

Pharmacological Treatment of Dementia

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
540 Gaither Road
Rockville, MD 20850
www.ahrq.gov

Contract No. 290-02-0020

Prepared by:

McMaster University Evidence-based Practice Center
McMaster University, Faculty of Health Sciences
Hamilton, Ontario, Canada

Investigators:

Pasqualina (Lina) Santaguida, PhD
Parminder Raina, PhD
Lynda Booker, BA
Christopher Patterson, MD
Fulvia Baldassarre, MSc
David Cowan, MD
Mary Gauld, BA
Mitch Levine, MD
Ayse Unsal, PhD

This document is in the public domain and may be used and reprinted without permission except those copyrighted materials noted for which further reproduction is prohibited without the specific permission of copyright holders.

Suggested Citation:

Santaguida PS, Raina P, Booker L, Patterson C, Baldassarre F, Cowan D, Gauld M, Levine M, Unsal A. Pharmacological Treatment of Dementia. Evidence Report/Technology Assessment No. 97 (Prepared by McMaster University Evidence-based Practice Center under Contract No. 290-02-0020). AHRQ Publication No. 04-E018-2. Rockville, MD: Agency for Healthcare Research and Quality. April 2004.

Preface

The Agency for Healthcare Research and Quality (AHRQ), through its Evidence-Based Practice Centers (EPCs), sponsors the development of evidence reports and technology assessments to assist public- and private-sector organizations in their efforts to improve the quality of health care in the United States. The reports and assessments provide organizations with comprehensive, science-based information on common, costly medical conditions and new health care technologies. The EPCs systematically review the relevant scientific literature on topics assigned to them by AHRQ and conduct additional analyses when appropriate prior to developing their reports and assessments.

To bring the broadest range of experts into the development of evidence reports and health technology assessments, AHRQ encourages the EPCs to form partnerships and enter into collaborations with other medical and research organizations. The EPCs work with these partner organizations to ensure that the evidence reports and technology assessments they produce will become building blocks for health care quality improvement projects throughout the Nation. The reports undergo peer review prior to their release.

AHRQ expects that the EPC evidence reports and technology assessments will inform individual health plans, providers, and purchasers as well as the health care system as a whole by providing important information to help improve health care quality.

We welcome written comments on this evidence report. They may be sent to: Director, Center for Outcomes and Evidence, Agency for Healthcare Research and Quality, 540 Gaither Road, Rockville, MD 20850.

Carolyn M. Clancy, M.D.
Director
Agency for Healthcare Research and Quality

Jean Slutsky, P.A., M.S.P.H.
Acting Director, Center for Outcomes and
Evidence
Agency for Healthcare Research and Quality

Name of Funding Government Agency Director
Director
Name of Funding Government Agency

The authors of this report are responsible for its content. Statements in the report should not be construed as endorsement by the Agency for Healthcare Research and Quality or the U.S. Department of Health and Human Services of a particular drug, device, test, treatment, or other clinical service.

Acknowledgements

The authors of this report would like to thank and acknowledge the assistance of the following people:

- Sarah Tanner-Bouma, Eileen Estrabillo, Connie Freeborn, Naomi Kirshenbaum, Elizabeth Simmons, Rob Stevens, and Hongmei Wu for assistance with data extraction and preparation of data tables;
- Carol Lefebvre for her assistance in developing the search strategies;
- Angela Eady for her assistance in developing the search strategies and editing the final version of the document;
- Bin Zhu, Gary Foster, and Ayaz Hyder for their assistance with the meta-analysis;
- Amy Steep and Tom Flemming of the McMaster University Health Sciences library for helping us with our searches, retrievals, and development of methods to access the library from off-site;
- Dan Ellington from ISI Researchsoft for answering questions regarding the management of our references and citations;
- Roxanne Cheeseman, Laurie Cocking, Susan Hanna, and Jessica Cheeseman for their assistance in the preparation and distribution of the manuscript.

Structured Abstract

Context. Dementia is a chronic progressive disease with no known cure. It affects cognition, behavior/mood, physical functions and activities of daily living, and caregiver burden. Therapeutic interventions for dementia aim to affect these domains.

Objectives. To review the evidence and answer the questions: Does pharmacotherapy for dementia syndromes improve cognitive symptoms and outcomes? Does pharmacotherapy delay cognitive deterioration or delay disease onset of dementia syndromes? Are certain drugs, including alternative medicines (non-pharmaceutical), more effective than others? Do certain patient populations benefit more from pharmacotherapy than others? What is the evidence base for the treatment of ischemic vascular dementia (VaD)?

Data sources. Studies were identified by searching the Cochrane Central trial registry, MEDLINE®, PreMedline®, EMBASE, AMED, CINAHL®, Ageline, and PsycINFO.

Study selection. English-language randomized controlled trials were selected if they evaluated pharmacological agents for adults with a diagnosis of dementia according to the criteria of International Classification of Diseases (ICD), Diagnostic and Statistical Manual of Mental Disorders (DSM) or National Institute of Neurological and Communicative Disorders and Stroke (NINCDS). Crossover trials and studies with a quality score < 3 on the Jadad Scale were excluded.

Data extraction. Data were extracted on type of dementia, severity of disease, setting, regimen of pharmacological agents, study duration, main outcome measures, adverse effects, and results. The quality of studies was assessed, and the quality of adverse effect reporting was assessed. Effect sizes were calculated and data were pooled when appropriate.

Data synthesis. (1) *Efficacy:* One hundred and eighty-six Randomized Controlled Trials (RCTs) evaluated 97 drugs. As expected the findings varied with the dementia population and the specific outcomes in the various domains. Those pharmacological agents that showed a consistent effect of benefit are as follows: A) Global assessment was improved by donepezil, galantamine, rivastigmine, velnacrine, cerebrolysin and idebenone; B) Cognition (general and specific) was improved by donepezil, galantamine, metrifonate (this drug has been withdrawn from use in North America because of safety concerns), nicergoline, physostigmine, rivastigmine, velnacrine, memantine, cerebrolysin, ginkgo biloba, idebenone and propentofylline; C) Behavior/mood was improved by haloperidol; D) Quality of life/Activities of Daily Living (ADL) was improved by donepezil, galantamine and posatirelin. In general, caregiver burden and quality of life/ADL were not frequently evaluated. (2) *Delay disease:* Cerebrolysin, selegiline plus vitamin E, and donepezil showed some significant effects in delaying disease progress in patients with mild to moderate and moderately severe Alzheimer's disease. (3) *Head to head comparisons:* Superiority was seen for sulphomucopolysaccharides over CDP-choline, donepezil over vitamin E, antagonistic-stress over nicergoline, antagonistic-stress over meclofenoxate, posatirelin over citicoline, and pyritinol over hydergine. (4) *Patient populations:* Stratified analyses included: age, gender, Apolipoprotein E (APOE) genotype,

disease type, disease severity, race by location, care dependence, and presence of depression. Single populations of dementia subjects with Down's syndrome, and depression were evaluated. Evidence was inconclusive for this question. (5) *Ischemic VaD*: A total of 20 pharmacological interventions in 29 studies were applied to vascular dementias. Differences were suggested between multi-infarct dementia (MID) and Alzheimer's disease (AD) for 5'-MTHF-trazodone, AD and VaD for citalopram, and AD and MID for Ginkgo biloba. Trials with VaD patients showed effects for memantine, nicergoline, pentoxifylline, idebenone, donepezil and cerebrolysin.

Conclusions. Pharmacotherapy for dementia can improve symptoms and outcomes. Adverse events should be more systematically reported. Few studies evaluated delay in either disease onset or progression, but there was some evidence suggesting delay in progression. Few studies compared drugs with other drugs. Due to poor evaluation, data was limited to consider efficacy of pharmacotherapy in different subgroups of patients. Some agents have been shown to be effective in VaD patients.

Table of Contents

Evidence report	1
Chapter 1. Introduction.....	3
Diagnosis of dementia.....	4
Analytic framework: understanding therapeutic aims of pharmacological treatment	5
Understanding efficacy of pharmacological interventions in dementia trials	5
Understanding efficacy of pharmacological interventions in dementia trials	6
Efficacy as measured by clinical versus statistical significance	6
Efficacy and outcome measures used in pharmacological intervention trials	7
Efficacy and potential risk of adverse events	8
Efficacy and intention to treat analysis	9
PRIMARY OBJECTIVES AND SCOPE OF SYSTEMATIC REVIEW	9
The questions	10
Chapter 2. Methods.....	11
THE RESEARCH TEAM.....	11
TOPIC ASSESSMENT AND REFINEMENT	11
Refinement of questions	11
Search strategy	11
Eligibility criteria.....	12
Data collection and reliability of study selection.....	16
Summarizing results: descriptive and analytic approaches	17
Meta-analysis	17
Power analyses.....	18
Peer review process.....	18
Chapter 3. Results.....	19
ELIGIBLE STUDIES	19
QUESTION 1: DOES PHARMACOTHERAPY FOR DEMENTIA SYNDROMES IMPROVE	
COGNITIVE SYMPTOMS AND OUTCOMES?	19
Interpretation of the results in the overall summary tables (ost) for individual studies	22
Statistical analysis.....	23
Quantitative and descriptive analyses	23
Results of cholinergic neurotransmitter modifying agents (cnma).....	23
Results of non-cholinergic neurotransmitter/neuropeptide modifying agents	
(NCNMA)	43
Results of other agents	50
QUESTION 2: DOES PHARMACOTHERAPY DELAY COGNITIVE DETERIORATION OR DELAY	
DISEASE ONSET OF DEMENTIA SYNDROMES?	62
Delay of Onset of Dementia	62
Delay of Progression.....	62
Survival Analyses	63

Staggered Withdrawal.....	63
QUESTION 3: ARE CERTAIN DRUGS, INCLUDING ALTERNATIVE MEDICINES (NON-PHARMACEUTICAL), MORE EFFECTIVE THAN OTHERS?	65
Head to Head Comparisons	65
QUESTION 4: DO CERTAIN PATIENT POPULATIONS BENEFIT MORE FROM PHARMACOTHERAPY THAN OTHERS?.....	68
QUESTION 5: WHAT IS THE EVIDENCE FOR THE TREATMENT OF VAD?	70
Summary Evidence Tables.....	73
Summary Table 1. Carnitine.....	75
Summary Table 2. Donepezil.....	76
Summary Table 3. Galantamine.....	77
Summary Table 4. Metrifonate.....	78
Summary Table 5. Nicergoline.....	79
Summary Table 6. Physostigmine.....	80
Summary Table 7. Posatirelin.....	81
Summary Table 8. Rivastigmine.....	82
Summary Table 9. Tacrine.....	83
Summary Table 10. Velnacrine.....	84
Summary Table 11. Various cholinergic neurotransmitter modifying agents.....	85
Summary Table 12. Haloperidol.....	86
Summary Table 13. Memantine.....	87
Summary Table 14. Selegeline.....	88
Summary Table 15. Various non-cholinergic neurotransmitter/neuropeptide modifying agents.....	89
Summary Table 16. Cerebrolysin.....	91
Summary Table 17. Estrogens.....	92
Summary Table 18. Ginkgo Biloba.....	93
Summary Table 19. Idebenone.....	94
Summary Table 20. Oxiracetam.....	95
Summary Table 21. Pentoxifylline.....	96
Summary Table 22. Propentofylline.....	97
Summary Table 23. Additional pharmacological agents.....	98
Summary Table 24. Drug vs drug studies.....	100
Summary Table 25. Vad/MID Studies.....	102
Chapter 4. Discussion.....	105
STRENGTH OF THE EVIDENCE.....	105
QUESTION 1: DOES PHARMACOTHERAPY FOR DEMENTIA SYNDROMES IMPROVE COGNITIVE SYMPTOMS AND OUTCOMES?	106
Summary of the systematic review results.....	106
Summary of cholinergic neurotransmitter modifying agents	106
Summary of non-cholinergic neurotransmitter/neuropeptide modifying agents	112
Summary of other pharmacological agents.....	113
Methodological issues and limitations in assessing efficacy of dementia agents.....	117
QUESTION 2: DOES PHARMACOTHERAPY DELAY COGNITIVE DETERIORATION OR DELAY DISEASE ONSET OF DEMENTIA SYNDROMES?	121

Summary of systematic review results	121
Methodological issues.....	122
QUESTION 3: ARE CERTAIN DRUGS, INCLUDING ALTERNATIVE MEDICINES (INCLUDING NON-PHARMACEUTICAL) MORE EFFECTIVE THAN OTHERS?	125
Summary of systematic review results	125
Relative efficacy must be evaluated in direct comparison trials.....	125
QUESTION 4: DO CERTAIN PATIENT POPULATIONS BENEFIT MORE FROM PHARMACOTHERAPY THAN OTHERS?.....	126
Summary of systematic review results	126
Representativeness of populations in the drug trials.....	126
QUESTION 5: WHAT IS THE EVIDENCE-BASE FOR THE TREATMENT OF VASCULAR DEMENTIA?	126
Summary of systematic review results	126
Diagnosis classification of VAD	127
DETERMINING CLINICAL RELEVANCE	127
LIMITATIONS OF THE MCMASTER AHRQ REVIEW	129
FUTURE RESEARCH RECOMMENDATIONS	130
References.....	133
Acronyms and Abbreviation.....	149
Figures	
Figure 1. Pathway for the progression of dementia and the ideal application of drug interventions within this framework.	5
Figure 2. Flow diagram showing the final number of studies meeting the eligibility criteria.	20
Figure 3. Proportion of studies as a function of year of publication.....	21
Figure 4. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the MMSE comparing carnitine versus placebo.	25
Figure 5. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the MMSE comparing donepezil and placebo.	27
Figure 6. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing donepezil versus placebo	27
Figure 7. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the CIBC+ (continuous data) comparing donepezil versus placebo	28
Figure 8. Relative Risk (RR) from the Random Effects Model (Random) for the CIBC+ (dichotomous data probability of improving) for a 5 mg dose of donepezil.	28
Figure 9. Relative Risk (RR) from the Fixed Effect Model (fixed) for the CIBC+ (dichotomous data [improved versus not]) for a 10 mg dose of donepezil.	28
Figure 10. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the Clinical Dementia Rating (CDR) comparing donepezil versus placebo.	29
Figure 11. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing galantamine at 24 mg dose versus placebo.	30

Figure 12. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing galantamine at 32 mg dose versus placebo.	31
Figure 13. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC comparing galantamine at 24 mg dose versus placebo.	31
Figure 14. Relative Risk (RR) from the Fixed Effects Mode Fixed I for CIBIC comparing galantamine at 32 mg dose versus placebo.	32
Figure 15. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the DAD comparing galantamine at 24 mg dose versus placebo.	32
Figure 16. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the DAD comparing galantamine at 32 mg dose versus placebo.	32
Figure 17. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing rivastigmine versus placebo.	38
Figure 18. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC+ comparing rivastigmine versus placebo.	39
Figure 19. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing cerebrolysin versus placebo.	52
Figure 20. Odd Ratio (OR) from the Random Effects Model (Random) for the CGI comparing cerebrolysin versus placebo.	53
Figure 21. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the MMSE change score comparing propentofylline versus placebo.	59
Figure 22. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the DSST change score comparing propentofylline versus placebo.	60
Figure 23. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing donepezil versus placebo.	66
Figure 24. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing galantamine versus placebo.	66
Figure 25. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing rivastigmine versus placebo.	67
Figure 26. Weighted Mean Difference (WMD) from the Fixed Effects Model (Fixed) for the ADAS-cog comparing tacrine versus placebo.	67
Figure 27. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC comparing donepezil versus placebo.	67
Figure 28. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC comparing galantamine versus placebo.	68
Figure 29. Relative comparison of effect sizes for studies using the CIBIC rivastigmine versus placebo.	68
Figure 30. Relative comparison of effect sizes for studies using the CIBIC comparing tacrine versus placebo.	68
Figure 31. Delay of symptomatic treatment effects.	123
Figure 32. Delay in disease progression treatment effects.	124

Tables

Table 1. Databases searched for relevant RCTS.	12
Table 2. List of cholinergic neurotransmitter modifying agents and the number of studies vs. Placebo for each of these.	23

Table 3. List of non-cholinergic neurotransmitter/neuropeptide modifying agents and the number of studies vs. placebo for each of these. Asterisk (*) indicates report of a drug vs. drug trial [comparator drug(s) in brackets].	43
Table 4. List of other pharmacological agents and the number of studies vs. placebo for each of these.	50
Table 5. Studies that withdrew the treatment agent but maintained at least single blinding.	64
Table 6. Studies that withdrew treatment and did not specify if blinding for washout or extension was maintained.	64
Table 7. Studies with stratified analyses.	69
Table 8. Studies evaluating vascular dementia patients relative to other dementias.	71
Table 9: Guide to Overall Summary Tables – Outcome Measures Classified by Domain.	72

Appendixes and Evidence Tables cited in this report are provided electronically at <http://www.ahrq.gov/clinic/epcindex.htm>.



Pharmacological Treatment of Dementia

Summary

Introduction

The focus of this review is the pharmacological treatment of dementia. Pharmacotherapy is often the central intervention used to improve symptoms or delay the progression of dementia syndromes. The available agents vary with respect to their therapeutic actions, and are supported by varying levels of evidence for efficacy. This report is a systematic evaluation of the evidence for pharmacological interventions for the treatment of dementia in the domains of cognition, global function, behavior/mood, quality of life/activities of daily living (ADL) and caregiver burden.

Many medications have been studied in dementia patients. These agents can be classified into three broad categories:

1. Cholinergic neurotransmitter modifying agents, such as acetylcholinesterase inhibitors.
2. Non-cholinergic neurotransmitters/neuropeptide modifying agents.
3. Other pharmacological agents.

Although only five agents have been approved by the Food and Drug Administration (FDA) for the treatment of dementia, many other pharmacological agents have been evaluated in trials and may be prescribed in off-label use.

Given the range of pharmacological agents that have been tested in dementia, a systematic review of these interventions (using a consistent methodology) provides a meaningful contribution in this area. The key questions addressed in this systematic review are as follows:

1. Does pharmacotherapy for dementia syndromes improve cognitive symptoms and outcomes?
2. Does pharmacotherapy delay cognitive deterioration or delay disease onset of dementia syndromes?
3. Are certain drugs, including alternative medicines (non-pharmaceutical), more effective than others?
4. Do certain patient populations benefit more from pharmacotherapy than others?
5. What is the evidence base for the treatment of ischemic vascular dementia (VaD)?

This review considers different types of dementia populations (not just Alzheimer's Disease [AD]) in subjects from both community and institutional settings. The studies eligible in this systematic review were restricted to parallel RCTs of high methodological quality.

Methods

A team of content specialists was assembled from both international and local experts. The purpose of the expert panel was to assist in the topic assessment and refinement process; in addition, complex methodological issues were evaluated by this expert panel.

Search Strategy

Search strategies were developed and undertaken in the electronic databases including Cochrane Central, MEDLINE®, PreMEDLINE®, EMBASE, AMED, CINAHL®, AgeLine, and PsycINFO. In addition to the electronic databases, the bibliographies of retrieved papers were reviewed.



Eligibility Criteria:

Studies were included that met the following criteria:

- Populations included dementia patients who were 18 years or older in age.
- Diagnosis of dementia using criteria of International Classification of Diseases (ICD) 9 or 10, Diagnostic and Statistical Manual of Mental Disorders (DSM) III, III-R or IV, National Institute of Neurological and Communication Disorders and Stroke (NINCDS), Neurological and Communication Disorders and Stroke-Alzheimer's Disease and Related Disorders Association (NINCDS-ADRDA), or Neurological and Communication Disorders and Stroke-Association Internationale pour la Recherche et l'Enseignement en Neurosciences (NINCDS-AIREN).
- Potential populations at high risk of dementia conversion in order to address the issue of delay in onset. These populations included: Mild Cognitive Impairment (MCI), Cognitive Impairment not Dementia (CIND), Cognitive Loss No Dementia (CLoND).
- Interventions were restricted to pharmacological agents, including food supplements (as defined by the FDA) administered for at least 1 day.
- Parallel design randomized control trials (RCT) in the English language of any sample size.
- Score of 3 or greater on the modified Jadad quality scale.

All types of instruments were considered for this review within the outcome domains.

Populations of dementias caused by toxic agents (e.g., alcohol) and temporary dementia (e.g., side effect of anesthesia) were excluded.

Data Collection and Reliability of Study Selection

All studies meeting eligibility criteria were reviewed to assess quality and data abstracted according to predetermined criteria. The articles were grouped according to the pharmacological agent used in the intervention. A team of study assistants were trained in the criteria for eligibility and quality for the purposes of this systematic review. Standardized forms and a guide explaining the criteria were developed from previous templates.

Study outcomes were classified into the following domains:

1. General cognitive function.
2. Specific cognitive function.
3. Global clinical assessment.
4. Behavior/mood.
5. Quality of life/ADL.
6. Effects on primary caregiver (also referred to as caregiver burden).

7. Safety as measured by the incidence of adverse effects (particularly serious events).
8. Acceptability of treatment as measured by withdrawal rate from trial due to side effects of the medication.

Measurement of Benefits and Harms

Evaluation of efficacy is based upon reported changes for outcomes in the principal domains of interest. Evaluation of the potential for harm is considered within three main areas: 1) the most frequently reported adverse events across studies for a specific drug, 2) the overall withdrawal rate due to adverse events for both the control and treatment groups, and 3) the range of frequencies reported for a subset of specific symptoms (nausea, diarrhea, dizziness, agitation, eating disorder) selected *a priori* and evaluated for all pharmacological interventions.

Measure of Effect Size and Meta-analysis

Effect sizes (ES) for trials were conducted for those pharmacological interventions with the same outcomes. In studies with multiple dosage groups and where sufficient data were provided, each dose level had an ES estimated separately relative to placebo. Before calculating a pooled effect size measure, the reasonableness of pooling was assessed on clinical and biological grounds, in terms of clinical homogeneity and therefore statistical meta-analysis was not appropriate for all outcomes or interventions.

Results

Question 1: Does pharmacotherapy for dementia syndromes improve cognitive symptoms and outcomes?

Seventy-two studies examined cholinergic neurotransmitter modifying agents, 61 studies examined non-cholinergic neurotransmitter/neuropeptide modifying agents and 76 trials evaluated other agents used to treat dementia. Table 1 lists all the pharmacological agents and the number of trials (in brackets) eligible for review in this study. Twenty of these agents are detailed in this summary. All drug agents are detailed in the full report.

Summary of Cholinergic Neurotransmitter Modifying Agents

Carnitine. Six trials¹⁻⁶ evaluated carnitine in 925 subjects with mild to moderate severity, recruited predominately from the community. A dose of 2 to 3 g was compared to placebo for either 24 or 52 weeks.

Evidence of benefit is conflicting for the domains of general or specific cognition. Results were not statistically significant in any study but the lack of sufficient power may have influenced

these results. Similarly, no statistically significant differences were found in the domains of global assessment, behavior/mood, and quality of life/ADL. Statistical power could not be evaluated for the most of these outcomes.

Four of the six studies scored 3 for quality on reporting adverse events. Withdrawal rates due to adverse events varied from 0–3 percent (excluding results from one outlier trial⁷), and gastrointestinal symptoms were the most frequently reported types of adverse events.

Donepezil. Ten trials^{8–17} in 3239 subjects evaluated the efficacy of donepezil compared to placebo, and one trial¹⁸ compared donepezil with vitamin E. Eight of the studies evaluated AD patients, for which at least half were recruited from the community (other studies did not specify). The subjects had predominately mild to moderate disease and doses of 5 or 10 mg were used with study duration from 12 to 56 weeks.

There is consistent evidence of benefit in the domains of general cognitive function and global assessment; the combined effect sizes for the Alzheimer's Disease Assessment Scale-Cognitive Section (ADAS-cog) and the Clinician's Interview-Based Impression of Change (CIBIC) were estimated. Based on the three studies that evaluated two different doses (5 and 10 mg), there was no consistent dose response relationship as the benefit was of similar magnitude for global assessment outcomes. Two of the three studies that evaluated behavior/mood outcomes, using the Neuropsychiatric Inventory (NPI), showed no statistically significant changes relative to placebo but these trials lacked sufficient power to detect a difference. There is evidence of benefit in ADL outcomes, although this outcome was evaluated by a variety of instruments. Caregiver burden outcomes were measured in a single study that did not report the findings for this domain.

Adverse events quality scores were 3 or greater for the majority of studies (n=7). Four trials provided evidence of a dose response for adverse events. One study showed a statistical difference for balance-related problems and asthenia (neurological fatigue) between placebo and treatment groups. Withdrawal due to adverse events ranged from 0–18 percent for treatment groups and 0–11 percent for placebo. Four out of 6 studies testing for differences between groups were statistically significant for diarrhea, nausea and vomiting.

Galantamine. Six trials^{19–24} in 3530 subjects compared the efficacy of galantamine with placebo. Doses of 24 and 32 mg were evaluated in half of these studies. Five studies evaluated only AD patients and there was limited information regarding the subjects' residence (community or institutional settings). All

studies recruited subjects with mild to moderate disease and the drug was administered from 3 to 6 months duration.

Evidence of benefit is consistent in the domains of general cognitive function, global assessment and quality of life/ADL. Two of the three studies that evaluated behavior/ mood found statistically significant differences in favor of galantamine. A dose effect was evident in the ADL domain when comparing the pooled estimates of the Disability Assessment for Dementia (DAD); no dose effect was observed for outcomes in the global assessment domain, and this could not be evaluated for the general cognition domain. Caregiver burden was not evaluated in any trial.

Five of the six trials scored 3 out of 5 on our quality scale for rating adverse events. Withdrawal rates due to adverse events ranged from 4–9 percent for placebo and 8–27 percent for the treatment group. One study showed a dose response for adverse events. Although four trials did not report significance testing for differences between groups, two trials did report a statistically significant difference in weight loss between the placebo and treatment group. The most common adverse events were gastrointestinal symptoms (nausea and vomiting, diarrhea), eating disorders/weight loss, and dizziness.

Metrifonate. Nine studies^{25–33} compared metrifonate to placebo in 2759 subjects with mild to moderate AD (the majority of studies did not specify community settings). Metrifonate doses from 50 to 80 mg were given for 21 days to 26 weeks duration.

All but one study showed metrifonate to have a consistent positive effect on measures of general cognitive function; none of the studies evaluated specific cognitive function measures. Effects on global assessment were less consistent but suggested a positive effect in four of the eight studies. Evidence for effect in the domains of behavior/mood and quality of life/ADL were not statistically significant in the majority of studies that evaluated these domains; however these were primarily evaluated as secondary outcomes and likely lacked sufficient power.

With the exception of a single study, quality scores for reporting adverse events were greater than 3. However, only one trial tested for differences between groups and found nausea and vomiting, diarrhea, and muscle and joint disorder to have statistically significantly differences. Withdrawal due to adverse events varied from 0–9 percent for placebo and 0–12 percent for the treatment group. It was difficult to determine which types of reported adverse events had the potential to cause serious harm. This is noteworthy as metrifonate has been withdrawn from use in North America, and Bayer has suspended Phase III trials,³⁴ because some patients in clinical

trials have experienced serious muscle weakness. This decision was based on the results of an experimental study showing risk of respiratory paralysis with the use of metrifonate. Other adverse events of concern included severe leg cramps, dyspepsia, and bradycardia. None of the studies that we reviewed indicated that if present, these events differed with statistical significance between groups. It is not clear if this inconsistency is a function of the methods used to collect and report adverse events, or a limitation of RCTs as a source of detecting serious adverse events when the incidence is low.

Nicergoline. Four trials³⁵⁻³⁸ in 705 subjects compared nicergoline to placebo and one trial³⁹ compared it to a second drug (antagonic-stress) in mixed populations that included AD, Multi-Infarct Dementia (MID), Progressive Degenerative Dementia (PDD), Vascular Dementia (VaD), mixed dementia, and Senile Dementia of the Alzheimer's Type (SDAT), which were classified as mild to moderate in severity.

All placebo-controlled trials found a positive effect for general cognitive outcomes, but half the results were based on observed case (OC) analyses. The evidence for benefit was mixed in the domain of global assessments. No statistically significant differences were found for behavior/mood, nor quality of life/ADL outcomes but these were evaluated in few studies and as secondary outcomes (suggesting that sufficient power was an issue).

Quality scores for reporting adverse events varied from 2 to 5 for these four trials, and none tested for differences between groups. Withdrawal due to adverse events varied from 0–8 percent for placebo and 0–9 percent for the treatment group. With the exception of headache, which was reported in all four trials, it was difficult to determine which types of adverse events most characterized exposure to this pharmacological agent.

Physostigmine. Four studies⁴⁰⁻⁴³ in 1198 subjects with mild to moderate AD evaluated physostigmine administered in patch and oral form (30 to 60 mg dose) from 6 to 24 weeks duration. All subjects were recruited from the community.

There is evidence that physostigmine has a statistically significant positive effect on general cognitive function, as three of the four studies showed improvement. Evidence for an effect on global function was mixed with no consistent effect. Similarly, for quality of life/ADL outcomes, all three studies that evaluated this domain showed no statistically significant difference but these were secondary outcomes and may reflect a lack of power. Behavior/ mood and caregiver burden outcomes were not tested.

The quality scores for reporting adverse events were generally low, scoring 1 or 2 out of 5. Withdrawal rates due to adverse events varied from 1–5 percent for placebo and 12–55 percent

in the treatment group, with one study not reporting rates. The high withdrawal rates were in studies with sample sizes that varied from 181 to 475 subjects. A single study tested for differences between groups, and found that dizziness, tremor, weight loss, asthenia, confusion, delirium, and respiratory problems (not detailed) were significantly different statistically. The cluster of reported types of adverse events suggests that gastrointestinal problems (abdominal pain, diarrhea, nausea and vomiting and eating disorder) were most frequently reported.

Posatirelin. Four trials⁴⁴⁻⁴⁷ evaluated posatirelin in 931 subjects in a variety of mild to moderate dementia populations (AD, PDD, VaD) using 10 mg per day dose for 3 months duration.

Three of the four trials showed statistically significant improvement in general cognitive function and quality of life/ADL (as measured by Gottfries-Brane-Steen (GBS) subscales for these domains). The evidence remains inconsistent for benefit in global assessment (evaluated in only one trial) and behavior/mood (mixed results). Caregiver burden and specific cognitive function were not evaluated.

Quality scores for reporting adverse events varied from 2 to 4. Withdrawal rates due to adverse events ranged from 0–3 percent in placebo and 0–4 percent in the treatment group. None of the studies tested for statistically significant differences between groups for adverse events. At least three studies reported arrhythmia, nausea/vomiting, headache, rash/skin disorder, and sleep disorder.

Rivastigmine. Six studies⁴⁸⁻⁵³ evaluated 2071 subjects with three of these studies limited to AD patients. Doses of rivastigmine varied from 1 to 12 mg, given for 14 to 26 weeks and only one study specified a community sample.

Evidence shows that general cognitive function improves with rivastigmine at dose of 12 mg but there are mixed results for efficacy at lower doses. Two trials evaluated specific cognitive function but the results were not consistent within studies (between general and specific measures); similarly, the results were not consistent for general and specific cognition between studies. There is consistent evidence of benefit for global function but the dosage at which this occurs has statistically significant variation among studies. In the domains of behavior/mood, quality of life/ADL, the findings were neither statistically significant nor consistent; most of these analyses were not based on intention to treat analysis and lack of sufficient power cannot be ruled out. Caregiver burden outcomes were not evaluated.

Quality scores for reporting adverse events varied from 2 to 5. Withdrawal rates due to adverse events ranged from 4–11

percent in the placebo and 11–27 percent in the treatment group. Two trials demonstrated a dose response; however, one of these trials showed statistically significant differences for nausea and vomiting only, and the other trial showed statistically significant differences for all the adverse events reported. The majority of studies reported dizziness, nausea and vomiting, eating disorder/weight loss, and headache. It should be noted that one study allowed intentional prescribed anti-emetic drugs to increase the tolerance of subjects taking rivastigmine.

Tacrine. Six studies⁵⁴⁻⁵⁹ evaluated tacrine in 994 subjects predominately with mild to moderate AD at doses of 80 to 160 mg lasting from either 12/13 or 30/36 weeks in duration. Two other studies^{60,61} involving 425 patients were non-placebo controlled studies. The majority of studies recruited community-based subjects.

A single trial showed benefit for general cognitive function. The small effect size was based on a series of related publications. The five trials showing no benefit for general cognitive function comprised small sample sizes and much shorter study duration. Thus, the evidence for benefit in general cognitive function is limited to a single trial. There is evidence for benefit in global function in two of the three trials. Changes in behavior/mood, quality of life/ADL domains, specific cognitive function, and caregiver burden were all not statistically significant, but lack of sufficient power cannot be ruled out.

The quality scores for reporting adverse events varied from 1 to 3. The proportion of subjects withdrawing due to adverse events ranged from 0–12 percent for placebo and 0–55 percent in the treatment group. The higher rates of withdrawal were associated with higher doses. Elevated alanine transaminase (ALT) or hepatic abnormality (placebo=4–13 percent, all doses tacrine=7–67 percent) was reported in six studies, raising concerns for the potential for serious liver damage. None of these trials tested for differences between treatment and placebo with respect to adverse events. Five studies reported nausea and vomiting, gastrointestinal problems, and dizziness. There is evidence for potentially serious adverse events associated with liver dysfunction in six trials.

Velnacrine. Three studies⁶²⁻⁶⁴ evaluated the effects of velnacrine in 774 AD patients with a probable severity classification. Doses between 75 mg twice daily and 225 mg were given for 15 to 24 weeks duration. Location of recruitment was not specified.

Statistically significant positive effects were observed for general cognitive function, and global assessment in the two studies with sample sizes over 300 subjects. Behavior/mood and

caregiver burden showed some benefit in one trial⁶² at the highest dose only. Quality of life/ADL was tested as a secondary outcome and showed mixed findings.

Quality scores for reporting adverse events were 3 for all studies. Withdrawal rates varied from 0–22 percent for the placebo group and 5–33 percent for the treatment group. None of the studies reported a dose response. None of the studies tested for statistical differences between the placebo and treatment groups. Two studies reported aberrant hematology and hepatic abnormality^{62,64}; for these two studies the rates of occurrence were 2–21 percent for placebo, and 32–40 percent for all doses. The potential for serious effects is not well specified in these trials. All studies reported diarrhea and nausea and vomiting.

Summary of Non-cholinergic Neurotransmitter/Neuropeptide Modifying Agents

Haloperidol. Five studies⁶⁵⁻⁶⁹ evaluated the effect of haloperidol relative to placebo in a total of 622 subjects with mild to moderate disease that included AD patients and mixed populations (MID/VaD/ PDD). One trial had only 15 patients, and one trial⁶⁵ lasted only 3 weeks. Two studies recruited subjects from institutions; one from the community; and, two did not specify.

Mixed results were observed for improvement in global assessment. In three of the trials there was benefit in the domain of behavior/mood which reached statistical significance. Two trials evaluated caregiver burden and found no statistically significant differences but lack of sufficient power cannot be ruled out. Few studies evaluated outcomes in quality of life/ADL. Haloperidol did not affect general cognitive function in two trials and was not evaluated in the other studies.

The quality scores for reporting adverse events varied from 1 to 5 and only three of five studies reported withdrawal rates; the proportion of subjects withdrawing due to adverse events ranged from 5–17 percent for placebo and 17–33 percent in the treatment group. One trial showed a dose-response effect but the study lasted only 3 weeks. Three trials tested for differences between treatment and placebo with respect to extra-pyramidal symptoms (placebo=17–32 percent, all doses=34–97 percent), and two found statistically significant differences.^{65,66} One study⁶⁶ found statistically significant differences between groups for balance-related problems.

Memantine. Three trials⁷⁰⁻⁷² evaluated memantine in 1066 patients, primarily with VaD, with 10 or 20 mg doses for durations of 12 or 28 weeks. Disease severity was moderate to severe in a single study⁷⁰ and mild to moderate in the remaining two studies.^{71,72} One study included patients that were

institutionalized; one study included community subjects; and the other study did not report the source of patients.

Consistent evidence of benefit in general cognitive function was demonstrated in the two studies that evaluated this domain. Findings for global assessment are mixed. The only trial that evaluated mixed dementia populations (including some VaD) with moderate to severe dementia found statistically significant improvements in global function, behavior/mood, and quality of life/ADL outcomes, but did not evaluate general cognitive function. It should be noted that this trial with mixed populations used half the dose of memantine for half the study duration in patients with greater disease severity, and had approximately half the sample size of the other two trials evaluated in this systematic review. Despite a lower dose, a smaller number of more severely affected patients and a shorter duration, a statistically significant difference was found.

The quality scores for reporting adverse events varied from 3 to 4. Only two of three studies reported withdrawal rates; the proportion of subjects withdrawing due to adverse events ranged from 3–7 percent for placebo and 9–12 percent in the treatment group. A single trial tested for differences between treatment and placebo, and none of the comparisons were significantly different statistically.

Selegiline. Six trials⁷³⁻⁷⁸ evaluated selegiline in 733 patients with AD, PDD, and dementia Alzheimer's type (DA) with 10 mg per day and study duration of 60 days or 2 years.

All but one trial that evaluated general cognition showed no statistically significant changes. A single trial found statistical improvements in specific cognitive tests (Sternberg Memory tests); this trial also showed statistically significant improvements in global assessment and behavior/mood. Only this trial, which had the highest quality score (7), showed consistently positive findings across all domains tested. Three of the five trials that evaluated part or all of these domains had very small sample sizes and were likely underpowered, possibly accounting for the inconsistent findings. Based on a single trial there is evidence that selegiline and selegiline combined with vitamin E, delays the time to important functional decline milestones.

The quality scores for reporting adverse events varied from 0 to 3. The proportion of subjects withdrawing due to adverse events ranged from 0–4 percent for placebo and 0–9 percent in the treatment group. Only one trial tested for differences between the treatment and placebo groups and showed that balance and falls were statistically significantly different (worse) between groups (particularly the group with selegiline combined with vitamin E [22 percent] versus placebo [5

percent]). However, when adjusted for multiple comparisons, these were no longer statistically significant.

Summary of Other Pharmacological Agents

Cerebrolysin. Six studies⁷⁹⁻⁸⁴ evaluated the effect of cerebrolysin in a total of 819 subjects. All but one of the trials included only AD patients with mild to moderate disease. All of the studies used the same dose of cerebrolysin, 30 ml per day for 5 days per week for 4 to 24 weeks duration. Location of recruitment was not specified.

Cerebrolysin showed a statistically significant improvement in cognition in four of five studies that evaluated this domain. Although a pooled estimate for the ADAS-cog was calculated, the model was positive for heterogeneity and the overall estimate was not statistically significant. The results for specific cognitive tests for the three trials that evaluated this domain were inconsistent. Global assessment measures showed a statistically significant effect in five of the trials. A summary estimate for the Clinical Global Impression (CGI) was presented; this model was also positive for heterogeneity but statistically significant for an overall effect. Two out of three studies showed an effect for behavior/ mood, but none of the six studies showed an effect on quality of life/ADL. No study measured caregiver burden.

Two of the six trials scored 5 out of 5 on our quality scale for rating adverse events, but did not report any adverse events. Two studies scored 4, and the other two trials scored 3 and 2. All the studies with scores equal to 4 or less tested for statistical differences in adverse events between placebo and treatment groups. Withdrawals due to adverse events were not reported in one study, and were 1 percent in two studies and none withdrew in three studies. A statistically significant difference between treatment and control group was reported in one study for weight change, anxiety, and headache.

Estrogen. Five studies⁸⁵⁻⁸⁹ evaluated estrogens for dementia in 247 patients with primarily mild to moderate AD from the community, with the exception of one study that included moderate to severe dementia patients who were all institutionalized. One of the studies with AD patients provided 0.10 mg per day by skin patch for 8 weeks and the others used 1.25 mg per day for 12 to 52 weeks duration. The study including severe subjects used 2.5 mg per day for 4 weeks.

Three trials evaluated general cognitive function and all showed statistically non-significant findings; two trials lacked sufficient power to show changes on the ADAS-cog. Two other trials evaluated specific cognitive function but results were mixed. Most of the outcomes evaluated in the domains of global assessment, behavior/mood, and quality of life/ADL

were secondary outcomes and none showed statistically significant differences (but lack of power could be a factor).

One of the five trials scored 5 out of 5 on our quality scale for rating adverse events, but did not report any adverse event. Withdrawal rates due to adverse events ranged from 0–5 percent for placebo and 0–14 percent for the treatment group. The most frequently reported adverse event was vaginal bleeding and a single trial reported a statistically significant difference between placebo and treatment group for this symptom. It was not clear from the descriptions provided in the study if they had ascertained whether vaginal bleeding was present prior to the trial commencement.

Ginkgo biloba. Three trials⁹⁰⁻⁹² evaluated Ginkgo biloba, 120 to 240 mg per day for 3 to 12 months, in a total of 563 subjects with mixed dementias of mild to moderate severity. All were recruited from the community.

The largest trial had the longest treatment duration but the lowest daily dosage and reported a statistically significant impact for general cognitive function but had mixed findings for global assessment. A second large trial found positive changes for neuropsychological tests, global assessment, and behavior/mood outcomes with double the dosage of the previously described trial and half the treatment interval. In this RCT, clinical efficacy was assessed by using a responder analysis, with therapy response being defined as response in at least two of the three variables: CGI—global function, Syndrome Kurz test (SKT)—special cognitive function, and Nurnberger-Alters-Beobachtungs-Skala (NAB)—ADL. A single trial evaluated behavior/mood and the result was not statistically significant. No trial evaluated caregiver burden or quality of life/ADL.

All three trials scored 3 or greater on the quality scale for rating adverse events. Two studies had no withdrawals due to adverse events, and one trial had a withdrawal rate of 6 percent for both placebo and treatment groups. Two studies reported no adverse events. One study reported a statistically significant difference between the treatment and the placebo group for skin disorders. The same study reported gastrointestinal and headache adverse effects, but did not test for statistical differences between the placebo and the treatment group.

Idebenone. Four studies⁹³⁻⁹⁶ evaluated the drug idebenone in 1153 subjects of mixed dementia populations of mild to moderate severity; one of these trials evaluated idebenone relative to tacrine. Doses varied from 30 mg per day to 360 mg per day, and the treatment interval ranged from 90 days to 60 weeks.

There was evidence of benefit in general cognitive function and global assessment. Several studies evaluated behavior/mood

and quality of life/ADL and these outcomes were found to be statistically different. None of the trials evaluated caregiver burden.

Quality scores for reporting adverse events varied from 1 to 5. Rates of withdrawal due to adverse events varied from 0–5 percent for the placebo group and 0–5 percent in the treatment group; a single trial did not report withdrawal rates. Two trials tested for statistical differences between groups and found none. Although no clear pattern emerges, three studies identified at least one balance-related adverse event.

Oxiracetam. Five studies⁹⁷⁻¹⁰¹ evaluated oxiracetam in 554 subjects with different dementia syndromes of mild to moderate severity. All studies used 1600 mg daily, with one exception where the dose ranged between 1600-2400 mg per day. The treatment interval ranged from 90 days to 26 weeks duration.

All outcomes shown to be positive for this drug were based on Observed Cases (OC) evaluation. The two trials that evaluated general cognitive function showed benefit. The findings for specific cognitive function were mixed. A single trial evaluated global assessment and showed statistically significant change. Behavior/mood and quality of life/ADL outcomes showed mixed results. No study evaluated caregiver burden.

The quality scores for reporting adverse events varied from 2 to 5. The proportion of withdrawals due to adverse events varied from 0–9 percent for the placebo group and 0–6 percent for the treatment group. No clear pattern for adverse events is evident, but three of the five studies reported gastrointestinal related problems, primarily abdominal pain.

Pentoxifylline. Three placebo-controlled studies¹⁰²⁻¹⁰⁴ evaluated pentoxifylline and one study compared pentoxifylline to sulodexide, with a total of 482 subjects with predominately MID. The dose administered in all studies was 1200 mg per day but varied between once or three times daily. The treatment intervals ranged from 12 to 36 weeks.

All three placebo trials showed statistically non-significant findings for any primary outcome evaluated on all subjects in the study. Two of these trials had very small sample sizes (n=38, n=28) and employed Observed Cases (OC) analyses; this suggests that the trials lacked sufficient power to evaluate multiple outcomes. The remaining trial had a large sample size (n=289) and employed an Intention to Treat (ITT) analysis; all primary outcomes evaluated were not statistically significant.

The quality scores for reporting adverse events were generally low, varying from 1 to 3. Withdrawal rates due to adverse events varied from 0–25 percent in the placebo group and 0–22 percent in the treatment group. The two studies that

reported adverse events indicated the presence of gastrointestinal disturbances, including abdominal pain and nausea and vomiting.

Propentofylline. Four trials¹⁰⁵⁻¹⁰⁸ using propentofylline in 510 patients with AD and VaD were included. A dose of 900 mg per day was consistent across all studies, and the treatment duration ranged from 3 to 12 months.

Two studies with small sample sizes (n=30) showed no statistically significant results for any outcome evaluated but likely lacked power. There were two trials that found benefit in general cognitive function based on the Mini-Mental Status Exam (MMSE). The results for specific cognitive function as measured by the Digit Symbol Substitution Test (DSST) were mixed, as were those for global assessment. Behavior/mood outcomes were evaluated in a single trial and showed no statistically significant difference; this same trial evaluated quality of life/ADL and showed no statistically significant difference. No trial evaluated caregiver burden.

The quality scores for reporting adverse events varied from 1 to 4. The percentage of withdrawals varied from 0–13 percent for the placebo group and 0–12 percent for the treatment group. None of the trials tested for differences between groups. Three of the trials reported gastrointestinal events that included abdominal pain, constipation, and nausea and vomiting.

Question 2: Does pharmacotherapy delay cognitive deterioration or delay disease onset of dementia syndromes?

Delay of Onset of Dementia

The concept of “delay onset” was operationalized to imply conversion from a state of cognitive impairment, classified as MCI, CLoND or CIND, to a true dementia state. No studies with this population met the final eligibility criteria, although four trials¹⁰⁹⁻¹¹² advanced to the full text screening stage. The lack of studies eligible for evaluation in this systematic review points to a gap in the literature for pharmacological interventions (attempting to demonstrate a delay in disease onset) in MCI-type populations.

Delay of Progression of Dementia

The need for good evaluation of disease progression in trials was also identified. In general, few studies evaluated subjects in more severe states of the disease. This suggests that a bias exists towards evaluating mild to moderate disease in the trials eligible in this systematic review; this may reflect an underlying assumption that the less severe groups are most likely to benefit from drug trials. Since so few studies have evaluated the more severe groups, this assumption may require some empirical

justification in future research. A consensus is required regarding the diagnostic criteria to be used to establish levels of severity.

Three studies evaluating cerebrolysin, selegiline and vitamin E, and donepezil have shown statistically significant effects in delaying disease progress in mild to moderate and moderately severe disease in patients with AD. This delay in progress was expressed in terms of delay in days to primary event or statistical differences between placebo at a specified time interval. Although these trials coincidentally evaluated dementia patients over the longest time interval, their protocol did not withdraw the drug at the end of the study. Theoretically, conclusive evidence of disease delay would be demonstrated if the treatment groups did not return to the level of the placebo. Thus, distinguishing between symptomatic and disease modifying effects is not possible unless the drug is withdrawn and the treatment groups are observed for these changes.

When studies attempted to evaluate disease progression, long-term (1 year or greater) trials continued in an “open-label fashion,” where blinding was no longer maintained. This limits the confidence that bias did not affect the subsequent changes in the outcomes. It was observed that increasing levels of dropout (for a variety of reasons) also plagued these open-label phases of evaluation. From a practical perspective, maintaining adherence in longer-term trials in dementia patients is challenging, particularly for those in the placebo arm or for those with interventions that have a high proportion of adverse events. Although this practical challenge exists, the findings of this review suggest that there is a gap in the literature showing delay of the disease process of dementia related disorders.

Question 3: Are certain drugs, including alternative medicines (non-pharmaceutical) more effective than others?

Head to head comparisons of drugs in the treatment of dementia

A total of 26^{18,39,47,60,61,65,66,68,69,73,113-128} studies compared efficacy of the two or more pharmacological agents relative to each other. In general, few drugs showed statistically significant differences relative to each other. Those that did include (listed in declining order of performance):

1. Sulphomucopolysaccharides versus CDP-choline:¹¹⁷ Statistically significant differences were seen in favor of sulphomucopolysaccharides in measures of behavior and global assessment in 30 institutionalized patients with mild to moderate MID.

2. Donepezil and vitamin E:¹⁸ Statistically significant differences were seen in favor of donepezil in general cognitive function 54 patients with mild AD.
3. Antagonic stress versus nicergoline:³⁹ Statistically significant differences were seen in favor of antagonic stress in cognition as well as a global assessments in 62 subjects with mild to moderate AD.
4. Antagonic stress versus meclofenate:¹²⁴ Statistically significant differences were seen in favor of antagonic stress in measures of cognition and global assessment in 63 patients with mild to moderate AD.
5. Posatirelin versus citicoline:⁴⁷ Statistically significant differences were seen in favor of posatirelin in general cognitive measure and mood in 222 community living patients with mild to moderate AD.
6. Pyritinol versus hydergine:¹²⁵ A significant difference in favor of pyritinol in a global assessment measure in 102 Hispanic patients with mild to moderate AD.
7. Idebenone⁶¹ versus tacrine: Mixed results were observed; the Efficacy Index Score showing a statistically significant benefit over tacrine, while the global assessment showed no difference in 203 individuals with AD, 44 of whom completed the study.

Current drugs approved in the United States for the treatment of dementia

What may be most relevant to clinicians are head to head comparison of the cholinergic modifying neurotransmitter pharmacological agents, particularly those currently approved for the treatment of dementia (tacrine, rivastigmine, galantamine, donepezil) in the United States. The evidence for each of these drugs has been extensively detailed, and the relative merits and handicaps of each are outlined in the results section of the full report (Chapter 3). Relative effectiveness as demonstrated by effect sizes for the ADAS-cog and the CIBIC are also compared in Chapter 3. Although, the psychometric properties of these two outcomes are commonly accepted, comparison across the populations in these pooled estimates may not lend themselves to direct comparison across these four different specific drugs; populations may be different and reporting of adverse events is not consistent. Thus, inferences about the relative efficacy of these four medications specific for the treatment of dementia should be made cautiously as head to head comparisons were not undertaken.

Question 4: Do certain patient populations benefit more from pharmacotherapy than others?

In general, very few trials examined the efficacy of dementia drugs across different populations or described the population characteristics in sufficient detail. From the 15 studies ^{2,3,8,10-}

^{12,23,24,61,84,93,129-132} that reported stratified analyses, eight different variables were identified, which included age, gender, Apolipoprotein E gene (APOE) genotype, disease type, disease severity (as determined by MMSE/ ADAS-cog threshold levels), treatment center, care dependence, and presence of depression. Additionally, three trials were identified that evaluated efficacy in 1) patients with Down's syndrome and dementia, 2) different races as a function of treatment center of a multicenter trial, and 3) depressed patients. Given the relatively small number of trials evaluating these variables within different populations and different pharmacological interventions, the findings of this review are inconclusive with respect to these variables. A significant gap in the literature has been identified.

Question 5: What is the evidence-base for the treatment of ischemic vascular dementia?

A total of 20 pharmacological interventions in 29 studies ^{17,36,38,44,46,70-72,81,92,96,98,102-104,106,107,117,126,128,133-141} were applied specifically to VaD classified dementias. The majority of these pharmacological interventions (n=14) were represented by single trials, limiting the ability to judge the evidence; these interventions included ateroid, buflomedil, cerebrolysin, sulphomucopolysaccharides (CDP choline), citalopram, donepezil, Ginkgo biloba, idebenone, minaprine, nimodipine, oxiracetam, 5-THF (trazodone), vincamine, and xantinolnicotinate. Six interventions had more than a single trial, and these included Choto-san (n=2), memantine (n=3), nicergoline (n=2), pentoxifylline (n=4), posatirelin (n=2), and propentofylline (n=2). In general, when the drug interventions were shown to be effective, it was in the domains of cognitive function (both general and specific) and global assessment. Other domains were less frequently evaluated. Several trials attempted to test for differences between VaD groups and other dementia types.

Discussion

The findings of this report suggest several important areas for future research using pharmacological treatments for dementia and these include:

Analytic framework of the intended aim of the therapy on the disease

- Better conceptualization and research design to capture "delay in progression."
- Clearer consensus on defining efficacy (benefits and clinically important change).
- Longer term studies (> 12 months).

Potential for bias

- Clarification of the role of industry sponsorship; one recommendation should be that all studies are required to disclose such information in future, including who analyzed the results.
- More concerted effort to incorporate unpublished studies and negative trials in future reviews.

Population

- Inclusion of the spectrum of severity in the patient populations (nothing to suggest that severe patients may not benefit from pharmacotherapy aimed at cognitive function improvement).
- The need for validation of trials and testing processes within cultures other than the traditional white population.
- Examining the efficacy of interventions in different sub-populations (age, disease severity levels, etc.).
- Better measurement and reporting of important patient characteristics (including baseline cognition scores, co-morbid conditions, the use of other medications, etc.).
- Inclusion of MCI type groups of subjects to evaluate “delay of onset” (studies in progress).

Outcomes

- Expansion of outcomes collected to include more than just cognitive function, and especially include caregiver burden and quality of life/ADL.
- Clear operational definitions for determining critical outcomes (delay to onset, delay to progression, important effect size, etc.).

- Understanding of how therapies are addressed and what outcomes are produced in different cultures.
- Production of other testing tools to detect both onset and responses to therapies across varied cultural groups.
- Improvement in the reporting of adverse events to evaluate harm and risk vs. benefit.
- Improvement in detailing adverse events associated with the duration period and those occurring following this period.

Analysis

- Appropriate analytical strategies that take into account intention to treat (ITT)/ last observation carried forward (LOCF) analyses; where possible both observed case and ITT/LOCF analyses should be presented.
- Sufficient data to estimate effect size, taking into account variability in both treated and control populations on the primary measures.
- Reporting the power of the study when findings are statistically non-significant.

Intervention

- Undertake more studies with direct comparison of drugs to determine the relative efficacy of agents.
- Improved description of the titration process.
- Improved collection of adverse events undertaken in a systematic fashion with standardized instruments.

Table 1. Pharmacological interventions and the number of trials (#) evaluated in this systematic review.

Cholinergic neurotransmitter modifying agents	
Antagonic Stress (2)	Metrifonate (9)
Acetyl-L-Carnitine (6)	Nicergoline (5)
Donepezil (11)	Physostigmine (4)
Eptastigmine (2)	Posatirelin (4)
Galantamine (6)	Rivastigmine (6)
Huperzine-A (2)	Sabeluzole (1)
Linopirdine (2)	Tacrine (8)
Mexofenoxate (1)	Velnacrine (3)
Non-cholinergic neurotransmitter/neuropeptide modifying agents	
Alaproclate (1)	Memantine (3)
Alprazolam (1)	Mianserin (1)
Anapso (1)	Minaprine (1)
BMV (Nootropic) (1)	Moclobemide (1)
Carbamazepine (2)	Naftidrofuryl (1)
Citalopram (2)	Olanzapine (2)
Diphenhydramine (1)	Oxazepam (1)
Divalproex (2)	Paroxetine (1)
Fluoxetine (2)	Perphenazine (1)
Fluvoxamine (1)	Phosphatidylserine (2)
Haloperidol (8)	Risperidone (2)
Imipramine (1)	Selegiline (6)
Lisuride (1)	Sertraline (2)
Lorazepam (2)	Thioridazine (1)
Loxapine (2)	Tiapride (2)
Lu25-109 (1)	Trazodone (2)
Maprotiline (1)	Xanomeline (1)
Melperone (1)	
Other agents	
5'-MTHF (1)	Misoprostol (1)
Aniracetam (1)	Monosialotetrahexosylganglioside (GM-1) (1)
Amitriptyline (1)	N-Acetylcysteine (1)
Ateroid (1)	Nimesulide (1)
Buflomedil (1)	Nimodipine (2)
Cerebrolysin (6)	Nizatidine (1)
Choro-San (1)	Nootropic (1)
Choto-San (1)	ORG 2766 (2)
Citicoline (2)	Oxiracetam (5)
Cyclandelate (2)	Pentoxifylline (4)
Denbufylline (1)	Piracetam (1)
Desferrioxamine (1)	Prednisone (1)
Diclofenac (1)	Propentofylline (4)
Ergokryptine (CMB 36-733) (1)	Pyritinol (1)
Ergokryptine (Dek) (1)	Silymarin + Tacrine (1)
Estrogens (5)	Simvastatin (1)
Ginkgo Biloba (3)	Sulphomucopolysaccharides (1)
Glycosaminoglycan Polysulfate (1)	Sulodexide (1)
Guanfacine (1)	Thiamine (1)
Hydergine (1)	Vasopressin (DDAVP) (1)
Hydroxychloroquine (1)	Vincamine (1)
Idebenone (5)	Vitamin E (2)
Indomethacin (1)	Xantinolnicotinate (1)

Availability of the Full Report

The full evidence report from which this summary was taken was prepared for the Agency for Healthcare Research and Quality (AHRQ) by McMaster University Evidence-based Practice Center under Contract No. 290-02-0020. It is expected to be available in April 2004. At that time, printed copies may be obtained free of charge from the AHRQ Publications Clearinghouse by calling 800-358-9295. Requesters should ask for Evidence Report/Technology Assessment No. 97, *Pharmacological Treatment of Dementia*. In addition, Internet users will be able to access the report and this summary online through AHRQ's Web site at www.ahrq.gov.

Suggested Citation

Santaguida PS, Raina P, Booker L, Patterson C, Baldassarre F, Cowan D, Gauld M, Levine M, Unsal A. Pharmacological Treatment of Dementia. Summary, Evidence Report/Technology Assessment No. 97 (Prepared by McMaster University Evidence-based Practice Center under Contract No. 290-02-0020). AHRQ Publication No. 04-E018-1. Rockville, MD: Agency for Healthcare Research and Quality. April 2004.

References

1. Thal LJ, Calvani M, Amato A, et al. A 1-year controlled trial of acetyl-L-carnitine in early-onset AD. *Neurology* 2000a; 55(6):805-10.
2. Thal LJ, Carta A, Clarke WR, et al. A 1-year multicenter placebo-controlled study of acetyl-L-carnitine in patients with Alzheimer's disease. *Neurology* 1996a; 47(3):705-11.
3. Sano M, Bell K, Cote L, et al. Double-blind parallel design pilot study of acetyl levocarnitine in patients with Alzheimer's disease. *Arch Neurol* 1992 Nov; 49(11):1137-41.
4. Livingston GA, Sax KB, McClenahan Z, et al. Acetyl-L-carnitine in dementia. *Int J Geriatr Psychiatry* 1991; 6(12):853-60.
5. Spagnoli A, Lucca U, Menasce G, et al. Long-term acetyl-L-carnitine treatment in Alzheimer's disease. *Neurology* 1991 Nov; 41(11):1726-32.
6. Rai G, Wright G, Scott L, et al. Double-blind, placebo-controlled study of acetyl-L-carnitine in patients with Alzheimer's disease. *Curr Med Res Opin* 1990; 11(10):638-47.
7. Dunne MP, Hartley LR. Scopolamine and the control of attention in humans. *Psychopharmacologia* 1986; 89(1):94-7.
8. Prasher VP, Huxley A, Haque MS, et al. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Down syndrome and Alzheimer's disease: Pilot study. *Int J Geriatr Psychiatry* 2002 Mar; 17(3):270-8.
9. Mohs RC, Doody RS, Morris JC, et al. A 1-year, placebo-controlled preservation of function survival study of donepezil in AD patients. *Neurology* 2001 Aug 14; 57(3):481-8.
10. Winblad B, Engedal K, Soininen H, et al. A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology* 2001b; 57(3):489-95.
11. Feldman H, Gauthier S, Hecker J, et al. A 24-week, randomized, double-blind study of donepezil in moderate to severe Alzheimer's disease. *Neurology* 2001 Aug 28; 57(4):613-20.
12. Tariot PN, Cummings JL, Katz IR, et al. A randomised, double-blind, placebo-controlled study of the efficacy and safety of Donepezil in patients with Alzheimer's disease in the nursing home setting. *J Am Geriatr Soc* 2001a; 49(12):1590-9.
13. Burns A, Rossor M, Hecker J, et al. The effects of donepezil in Alzheimer's disease—results from a multinational trial. *Dement Geriatr Cogn Disord* 1999 May; 10(3):237-44.
14. Rogers SL, Farlow MR, Doody RS, et al. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. *Neurology* 1998b; 50(1):136-45.
15. Rogers SL, Doody RS, Mohs RC, et al. Donepezil improves cognition and global function in Alzheimer disease: a 15-week, double-blind, placebo-controlled study. *Arch Intern Med* 1998a; 158(9):1021-31.
16. Rogers SL, Friedhoff LT. The efficacy and safety of donepezil in patients with Alzheimer's disease: results of a US multicentre, randomized, double-blind, placebo-controlled trial. The Donepezil Study Group. *Dementia* 1996 Nov; 7(6):293-303.
17. Pratt RD, Perdomo CA. Donepezil-treated patients with probable vascular dementia demonstrate cognitive benefits. *Ann N Y Acad Sci* 2002 Nov; 977:513-22.
18. Thomas A, Iacono D, Bonanni L, et al. Donepezil, rivastigmine, and vitamin E in Alzheimer disease: a combined P300 event-related potentials/neuropsychologic evaluation over 6 months. *Clin Neuropharmacol* 2001 Jan; 24(1):31-42.
19. Erkinjuntti T, Kurz A, Gauthier S, et al. Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: a randomised trial. *Lancet* 2002 Apr 13; 359(9314):1283-90.
20. Rockwood K, Mintzer J, Truyen L, et al. Effects of a flexible galantamine dose in Alzheimer's disease: a randomised, controlled trial. *J Neurol Neurosurg Psychiatry* 2001; 71(5):589-95.
21. Wilkinson D, Murray J. Galantamine: a randomized, double-blind, dose comparison in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 2001 Sep; 16(9):852-7.
22. Tariot PN, Solomon PR, Morris JC, et al. A 5-month, randomized, placebo-controlled trial of galantamine in AD. The Galantamine USA-10 Study Group. *Neurology* 2000 Jun 27; 54(12):2269-76.
23. Wilcock GK, Lilienfeld S, Gaens E. Efficacy and safety of galantamine in patients with mild to moderate Alzheimer's disease: multicentre randomised controlled trial. *Galantamine International-1 Study Group. BMJ* 2000 Dec 9; 321(7274):1445-9.
24. Raskind MA, Peskind ER, Wessel T, et al. Galantamine in AD: a 6-month randomized, placebo-controlled trial with a 6-month extension. The Galantamine USA-1 Study Group. *Neurology* 2000 Jun 27; 54(12):2261-8.
25. Schulz R, O'Brien A, Czaja S, et al. Dementia caregiver intervention research: in search of clinical significance. *Gerontologist* 2002 Oct; 42(5):589-602.
26. Jann MW, Cyrus PA, Eisner LS, et al. Efficacy and safety of a loading-dose regimen versus a no-loading-dose regimen of metrifonate in the symptomatic treatment of Alzheimer's disease: a randomized, double-masked, placebo-controlled trial. *Clin Ther* 1999; 21(1):88-102.
27. Raskind MA, Cyrus PA, Ruzicka BB, et al. The effects of metrifonate on the cognitive, behavioral, and functional performance of Alzheimer's disease patients. *J Clin Psychiatry* 1999; 60(5):318-25.

28. Becker RE, Colliver JA, Markwell SJ, et al. Effects of metrifonate on cognitive decline in Alzheimer's disease: a double-blind, placebo-controlled, 6-month study. *Alzheimer Dis Assoc Disord* 1998 Mar; 12(1):54-7.
29. Morris JC, Cyrus PA, Orazem J, et al. Metrifonate benefits cognitive, behavioral, and global function in patients with Alzheimer's disease. *Neurology* 1998 May; 50(5):1222-30.
30. Cummings JL, Cyrus PA, Bieber F, et al. Metrifonate treatment of the cognitive deficits of Alzheimer's disease. *Neurology* 1998b; 50(5):1214-21.
31. Pettigrew LC, Bieber F, Lettieri J, et al. Pharmacokinetics, pharmacodynamics, and safety of metrifonate in patients with Alzheimer's disease. *J Clin Pharmacol* 1998 Mar; 38(3):236-45.
32. Becker RE, Colliver JA, Markwell SJ, et al. Double-blind, placebo-controlled study of metrifonate, an acetylcholinesterase inhibitor, for Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1996; 10(3):124-31.
33. Cummings J, Bieber F, Mas J, et al. Metrifonate in Alzheimer's disease: results of a dose finding study. *Alzheimers Dis Biol Diagn Ther* 1997:665-9.
34. Nordberg A, Svensson AL. Cholinesterase inhibitors in the treatment of Alzheimer's disease: a comparison of tolerability and pharmacology. *Drug Saf* 1998 Dec; 19(6):465-80.
35. Winblad B, Bonura ML, Rossini BM, et al. Nicergoline in the treatment of mild-to-moderate Alzheimer's disease: a European multicentre trial. *Clin Drug Investig* 2001a(9):621-32.
36. Herrmann WM, Stephan K, Gaede K, et al. A multicenter randomized double-blind study on the efficacy and safety of nicergoline in patients with multi-infarct dementia. *Dement Geriatr Cogn Disord* 1997 Jan; 8(1):9-17.
37. Nappi G, Bono G, Merlo P, et al. Long-term nicergoline treatment of mild to moderate senile dementia. Results of a multicentre, double-blind, placebo-controlled study. *Clin Drug Investig* 1997(6):308-16.
38. Saletu B, Paulus E, Linzmayer L, et al. Nicergoline in senile dementia of Alzheimer type and multi-infarct dementia: a double-blind, placebo-controlled, clinical and EEG/ERP mapping study. *Psychopharmacologia* 1995 Feb; 117(4):385-95.
39. Schneider F, Popa R, Mihalas G, et al. Superiority of antagonistic-stress composition versus nicergoline in gerontopsychiatry. *Ann N Y Acad Sci* 1994 Jun; 717:332-42.
40. Moller HJ, Hampel H, Hegerl U, et al. Double-blind, randomized, placebo-controlled clinical trial on the efficacy and tolerability of a physostigmine patch in patients with senile dementia of the Alzheimer type. *Pharmacopsychiatry* 1999 May; 32(3):99-106.
41. Thal LJ, Schwartz G, Sano M, et al. A multicenter double-blind study of controlled-release physostigmine for the treatment of symptoms secondary to Alzheimer's disease. *Neurology* 1996b; 47(6):1389-95.
42. Van Dyck CH, Newhouse P. Extended-release physostigmine in Alzheimer's disease: a multicenter, double-blind, 12-week study with dose enrichment. *Arch Gen Psychiatry* 2000; 57(2):157-64.
43. Thal LJ, Ferguson JM, Mintzer J, et al. A 24-week randomized trial of controlled-release physostigmine in patients with Alzheimer's disease. *Neurology* 1999 Apr 12; 52(6):1146-52.
44. Ferrari E, Cucinotta D, Albizzati MG, et al. Effectiveness and safety of posatirelin in the treatment of senile dementia: a multicenter, double-blind, placebo-controlled study. *Arch Gerontol Geriatr* 1998; 27(Suppl 6):163-74.
45. Gasbarrini G, Stefanini G, Addolorato G, et al. Posatirelin for the treatment of degenerative and vascular dementia: results of explanatory and pragmatic efficacy analyses. *Arch Gerontol Geriatr* 1997; 26(1):33-47.
46. Parnetti L, Ambrosoli L, Agliati G, et al. Posatirelin in the treatment of vascular dementia: a double-blind multicentre study vs placebo. *Acta Neurol Scand* 1996 Jun; 93(6):456-63.
47. Parnetti L, Ambrosoli L, Abate G, et al. Posatirelin for the treatment of late-onset Alzheimer's disease: a double-blind multicentre study vs citicoline and ascorbic acid. *Acta Neurol Scand* 1995 Aug; 92(2):135-40.
48. Potkin SG, Anand R, Fleming K, et al. Brain metabolic and clinical effects of rivastigmine in Alzheimer's disease. *Int J Neuropsychopharmacol* 2001 Sep; 4(3):223-30.
49. McKeith I, Del Ser T, Spano P, et al. Efficacy of rivastigmine in dementia with Lewy bodies: a randomised, double-blind, placebo-controlled international study. *Lancet* 2000 Dec 16; 356(9247):2031-6.
50. Forette F, Anand R, Gharabawi G. A phase II study in patients with Alzheimer's disease to assess the preliminary efficacy and maximum tolerated dose of rivastigmine (Exelon registered). *Eur J Neurol* 1999; 6(4):423-9.
51. Rosler M, Anand R, Cicin-Sain A, et al. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: international randomised controlled trial. *BMJ* 1999 Mar 6; 318(7184):633-8.
52. Agid Y, Dubois B, Anand R, et al. Efficacy and tolerability of rivastigmine in patients with dementia of the Alzheimer type. *Curr Ther Res Clin Exp* 1998; 59(12):837-45.
53. Corey-Bloom JR, Anand JV, Veach J, et al. A randomized trial evaluating the efficacy and safety of ENA 713 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe Alzheimer's disease. *Int J Geriatr Psychopharmacol* 1998; 1:55-65.
54. Knapp MJ, Knopman DS, Solomon PR, et al. A 30-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. The Tacrine Study Group. *JAMA* 1994b; 271(13):985-91.
55. Maltby N, Broe GA, Creasey H, et al. Efficacy of tacrine and lecithin in mild to moderate Alzheimer's disease: double blind trial. *BMJ* 1994 Apr 2; 308(6933):879-83.
56. Prentice N, Van BM, Dougall NJ, et al. A double-blind, placebo-controlled study of tacrine in patients with Alzheimer's disease using SPET. *J Psychopharmacol (Oxf)* 1996; 10(3):175-81.
57. Weinstein HC, Teunisse S, Van Gool WA. Tetrahydroaminoacridine and lecithin in the treatment of Alzheimer's disease. Effect on cognition, functioning in daily life, behavioural disturbances and burden experienced by the carers. *J Neurol* 1991 Feb; 238(1):34-8.
58. Wong WJ, Liu HC, Fuh JL, et al. A double-blind, placebo-controlled study of tacrine in Chinese patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 1999 Jul; 10(4):289-94.
59. Wood PC, Castleden CM. A double-blind, placebo controlled, multicentre study of tacrine for Alzheimer's disease. *Int J Geriatr Psychiatry* 1994; 9(8):649-54.
60. Allain H, Schuck S, Lebreton S, et al. Aminotransferase levels and silymarin in de novo tacrine-treated patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 1999 May; 10(3):181-5.
61. Gutzmann H, Kuhl KP, Hadler D, et al. Safety and efficacy of idebenone versus tacrine in patients with Alzheimer's disease: results of a randomized, double-blind, parallel-group multicenter study. *Pharmacopsychiatry* 2002 Jan; 35(1):12-8.
62. Antuono PG. Effectiveness and safety of velnacrine for the treatment of Alzheimer's disease. A double-blind, placebo-controlled study. Mentane Study Group. *Arch Intern Med* 1995 Sep 11; 155(16):1766-72.

63. Huff FJ, Antuono P, Murphy M, et al. Potential clinical use of an adrenergic/cholinergic agent (HP 128) in the treatment of Alzheimer's disease. *Ann N Y Acad Sci* 1991; 640:263-7.
64. Zemlan FP, Folks DG, Goldstein BJ, et al. Velnacrine for the treatment of Alzheimer's disease: a double-blind, placebo-controlled trial. *J Neural Transm Gen Sect* 1996; 103(8-9):1105-16.
65. Allain H, Dautzenberg PH, Maurer K, et al. Double blind study of tiapride versus haloperidol and placebo in agitation and aggressiveness in elderly patients with cognitive impairment. *Psychopharmacologia* 2000 Mar; 148(4):361-6.
66. Teri L, Logsdon RG, Peskind E, et al. Treatment of agitation in AD: a randomized, placebo-controlled clinical trial. *Neurology* 2000 Nov 14; 55(9):1271-8.
67. Petrie WM, Ban TA, Berney S, et al. Loxapine in psychogeriatrics: a placebo- and standard-controlled clinical investigation. *J Clin Psychopharmacol* 1982 Apr; 2(2):122-6.
68. De Deyn PP, Rabheru K. A randomized trial of risperidone, placebo, and haloperidol for behavioral symptoms of dementia. *Neurology* 1999 Sep 22; 53(5):946-55.
69. Auchus AP, Bissey-Black C. Pilot study of haloperidol, fluoxetine, and placebo for agitation in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1997; 9(4):591-3.
70. Winblad B, Poritis N. Memantine in severe dementia: results of the 9M-Best Study (benefit and efficacy in severely demented patients during treatment with memantine). *Int J Geriatr Psychiatry* 1999 Feb; 14(2):135-46.
71. Orgogozo J, Rigaud AS, Stoffler A, et al. Efficacy and safety of memantine in patients with mild to moderate vascular dementia. *Stroke* 2002; 33:1834-9.
72. Wilcock G, Mobius HJ, Stoffler A. A double-blind, placebo-controlled multicentre study of memantine in mild to moderate vascular dementia (MMM500). *Int Clin Psychopharmacol* 2002; 17(6):297-305.
73. Sano M, Ernesto C, Thomas RG, et al. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *N Engl J Med* 1997; 336(17):1216-22.
74. Mangoni A, Grassi MP, Frattola L, et al. Effects of a MAO-B inhibitor in the treatment of Alzheimer disease. *Eur Neurol* 1991; 31(2):100-7.
75. Agnoli A, Fabbrini G, Fioravanti M, et al. CBF and cognitive evaluation of Alzheimer type patients before and after IMAO-B treatment: a pilot study. *Eur Neuropsychopharmacol* 1992 Mar; 2(1):31-5.
76. Burke WJ, Roccaforte WH, Wengel SP, et al. L-deprenyl in the treatment of mild dementia of the Alzheimer type: results of a 15-month trial. *J Am Geriatr Soc* 1993a; 41(11):1219-25.
77. Filip V, Kolibas E. Selegiline in the treatment of Alzheimer's disease: a long-term randomized placebo-controlled trial. Czech and Slovak Senile Dementia of Alzheimer Type Study Group. *J Psychiatry Neurosci* 1999 May; 24(3):234-43.
78. Freedman M, Rewilak D, Xerri T, et al. L-deprenyl in Alzheimer's disease: cognitive and behavioral effects. *Neurology* 1998; 50(3):660-8.
79. Ruether E, Ritter R, Apecechea M, et al. Efficacy of the peptidergic nootropic drug cerebrolysin in patients with senile dementia of the Alzheimer type (SDAT). *Pharmacopsychiatry* 1994 Jan; 27(1):32-40.
80. Bae CY, Cho CY, Cho K, et al. A double-blind, placebo-controlled, multicenter study of Cerebrolysin for Alzheimer's disease. *J Am Geriatr Soc* 2000 Dec; 48(12):1566-71.
81. Xiao S, Yan H, Yao P, et al. The efficacy of cerebrolysin in patients with vascular dementia: results of a Chinese multicentre, randomised, double-blind, placebo-controlled trial. *Hong Kong Journal of Psychiatry* 1999(2):13-9.
82. Xiao S, Yan H, Yao P, et al. Efficacy of PFP 1070 (cerebrolysin) in patients with Alzheimer's disease: a multicentre, randomised, double-blind, placebo-controlled trial. *Clin Drug Investig* 2000(1):43-53.
83. Ruether E, Husmann R, Kinzler E, et al. A 28-week, double-blind, placebo-controlled study with Cerebrolysin in patients with mild to moderate Alzheimer's disease. *Int Clin Psychopharmacol* 2001 Sep; 16(5):253-63.
84. Panisset M, Gauthier S, Moessler H, et al. Cerebrolysin in Alzheimer's disease: a randomized, double-blind, placebo-controlled trial with a neurotrophic agent. *J Neural Transm Gen Sect* 2002; 109(7-8):1089-104.
85. Asthana S, Baker LD, Craft S, et al. High-dose estradiol improves cognition for women with AD: results of a randomized study. *Neurology* 2001 Aug 28; 57(4):605-12.
86. Wang PN, Liao SQ, Liu RS, et al. Effects of estrogen on cognition, mood, and cerebral blood flow in AD: a controlled study. *Neurology* 2000 Jun 13; 54(11):2061-6.
87. Henderson VW, Paganini-Hill A, Miller BL, et al. Estrogen for Alzheimer's disease in women: randomized, double-blind, placebo-controlled trial. *Neurology* 2000 Jan 25; 54(2):295-301.
88. Mulnard RA, Cotman CW, Kawas C, et al. Estrogen replacement therapy for treatment of mild to moderate Alzheimer disease: a randomized controlled trial. *JAMA* 2000(8):1007-15.
89. Kyomen HH, Satlin A, Hennen J, et al. Estrogen therapy and aggressive behavior in elderly patients with moderate-to-severe dementia: results from a short-term, randomized, double-blind trial. *Am J Geriatr Psychiatry* 1999; 7(4):339-48.
90. Le Bars PL, Katz MM, Berman N, et al. A placebo-controlled, double-blind, randomized trial of an extract of Ginkgo biloba for dementia. *JAMA* 1997(16):1327-32.
91. Maurer K, Ihl R, Dierks T, et al. Clinical efficacy of Ginkgo biloba special extract EGb 761 in dementia of the Alzheimer type. *J Psychiatr Res* 1997 Nov; 31(6):645-55.
92. Kanowski S, Herrmann WM, Stephan K, et al. Proof of efficacy of the Ginkgo biloba special extract EGb 761 in outpatients suffering from mild to moderate primary degenerative dementia of the Alzheimer type or multi-infarct dementia. *Pharmacopsychiatry* 1996 Mar; 29(2):47-56.
93. Weyer G, Babej-Dolle RM, Hadler D, et al. A controlled study of 2 doses of idebenone in the treatment of Alzheimer's disease. *Neuropsychobiology* 1997; 36(2):73-82.
94. Gutzmann H, Hadler D. Sustained efficacy and safety of idebenone in the treatment of Alzheimer's disease: update on a 2-year double-blind multicentre study. *J Neural Transm Suppl* 1998; 54:301-10.
95. Bergamasco B, Scarzella L, La Commare P. Idebenone, a new drug in the treatment of cognitive impairment in patients with dementia of the Alzheimer type. *Funct Neurol* 1994 May; 9(3):161-8.
96. Marigliano V, Abate G, Barbagallo-Sangiorgi G, et al. Randomized, double-blind, placebo controlled, multicentre study of idebenone in patients suffering from multi-infarct dementia. *Arch Gerontol Geriatr* 1992; 15(3):239-48.
97. Bottini G, Vallar G, Cappa S, et al. Oxiracetam in dementia: a double-blind, placebo-controlled study. *Acta Neurol Scand* 1992 Sep; 86(3):237-41.

98. Maina G, Fiori L, Torta R, et al. Oxiracetam in the treatment of primary degenerative and multi-infarct dementia: a double-blind, placebo-controlled study. *Neuropsychobiology* 1989; 21(3):141-5.
99. Mangoni A, Perin C, Smirne S, et al. A double-blind, placebo-controlled study with oxiracetam in demented patients administered the Luria-Nebraska Neuropsychological Battery. *Drug Dev Res* 1988; 14(3-4):217-4.
100. Rozzini R, Zanetti O, Bianchetti A. Effectiveness of oxiracetam therapy in the treatment of cognitive deficiencies secondary to primary degenerative dementia. *Acta Neurol (Napoli)* 1992 Apr; 14(2):117-26.
101. Burgio LD, Reynolds CFI, Janosky JE, et al. A behavioral microanalysis of the effects of haloperidol and oxazepam in demented psychogeriatric inpatients. *Int J Geriatr Psychiatry* 1992; 7(4):253-62.
102. Ghose K. Oxpentifylline in dementia: a controlled study. *Arch Gerontol Geriatr* 1987 Apr; 6(1):19-26.
103. Black RS, Barclay LL, Nolan KA, et al. Pentoxifylline in cerebrovascular dementia. *J Am Geriatr Soc* 1992 Mar; 40(3):237-44.
104. Knezevic S. European Pentoxifylline Multi-Infarct Dementia Study. *Eur Neurol* 1996; 36(5):315-21.
105. Mielke R, Ghaemi M, Kessler J, et al. Propentofylline enhances cerebral metabolic response to auditory memory stimulation in Alzheimer's disease. *JNS* 1998 Jan 21; 154(1):76-82.
106. Mielke R, Kittner B, Ghaemi M, et al. Propentofylline improves regional cerebral glucose metabolism and neuropsychologic performance in vascular dementia. *JNS* 1996 Sep 15; 141(1-2):59-2.
107. Marcusson J, Rother M, Kittner B, et al. A 12-month, randomized placebo-controlled trial of propentofylline (HWA 285) in patients with dementia according to DSM III-R. *Dement Geriatr Cogn Disord* 1997; 8(5):320-8.
108. Saletu B, Moller HJ, Grunberger J, et al. Propentofylline in adult-onset cognitive disorders: double-blind, placebo-controlled, clinical, psychometric and brain mapping studies. *Neuropsychobiology* 1990 Sep; 24(4):173-84.
109. Convit A, de Asis J, de Leon MJ, et al. Atrophy of the medial occipitotemporal, inferior, and middle temporal gyri in non-demented elderly predict decline to Alzheimer's disease. *Neurobiol Aging* 2000 Jan; 21(1):19-26.
110. Rapp S, Brenes G, Marsh AP. Memory enhancement training for older adults with mild cognitive impairment: a preliminary study. *Aging Ment Health* 2002 Feb; 6(1):5-11.
111. Johnson SA, Simmon VF. Randomized, double-blind, placebo-controlled international clinical trial of the AMPAKINE CX516 in elderly participants with mild cognitive impairment. A progress report. *J Mol Neurosci* 2002; 19(1-2):197-200.
112. Sherwin BB. Estrogen and cognitive functioning in men with mild cognitive impairment. *J Mol Neurosci* 2002; 19(1-2):219-23.
113. Meehan KM, Wang H, David SR, et al. Comparison of rapidly acting intramuscular olanzapine, lorazepam, and placebo: a double-blind, randomized study in acutely agitated patients with dementia. *Neuropsychopharmacology* 2002 Apr; 26(4):494-504.
114. Carlyle W, Ancill RJ, Sheldon L. Aggression in the demented patient: a double-blind study of loxapine versus haloperidol. *Int Clin Psychopharmacol* 1993; 8(2):103-8.
115. Ancill RJ, Carlyle WW, Liang RA, et al. Agitation in the demented elderly: a role for benzodiazepines? *Int Clin Psychopharmacol* 1991; 6(3):141-6.
116. Coccaro EF, Kramer E, Zemishlany Z, et al. Pharmacologic treatment of noncognitive behavioral disturbances in elderly demented patients. *Am J Psychiatry* 1990 Dec; 147(12):1640-5.
117. Cucinotta D, Romagnoli S, Godoli G, et al. Comparison of sulfomucopolysaccharides and cytidine diphosphocholine in the treatment of multi-infarct dementia: a randomized double-blind test. *Curr Ther Res Clin Exp* 1988; 43(1):12-20.
118. Karlsson I, Godderis J, Augusto De Mendonca LC, et al. A randomised, double-blind comparison of the efficacy and safety of citalopram compared to mianserin in elderly, depressed patients with or without mild to moderate dementia. *Int J Geriatr Psychiatry* 2000 Apr; 15(4):295-305.
119. Pollock BG, Mulsant BH, Rosen J, et al. Comparison of citalopram, perphenazine, and placebo for the acute treatment of psychosis and behavioral disturbances in hospitalized, demented patients. *Am J Psychiatry* 2002; 159(3):460-5.
120. Barnes R, Veith R, Okimoto J. Efficacy of antipsychotic medications in behaviorally disturbed dementia patients. *Am J Psychiatry* 1982; 139(9):1170-4.
121. Chan WC, Lam LC, Choy CN, et al. A double-blind randomised comparison of risperidone and haloperidol in the treatment of behavioural and psychological symptoms in Chinese dementia patients. *Int J Geriatr Psychiatry* 2001 Dec; 16(12):1156-62.
122. Katona CL, Hunter BN, Bray J. A double-blind comparison of the efficacy and safety of paroxetine and imipramine in the treatment of depression with dementia. *Int J Geriatr Psychiatry* 1998 Feb; 13(2):100-8.
123. Taragano FE, Lyketsos CG, Mangone CA, et al. A double-blind, randomized, fixed-dose trial of fluoxetine vs. amitriptyline in the treatment of major depression complicating Alzheimer's disease. *Psychosomatics* 1997 May; 38(3):246-52.
124. Popa R, Schneider F, Mihalas G, et al. Antagonic-stress superiority versus meclofenoxate in gerontopsychiatry. *Arch Gerontol Geriatr* 1994; 18(Suppl 4):197-206.
125. Spilich GJ, Wannenmacher W, Duarte A, et al. Efficacy of pyritinol versus hydergine upon cognitive performance in patients with senile dementia of the Alzheimer's type: a double-blind multi-center trial. *Alzheimers Res* 1996(3):79-84.
126. Parnetti L, Mari D, Abate G, et al. Vascular dementia Italian sulodexide study (VA.D.I.S.S.). Clinical and biological results. *Thromb Res* 1997 Jul 15; 87(2):225-33.
127. Gutzmann H, Kuhl KP, Kanowski S, et al. Measuring the efficacy of psychopharmacological treatment of psychomotoric restlessness in dementia: clinical evaluation of tiapride. *Pharmacopsychiatry* 1997 Jan; 30(1):6-11.
128. Passeri M, Cucinotta D, Abate G, et al. Oral 5'-methyltetrahydrofolic acid in senile organic mental disorders with depression: results of a double-blind multicenter study. *Aging (Milano)* 1993 Feb; 5(Milano):63-71.
129. Alvarez XA, Pichel V, Perez P, et al. Double-blind, randomized, placebo-controlled pilot study with anapso in senile dementia: effects on cognition, brain bioelectrical activity and cerebral hemodynamics. *Methods Find Exp Clin Pharmacol* 2000 Sep; 22(7):585-94.
130. Ruether E, Alvarez XA, Rainer M, et al. Sustained improvement of cognition and global function in patients with moderately severe Alzheimer's disease: a double-blind, placebo-controlled study with the neurotrophic agent Cerebrolysin. *J Neural Transm Suppl* 2002(62):265-75.
131. Schellenberg R, Todorova A, Wedekind W, et al. Pathophysiology and psychopharmacology of dementia: a new study design. *Neuropsychobiology* 1997; 35(3):132-42.

132. Reifler BV, Teri L, Raskind M, et al. Double-blind trial of imipramine in Alzheimer's disease patients with and without depression. *Am J Psychiatry* 1989 Jan; 146(1):45-9.
133. Ban TA, Morey LC, Santini V. Clinical investigations with ateroid in old-age dementias. *Semin Thromb Hemost* 1991b; 17(Suppl 2):161-3.
134. Cucinotta D, Aveni Casucci MA, Pedrazzi F, et al. Multicentre clinical placebo-controlled study with buflomedil in the treatment of mild dementia of vascular origin. *J Int Med Res* 1992 Apr; 20(2):136-49.
135. Shimada Y, Terasawa K, Yamamoto T, et al. A well-controlled study of Choto-san and placebo in the treatment of vascular dementia. *J Tradit Med* 1994; 11:246-55.
136. Terasawa K, Shimada Y, Kita T, et al. Choto-san in the treatment of vascular dementia: a double-blind, placebo-controlled study. *Phytomedicine* 1997; 4(1):15-22.
137. Nyth AL, Gottfries CG. The clinical efficacy of citalopram in treatment of emotional disturbances in dementia disorders: a Nordic multicentre study. *Br J Psychiatry* 1990 Dec; 157:894-901.
138. Passeri M, Cucinotta D, de Mello M, et al. Comparison of minaprine and placebo in the treatment of Alzheimer's disease and multi-infarct dementia. *Int J Geriatr Psychiatry* 1987; 2(2):97-103.
139. Saletu B, Anderer P, Semlitsch HV. Relations between symptomatology and brain function in dementias: double-blind, placebo-controlled, clinical and EEG/ERP mapping studies with nicergoline. *Dement Geriatr Cogn Disord* 1997; Vol 8(Suppl 1):12-21.
140. Pantoni L, Rossi R, Inzitari D, et al. Efficacy and safety of nimodipine in subcortical vascular dementia: a subgroup analysis of the Scandinavian Multi-Infarct Dementia Trial. *JNS* 2000b; 175(2):124-34.
141. Fischhof PK, Moslinger-Gehmayr R, Herrmann WM, et al. Therapeutic efficacy of vincamine in dementia. *Neuropsychobiology* 1996; 34(1):29-35.



www.ahrq.gov
AHRQ Pub. No. 04-E018-1
April 2004

ISSN 1530-440X

Pharmacological Treatment of Dementia

Evidence Report

Chapter 1. Introduction

This review focuses on the pharmacological treatment of dementia. Dementia is a syndrome of acquired cognitive defects sufficient to interfere with social or occupational functioning, which results from various central neurodegenerative and ischemic processes. Dementia has become a major public health problem due to its increasing prevalence, long duration, caregiver burden, and high financial cost of care. The prevalence of dementia varies as a function of the defining criteria as shown by Erkinjuntti et al. (1997),¹ who showed a range from 3.1% using the International Classification of Diseases, Tenth Revision (ICD-10) criteria, up to 29.1% using the Diagnostic and Statistical Manual, Third Edition (DSM-III) criteria. Jorm et al. (1987)² conducted a meta-analysis based on 22 international studies and found that the actual prevalence rates differed significantly from study to study. However, this meta-analysis demonstrated that the prevalence increased exponentially with age. The prevalence ranged from 0.7% for 60 – 64 year olds to 24% for people over the age of 85 years. In the United States, the prevalence of Alzheimer's disease (AD) is projected to quadruple to one in 45 Americans in the next 50 years³ across all ages. The Canadian Study of Health & Aging (CSHA)⁴ estimated the prevalence of dementia in Canada at 8% (approximately 252,600 cases) in 1991 among seniors over the age of 65 years. The prevalence of dementia increases to 34% among those aged 85 years or more.⁴ The age-standardized incidence of dementia in Canada has been estimated at 21.8 per thousand for females and 19.1 per thousand for males.⁵ The prevalence is expected to double to half a million cases in Canada by 2013.⁶ Because the world's population is progressively aging, especially in the developed nations, more people are falling into age groups where the prevalence of dementia is highest. From a clinical perspective, dementia predominately affects 1) cognition, 2) behavior/mood, 3) physical functions and activities of daily living, and 4) caregiver burden. Therapeutic interventions for dementia aim to affect these four primary domains.

Pharmacotherapy is often the primary intervention used to improve symptoms or delay the progression of dementia syndromes. The pharmacological agents used vary significantly with respect to their therapeutic actions. The most common pharmacological agents used in North America modify the activity of cholinesterases—enzymes, which degrade acetylcholine, a neurotransmitter that is critical to the neurons involved in cognition (e.g. memory, thought, and judgment). Other approaches include the use of anti-oxidants, which work by minimizing the effects of free radicals that are released through normal oxidative metabolism. These free radicals may cause neuronal damage and play a role in the development of dementia. Similarly, it is believed that inflammation contributes to nerve cell damage and dementia; hence anti-inflammatory drugs may act by decreasing inflammation, potentially reducing nerve degeneration, which may in turn slow or even prevent dementia illnesses.

Other pharmacological interventions that have been studied include cholesterol-lowering agents, anti-hypertensives, folic acid, hormones (e.g. estrogen), behavior and mood altering drugs, anti-amyloid strategies (e.g. immunization, aggregation inhibitors, and secretase inhibitors), transition metal chelators, nerve growth factors, and agents that target neurotransmitters other than acetylcholine and its receptors. The various pharmacotherapeutic agents available to treat problems associated with dementia have varying levels of evidence to support their efficacy. This report is a systematic evaluation of the evidence for pharmacological

interventions in the treatment of dementia in the domains of cognition, global function, behavior/mood, quality of life/ADL, and caregiver burden.

Diagnosis of Dementia

Determination of disease onset presents considerable difficulty, as dementia, by definition, has an insidious and gradual progression. A number of diagnostic models have been used to classify dementia. In 1988 the National Institute of Neurological and Communicative Disorders and Stroke - Alzheimer's Disease and Related Disorders Association (NINCDS/ADRDA) research on diagnostic criteria were published for AD, which served to increase the validity and reliability of the clinical diagnosis.^{7,8} Trials published prior to this time may reflect mixed populations other than AD. Other models used to diagnose dementia include: the International Classification of Diseases (ICD) version 9 or 10, the Diagnostic and Statistical Manual of Mental Disorders (DSM) III, III-R, and IV (American Psychiatric Association), and the NINCDS/ADRDA.⁹ The difficulty with these different diagnostic criteria for dementia is that they are not interchangeable. Erkinjuntti et al. (1997)¹ compared six commonly used classification schemes (DSM-III, DSM-III-R, DSM-IV, ICD-9, ICD-10, and the Cambridge Examination for Mental Disorders in the Elderly (CAMDEX)). They showed that the prevalence of dementia can differ by a factor of 10 depending on the diagnostic criteria used. Two other studies have demonstrated that the prevalence of vascular dementia (VaD) varies with the classification system and therefore these criteria for diagnosis are not interchangeable.^{10,11} Furthermore, there is controversy about the validity of the clinical classification of VaD, as autopsy confirmation often does not substantiate the clinical diagnosis.^{12,13} The majority of dementias were actually AD with co-existing vascular and Parkinson's disease lesions.¹⁴ In contrast, the clinical accuracy of AD diagnosis is relatively high.⁷ The discovery that a long pre-clinical period precedes AD has led to the establishment of early diagnostic indices of dementia. This border zone between normality and dementia has been given numerous names and definitions, which include: benign senescent forgetfulness (BSF), age associated memory impairment (AAMI), age-consistent memory impairment (ACMI), age-associated cognitive decline (AACD), mild cognitive impairment (MCI), cognitive loss no dementia (CLOND), and cognitive impairment but not dementia (CIND). The prevalence for this pre-clinical or mild form of cognitive decline varies with the classification system used.¹⁵ Unfortunately, the classifications used to diagnose early mild cognitive decline are not interchangeable. MCI¹⁶ is emerging as the preferred term for this condition¹⁷ using the criteria of Petersen et al.¹⁶ Ritchie et al.¹⁸ (2001) estimated the prevalence of MCI to be 3.2% with an 11.1% conversion rate to dementia within a 3 year period.

Analytic Framework: Understanding Therapeutic Aims of Pharmacological Treatment

Dementia is a chronic progressive disease for which no known cure exists. Pharmacological interventions used to treat dementia are intended to achieve at least one of the following broad therapeutic aims:


Prevention of onset of the disease. In the context of this review, this applies to those at greatest risk (such as those with the clinical diagnosis of MCI) of conversion to a dementia syndrome.

Symptomatic treatment of the disease. Symptomatic benefit can be described as maintenance (or stabilization) or improvement of the current cognitive, behavioral, functional, or caregiver status only while on active treatment with the pharmacological intervention. Withdrawal of the pharmacological therapy results in a decline towards baseline or placebo levels of relevant outcomes.

Delay in the progression of the disease. A therapeutic intervention that brings about delay in the progression of the disease can be described as either 1) one that maintains (or stabilizes) or improves current cognitive, behavioral, functional, or caregiver status, which is sustained, or 2) one that can be shown to alter the rate of decline of the disease progression, even when the drug is withdrawn.

Figure 1 details the analytic framework for the progression of dementia and shows when various pharmacological interventions would ideally be administered for the intended therapeutic benefit within this pathway. The scope of this review did not include the evaluation of normal healthy aging populations. Rather, pharmacological agents intended for populations at increased risk of conversion to dementia syndromes, such as MCI, were eligible for evaluation in this systematic review.

Figure 1. Pathway for the progression of dementia and the ideal application of drug interventions within this framework.



Understanding Efficacy of Pharmacological Interventions in Dementia Trials

It has been suggested that dementia does not have uniformly accepted criteria for disease progression or consensus regarding the magnitude of clinically important changes.^{19,20} With respect to the therapeutic aim, the practical consequences of these unresolved issues are that the same efficacy variables have been used to both show evidence of symptomatic benefit and demonstrate the effects on disease progression. Thus, the design of a clinical trial (rather than the outcome) is critical to demonstrating which of these two therapeutic outcomes (symptomatic benefit or delay in progression) is being achieved with the pharmacological agent.²¹

Irrespective of which therapeutic aim is being achieved by the pharmacological agent, the lack of consensus on these two issues has even more important implications when considering the definition of “efficacy” for either treatment goal. A change in a relevant outcome measure that is due to factors other than chance is deemed statistically significant. The criteria to determine efficacy solely on statistical significance have long been recognized as problematic from an interpretation perspective. Clinically meaningful change reflects a different level of “significance” and often requires consensus among experts within the field to establish what magnitude of change is important.²⁰

Efficacy as Measured by Clinical versus Statistical Significance

The dementia literature is not consistent in the criteria used for establishing efficacy, and there is no consensus on the meaning of clinical significance in the changes observed.^{20,22} In general, attempts are made to select an outcome measuring an important dementia attribute (such as cognition) and an additional outcome evaluating global change as observed by a clinician (with or without input from the caregiver). The outcomes selected to reflect these two domains vary, as do the number of attributes that are selected for evaluation.

The United States Food and Drug Administration (FDA) has established criteria for efficacy of dementia (specifically AD) drug interventions,²³ which require the following: 1) a double blind, placebo-controlled trial, 2) subjects who meet established criteria for AD, 3) sufficient length of follow-up to appreciate a meaningful effect of the drug on cognition, and 4) a clinical change of sufficient magnitude to be recognized by a clinician. In establishing these criteria, it was assumed that the outcome measuring cognition was the primary change of interest, and that the global clinical evaluation would mirror the changes in the primary variable.²⁴ In 1997, the European Medicine Evaluation Agency (EMA) issued new guidelines that incorporated two new concepts for the treatment of AD.²⁵ Firstly, the EMA guidelines suggested a preference for a measure of functional abilities in addition to a global measure, and noted that behavioral outcomes were important from a clinical perspective. Secondly, a definition of “responders” should be included in all trials, such that the degree of improvement in their cognition (or stabilization) was pre-specified. However, the magnitude of the change reflecting a clinically meaningful change was not specifically stated in either of these two guidelines. Sufficient magnitude of the change would reflect a clinically important difference, and this would vary with the type of outcome selected.

Several authors have attempted to define “clinically” relevant change. Gutzmann et al. (2002)²⁶ developed an Efficacy Index Score (EIS), which is a checklist that combines dropout as well as the relevant improvements individually across the three levels of assessment (cognitive function, activities of daily living and global function). Although, this summary score has not been validated relative to other traditional outcomes, it does present a unique example of determining efficacy in the context of anti-dementia drug interventions. Mayeux and Sano (1999)²⁷ in reviewing drug interventions for dementia, evaluated efficacy as a percent of the change in the treatment group relative to baseline (corrected for any change in the placebo group) and contrasted this with the percent of dropouts related to adverse events. Disease progression was considered with respect to the outcomes of 1) time until death, 2) nursing home placement, 3) loss of ability to perform Activities of Daily Living (ADL), or 4) severe dementia. In the context of clinical trials seeking to establish efficacy of pharmacological interventions, the latter outcomes may be problematic to ascertain.

Evaluation of the natural history of AD established some threshold values for expected decline or progression of the disease. Using the Alzheimer’s Disease Assessment Scale-Cognitive Section (ADAS-cog),²⁸ Rosen demonstrated that a decline of 1.28 points occurred within 12 weeks, a decline of 3.5 points within 6 months, and Stern et al. (1994)²⁹ showed a decline of 9 – 11 points by 1 year. Clinical experience would also suggest that the decline is not linear, with less deterioration in the early and later stages and the greatest acceleration in the middle severity category. The characteristics of the natural history of AD and other dementia types are best derived from longitudinal studies. Although, more details on the natural history of dementia are being reported, the fundamental difficulty still remains concerning the diversity of the outcome measures used to describe these changes. The picture of cognitive, behavioral, and functional decline will therefore vary with the outcome measure selected to describe it. Additionally, the diversity has a negative impact on comparisons of drug efficacy that can be made across trials.³⁰

Efficacy and Outcome Measures Used in Pharmacological Intervention Trials

No specific set of commonly accepted outcomes that define efficacy or “clinical relevance” applies to all the pharmacological interventions that have been used to treat dementia. More than 175 outcome measures are listed in Appendix E. EMEA guidelines acknowledge that no single test encompasses the broad range of disease characteristics associated with AD; nor has there been convincing evidence that an ideal (or reference) instrument exists to capture cognitive, behavioral, functional, or caregiver status.²⁵ The FDA has recommended that “dual efficacy” of dementia drug interventions be established by significant change in both a psychological measure and a global change measure. The outcomes used to measure these attributes within these two domains were not specified. In practice, there has been a general trend in North America toward using the outcomes ADAS-cog, the Mini-Mental State Examination (MMSE), and the Clinicians’ Interview-Based Impression of Change-Plus (CIBIC+) to capture the two domains when evaluating drugs for AD populations. However, these frequently used outcome measures may not be the best choice with respect to capturing “clinically relevant change”. The psychometric instrument properties must also be taken into consideration. For example, it has

been suggested that the ADAS-cog is weighted predominately to evaluate memory loss at the expense of other cognitive domains (especially executive control functions),³¹ which suggests that the face validity of this instrument may be in question. The generalizability of these results may be limited to dementia in which memory impairment is a key feature as the instrument is less sensitive to personality and executive dysfunction changes seen in a less typical dementia, such as frontotemporal dementia. The responsiveness (ability to detect change) of the CIBIC+ has not been well established.³² This suggests that some of the most established outcomes used to evaluate efficacy of pharmacological interventions are far from ideal.

Demers et al. (2000)³³ critically appraised some of the most commonly used scales evaluating global assessment,³² quality of life/ADL,³⁰ and behavior/mood³⁴ with respect to the quality of their psychometric properties. Several important limitations were identified in these reviews for the measures they evaluated, and these include 1) a lack of responsiveness data, 2) diversity in the content of the scales (capturing various aspects of a domain, for example, behavior), and 3) limited studies on reliability and validity (which are sample specific). The literature evaluating outcome measures used in dementia trials would suggest that most instruments have significant limitations, or at least more data are required to establish the required properties for acceptability of the scales.

Given the current state of development of research on outcome measures used in dementia trials for determining efficacy, a dilemma is clearly at hand. Ideally, all outcomes used to evaluate efficacy should have demonstrated acceptable psychometric properties, such as reliability, validity (construct), and responsiveness. However, since none of these outcomes have been accepted as standards, the selection of the most appropriate outcome is purely arbitrary. Similarly, establishing a rationale to exclude studies based on the specific type of outcome measure would be arbitrary. For this reason, no exclusion criteria based on outcome measures were used as eligibility criteria for this study.

Efficacy and Potential Risk of Adverse Events

Increasing attention has been given to the potential for harm, and not just benefits, when considering the efficacy of drug interventions. Empirical evidence across diverse medical fields indicates that reporting of safety information, including milder adverse events, receives much less attention than the positive efficacy outcomes.³⁵ Thus, an evaluation of the benefits of anti-dementia pharmacological agents alone may present a biased view of the overall benefit of the intervention. In the context of this systematic review, the type and frequency of adverse events associated with the use of a drug intervention will be scrutinized to a greater extent than previous reviews of anti-dementia drugs.

Capturing and evaluating adverse events is problematic. Typical randomized controlled trial (RCT) dose finding studies should consist of the comparison of several doses of a drug versus placebo; efficacy is demonstrated relative to a placebo group or relative to a different dose group. Ideally, the goal of early phase trials is to estimate the minimum effective dose or the maximum safe dose (or both). However, it is misleading to assume that drugs shown to be safe and effective in trials are safe and effective in all other circumstances.³⁶ The nature of pre-market clinical trials makes it difficult to evaluate the benefits of drugs for the universe of potential

users, as criteria restricting entry into the trial do not necessarily reflect dementia patients in general. By their nature, some adverse events are not easily anticipated, and therefore are not screened for in some trials. The implementation of pharmaco-vigilance systems attests to the need for further capture of potential adverse events not captured in trials. Adverse events may be hard to predict or anticipate and are captured only if a trial protocol was designed to measure these events. A limited number of standardized instruments exist to capture these events reliably. Unique to individuals with cognitive decline is the potential problem of validity of the self-report instrument, even if completed by the caregiver. Furthermore, many trials may be underpowered to detect adverse events with an incidence of 1/1000.³⁷ Despite these limitations, quality criteria for the collection and reporting of adverse events have been identified.^{35,37} An instrument to evaluate the quality of reporting adverse events has been developed and used in this report to determine the strength of the evidence for adverse events in the context of determining efficacy.

Efficacy and Intention to Treat Analysis

Determining efficacy in dementia trials evaluating pharmacological interventions may vary depending on the selection of the analysis type. In general, the types of analyses of primary data in trials fall into two main categories: 1) intention to treat analyses (ITT) or last observation carried forward (LOCF), and 2) observed case (OC) or completed trial (CT). The advantages of ITT over OC analyses have been well explicated.³⁸ It is recognized that non-compliance is not a random event; thus, ITT analyses should be used to base principal conclusions of efficacy.³⁹ In the context of some anti-dementia drug therapies, where dropout rates due to adverse events and other non-compliance reasons may be high, the ITT analysis minimizes bias and the potential for type I errors when considering treatment efficacy. However, the ITT analysis, while less biased, does tend to reduce treatment effects to the extent that there are dropouts and crossover patients. The optimal analysis, when there is a large loss to follow-up, is to conduct the analysis both ways and look for consistency.

Primary Objectives and Scope of Systematic Review

A large number of pharmacological interventions have been studied in dementia patients. These agents can be classified into three broad categories: 1) cholinergic neurotransmitter modifying agents, such as acetylcholinesterase inhibitors, 2) non-cholinergic neurotransmitter/neuropeptide modifying agents, and 3) other pharmacological agents. Although only four agents have been approved by the FDA for the treatment of dementia, many other pharmacological agents are being evaluated in trials in off-label use. In both these circumstances, there was a need to determine the evidence to support claims of efficacy and to describe adverse events.

The Questions

Given the range of pharmacological agents that have been used to treat dementia, evaluation of all of these interventions in a systematic review (which afforded a consistent methodology) should serve as a meaningful contribution in this area. The purpose of this systematic review is to answer the following questions:

- 1) *Does pharmacotherapy for dementia syndromes improve cognitive symptoms and outcomes?*
- 2) *Does pharmacotherapy delay cognitive deterioration or delay disease onset of dementia syndromes?*
- 3) *Are certain drugs, including alternative medicines (non-pharmaceutical), more effective than others?*
- 4) *Do certain patient populations benefit more from pharmacotherapy than others?*
- 5) *What is the evidence base for the treatment of VaD?*

This review considers different dementia populations (not just AD) and subjects from both community and institutional settings. The interventions were limited to pharmacological agents (including nutraceuticals), and these were not restricted to those that have received official approval in North America. The studies eligible in this systematic review were restricted to parallel RCTs, but the study outcomes were not limited to specific types.

The review will serve to evaluate the quality of the evidence and identify important gaps in the literature. Future recommendations will serve the dementia research community specifically. This evidence report will support the American College of Physicians (ACP) and the American Academy of Family Physicians (AAFP) in developing “best practices” and practice guidelines for the evidence-based treatment of dementia for providers, patients and the public.

Chapter 2. Methods

The Research Team

A multidisciplinary local research team representing geriatric and dementia epidemiology/ systematic review methods (P. Raina, PhD), pharmaco-epidemiology (M. Levine, MD, PhD), geriatric medicine/ dementia (D. Cowan, MD; C. Patterson, MD), rehabilitation/ systematic review methods (P. Santaguida, PT, PhD), and neuropsychology (A. Unsal, PhD) was assembled. The core research team, including experienced staff at the McMaster Evidence-based Practice Center (EPC) (F. Baldassarre, MSc; L. Booker, BA; M. Gault, BA) participated in regular meetings and reached consensus on key methodological issues. An international Technical Expert Panel (TEP) was assembled to provide high-level content expertise in dementia and participated in conference calls on an as-needed basis. Participants in this panel were: Larry W. Chambers, PhD. Ottawa, ON, Canada; Thomas Cook, MD. (ACP appointee) Colorado Springs, CO, USA; Rachelle Doody, MD, PhD. Houston, TX, USA; John Feightner, MSc, MD. London, ON, Canada; Rodney Hornbake, MD. (ACP appointee) Hadlyme, CT, USA; David Hogan, MD. Calgary, AB, Canada; Roy Jones, MD. Bath, UK; and Holly Tuokko, PhD. Victoria, BC, Canada.

Topic Assessment and Refinement

Refinement of Questions

The first step during the topic assessment and refinement process was to organize a teleconference with the partner organization, the Task Order Officer (TOO), invited topic experts, and the McMaster team in order to define the magnitude of the topic addressed and to refine/clarify the preliminary research questions for this evidence report. It was agreed that this evidence report would focus on addressing the efficacy of pharmacotherapies for dementia syndromes. Regular teleconferences were held with the TOO, the partner, and technical experts throughout the data refinement and extraction phase.

Search Strategy

Search strategies were developed and undertaken in the electronic databases listed in Table 1 for the time periods specified. The order of the databases in Table 1 also represents the sequence that the databases were searched. Appendix A details the search terms for all databases.

Table 1. Databases searched for relevant RCTs.

Database searched	Search date	Period searched
Cochrane Central	February 3, 2003	1st Quarter 2003
MEDLINE® & PreMedline®	February 4, 2003	1998 to 2003 week 4
EMBASE	February 6, 2003	1998 to 2003 week 5
AMED	March 4, 2003	1985 to 2003 February
CINAHL®	March 5, 2003	1982 to February 2003 week 3
Ageline	March 6, 2003	1978 to 2002 December
PsycINFO	March 7, 2003	1967 to 2002 December

Expert opinion was sought on the most efficient search strategies to minimize noise in the collection of citations. Some of the medical subject headings (MeSH) used to select RCTs yielded a large number of non-RCT literature due to misclassification of the study design terms. For example, in previous indexing, terms like “longitudinal study” or “comparative study” were applied to RCTs; conversely, the MeSH terms “random” or “randomized” in the title or abstract were not consistently used. However, some recent methodological work has suggested that more specific search term approaches can be used, which increases the sensitivity and specificity of the search results.⁴⁰ The Cochrane Central Trial Registry contains correctly re-classified RCT/Controlled Clinical Trial (CCT) trials that were misclassified in MEDLINE® and EMBASE from 1966 to 1998. All published RCTs to 1998 are contained within this database. Hence we commenced our search with the Cochrane Central Trial Registry database. For this reason, MEDLINE® and EMBASE were searched from 1998 forward for relevant studies, and all the other databases from their inception.

Specific drug names and manufacturer brands were considered as potential search terms. However, the local research team was in agreement that listing specific drug names would bias the yield to include only those pharmacological agents searched and would not capture newer drug therapies. Thus the recommendation was to not restrict the search to known pharmacological agents but to include whatever agents were in the literature.

In addition to the electronic databases, the bibliographies of retrieved papers were retrieved. Any citations recommended by the local research team, the TEP, or the peer reviewers were retrieved and screened.

Eligibility Criteria

Inclusion. Studies were included that contained the following criteria:

- 1) Age: Studies involving dementia patients who were 18 years or older in age
- 2) Diagnostic model used to determine dementia: The diagnosis of dementia using any of these criteria:
 - i) ICD 9 or 10.^{41,42}
 - ii) DSM III, III-R, and IV.^{43,44,45}
 - iii) NINCDS.⁹
 - iv) NINCDS-ADRDA⁹ or NINCDS-AIREN.⁴⁶

- 3) Diagnostic criteria used to determine cognitive impairment (pre-dementia): In the case of not yet diagnosed dementia, specific diagnostic categories were accepted for the following:
 - i) mild cognitive impairment (MCI)..⁴⁷
 - ii) cognitive impairment not dementia (CIND).⁴⁸
 - iii) cognitive loss no dementia (CLoND).⁴⁹
- 4) Disease classifications for dementia: These included AD, senile dementia of the Alzheimer's type (SDAT), Lewy body disease, VaD, multi-infarct dementia (MID), AIDS/HIV dementia, Parkinson's disease dementia (PDD), progressive supranuclear palsy (PSP), mixed diagnosis dementia, encephalopathy, Mesulam syndrome, progressive non-fluent aphasia, Binswanger disease, subcortical leukoencephalopathy, circumscribed lobar brain atrophy, Pick disease, amyloid beta-protein (not Down's syndrome or trisomy), cerebral amyloid angiopathy, neurofibrillary tangles, threads, senile plaques, corticobasal ganglionic degeneration, cerebral autosomal dominant ischemia with subcortical leukoencephalopathy (CADASIL), Huntington's disease with dementia, hydrocephalus (for additional terms used in the search strategy, see Appendix A).
- 5) Severity classification: This was accepted in whichever classification system the studies specified. The majority of studies specified threshold criteria using the MMSE as follows: mild > 22, moderate 14 – 21, and 10 – 14 as severe. Many studies used the definition of mild to moderate as a range from 10 to 26 based on criteria established by Folstein et al.⁵⁰ Some studies specified a category (i.e. mild to moderate) but did not report the baseline MMSE values for the groups compared.

Some studies specified two categories (mild to moderate) and (moderate to severe) based on the DSM-III-R criteria. Cambridge Examination for Mental Disorders in the Elderly (CAMDEX) specifies levels of severity (minimal, mild, moderate, severe). Similarly, some studies reported a category of severity without stating which method was used. In these instances, the category of severity specified was accepted as reported by the study authors.

Exclusion. Studies that had populations with any of the characteristics listed below were excluded.

- 1) Dementia disease classification: i) alcohol caused dementia/ Korsakoff's syndrome, ii) Creutzfeldt-Jakob syndrome, c) spongiform encephalopathy, iii) hypothyroidism, iv) vitamin B12 deficiency, v) neurosyphilis.
- 2) Dementia diagnosed using only Lowb, Hachinski (specific for VaD) criteria.⁵¹
- 3) All organically caused dementias which includes "Delirium, Dementia, Amnesic Disorders, and Cognitive Disorder Otherwise Specified. The predominant disturbance is a clinically significant deficit in cognition that represents a significant change from a previous level of functioning. For each disorder in this section, the etiology is either a general medical condition (although the specific general medical condition may not be identifiable) or a substance (i.e., a drug of abuse, medication, or toxin), or a combination of these factors."⁴³
- 4) Temporary dementia (e.g. side effect of anesthesia) classified as follows: Delirium: a delirium is characterized by a disturbance of consciousness and a change in cognition

- that develop over a short period of time. The disorders included in the "Delirium" section are listed according to presumed etiology: delirium due to a general medical condition, substance-induced delirium (i.e. due to a drug of abuse, a medication, or toxin exposure), delirium due to multiple etiologies, or delirium not otherwise specified (if the etiology is indeterminate).
- 5) Normal or healthy volunteers: studies that deal with healthy people (i.e. prevention is limited to people who have any form of the above); volunteer study population
 - 6) General population of elderly persons.
 - 7) Study subjects selected for depression (some patients may have dementia but not all) and where there is no stratified analysis by disease subgroup (i.e. the dementia subjects).

Study Design. Eligible studies included parallel design RCTs only. Although crossover trials are suitable for chronic diseases, they may be prone to period effects or period by treatment interactions. Period effects are systematic changes in the outcome that apply to all patients due to temporal changes in the disease or to the measurement instrument. Period by treatment interactions occur when the efficacy of the intervention varies by period. Additionally, a carry-over effect may occur if there is not an adequate washout period. Apart from the weaknesses of this design, some limitations arise when considering the potential for meta-analytic analyses. Traditionally, first period data from a crossover trial are abstracted and can be potentially combined with parallel trials for analyses of a pooled estimate; the reporting of the study results (positive or negative) would also be based on this first period data. In a preliminary phase of the review, several crossover trials were examined. Many did not report first period data, which precluded any potential for combining with parallel trials; many trials also did not undertake statistical tests during the first experience, thus making it difficult to report the direction and significance of the first period findings. Finally, because this systematic review was considering a variety of drug interventions administered over differing time intervals, period effects might be an important source of bias. For all these reasons, the decision was made to exclude crossover trials from this systematic review.

Language of Publication. Studies published in the English language were eligible. The scope and resources of this review did not permit translation of studies published in other languages.

Sample Size. No sample size restrictions were applied.

Treatment Interventions. Drug interventions were eligible in the following manner:

- 1) Pharmacological agents: all types of pharmacological treatment were considered in this review, including food supplements (as defined by the FDA). Government approval was not a requirement, and as such, off-label use of drugs (i.e. drugs approved for other conditions but used in the treatment of dementia) were eligible for this review.
- 2) Dose: all doses and dosing schedules and any mode of administration (oral, subdermal, transdermal, intravenous, suppository, or intra-muscular injection) were considered.
- 3) Treatment period: the period of treatment must equal or exceed 1 day.

- 4) Follow-up length: Any duration of follow-up was eligible. Different drugs require different time periods to show an effect. For example, antidepressant and anti-psychotic medications may take a month or more to be effective. Some dementia drugs take a minimum of 2 months. For interventions such as vitamin E or Ginkgo biloba, the time to effect is not well established. Thus, an absolute limit to the minimum number of months of follow-up could not be applied to all potential interventions. It was anticipated that many studies with some of the most recent pharmacological agents (i.e. donepezil) would have a minimum follow-up of 24 weeks.

Study Outcomes. No specific set of commonly accepted outcomes that define efficacy or “clinical relevance” were applicable to all the pharmacological interventions that have been used to treat dementia. The literature evaluating outcome measures in dementia trials would suggest that most instruments have significant limitations or at least more data are required to establish the required properties for acceptability of the scales. Since none of the outcomes used in dementia trials have been accepted as standards (no consensus), the selection of the most appropriate or clinically relevant outcome is purely arbitrary. Similarly, establishing a rationale to exclude studies based on the specific type of outcome measure would be arbitrary. For this reason, no exclusion criteria based on outcome measures were used as eligibility criteria for this study; rather the domains of interest for inclusion have been identified.

Studies with the following outcomes were included:

- General cognitive function (e.g., ADAS-cog).
- Specific cognitive function (e.g., Weschler Memory Tests).
- Global clinical assessment (e.g., CIBIC).
- Behavior/mood (disturbances characterized by agitation, wandering, sleep cycle disturbance, depression, obsessive compulsive activities) (e.g., Behavioral Pathology in Alzheimer’s Disease Rating Scale (BEHAVE_AD)).
- Quality of life/ADL (e.g., Instrumental Activities of Daily Living (IADL)).
- Effects on primary caregiver (also referred to as caregiver burden).
- Safety as measured by the incidence of adverse effects (e.g., particularly serious adverse events).
- Acceptability of treatment as measured by withdrawal rate from trial due to side effects of the medication. (e.g., dropouts due to adverse events).
- Mortality.
- Dependency or Rate of Institutionalization/ or continued residence in own home.
- Use of services.

Studies with the following outcomes were excluded as follows:

- Studies which reported only biological/ physiological outcomes, such as plasma levels, changes on functional imaging, or electroencephalography (EEG) activity, were noted but not assessed as efficacy measures.

- Outcomes reported in the trials should reflect changes in the person with dementia. If the study population did not all have dementia, only data subgrouped for dementia was examined.

Minimum quality threshold score for eligibility.

Exclusion part I: Pre-Jadad score. Studies were also screened to determine a minimum threshold for quality, sometimes described as “fatal flaws” in the trial design. Specifically, all studies had to include at least some mention of the term “randomization” or “withdrawal(s)” in the text of the paper. Trials that did not at least mention these components were excluded, as they possessed a fatal flaw.

Exclusion part II: Post-Jadad score. The methodological quality of the primary studies was assessed using the modified Jadad scale for RCTs⁵² (Appendix B). The reliability of this modified scale was shown to be high, as measured by the intraclass correlation coefficient (ICC = 0.90).⁵² Each study was evaluated by two reviewers, and the level of agreement was determined statistically. The first three items on the scale rate elements that have been shown to bias meta-analytic results. These include randomization, blinding, and withdrawal. If these items alone are considered, the maximum score is 5. Any study that did not score 3 or more on the scale was excluded from the review. Therefore, this review abstracts detailed data only from studies that achieved moderate to high ratings on the quality scale.

Evaluating the methodological quality of studies and rating the strength of the evidence.

Quality of the RCT. The methodological quality of the primary studies was assessed using the modified Jadad scale for RCTs.⁵²

Quality of reporting adverse events. The potential for risk, or adverse events, was an important component to consider with respect to efficacy. The Jadad scale for quality does not take into account factors associated with adequate collection and reporting of adverse events as detailed by Ioannidis and Lau (2002).³⁵ Therefore, a summary checklist was developed to determine the potential quality in the collection and reporting of adverse events (Appendix B). This score was used to evaluate the relative quality of the adverse events reported.

Data Collection and Reliability of Study Selection

During the identification phase, two independent reviewers evaluated the title and abstract for eligibility; those meeting the criteria were retrieved as well as those that reported insufficient information to determine eligibility. Two independent reviewers examined the full text of these articles (passing from the title and abstract phase). All studies meeting eligibility criteria were reviewed to assess quality and abstracted according to predetermined criteria. The articles were grouped according to the pharmacological agent used in the intervention.

A team of study assistants was trained in the eligibility criteria for the purposes of this systematic review. Standardized forms and a guide explaining the criteria were developed from previous templates (Appendix B). Two reviewers were used for the identification, selection,

validity, and abstraction phases of the systematic review. Disagreements were resolved by consensus. The reviewers were experienced EPC staff with post-graduate training in research methods. The reviewers and abstractors would consult with more senior members of the TEP for content expertise or methods-related issues.

Summarizing Results: Descriptive and Analytic Approaches

It was expected that studies of the pharmacological agents used in the management of dementia would be quite diverse with respect to the intended therapeutic effect. For these studies, evidence and summary tables (Appendix C) were constructed to describe the more salient characteristics of the included studies.

Meta-analysis

Statistical meta-analysis was not appropriate for all outcomes or interventions. Before calculating a pooled effect measure, the reasonableness of pooling was assessed on clinical and biological grounds, in terms of clinical homogeneity. Tabular summaries of key characteristics, participants, interventions, and outcomes were considered. A priori, it was decided that pooled estimates would be undertaken for studies with the same pharmacological intervention and the same outcome measure and that a minimum of three studies was necessary for pooling for a specific outcome*. Consideration was given to the similarity of study populations when selecting studies to be included in the pooled estimates. Although many studies evaluated multiple outcomes, data necessary for meta-analysis were not provided in all eligible trials. When sufficient data were provided to estimate the weighted mean difference (WMD), then a meta-analysis was undertaken. WMD was selected as the pooled estimate (versus the standardized mean difference) because the outcome measures did not differ between studies eligible for pooled estimates. For WMD, the difference between the treated and control groups are weighted by the inverse of the variance.

Analysis was undertaken in RevMan 4.2 (Review Manager, Cochrane Collaboration, 2003), and the random-effects model was used to conduct our analyses. In cases where heterogeneity existed, the results of the random-effects model only were considered for interpretation of the results of the pooled estimate. RevMan 4.2 automatically tests the homogeneity of the results of the individual studies for each comparison of dichotomous or continuous data. Tests of homogeneity are formal statistical analyses for examining whether the observed variation in study results is compatible with the variation expected by chance alone. The more significant the results of the test (the smaller the p-value), the more likely that the observed differences were due to unknown factors likely not controlled for in the study. Sensitivity analysis or meta-regression was not undertaken to assess the extent to which the methodological quality of studies, population characteristics, dose, etc., accounted for variation in the primary outcome.

* A priori it was decided that a minimum of three studies would be required for undertaking pooled estimates. It was assumed that if two studies were meta-analyzed, theoretically the estimates could be in opposite directions leading to un-interpretable estimates. In this same situation, a third study would allow for interpretation of the direction of the effect.

Power Analyses

Power analyses were conducted for select pharmacological interventions reporting non-significant findings for all primary outcomes reported in the paper. In addition, if the trial reported the outcomes of MMSE, ADAS-cog, or the CIBIC+, the power for these was also estimated. It was assumed that the desired level of significance was set to alpha equal to 0.05. Adequate power was defined as at least 80% power.

Peer Review Process

A list of potential peer reviewers was created at the outset of the study. During the course of the project, additional names were added to this list by the McMaster Center and Agency for Healthcare Research and Quality (AHRQ). In May 2003, the individuals on the list were approached by the McMaster team and asked if they would act as peer reviewers of this evidence report. A total of 26 experts agreed and received a copy of the draft report and a copy of the “Structured Format for Referee’s Comments” (Appendix D). A list of the reviewers’ names and their affiliation is provided in Appendix D. In addition, a criticism editor, Dr. Patricia Huston, who is external to the McMaster EPC, was asked to review the draft report and synthesize the peer review comments. The report from the criticism editor was then used to prioritize the incorporation of peer review comments into the final version of this evidence report.

Chapter 3. Results

In this chapter, the presentations of the main results of the systematic review are organized according to the five questions that were addressed. The first question, concerning efficacy of the pharmacological interventions, contains results from all eligible studies. Subsets of trials were then selected from this larger set to address the remaining four questions (see Chapter 2 Methods).

Eligible Studies

Figure 2 shows the final yield of eligible studies for evaluation, and the inclusion/exclusion criteria are listed in Chapter 2. Approximately 10.5% of identified studies met the eligibility criteria in the title and abstract phase. Similarly, 14.7% of the full text screened citations were eligible for full data abstraction. Several trials were identified as “companion papers”, indicating that results for these related studies were based on the same study subjects. These related studies were evaluated and a main publication was selected (usually the first chronological publication), and the remaining trials were searched for any additional data for abstraction; the “companion papers” were not considered as unique studies. English-language reports only were included in this review.⁵³ Although this is acknowledged as a possible source of bias, the overall proportion of potentially eligible non-English studies for review in title and abstract was small (7%).

Figure 3 indicates the distribution of eligible studies as a function of publication year grouped into approximately 5-year intervals. The largest proportion of studies (83%) was published within the last 11 years, with the greatest number from 1997 forward. This may have some implications for future systematic reviews with respect to the years searched.

This systematic review yielded a total of 97 pharmacological agents used in the treatment of dementia from 186 unique studies. These 97 interventions have been classified according to three broad categories: 1) cholinergic neurotransmitter modifying agents, 2) non-cholinergic neurotransmitter/ neuropeptide modifying agents, and 3) other agents.

Question 1: Does pharmacotherapy for dementia syndromes improve cognitive symptoms and outcomes?

The largest number of eligible citations evaluated was cholinergic neurotransmitter modifying agents (n = 72). The remaining citations were distributed amongst the non-cholinergic neurotransmitter/neuropeptide modifying agents (n = 61) and other agents (n = 76) categories. Some studies evaluated agents in more than one category. The results for all pharmacological agents are presented in this chapter in summary format with descriptive text and an overall summary table (OST) for each drug located at the end of this chapter. The specific details abstracted from each individual study are presented within Appendix C (guide to the results tables) and organized into these same three therapeutic effect classification groups and by pharmacological agent. These Evidence Tables are available on-line at <http://www.ahrq.gov/clinic/epcindex.htm>.

Figure 2. Flow diagram showing the final number of studies meeting the eligibility criteria.





Figure 3. Proportion of studies as a function of year of publication.



Appendix C contains three sets of tables with key study descriptors as follows:

Key characteristics. Summarizes the following aspects of each study: features (author, year published, funding source, modified Jadad scale quality score, number randomized, number completing the trial, subgroup analysis), population characteristics (diagnosis, criteria for diagnosis, disease severity, percent male, age, dwelling, and differentiating demographics), intervention (doses, titration scheme, and intervention period), and a complete list of outcomes administered in the study protocol.

Study results. Details the changes observed (the magnitude of these changes, the comparison groups analyzed, and the findings of any statistical testing) for those outcomes for which appropriate data was reported (for up to three time periods if available). When reported in studies, baseline measures, particularly MMSE score, were also detailed in these tables.

Study adverse events. Lists the specific types of adverse events (side effects, adverse reactions, and serious events) reported, any statistically significant differences between groups, the proportion of withdrawals due to adverse events, and the quality rating score (based on a checklist devised at the McMaster EPC and on the work of Ioannidis and Lau (2002)³⁵) specific to the collection and reporting of these adverse events.

Interpretation of the Results in the Overall Summary Tables (OST) for Individual Studies

To facilitate the presentation of information within the OST (found at the end of this chapter), the outcomes reported in eligible studies were classified into seven domains: 1) general cognition scales, 2) specific cognition tests (neuropsychological tests evaluating specific attributes of cognition, such as short- and long-term memory, word fluency, etc.), 3) global assessment, 4) behavior/mood, 5) quality of life/ADL, 6) caregiver burden, and 7) other. The EPC research team reached consensus on the classification of the various outcome measures within these seven domains. For example, the ADAS-cog and MMSE were classified as “general cognition scales”, and the BEHAVE-AD and NOSGER were placed in the “behavior and mood” domain (see Appendix C guide to the results tables). The complete list of outcomes that were reported in the studies evaluated in this review and the domains that they were classified within is found in Appendix E. Table 9 in the report presents a guide to the overall summary tables by domain.

For each of the outcomes reported by a study, four interpretations of the results were possible. The four options for interpretation are as follows:

- SC* = *significant change*. Demonstrated by statistical significance ($p \leq 0.05$) for the primary outcomes from an ITT analysis comparing treatment and placebo groups, or comparing differences among dose groups.
- NS* = *not significant*. The corollary of SC indicating no statistical significance.
- MX* = *mixed results*. Primary outcomes within the same domain show opposite or inconclusive statistical significance; for example, in the general cognition domain, half the studies show significant change and the other half show no significance).
- NR* = *not reported*. Outcome was collected but not statistically evaluated or not reported in the publication.
- NT* = *not tested*. No outcomes in this domain were tested.

Secondary outcome results were reported in the absence of any primary outcome data (for the domain of interest) and were demarcated with a (2°) in the OST. Similarly, analyses other than ITT were denoted with an asterisk (*) in the OST. If the report describes my subgroup analyses, the word SUBGROUP appears in the “other” column.

Adverse events were not always clearly described in many studies. A priori, we selected 5 generic symptoms (nausea, dizziness, agitation, eating disorder, and diarrhea) and selected to detail the ranges amongst studies for both placebo and treatment groups for these symptoms. The percent of withdrawals for both groups due to adverse events was reported. Adverse events reported to be statistically significant are highlighted for the reader. The details in addition to the quality score rating will assist the reader in evaluating the potential for harm.

Statistical Analysis

Power Analyses and Measures of Effect for combined studies. Power analyses (PW) were for individual trials for select pharmacological interventions (donepezil, galantamine, tacrine, rivastigmine, memantine, estrogen, carnitine, ginkgo biloba, selegiline) for all primary outcomes. In addition, if the trial reported the outcomes of the MMSE, ADAS-cog, or CIBIC+, power was also estimated (for individual trials of pharmacological interventions that had a minimum of three trials with a common outcome). Quantitative meta-analyses were undertaken in interventions that had a minimum of three trials using the same outcome scales and which provided sufficient data to permit calculation of effect sizes (as an Odds Ratio (OR), Relative Risk (RR) and Weighted mean difference (WMD)). The random-effects model results are presented to the reader.

Quantitative and Descriptive Analyses

Results of cholinergic neurotransmitter modifying agents (CNMA)

A total of 70 studies evaluating 16 cholinergic neurotransmitter modifying agents were eligible for review (Table 2). Six studies directly compared different drugs, and these trials are considered separately in the section that addresses question three. Overall results for each of the trials for each of the interventions are detailed in the OST located at the end of this chapter and organized by drug. All other study details are available in Evidence Tables 1 through 93 in Appendix C.

Table 2. List of Cholinergic neurotransmitter modifying agents and the number of studies vs. placebo for each of these. Asterisk (*) indicates report of a drug vs. drug trial [comparator drug(s) in brackets].

Drug	Number of studies vs. placebo	Drug	Number of studies vs. placebo
Antagonic Stress * [Meclofenoxate] *[Nicergoline]	0**	Metrifonate	9
Carnitine	6	Nicergoline*[Antagonic Stress]	4*
Donepezil *[Vitamin E]	10*	Physostigmine	4
Eptastigmine	2	Posatirelin *[Citicoline]	4*
Galantamine	6	Rivastigmine	6
Huperzine-A *[Tablet Capsule]	1*	Sabeluzole	1
Linopirdine	2	Tacrine *[Idebenone] *[Silymarin]	6**
Mexofenoxate *[Antagonic stress]	0*	Velnacrine	3

Carnitine (also known as acetyl-L-carnitine, gamma-trimethyl- β -acetylbutyrobetaine (Alcar). See Evidence Tables 1 through 8 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. A total of six studies^{54,55,56,57,58,59} evaluating carnitine were included in this review. Four of the reports were published from 1990 to 1992^{56,57,58,59} while the remaining two, both by same authors, were published in 1996 and 2000.^{55,54}

Design/methodology. A total of 925 subjects were evaluated in these six studies comparing carnitine and placebo. The range of study sample sizes was from 30 to 431 subjects. Quality scores (out of 8 points) ranged from moderate⁵⁷ (5) to high^{58,59} (7), and all of the studies were partially or totally funded by industry.

Populations. All trials were conducted on AD patients, and all but one study used the NINCDS criteria for diagnosis. None of the trials reported including patients with severe dementia; all were classified as mild to moderate.

One trial⁵⁸ had a mix of community and institutional patients, and one study reported using a community sample.⁵⁶ The mean age of the samples ranged from 59⁵⁴ to 79 years,⁵⁹ with the majority reporting mean age greater than 70 years. One study⁵⁶ did not report mean age. Three studies specified the baseline MMSE^{54,55,57} (range 16.1 to 20.6) and one trial specified the modified MMSE⁵⁶ (mean 35) demonstrating no differences between placebo and treatment groups.


Intervention. The dose varied from 2 to 3 grams per day, and treatment duration was either 24 weeks^{56,57,59} or 52 weeks.^{54,55,58} One study⁵⁷ did not report the dose used. No titration period was used for this drug in any of the studies.

Primary outcomes. All of the studies measured cognition as a main outcome; half of the trials also measured outcomes in the behavior/mood domain.^{58,54,55} All but one of the studies⁵⁸ used a quality of life/ADL or functional status measure. Only one study⁵⁵ evaluated caregiver burden. Of the four studies^{59,58,54,55} reporting the findings from a global measure, only one⁵⁵ used the Caregiver-rated Global Impression of Change (CGIC).

Analysis. Half of the studies reported ITT analyses^{58,54,55} and the remaining trials reported OC results.^{57,59,56} The ability to combine results was limited with only three studies having the common outcomes of ADAS-cog, modified MMSE (mMMSE), and Clinical Dementia Rating (CDR).

Results and interpretation. See Summary Table 1. The four studies that evaluated “general cognitive function” did not find statistically significant differences in this domain. For those trials that provided sufficient data to estimate power, three trials^{54,55,56} were underpowered for the MMSE (PW = 0.15 to 0.19), and two trials^{54,55} for the ADAS-cog (PW = 0.08 to 0.09) and the CDR (PW = 0.06 to 0.11). Meta-analysis was undertaken for the MMSE scores; although favoring a treatment effect, the pooled effect size (WMD = 0.55) was modest and zero was contained within the confidence interval (Figure 4) for the random effects models. The pooled estimate favoring treatment may suggest some potential for benefit in general cognitive function, but this must be verified in future research.

Figure 4. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the MMSE comparing carnitine versus placebo.



For “specific cognitive tests”, two out of four studies did not detect statistical differences relative to placebo, and the remaining two showed mixed results (see Summary Table 1). No significant differences were found in the domains of global assessment, behavior/mood, and quality of life/ADL; power could not be evaluated for the majority of these outcomes in the trials (insufficient data reported to permit calculation).

Four^{57,59,56,58} of the six trials scored 3 out of 5 on our quality scale for rating adverse events. Two trials^{55,54} did not adequately report adverse events (score = 1), but tested for statistical differences between groups. Withdrawal rates due to adverse events ranged from 0 – 3% in all studies, with the exception of a single trial⁵⁹ where the percentage was 22% (placebo) and 44% (treatment). The high rates in this trial are likely related to the small sample size ($n = 36$). This same trial⁵⁹ was also the sole study reporting dizziness and anxiety (confusion, depression). In general, gastrointestinal symptoms (Evidence Table 8) were the most frequently reported adverse events, but most studies did not test for statistical differences in the rates between the groups. The percent of subjects reporting of a priori selected symptoms across all studies are as follows: 1) nausea (placebo = 6 - 14%, all doses carnitine = 28%), 2) dizziness (not reported as an event for either placebo or treatment group), 3) agitation (placebo = 6%, all doses carnitine = 7%), 4) diarrhea (not reported as an event for either placebo or treatment group), and 4) eating disorder (not reported as an event for either placebo or treatment group).

Donepezil. See Evidence Tables 9 through 21 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. A total of 11 studies^{60,61,62,63,64,65,66,67,68,69,70} evaluating donepezil were eligible for this systematic review. One study⁷⁰ compared donepezil to vitamin E rather than placebo. All were published within the last 6 years ($n = 1$, 1996), ($n = 2$, 1998), ($n = 1$, 1999), ($n = 5$, 2001), ($n = 2$, 2002). Three of these studies^{66,67,69} were undertaken by the same research group at different time periods and had unrelated cohorts of patients.

Design/methodology. A total of 3239 subjects (range of study sample size, 30 – 893) were included in these trials. The modified Jadad scale quality scores ranged from 5⁶¹ to 8.^{63,64} All studies were funded by industry sponsors with the exception of a single trial⁷⁰ that did not specify their source of support.

Populations. All but one study⁶⁰ used the NINCDS criteria to diagnose dementia. Eight studies included only AD patients, one study included only VaD,⁶⁸ and the remaining two included Parkinson’s disease dementia (PDD⁶⁵) and AD patients with cardiovascular disease.⁶⁴ A single

trial included subjects with Down's syndrome and AD.⁶⁰ The severity of the dementia patients was described as mild to moderate in five studies,^{65,66,60,62,70} mild to moderately severe in two studies,^{69,67} probable in two studies,^{61,68} and moderate to severe in two studies.^{63,64}

Some studies specified that the dementia patients were recruited from the community,^{68,63,60} one study from institutional setting,⁶⁴ and the remaining did not specify the living arrangements. Mean ages of the study subjects ranged from 54 to 85.7 years with most studies representing ages in the upper to mid 70s.

Six studies^{61,62,65,66,67,69} specified the race of the subjects, and of these, the overwhelming sample was Caucasian (range from 92 - 100%). All but one study⁶⁸ specified the proportion of men recruited, the range being from 18 - 46%. Four of these studies presented some results stratified by gender,⁶² age,⁶⁴ APOE genotype,⁶² baseline MMSE,^{63,64} patients with Down's syndrome,⁶⁰ and the use of psychoactive drugs.⁶³ Three studies specified the baseline MMSE^{61,64,70} demonstrating no differences between placebo and treatment group, and the mean values varied from 14 to 16.

Intervention. Five studies evaluated a 10 mg dose given once daily,^{60,61,62,63,66} two studies 5 mg daily,^{64,69} and four studies compared 5 mg and 10 mg dose groups.^{65,68,67,70} Titration periods observed included 7 days^{65,67} and 4 weeks,^{70,68} these were not specified in the remaining studies. The total duration of the drug (including titration) varied from 12,⁶⁵ 15,⁶⁷ 23/24,^{60,63,64,65,66,68} and 54/56^{61,62} weeks.

Primary outcomes. Specific cognitive tests and caregiver burden were not evaluated in these studies. Nine studies used the MMSE, and six studies the ADAS-cog.

Analysis. All but one of the studies⁶⁰ comparing donepezil to placebo used ITT analysis. The study using OC analysis showed no statistical difference between treatments. It had the smallest sample size of 30 subjects and was underpowered (PW = 0.16) for the behavioral measure used, NPI.

Results and interpretation. See Summary Table 2. For the 10 trials^{60,61,62,63,64,65,66,67,69,68} comparing donepezil to placebo, two studies^{60,64} did not show a positive effect for the domain of general cognition. However, both of these studies evaluated the outcomes in this domain as secondary outcomes, with one trial⁶⁴ lacking sufficient power for the MMSE (PW = 0.69); for the other trial, power could not be evaluated. For the eight trials^{61,62,63,64,66,69,67,68} showing a positive effect on general cognition, all but one trial used the MMSE as an outcome, which allowed for a pooled effect size estimate (Figure 5); we assumed that the VaD patients in one trial⁶⁵ could be combined with the other dementia populations. Figure 5 shows a consistent treatment effect for improvement in general cognitive function as measured by the MMSE, and the overall effect was statistically significant. Figure 6 shows the four trials^{65,66,67,68} that used the ADAS-cog to measure general cognitive function change. A consistent effect favoring treatment was evident, and the test for overall effect was statistically significant. It should be noted that some of the values used in the pooled estimates for the MMSE and the ADAS-cog were derived from figures showing means and confidence intervals in the trial reports, thus introducing some imprecision into these estimates.

Figure 5. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the MMSE comparing donepezil and placebo.





Figure 6. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing donepezil versus placebo



Ten studies^{60,62,63,64,70,65,66,67,69,68} evaluated global assessment, and with the exception of three trials,^{60,69,70} all studies showed a statistically significant difference in this domain. The overall effect for the CIBIC (Figure 7), and the CIBIC + (Figure 8, expressed as a proportion of improved versus not improved) were estimated for the 5 mg dose of donepezil. Figure 9 shows the summary estimate for the three studies^{65,66,68} that evaluated the 10 mg dose of donepezil; heterogeneity was significant (p = 0.007) in this meta-analysis, but the overall effect was significant (p = 0.002). Similarly, Figure 10 shows the summary estimate for the Clinical Dementia Rating (CDR) global assessment measure. A radial plot of these three studies was undertaken and suggests that one trial⁶⁴ could be an important source of the heterogeneity. Summary estimate was also calculated for the Neuropsychiatric Inventory (NPI), which was classified in the behavior domain. It was noted that two of the three studies that evaluated this outcome were lacking sufficient power^{60,64} (PW = 0.11, PW = 0.16). The test for heterogeneity was significant but the test for overall effect was not. Thus, the results of the summary estimates for the NPI outcome are problematic. Two global assessment outcomes, the CIBIC+ and the CDR, show a consistent effect favoring the drug treatment at 5 mg; the evidence is inconsistent for the 10 mg dose.

Figure 7. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the CIBC+ (continuous data) comparing donepezil versus placebo




Figure 8. Relative Risk (RR) from the Random Effects Model (Random) for the CIBC+ (dichotomous data probability of improving) for a 5 mg dose of donepezil.




Figure 9. Relative Risk (RR) from the Fixed Effect Model (fixed) for the CIBC+ (dichotomous data [improved versus not]) for a 10 mg dose of donepezil.





Figure 10. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the Clinical Dementia Rating (CDR) comparing donepezil versus placebo.



Five of the eight studies^{60,61,63,64,65,66,67,69} measuring Activities of Daily Living (ADL) found a significant difference in the various outcomes used to assess ADL, but none of these could be combined into a summary estimate. It should be noted that the majority of trials selected these ADL variables as secondary outcomes. Behavior outcomes were not significant or showed mixed results for the three studies that evaluated this domain but these lacked sufficient power. Only one study collected caregiver stress and health service utilization outcomes⁶³ but did not report these data.

Quality scores for reporting adverse events varied from 1 to 4 but the majority of trials scored 3 or greater ($n=7$). One recently published study⁶⁸ scored 1, with no events detailed. Withdrawal due to adverse events ranged from 0 - 18% for treatment groups and 0 - 11% for placebo (see Evidence Table 21). Four studies^{65,68,67,66} were able to demonstrate a dose effect, with increasing frequency of events as dosage increased. One study⁶² reported significant differences between treatment and placebo for balance problems and asthenia (neurological fatigue). Fatigue was shown to be significant in two other studies.^{61,66} Four^{65,61,66,67} out of six studies testing differences between groups were statistically significant for diarrhea (placebo = 3 - 21%, all doses donepezil = 0 - 38%), nausea and vomiting (placebo = 4 - 9%, all doses donepezil = 4 - 25%). The other a priori symptom reported was agitation and frequencies for placebo varied from 0 - 8% and for all doses from 3 - 19%; but these were not shown to be statistically different. No serious adverse events requiring hospitalization were reported or shown to differ statistically between groups.

Galantamine. See Evidence Tables 22 through 29 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Six studies of galantamine^{71,72,73,74,75,76} were included in this review. All compared galantamine with placebo and were published between 2000 and 2002 (from six different authors).

Design/methodology. The sample sizes for subjects ranged from 285⁷³ to 978⁷⁴ with 3530 subjects evaluated in total. All quality scores were high, with either 7 or 8 on the Jadad scale. Funding sources for these studies varied; one study did not report funding source,⁷⁴ one was funded by a non-industry source,⁷⁵ one was partially funded by industry,⁷¹ and three were funded by industry.^{72,73}

Populations. All but one of the studies⁷¹ included AD patients only, and this single study mixed VaD and AD patients. All subjects in these studies were classified as mild to moderate.

One study specified that the subjects were from the community.⁷³ All studies included from 36 - 53% male subjects, and the mean age ranged from 72.2 to 76.8 years. Three studies^{71,74,76} specified race, and with the largest proportions being white (range from 91.5 - 99.9%). Two studies evaluated subgroups, based on baseline MMSE⁷⁵ and APOE genotype.^{75,76}

Intervention. All studies had a titration period, starting at 4 mg per day^{71,73} or 8 mg per day.^{72,74,75,76} Four studies increased the dose weekly,^{71,72,75,76} and one study increased every 2 to 3 days.⁷³ All studies had a treatment dose of 24 mg per day. Three studies^{73,75,76} included other doses ranging from 16 mg per day to 36 mg per day. Trials were undertaken for a minimum of 3 and maximum of 6 months.

Primary outcomes. All domains were measured except for specific cognitive tests and caregiver burden. All studies used the ADAS-cog and CIBIC or CGIC measures as primary outcomes. None reported baseline mean MMSE values.

Analysis. All but one⁷¹ of the studies reported ITT analysis, and the results of this study did not differ from the others.

Results and interpretation. See Summary Table 3. Five of the six trials^{71,72,73,74,75,76} that evaluated general cognitive function showed a significant effect. One trial⁷³ showed mixed effects with the ADAS-cog showing some improvement at the 24 mg but not the 32 mg dose level. Figures 11 and 12 show the pooled estimate for the ADAS-cog for five studies for 24 and 32 mg doses; one trial⁷¹ was excluded from the pooled estimate as the population of this study was thought to be a source of heterogeneity. However, the test for heterogeneity for the 24 mg dose (Figure 11) was significant ($p = 0.001$) despite omitting this study, but the overall effect was significant ($p = 0.0005$); this estimate should be interpreted with caution. The pooled estimate for the 32 mg dose (Figure 12) showed a consistent effect favoring treatment and was significant ($p < 0.00001$).

Figure 11. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing galantamine at 24 mg dose versus placebo.





Figure 12. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing galantamine at 32 mg dose versus placebo.



All studies evaluated global assessment with the CIBIC+ with one exception.⁷³ All studies with the exception of this study showed a significant difference between placebo and treatment for both the 24 and 32 mg doses. Figures 13 and 14 show the pooled estimates for the CIBIC+ for these two dosages and suggest an overall effect size of equivalent magnitude for either dose. Similarly, all studies evaluated quality of life/ADL with a variety of different outcome measures; four studies^{71,72,74,75} showed statistically significant differences between groups, and two studies did not.^{76,73} Figures 15 and 16 shows the results of pooling the estimates for the Disability Assessment for Dementia (DAD)⁷⁷ outcome in those studies that provided sufficient data. In this instance, we included the trial with mixed dementia populations⁷¹ to have a minimum of three studies required for a pooled estimate. A consistent, statistically significant effect favoring treatment is evident; the higher dose of 32 mg shows a slightly larger effect size relative to 24 mg. Two of the three studies that reported on behavior/mood outcomes showed statistically significant differences (Summary Table 3).

Figure 13. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC comparing galantamine at 24 mg dose versus placebo.




Figure 14. Relative Risk (RR) from the Fixed Effects Mode Fixed I for CIBIC comparing galantamine at 32 mg dose versus placebo.




Figure 15. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the DAD comparing galantamine at 24 mg dose versus placebo.





Figure 16. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the DAD comparing galantamine at 32 mg dose versus placebo.



Five^{71,73,74,75,76} of the six trials scored 3 out of 5 on our quality scale for rating adverse events. One trial⁷² scored 4. Withdrawal rates due to adverse events ranged from 4 – 9% for placebo and 8 – 27% for the treatment group. One study⁷³ showed a dose response for adverse events. Although, most trials did not report testing for differences between groups, two trials^{76,75} reported a statistical significant difference in weight loss between the placebo and treatment group. Statistical differences for aberrant hematology were not significant in any of the five studies that evaluated this (Evidence Table 29). The most common types of adverse events reported were gastrointestinal symptoms (nausea and vomiting, diarrhea), eating

disorders/weight loss, and dizziness (four studies, see Evidence Table 29). The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea and vomiting (placebo = 3 - 13%, all doses = 6 - 44%), 2) dizziness (placebo = 3 - 11%, all doses = 4 - 19%), 3) diarrhea (placebo = 2 - 10%, all doses = 4 - 19%), 4) agitation (placebo = 1 - 9%, all doses = 6 - 15%), and 5) eating disorder (placebo = 0 - 6%, all doses = 4 - 20%).

Metrifonate. See Evidence Tables 30 through 40 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Nine studies^{78,79,80,81,82,83,84,85,86} were eligible for this systematic review. Studies were published from 1996 to 1999, and all studies compared metrifonate to placebo.

Design/methodology. Overall, 2759 subjects were evaluated. Sample sizes randomized ranged from 27 to 605 subjects. Quality of the studies ranged from 5⁸⁰ to 8,⁸¹ with the majority of studies earning 6 points.^{82,83,79,85,78} Two studies were non-industry-funded^{78,79} and one did not specify the source of funding,⁸⁰ while the other six studies were partially or wholly funded by industry.

Populations. The subjects in all included trials were classified as having mild to moderate AD. Not all trials specified the source of recruitment or the racial composition of subjects. Three studies specified a community sample,^{86,79,85} and three trials reported the racial composition 1,^{83,86,84} which was greater than 90% white in all cases. Mean age for all of the studies ranged from 71.4 to 75.0 years, with one study not reporting the mean age.⁸⁰

Intervention. All but one study⁸⁶ reported the loading dose, which varied from 0.5 mg per kg to 5.0 mg per kg. Following this initial loading period, the maintenance dose varied from 0.65 mg per kg to 4 mg per kg and 50 mg per day. The duration of the study treatments varied from 21 days to 26 weeks.

Primary outcomes. All outcome domains were evaluated with the exception of caregiver burden. ADAS-cog, CIBIC+, and MMSE were most frequently used as outcomes.

Analysis. Four trials reported OC analyses^{78,79,80,85} and the remaining reported ITT analyses.

Results and interpretation. See Summary Table 4. A consistent positive change in cognitive function was found in all studies that reported this outcome (n = 8). One study⁸⁵ tested cognitive function, global assessment, and behavioral outcomes and reported the baseline endpoint scores but did not test for differences between treatment and placebo groups. Four of the eight studies reporting global assessment outcomes showed statistically significant differences between groups, and two^{83,86} showed mixed results. The remaining two trials showed no significant results for global assessment, but they were secondary rather than primary outcomes. All studies that evaluated behavior/mood and quality of life/ADL outcomes (with the exception of one trial⁸⁶) showed no significant findings or mixed findings; it should be noted that all were secondary outcomes. There were not enough similar outcomes reported to complete a pooled analysis for metrifonate.

With the exception of a single study, quality scores for reporting adverse events were greater than 3 and generally well reported. However, only one trial⁸³ tested for differences between

groups and found nausea and vomiting, diarrhea, and muscle and joint disorder to be significantly different. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea and vomiting (placebo = 3 - 14%, all doses = 2 - 50%), 2) dizziness (placebo = 1%, all doses = 3 - 4%), 3) diarrhea (placebo = 4 - 14%, all doses = 11-19%), 4) agitation (placebo = 2 - 14%, all doses = 8 - 33%), and none reported eating disorder as an adverse event. Withdrawal rates due to adverse outcomes varied from 0 – 9% for placebo and 0 – 12% for treatment groups. Some studies indicated arrhythmia^{80,82,83,85} and hypotension^{82,85} and hematological abnormality⁸² but did not test for differences between groups. The majority of studies reported that laboratory tests including liver function and hematology were within normal limits. Overall, it was difficult to determine which types of adverse events reported had the potential to cause serious harm. This is some concern as metrifonate is no longer used as a therapy for dementia due to its potential for serious adverse events that include: respiratory paralysis, bradychardia, severe leg cramps and dyspepsia.⁸⁷

Nicergoline. See Evidence Tables 41 through 47 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Four studies^{88,89,90,91} compared the effect of nicergoline to placebo, and one study,⁹² published in 1994, compared nicergoline to antagonistic-stress. The four placebo comparison studies were published in 1995,⁹¹ 1997,^{89,90} and 2001.⁸⁸

Design/methodology. Sample sizes in the controlled studies varied from 108 to 346, with the total number of subjects included totaling 705. The drug versus drug study had only 62 subjects. The placebo trials all had quality scores of 6 points, while the non-placebo trial had a quality score of 5. Funding sources were reported only in two studies,^{88,90} and both were industry-funded.

Populations. These studies had a very mixed population of dementia patients. Two included AD only,^{88,92} one trial MID only,⁸⁹ one trial included both senile dementia of Alzheimer type (SDAT) and MID,⁹¹ and one trial included PDD, VaD, and mixed dementia.⁹⁰ All subjects had mild to moderate dementia. Studies included 38 - 55% male subjects; one study⁹¹ did not report the gender proportions. Mean age of subjects ranged from 69.3 to 73.7 years with one study not reporting⁹¹ this value. One study⁹¹ compared SDAT patients to MID patients.

Intervention. All trials versus placebo used 60 mg per day, but duration varied from 2 months,⁹¹ 6 months,^{88,89,92} and 12 months.⁹⁰

Primary outcomes. Caregiver burden was the only domain not evaluated by at least one of the studies. Three trials^{89,90,91} specified baseline MMSE, and this varied from 20 to 22.

Analysis. Two of the trials reported OC analyses^{90,91} and two reported ITT analyses^{89,88}; the trial comparing nicergoline to antagonistic-stress presented OC analysis only.

Results and interpretation. See Summary Table 5. There was a consistent positive effect for improvement in general cognitive function as all four studies showed a statistically significant difference. The evidence for benefit in the global assessment domain is inconclusive as only two of the trials^{89,90} found significant differences and two trials^{88,93} had mixed results (see Summary Table 5). Two trials^{91,88} measured behavior/mood and found no significant difference. A single

trial⁸⁸ evaluated quality of life/ADL; although two outcomes in this domain were used (both as secondary measures), none was significant relative to placebo. There were not enough similar outcomes reported to complete a pooled analysis for nicergoline.

Quality scores for reporting adverse events varied from 2 to 5 for these four trials, and none tested for differences between groups. Withdrawal due to adverse events varied from 0 – 8% for placebo and 0 to 9% for the treatment group. The trial with the lowest quality reporting score⁸⁹ reported the most number of different events (up to 23 event types). With the exception of headache, which was reported in all four trials, it was difficult to determine which types of adverse events most characterized exposure to this pharmacological agent. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = 3%), 2) dizziness (placebo = 1 – 2%, all doses = 0% or not reported), 3) diarrhea (placebo = 2 – 6%, all doses = 2 – 4%), 4) agitation (placebo = 5%, all doses = not reported), and none reported eating disorder as an adverse event.

Physostigmine. See Evidence Tables 48 through 53 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Four studies were eligible for our review,^{94,95,96,97} all comparing physostigmine to placebo only. The studies were from 1996, 1999, and 2000 with two of the studies being by the same author.^{95,96}

Design/methodology. Sample size ranged from 176⁹⁷ to 475⁹⁶ with an overall total of 1198 subjects. Quality scores were 5,⁹⁶ 6,^{94,95} and 7⁹⁷ out of 8 possible points. Two studies were industry-funded,^{97,96} one was partially funded by industry,⁹⁵ and one did not report funding source.⁹⁴

Populations. All subjects had a diagnosis of mild to moderate or probable AD. Only one study⁹⁶ reported that all subjects were drawn from the community. Mean age ranged from 68.6 to 73.4 years, and the proportion of male subjects varied from 39.8 – 63%.

Intervention. Treatment schedules varied across the studies. One study⁹⁴ used a patch (30 and 60 mg), one trial used 30 mg (15 mg twice daily),⁹⁷ one trial had a washout and titration every 3 weeks to 30 or 36 mg per day,⁹⁶ and another trial titrated weekly to 15 mg twice daily.⁹⁵ Duration of treatment ranged from 6 weeks⁹⁵ to 24 weeks.^{94,96}

Primary outcomes. All studies used the ADAS-cog as a primary outcome, and none reported caregiver burden or behavior/mood measures. No studies reported baseline MMSE scores.

Analysis. All but one trial⁹⁴ used ITT analysis.

Results and interpretation. See Summary Table 6. Although all four trials measured general cognitive function, only three reported the results. All of these were statistically significant (see Summary Table 6) for the ADAS-cog, with change scores varying from 0.95 to 2.9 (change from baseline). Two trials^{95,96} found significant change for global assessment outcomes; the remaining two showed mixed results⁹⁷ and non-significance.⁹⁴ Behavior/mood was only measured in one trial,⁹⁴ and the effect was not reported. Three of the trials included measures of

quality of life/ADL as secondary outcomes^{95,96,97} and all found no significant difference from placebo but these were secondary outcomes and may reflect a lack of power. There were not enough similar outcomes reported to complete a pooled analysis for physostigmine.

The quality scores for reporting adverse events were generally low, scoring 1 or 2 out of 5. Withdrawal rates due to adverse events varied from 1 - 5% for placebo and 12 - 55% in the treatment group, with one study⁹⁷ not reporting rates. The high withdrawal rates were in studies with sample sizes that varied from 181 to 475 subjects. A single study⁹⁷ tested for differences between groups, and found that dizziness, tremor, weight loss, asthenia (varying from 6 - 22% for all doses), confusion, delirium, and respiratory problems were significantly different. The cluster of reported types of adverse events suggests that gastrointestinal problems (abdominal pain, diarrhea) (placebo = 1 - 9%, all doses = 13 - 28%), nausea and vomiting (placebo = 1 - 9%, all doses = 9 - 75%) and eating disorder (placebo = 2 - 6%, all doses = 5 - 16%) were most frequently reported. Dizziness (placebo = 4 - 13%, all doses = 11 - 38%) and agitation (placebo = 6 - 16%, all doses = 4 - 8%) were also reported. No events deemed serious enough for hospitalization were reported.

Posatirelin. See Evidence Tables 54 through 59 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Four studies^{98,99,100,101} compared posatirelin to placebo, and one of these¹⁰¹ also compared it to citicoline. One study was published in each of the years from 1995 to 1998, and all studies were conducted in Italy. Two studies were by the same author.^{100,101}

Design/methodology. Populations randomized in the studies varied from 136¹⁰⁰ to 360⁹⁹ with a total of 931 in all trials. Quality scores ranged from 5⁹⁸ to 7¹⁰⁰ out of a possible 8 points. Three of the four studies did not report the source of their funding, but one trial¹⁰⁰ reported partial funding by industry.

Populations. No two studies included exactly the same populations; one trial had only AD,¹⁰¹ one trial had only VaD,¹⁰⁰ one trial had mixed AD and VaD,⁹⁹ and another trial had mixed AD, VaD, and PDD.⁹⁸ This latter trial⁹⁸ compared populations in a subgroup analysis of AD versus VaD. All studies evaluated populations with mild to moderate disease.

The mean age of the subjects ranged from 69.4¹⁰⁰ to 78.8⁹⁸ with the percentage of male subjects varying from 34¹⁰¹ to 66%.¹⁰⁰ One study¹⁰⁰ included a dementia population who also had hypertension.

Intervention. A dose of 10 mg per day was used in all studies, and treatment interval varied from 3 to 4 months.

Primary outcomes. General cognitive function was evaluated in all studies using the intellectual impairment Gottfries-Br  ne-Steen (GBS) subscale and the MMSE was used in one trial.¹⁰¹ Specific cognitive function was evaluated in one trial.¹⁰⁰ Quality of life/ADL was evaluated with the ADL subscale of the GBS, and behavior/mood was evaluated with the emotional impairment subscale of the GBS. GBS total score was assumed to be a measure of quality of life/ADL rather than global assessment. None of the trials reported baseline MMSE scores.

Analysis. Two of the studies^{100,101} used OC analysis to report outcomes.

Results and interpretation. See Summary Table 7. All four studies evaluated general cognitive function, and three of these trials^{100,99,98} reported significant differences using the intellectual impairment subscale of the GBS as a measure of this attribute. One study¹⁰⁰ measured this same outcome and the MMSE, but did not report results for the latter outcome. One study¹⁰¹ measured reported changes within a treatment relative to baseline and not relative to placebo; this study demonstrated superiority for posatirelin relative to citicoline (a third comparison group), but did not test for differences between the placebo group. Showing non-inferiority of citicoline in this trial does not establish efficacy with respect to placebo. Statistically significant changes were also shown for the domain of quality of life/ADL as measured by the GBS total score or GBS ADL (FactorII) subscales in three of the studies.^{98,99,100} A single trial¹⁰⁰ evaluated global assessment using the TP Global scale in VaD subjects. There were not enough similar outcomes reported to complete a pooled analysis for posatirelin.

Quality scores for reporting adverse events varied from 2 to 4. Withdrawal rates due to adverse events ranged from 0 - 3% in placebo and 0 - 4% in the treatment group. One trial¹⁰⁰ did not report the rate of withdrawal. None of the studies tested for significant differences between groups. All studies reported the presence of agitation, and at least three studies reported arrhythmia, nausea/vomiting, headache, rash/skin disorder, and sleep disorder; there is no evidence to suggest that these differed between placebo and treatment group. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = 1 - 4%), 2) dizziness (placebo = not reported, all doses = 1%), 3) diarrhea (placebo = 2%, all doses = 2%), 4) agitation (placebo = 1 - 5%, all doses = 1 - 5%), and none reported eating disorder as an adverse event. No serious adverse events were reported.

Rivastigmine. See Evidence Tables 60 through 67 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Six studies were eligible for this review, all comparing rivastigmine to placebo. Studies were published in 1998,^{102,103} 1999,^{104,105} 2000,¹⁰⁶ and 2001.¹⁰⁷

Design/methodology. Six studies evaluated 2071 subjects in total, with studies ranging from 27¹⁰⁷ to 725¹⁰⁴ subjects. The quality of studies varied from 5 to 8, with three studies^{106,104,103} scoring 8 points and one¹⁰⁷ earning 5 points. All studies were funded by industry sponsors.

Populations. Four studies were evaluated in AD patients,^{107,105,104,103} one trial¹⁰² dementia of the Alzheimer's type (DAT), and one study¹⁰⁶ Lewy body dementia subjects for mild to moderately severe subjects. One study¹⁰³ reported a subgroup analysis by vascular risk. One study reported a community sample in their trial.¹⁰⁴ Mean age for the studies ranged from 69.4¹⁰² to 75.9 years.¹⁰⁷ Two studies^{107,105} did not report the ratio of male subjects in their study, and the other four varied from 39 - 56%. One trial¹⁰⁴ reported co-morbidity of diabetes, hypertension and arthritis; one trial¹⁰³ reported concurrent medication use for cardiovascular, gastrointestinal and analgesic aids.


Intervention. Doses for rivastigmine varied from 1 mg¹⁰³ to 12 mg,^{106,105,104} and treatment duration varied from 14¹⁰⁴ to 26^{107,103} weeks. All studies titrated the dose of drug over a period ranging from 2 weeks¹⁰⁶ to 12 weeks.¹⁰⁷

Primary outcomes. The ADAS-cog and CIBIC+ were evaluated in half the studies. Baseline MMSE was reported in two trials,^{103,106} and the mean scores varied from 18 to 20. Specific cognitive function, behavior/mood, and quality of life/ADL were infrequently evaluated, and caregiver burden was not evaluated in the trials.

Analysis. Trials were evenly divided between ITT analyses^{106,104,103} and OC.^{107,105,102}

Results and interpretation. See Summary Table 8. Although, general cognitive function was evaluated in five studies, only four reported findings, which were all statistically significant; one of these was borderline significant ($p = 0.054$) (see Summary Table 8). Two of these studies^{104,103} represented two sites (North American and European) of the same protocol. Although the same protocol was used, one study¹⁰³ found significance for both high (6 – 12 mg) and lower (1 – 4 mg) dosages, but the other trial did not show significance for the lower dose, likely due to lack of power for this outcome ($PW = 0.67$). Interestingly, at the 12-week midpoint, the low dose groups in both these studies appeared to be worse than placebo for all primary outcome measures, but then migrated to improvement at the 26-week endpoint. For those studies^{105,104,103} that reported ADAS-cog change scores from baseline for the treatment group, mean change values varied from -2.75 to 0.26. Figure 17 shows the pooled estimate for those trials that provided sufficient data and represents the 12 mg dose. However, the test for homogeneity was significant suggesting that the pooled estimate should be interpreted with caution (albeit a significant overall effect). For those studies that reported MMSE change scores from baseline for the treatment group, mean values varied from 0.0 to 0.6; a single trial¹⁰³ showed a decline of 7.9 points relative to baseline for the placebo after 26 weeks (other trials reporting MMSE scores did not report such marked change).

Figure 17. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing rivastigmine versus placebo.




Two trials evaluated specific cognitive tests; one trial¹⁰⁶ showed statistical significance for the CCASSS but not for the MMSE (general cognitive test). Similarly, another trial¹⁰⁵ found the Weschler Logical memory (instant recall) to be significant but the ADAS-cog (general cognitive outcome) was borderline significant.

With respect to global changes, five of six studies showed significant changes, and from these, three studies^{104,103,102} were for the high dose only. One of these studies¹⁰² defined the high dose as 6 mg per day, which was the minimum dose level for the other two studies. One study¹⁰⁷ showed a statistical difference for the two deterioration categories (5 – 7) in the CIBIC+, but not in the improvement categories when comparing treatment and placebo groups. Figure 18 shows

the pooled estimate for the CIBIC+ in those studies that provided sufficient information. A consistent effect favoring treatment is shown, but the two smaller trials display large confidence intervals (Figure 18). There was no clear trend in the domains of behavior/mood and quality of life/ADL as not all studies evaluated these domains.

Figure 18. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC+ comparing rivastigmine versus placebo.



Quality scores for reporting adverse events varied from 2 to 5. Withdrawal rates due to adverse events ranged from 4 - 11% in the placebo and 11 - 27% in the treatment group. One trial¹⁰⁷ did not report the withdrawal rates or the types of adverse events observed. Two trials^{103,104} demonstrated a dose response; however, one of these trials¹⁰⁴ showed significant differences for nausea and vomiting only, and the other trial¹⁰³ showed significant for all the adverse events reported. With respect to the types of adverse events, the majority of studies reported dizziness, nausea and vomiting, eating disorder/weight loss, and headache. It should be noted that one study¹⁰⁵ allowed intentional prescribed anti-emetic drugs to increase the tolerance of subjects taking rivastigmine. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3 - 10%, all doses = 8 - 58%), 2) dizziness (placebo = 0 - 7%, all doses = 6 - 20 %), 3) diarrhea (placebo = 2 - 9%, all doses = 7 - 17%), 4) eating disorder (placebo = 4 - 8%, all doses = 4 - 19%), and 5) agitation was not reported. No serious adverse events were reported.

Tacrine. See Evidence Tables 68 through 77 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Eight studies evaluating tacrine were eligible for this review; tacrine was compared to placebo in six trials^{108,109,110,111,112,113} (lecithin was assumed to be like placebo) and to other drugs in two trials.^{114,26} One study¹¹⁴ compared two arms with tacrine, one with silymarin added and one placebo arm. The other non-placebo controlled trial compared tacrine with idebenone.²⁶ These drug versus drug trials were published in 1999 and 2002. The placebo controlled trials were published 1991,¹¹¹ 1994,^{108,109,113} 1996,¹¹⁰ and 1999.¹¹²

Design/methodology. The placebo studies evaluated 994 patients in total with a range of 13¹¹¹ to 663¹⁰⁸ subjects per study. Quality scores out of 8 points were evenly distributed with two studies each having scores of 5, 6 and 7. Both drug versus drug trials had scores of 7. One study¹⁰⁹ was not funded by industry, and one trial¹¹⁴ did not report its source of funding; the other six studies had at least partial, if not full, industry support.

Populations. All studies included AD subjects; one trial²⁶ also included PDD patients. Subjects in all studies had mild to moderate or probable disease. Five of the studies^{113,109,111,114,26} reported that their sample was from the community. The mean age ranged from 68¹¹⁰ to 75 years,¹¹³ and the percentage of male subjects in the studies varied from 13¹¹⁰ - 54%.¹¹³ One study reported race with 100% white subjects.²⁶

Intervention. See Summary Table 9. All placebo-controlled trials used a titration period to get to maximum dose, from 11 days¹¹³ to 18 weeks.^{112,108} Treatment doses varied from 80 mg per day¹¹⁰ to 160 mg per day.¹⁰⁸ Treatment duration was either 12/13 weeks,^{110,113,111} or 30/36 weeks^{112,108,109} for all placebo-controlled studies. The trial versus Idebenone²⁶ was for 60 weeks and the trial with silymarin¹¹⁴ was for 15 weeks.

Primary outcomes. All six of our identified domains were evaluated by at least one trial. All trials measured cognition; however, sufficient data to permit pooled analyses could not be adequately abstracted from all these studies that had similar outcomes. Baseline MMSE varied from 14 to 18 in the four trials^{112,110,109,114} that reported this.

Analysis. Five studies^{108,112,113,114,26} reported ITT analysis and three did not.^{109,111,110}

Results and interpretation. Of the six placebo-controlled studies, only one trial¹⁰⁸ showed statistical significance for general cognitive function as measured by the ADAS-cog (ES = -0.268). Three doses (80 mg, 120 mg, 160 mg) were compared in this trial, and the 120 and 160 mg per day were shown to be statistically significant (approximately a mean change of 2 points on the ADAS-cog). One trial¹¹² showed mixed results for the two outcomes used (CASI and MMSE) to evaluate general cognitive function; this trial was underpowered for both these outcomes (PW = 0.22 and 0.26, respectively). Three trials^{109,110,111} found no statistical differences between treatment and placebo but had small sample sizes, ranging from 12 to 32 subjects, and were likely underpowered (insufficient reporting to estimate power). A fourth trial¹¹³ also found no statistical difference (p = 0.55) for general cognitive function, but the study duration was 12 weeks. It should be noted that this study used an 80 mg dose, which was shown to have no benefit relative to higher doses of 120 and 160 mg.¹⁰⁸ A single trial¹⁰⁹ of small sample size evaluated specific cognitive tests and did not show statistical differences.

Three studies evaluated global assessment, and two^{113,108} found statistical significance; the trial showing no benefit¹¹² also showed inconclusive findings for general cognitive function as well (PW = 0.05 for the CGIC). Four trials^{108,109,110,113} evaluated behavior/mood and showed no difference between groups. Two trials^{109,111} with small sample sizes measured quality of life/ADL and showed no significant changes; lack of sufficient power cannot be ruled out. There were not enough similar outcomes reported to complete a pooled analysis for tacrine.

The quality scores for reporting adverse events varied from 1 to 3. The proportion of subjects withdrawing due to adverse events ranged from 0 - 12% for placebo and 0 - 55% in the treatment group. The higher rates of withdrawal were associated with higher doses. Elevated alanine transaminase (ALT) or hepatic abnormality (placebo = 4 - 13%, all doses = 7 - 67%) was reported in six studies, suggesting the potential for serious liver damage. None of these trials tested for differences between treatment and placebo with respect to adverse events. Five of the

studies reported nausea and vomiting (placebo = 0 - 9%, all doses = 9 - 37%); gastrointestinal problems and dizziness (placebo = 0 - 16%, all doses = 4 - 14%) was also noted in several studies. Frequencies of other a priori symptoms of interest are as follows: 1) agitation (placebo = 5 - 12%, all doses = 5 - 9%), and 2) diarrhea (placebo = 0 - 13%, all doses = 4 - 18%).

Velnacrine. See Evidence Tables 78 through 82 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Three studies evaluated velnacrine versus placebo, and these were published in 1991,¹¹⁵ 1995,¹¹⁶ and 1996.¹¹⁷

Design/methodology. A total of 774 subjects were studied with sample sizes ranging from 16¹¹⁵ to 449.¹¹⁶ Quality scores out of a possible 8 points varied from scores of 6^{117,115} or 7.¹¹⁶ All studies were sponsored by industry.

Populations. The characteristics of the populations all included probable AD subjects. The mean age of the participants ranged from 70.5 to 72.8 years and the percentage of male subjects ranged from 31 - 41%. Location of recruitment was not specified.

Intervention. The doses given for this drug overlapped between the studies, but they were on different schedules (once, twice, or three times per day). None of the studies had a titration period. One study¹¹⁷ compared four doses (three daily doses of 10 mg, 25 mg, 50 mg, and 75 mg). One study¹¹⁶ compared doses of 150 mg or 225 mg per day. The other study used a dose of 100 mg twice daily and had the smallest sample size.

Primary outcomes. General cognitive function was evaluated in all studies with the ADAS-cog, and none specified baseline MMSE values. A variety of outcomes were used to evaluate global assessment. At least one of these trials evaluated the other outcomes domains

Analysis. Only one of the trials¹¹⁶ used an ITT analysis.

Results and interpretation. See Summary Table 10. One¹¹⁵ trial was of very small sample size (n = 16) and of a 2 weeks duration, compared to the other two studies^{117,116} with 15 or 24 weeks. Similarly, this trial evaluated two outcomes (specific cognitive function and global assessment) and showed mixed or non-significant results, likely a function of being underpowered. The two remaining studies^{117,116} had sample sizes over 300 subjects and showed statistical significance for the domain of general cognitive function using the ADAS-cog. The magnitude of the change reported varied from -2.0 at 12 weeks and then -1.0¹¹⁶ at 24 weeks for the 225 mg dose group only; a mean change of 2.15¹¹⁷ for the 75 mg (three times daily) as observed at the study endpoint of 15 weeks (no other dosage group was reported for this study). The trial¹¹⁶ evaluating doses of 150 and 225 given once daily showed significant changes for 225 mg per day but not for 150 mg per day at endpoint (24 weeks), whereas, the trial with 75 mg twice daily did show significant change for general cognitive function.

All studies included assessment of global functioning, for which two^{117,116} found significant differences, and one¹¹⁵ had mixed results. Behavior/mood was evaluated in only one study¹¹⁷ as a secondary outcome with an OC analysis, and no significant effect was found. Similarly, quality of life/ADL was measured in two studies as secondary outcomes, which produced opposite

results (not significant¹¹⁷ and significant¹¹⁶). One of the trials¹¹⁶ measured effects on caregiver burden as a secondary outcome and found a significant effect. There were not enough similar outcomes reported to complete a pooled analysis for velnacrine.

Quality scores for reporting adverse events were 3 for all studies. Withdrawal rates varied from 0 - 22% for the placebo group and 5 - 33% for the treatment group. None of the studies reported a dose response. None of the studies tested for statistical differences between the placebo and treatment groups. Two studies reported aberrant hematology and hepatic abnormality^{116,117}; for these two studies the rate of occurrence were 2 - 21% for placebo, and 32 - 40% for all doses. All studies reported diarrhea and nausea and vomiting. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 0 - 4%, all doses = 3 - 8%), 2) dizziness (placebo = 3%, all doses = 0 - 8%), 3) diarrhea (placebo = 3%, all doses = 2 - 33%), 4) agitation (placebo = 4%, all doses = 1 - 4%), and 5) eating disorder (placebo = 1%, all doses 2 - 4%).

Various cholinergic neurotransmitter modifying agents. See Evidence Tables 83 through 93 at <http://www.ahrq.gov/clinic/epcindex.htm>. See Summary Table 11.

The remaining agents classified as cholinergic neurotransmitter modifying agents were grouped according to the number of studies for the purposes of presentation:

Cholinergic pharmacological agents that had two trials eligible for this review and were compared to placebo.

Eptastigmine (Evidence Tables 83, 84, 85, 93). Two trials^{118,119} evaluated eptastigmine in patients with mild to moderate AD (103 patients for 4 weeks and in 491 patients for 24 weeks). Both trials were industry-funded. The trial that used an ITT analysis¹¹⁸ and had the longer duration (24 weeks) showed significant change in the three domains: general cognitive function, global assessment, and quality of life/ADL. The OC analysis¹¹⁹ of the patients treated for 4 weeks showed no significant effect. One trial used the ADAS-cog as a primary outcome¹¹⁸ and showed a small increase of 1.05 and 0.41 for the 15 and 20 mg thrice daily doses, respectively, relative to the placebo group (which increased by 2.6 points). The CIBIC+ was significant for the higher dose group only in this same trial. The evidence of benefit for eptastigmine remains inconclusive given the lack of consistency between studies.

Linopirdine (Evidence Tables 83, 87, 88, 93). Two 1997 trials^{120,121} evaluated linopirdine in patients with mild to moderate AD patients for 4 or 6 weeks at 40 or 30 mg thrice daily. Both were at least partially industry-funded. One trial¹²⁰ included 382 patients on 30 mg dose during a 6 month trial and used an ITT analysis; this study showed statistically significant findings for general cognitive function alone as measured with the ADAS-cog (mean change 2.0 points); global assessment, quality of life/ADL, and behavior/mood were not significant. All outcomes evaluated in the second trial¹²¹ were not significant, even though OC analysis was used.

Cholinergic pharmacological agents with only one trial eligible for this systematic review.

Huperzine-A (Evidence Tables 83, 92, 93). This study¹²² showed a statistically significant benefit relative to placebo in an OC analysis of all domains that were evaluated: general cognitive function, behavior/mood, and quality of life/ADL. The study population was 103 Asian patients with mild to moderate AD, who were treated for 8 weeks.

Sabeluzole (Evidence Tables 83, 89, 93). This study¹²³ included 39 patients with mild to moderate AD and lasted 48 weeks. General cognitive function as measured by the ADAS-cog showed approximately a 5 point increase compared to a 7 point increase for placebo. The OC analysis showed no significant difference from placebo in general cognition.

Results of non-cholinergic neurotransmitter/neuropeptide modifying agents (NCNMA)

A total of 35 drugs in 50 studies were classified as non-cholinergic neurotransmitter/neuropeptide modifying agents. These pharmacological agents can be seen in Table 3. Sixteen of these studies involved direct comparisons to other drugs and these are considered separately in the section addressing Question Three. Overall results for each of the trials each intervention are detailed in OST located at the end of this chapter and organized by drug. All other study details are available in Evidence Tables 95 through 161 in the Appendices.

Table 3. List of Non-cholinergic neurotransmitter/neuropeptide modifying agents and the number of studies vs. placebo for each of these. Asterisk (*) indicates report of a drug vs. drug trial [comparator drug(s) in brackets].

Drug	Number of studies vs. placebo	Drug	Number of studies vs. placebo
Alaproclate	1	Memantine	3
Alprazolam *[Lorazepam]	0*	Mianserin *[Citalopram]	0*
Anapsos	1	Minaprine	1
BMV (Nootropic)	1	Moclobemide	1
Carbamazepine	2	Naftidrofuryl	1
Citalopram *[Mianserin] *[Perphenazine]	2**	Olanzapine *[Lorazepam]	2*
Diphenhydramine *[Haloperidol, Oxazepam]	0*	Oxazepam *[Diphenhydramine Haloperidol]	0*
Divalproex	2	Paroxetine *[Imipramine]	0*
Fluoxetine *[Haloperidol] *[Amitriptyline]	2**	Perphenazine *[Citalopram]	1*
Fluvoxamine	1	Phosphatidylserine	2
Haloperidol **[Risperidone] *[Loxapine] *[Diphenhydramine Oxazepam] *[Fluoxetine] *[Tiapride] *[Trazodone]	4*****	Risperidone **[Haloperidol]	1**

Table 3. List of Non-cholinergic neurotransmitter/neuropeptide modifying agents and the number of studies vs. placebo for each of these. Asterisk (*) indicates report of a drug vs. drug trial [comparator drug(s) in brackets] (continued).

Drug	Number of studies vs. placebo	Drug	Number of studies vs. placebo
Imipramine *[Paroxetine]	1*	Selegiline *[Vitamin E]	7*
Lisuride	1	Sertraline	2
Lorazepam *[Alprazolam] *[Olanzapine]	1**	Thioridazine *[Loxapine]	1*
Loxapine *[Haloperidol] *[Thioridazine]	1**	Tiapride *[Haloperidol] *[Melperone]	1**
Lu25-109	1	Trazodone *[Haloperidol] *[5'-MTHF]	1**
Maprotiline	1	Xanomeline	1
Melperone *[Tiapride]	0*		

Haloperidol. See Evidence Tables 94 through 103 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Five studies^{124,125,126,127,128} evaluating haloperidol relative to placebo as well as another drug were included in this review. Three additional trials^{129,130,131} (from 1990, 1993, and 2001) compared haloperidol to another drug and did not include a placebo group (these are detailed in question 3). One study was published in each of 1982, 1997, and 1999; two were from 2000.

Design/methodology. Sample sizes for the placebo-controlled studies were generally small with samples of 15,¹²⁸ 64,¹²⁶ 149,¹²⁵ 344,¹²⁷ and 306¹²⁴ for an overall total sample size of 622 subjects. All but one of the studies had a quality score of 6 out of a possible 8 points; the other study¹²⁷ had a score of 7 points. One study¹²⁴ did not indicate a funding source, three studies^{125,126,127} indicated some industry funding, although none showed total industry funding, and one study¹²⁸ had no industry funding.

Populations. Populations evaluated in the studies included three with only mild to moderate or probable AD,^{124,125,128} one with PDD and MID,¹²⁶ and one with PDD, VaD, and mixed dementia¹²⁷ (which reported subgroup information about VaD versus all subjects). Two placebo studies¹²⁶ reported the presence of subjects with severe disease. Two trials^{126,127} studied institutionalized patients while one¹²⁸ looked at community subjects. Ages in the studies ranged from a mean of 72.7 to 81.0 years, and 33 - 49% of subjects were male.

Intervention. Haloperidol doses ranged from 3 mg to 20 mg per day for a treatment period of 3 weeks,¹²⁴ 6 weeks,¹²⁸ 10 weeks,¹²⁶ 12 weeks¹²⁷ or 16 weeks.¹²⁵ The other drugs that haloperidol was compared to included fluoxetine,¹²⁸ loxapine,¹²⁶ risperidone,¹²⁷ tiapride,¹²⁴ trazodone & BMT¹²⁵ in the placebo controlled studies; loxapine,¹²⁹ risperidone,¹³¹ oxazepam & diphenhydramine¹³⁰ were evaluated in the head to head comparisons.

Primary outcomes. All studies evaluated behavioral outcomes, and at least one study evaluated the effect of haloperidol in each of the other domains included in this review with the exception of specific cognitive function. None of the studies reported baseline MMSE values.

Analysis. Two studies^{124,127} reported ITT analysis and three^{128,126,125} did not.

Results and interpretation. See Summary Table 12. Of the five studies that had a placebo group, only three trials evaluated general cognitive function. One trial¹²⁷ did not report the results for this domain and two showed no significant difference.^{124,125} Three trials^{124,126,127} found statistical differences for outcomes in the behavior/mood domain, and two trials^{128,125} showed no change. One of these non-significant trials¹²⁸ evaluating behavior had a very small sample size (n = 12) and was likely underpowered. Four trials evaluated global function, and the two studies^{124,125} that reported findings based on the CGIC and CGI showed both improvement and no benefit, suggesting inconsistent evidence for this domain; it should be noted that one of these trials lasted for only 3 weeks.¹²⁴ One trial evaluated quality of life using the IADL and showed statistical difference in favor of the placebo. Two trials^{125,128} evaluated caregiver burden and showed no effect; one of these studies¹²⁸ had very small sample size and was likely underpowered. There were not enough similar outcomes reported to complete a pooled analysis for haloperidol.

The quality scores for reporting adverse events varied from 1 to 5. Only three of five studies reported withdrawal rates; the proportion of subjects withdrawing due to adverse events ranged from 5 – 17% for placebo and 17 – 33% in the treatment group. One trial¹²⁴ showed a dose response effect, but the study only lasted for 3 weeks. Three trials tested for differences between treatment and placebo with respect to extrapyramidal symptoms (placebo = 17 - 32%, all doses = 34 - 97%), and two^{124,125} found significant differences. One study¹²⁵ found significant differences between groups for balance-related problems. Although reported by only two trials, the range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = not reported), and 2) dizziness (placebo = 24%, all doses = 21%), 3) no frequencies were reported for agitation, diarrhea, or eating disorder.

Memantine. See Evidence Tables 104 through 108 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Three studies comparing memantine to placebo were eligible for review. One study was published in 1999¹³² and the other two^{133,134} in 2002.

Design/methodology. Sample sizes ranged from 166 to 579 for a total population evaluated of 1066 subjects. Two studies^{132,133} earned 6 points out of 8 for the quality score while the other¹³⁴ earned 7 points. One report¹³² did not indicate the source of funding, and the other two had industry support or funding.

Populations. Two studies^{133,134} included VaD patients only, one of which¹³⁴ analyzed subgroups based on MMSE, type of VaD, and gender. The other study¹³² included VaD, DAT and PDD patients and did subgroup analysis comparing VaD to DAT and grouping for care dependence. One trial¹³² included patients with severe disease and was the only study to report that all of their subjects were institutionalized. One study¹³⁴ included only community subjects and the other study¹³³ did not report source of patients. Study subjects had a mean age of 71.2,¹³² 76.4,¹³³ and 77.4¹³⁴ years, and 42 - 53% were male.

Intervention. Two studies^{133,134} had a 4 week titration period with a final dose of 20 mg per day for the remaining 24 week study duration. The third study used a 2-week titration period with a final dose of 10 mg per day for the remaining 10 weeks of the study.

Primary outcomes. All studies evaluated global function. The ADAS-cog was evaluated in two studies^{134,133} and showed smaller changes of decline relative to placebo by approximately 1.5 points. All studies measured global function with the CGI-C but did not provide variance data to permit the calculation of the pooled estimates. Although, all trials measured MMSE, none reported baseline values for this outcome. Only one trial¹³² evaluated behavior/mood and quality of life/ADL. No study evaluated specific cognitive function or caregiver burden.

Analysis. All studies performed ITT analysis.

Results and interpretation. See Summary Table 13. Two studies^{134,133} in subjects with mild to moderate VaD showed significant findings for general cognitive function but not global assessment. The power could be estimated for one of these trials¹³³ and was found to be below acceptable levels (PW= 0.60). The third memantine trial¹³² in this review evaluated mixed dementia populations (including some VaD) with moderate to severe dementia and found significant differences for global function, behavior/mood, and quality of life/ADL, but did not evaluate general cognitive function. It should be noted that this trial¹³² used half the dose of memantine for half the study duration in patients with greater disease severity, and had approximately half the sample size of the other two trials evaluated in this systematic review. There were not enough similar outcomes reported to complete a pooled analysis for Memantine.

The quality scores for reporting adverse events varied from 3 to 4. Only two of three studies reported withdrawal rates; the proportion of subjects withdrawing due to adverse events ranged from 7 – 13% for placebo and 9 – 12% in the treatment group. One trial¹³³ tested for differences between treatment and placebo, and none of the comparisons were significant. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = 5%), 2) dizziness (placebo = 3 - 8%, all doses = 6 - 11%), 3) diarrhea (placebo = 4%, all doses = 4%), 4) agitation (placebo = 7 - 8%, all doses = 4 - 5%), and none reported eating disorder as an adverse event.

Selegiline. See Evidence Tables 109 through 116 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Six studies evaluated the effect of selegiline compared to placebo.^{135,136,137,138,139,140} A single study¹³⁵ compared selegiline to vitamin E, placebo and the combination of selegiline plus vitamin E. The studies were published in 1991,¹³⁶ 1992,¹³⁷ 1993,¹³⁸ 1997,¹³⁵ 1998,¹⁴⁰ and 1999.¹³⁹

Design/methodology. Sample sizes ranged from 10¹³⁷ to 341¹³⁵ with a total population evaluated of 733 subjects. Study quality scores were 5,^{140,135,137} 6,^{139,138} and 7.¹³⁶ Three trials^{139,137,136} did not report the source of funding, and the other three^{140,135,138} had some industry support.

Populations. Studies included patients with mild to moderate PDD, DAT, and AD. Two subgroup analyses based on the results of the clock drawing test¹³⁹ and the GDS result¹³⁶ were

reported. One study reported that the included patients were institutionalized.¹³⁹ Mean age of the subjects in the trials ranged from 68.6 to 83.0 years and all had male subjects (29 – 74%).

Intervention. All trials used the same dose, 10 mg per day, with three of the trials^{135,138,137} giving the drug in two 5 mg doses. One trial¹⁴⁰ reported a titration period of 7 days. The duration of the trials varied with treatment times of 2 months,¹³⁷ 3 months,¹³⁶ 6 months,^{139,140} 15 months,¹³⁸ and 24 months.¹³⁵

Primary outcomes. Quality of life/ADL and caregiver burden were not evaluated in any of the studies.

Analysis. Two studies^{135,140} carried out ITT analysis.

Results and interpretation. See Summary Table 14. Five of the six trials evaluated general cognitive function, and of these, only four reported their findings. Two of the trials^{138,140} showed non-significant findings, but these had very small sample sizes (10 and 41 subjects) and were likely underpowered. Two trials^{137,139} showed mixed results, and one of these was likely underpowered.¹³⁷ One trial¹³⁶ found significant changes for specific cognitive tests (Sternberg Memory tests). Similarly, this same trial showed significant differences for global assessment and behavior/mood. This is the only trial that showed consistently positive findings across domains tested, and it also had the highest quality score (7). However, the other studies evaluating specific cognitive functions, global assessment, and behavior/mood did not show consistent results (non-significant or mixed findings). There were not enough similar outcomes reported to complete a pooled analysis for selegiline.

There is some evidence that shows that selegiline and selegiline combined with vitamin E, increases the time to important functional decline milestones¹³⁵ using time to event in the survival analysis. The results of this study showed that the vitamin E, selegiline, and combined groups were statistically different (i.e., declined less) from the placebo group in analyses that included baseline MMSE score as a covariate (not significant when excluded). The median survival was 230 days (vitamin E), 215 days (selegiline), and 145 days (combined group). Moreover, the vitamin E group showed a statistically significant difference for the endpoint of institutionalization, and the other treatment groups did not. Thus, the findings of this study suggest that selegiline and vitamin E may delay clinically important deterioration in patients with moderately severe AD; this delay varied from 20 to 32 weeks. It should be noted that this study evaluated subjects over a 2 year period, the longest of any dementia trial; moreover, the population was moderate to severe with respect to severity.

The quality scores for reporting adverse events varied from 0 to 3. The proportion of subjects withdrawing due to adverse events ranged from 0 - 4% for placebo and 0 – 9% in the treatment group. Two trials^{137,138} did not report any adverse events. Only one trial¹³⁵ tested for differences between the treatment and placebo groups and showed that balance (worse) and falls were significantly different between groups (particularly the group with selegiline combined with vitamin E (22%) versus placebo (5%)). However, when adjusted for multiple comparisons, these were no longer significant. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 2%, all doses = 0%), 2) dizziness (placebo = 2 - 20%, all

doses = 0 - 30%), and 3) agitation (placebo = 4 - 16%, all doses = 4 - 23%); no trial diarrhea or reported eating disorder as an adverse event.

Various non-cholinergic neurotransmitter/neuropeptide modifying agents. See Evidence Tables 117 through 160 at <http://www.ahrq.gov/clinic/epcindex.htm>. See Summary Table 15.

Ten non-cholinergic neurotransmitter neuropeptide modifying agents versus placebo were studied in only two included trials:

Anapsos (Evidence Tables 117, 119, 120, 160). Anapsos versus placebo was reported for a total of 114 patients with AD or VaD in reports published 1993¹⁴¹ and 2000.¹⁴² Both studies were partially funded by industry, and varied in the drug dose and duration; one trial¹⁴² used 360 mg per day or 720 mg per day for 4 weeks, and the other trial¹⁴¹ used 300 mg three times a day for 12 weeks. They each reported only one domain, and showed a significant change for general cognitive function¹⁴² and no significant results for global assessment.¹⁴¹

Carbamazepine (Evidence Tables 117, 121, 122, 160). Two trials^{143,144} evaluated carbamazepine in a total of 72 patients. The 1998 study¹⁴⁴ included a mixed severity population of institutionalized patients with non-industry-funding but also some financial support from industry. Both studies titrated up from 100 mg per day to 300 mg per day for 6 weeks. They evaluated all domains except caregiver burden. The trial using OC¹⁴³ population showed no significant effect for all outcomes tested but was likely underpowered (n = 16). The trial using ITT¹⁴⁴ showed a significant change in global assessment and behavior/mood. The evidence for benefit remains inconclusive given the lack of consistency between trials.

Citalopram (Evidence Tables 117, 123, 124, 160). Citalopram was evaluated in a total of 183 patients with mixed dementias including AD, VaD, mixed, MID, PDD. One trial¹⁴⁵ was non-industry-funded and the other¹⁴⁶ did not report funding source. Treatment was 20 mg per day for two weeks in both trials, with one continuing for 2 more weeks with 30 mg per day. One trial¹⁴⁶ measured the global effect and had mixed results. Both studies measured behavior/mood: one¹⁴⁵ showing significant change and the other¹⁴⁶ showing no significant change.

Divalproex sodium (Evidence Tables 117, 125, 126, 160). Divalproex sodium was evaluated in 229 subjects with mixed populations of VaD and AD who were treated for 6 weeks with increasing dosages until 20 mg per kg daily¹⁴⁷ or until side effects appeared.¹⁴⁸ These trials were both industry-supported or funded and included 56 or 173 institutionalized patients with probable or possible disease. Both trials showed no significant change in cognition and behavior/mood, while only one study¹⁴⁸ measured quality of life/ADL and found no significant difference. Both trials did a global assessment; one study found no significant difference,¹⁴⁸ and the other¹⁴⁷ found a significant change in favor of placebo.

Fluoxetine (Evidence Tables 117, 140, 141, 160). Fluoxetine was studied in a total of 56 AD patients using 3 or 20 mg per day¹²⁸ or a titration from 10 to 40 mg per day¹⁴⁹ for 6 weeks. All patients included in one study¹⁴⁹ also had major or minor depression. One study¹²⁸ was not industry-funded, and the other¹⁴⁹ did not indicate the funding source. Overall, the two studies

evaluated general cognition, behavior/mood, quality of life/ADL, and caregiver burden; no significant differences between the drug and placebo were found.

Loxapine (Evidence Tables 117, 147, 159, 160). Loxapine was evaluated in two trials^{150,126} from 1982 and included a total of 124 patients with MID and PDD. One trial¹²⁶ reported moderate to severe disease. The mean age in the other trial¹⁵⁰ was 83.0 years compared to 72.7 years in the trial with severe patients. Both studies were partially funded by industry and lasted 8 or 10 weeks. Only two domains were evaluated: global assessment and behavior /mood. No significant difference was shown in one trial,¹⁵⁰ while the other trial¹²⁶ showed a significant difference for behavior/mood.

Olanzapine (Evidence Tables 117, 135, 146, 160). Olanzapine was evaluated by two industry-funded trials in a total of 478 institutionalized patients with AD, VaD, and mixed dementia. One study¹⁵¹ used 10 or 15 mg per day for 6 weeks and the other¹⁵² used 12.5 mg maximum for one day. Both studies showed no significant change in general cognition. Both showed a significant change in measures of behavior/mood. One study¹⁵² evaluated global assessment and found no significant differences.

Phosphatidylserine (Evidence Tables 117, 136, 137, 160). Two industry-funded trials studied a total of 193 patients with AD or PDD. One study¹⁵³ included institutionalized patients with mild to severe AD and a mean age of 62.1 years, and the other¹⁵⁴ included community patients with mild to moderate AD or PDD and a mean age of 71.0 years. Both studies did subgroup analysis based on severity of illness. The study of institutionalized patients¹⁵³ found significant change in the domain of general cognition and global assessment. The study with community patients found significant change in a global assessment but no significant change in a measure of quality of life/ADL.

Risperidone (Evidence Tables 117, 142, 144, 160). Two studies evaluated risperidone for 12 weeks in 625 AD, VaD, or mixed dementia patients with moderate to severe disease¹⁵⁵ and in 344 PDD, VaD, or mixed dementia patients with severe disease.¹²⁷ The studies were industry-funded or supported, and both did subgroup analysis: one by disease and the other by gender, age, race, and diagnosis. Both trials showed a significant change in a global assessment. One study¹⁵⁵ found a significant change in behavior/mood, and the other study¹²⁷ had mixed results for that domain. There was no significant change in cognition or quality of life/ADL according to one of the trials.¹²⁷

Sertraline (Evidence Tables 117, 138, 139, 160). Sertraline was evaluated in two studies: one trial¹⁵⁶ for 8 weeks in 31 late-stage institutionalized AD patients with major depression (mean age 89.0 years), and the other trial¹⁵⁷ for 13 weeks in a community sample of 22 patients with mild to moderate AD and depression (mean age 77.0 years). Both studies found no significant differences in cognition. The trial¹⁵⁶ in subjects with severe disease found no significant difference in behavior/mood; the second trial¹⁵⁷ had mixed results for this same domain. The study in patients with mild to moderate disease showed significant change for a global assessment and no significance for quality of life/ADL.

Non-cholinergic neurotransmitter/neuropeptide modifying interventions (NCNMA).

Fifteen drugs in this drug grouping were compared to placebo in only one included trial. Eight of these trials showed a significant difference from placebo (See Evidence Tables 117, 120, 128, 129, 133, 134, 143, 145, 149, 160 at <http://www.ahrq.gov/clinic/epcindex.htm>): Alaproclate¹⁵⁸ for 4 weeks in 43 institutionalized patients with mild to severe PDD, MID and mixed dementia was better for quality of life/ADL. Imipramine¹⁵⁹ for 8 weeks in a community sample of 61 PDD and AD patients with depression was better for global assessment. Lisuride¹⁶⁰ for 8 weeks in 22 patients with mild to moderately severe AD was better for cognition. Minaprine¹⁶¹ for 12 weeks in an institutionalized sample of MID or SDAT patients showed mixed results for behavior. Moclobemide¹⁶² for 6 weeks in 511 patients with mild to moderate AD who were from both the community and institutions was better for cognition and behavior/mood. Naftidrofuryl¹⁶³ for 6 months in 378 patients with mild to severe VaD or mixed dementia was better for cognition and global assessment. Tiapride¹²⁴ for 3 weeks in 306 institutionalized AD patients with aggressiveness or irritability was better for behavior/mood. Trazodone¹²⁵ for 16 weeks in 149 AD patients from the community was better for quality of life/ADL. Xanomeline¹⁶⁴ in 343 community AD patients for 6 months was better for cognition, global assessment, and quality of life/ADL.

Seven trials (See Evidence Tables 119, 127, 130, 131, 132, 146, 147, 148, 160 at <http://www.ahrq.gov/clinic/epcindex.htm>) found no significant differences from placebo when evaluating perphenazine,¹⁴⁵ thioridazine,¹⁵⁰ fluvoxamine,¹⁶⁵ lorazepam,¹⁵² LU25,¹⁶⁶ maprotiline,¹⁶⁷ and minaprine¹⁶¹.

Results of other agents

A total of 72 studies representing 46 different other agents were eligible for this review and these can be seen in Table 4. Twenty-two of these interventions were evaluated in a single trial and only briefly summarized in this chapter; greater detail is provided in Evidence Tables 161 through 249.

Table 4. List of Other pharmacological agents and the number of studies vs. placebo for each of these. Asterisk (*) indicates report of a drug vs. drug trial [comparator drug(s) in brackets].

Drug	Number of studies vs. placebo	Drug	Number of studies vs. placebo
Aniracetam	1	Misoprostol *[Diclofenac]	0*
5'-MTHF *[Trazodone]	0*	Monosialotetrahexosylganglioside (GM-1)	1
Amitriptyline *[Fluoxetine]	0*	N-Acetylcysteine	1
Ateroid	1	Nimesulide	1
Buflomedil	1	Nimodipine	2
Cerebrolysin	6	Nizatidine	1
Choro-San	1	Nootropic	1
Choto-San	1	ORG 2766	2

Table 4. List of Other pharmacological agents and the number of studies vs. placebo for each of these. Asterisk (*) indicates report of a drug vs. drug trial [comparator drug(s) in brackets] (continued).

Drug	Number of studies vs. placebo	Drug	Number of studies vs. placebo
Citicoline *[Posatirolin] *[Sulphomucopolysaccharides]	0**	Oxiracetam	5
Cyclandelate	2	Pentoxifylline *[Sulodexide]	3*
Denbufylline	1	Piracetam	1
Desferrioxamine	1	Prednisone	1
Diclofenac	1	Propentofylline	4
Ergokryptine (CMB 36-733)	1	Pyritinol *[Hydergine]	0*
Ergokryptine (Dek)	1	Silymarin + Taurine *[Placebo + Taurine]	0*
Estrogens	5	Simvastatin	1
Ginkgo Biloba	3	Sulphomucopolysaccharides *[Citicoline]	0*
Glycosaminoglycan Polysulfate	1	Sulodexide *[Pentoxifylline]	0*
Guanfacine	1	Thiamine	1
Hydergine *[Pyritinol]	1*	Vasopressin (DDAVP)	1
Hydroxychloroquine	1	Vincamine	1
Idebenone *[Taurine]	4*	Vitamin E *[Donepezil] *[Selegiline]	1**
Indomethacin	1	Xantinolnicotinate	1

Cerebrolysin. See Evidence Tables 161 through 168 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Six included studies^{168,169,170,171,172,173} compared cerebrolysin to placebo. One report was from 1994,¹⁷² one from 1999,¹⁷¹ two from 2000,^{169,170} one from 2001,¹⁶⁸ and one from 2002.¹⁷³

Design/methodology. The sample size in the studies ranged from 53¹⁶⁹ to 192¹⁷³ with a total of 819 subjects. The quality of studies varied from scoring 6¹⁷¹ to 8^{168,173} points out of a possible 8 points. One study¹⁷² did not indicate the source of funding, one trial had non-industry funding,¹⁷³ and the four remaining trials were funded by industry.^{168,169,170,171}

Populations. All but one of the six studies included AD patients; one study¹⁷¹ evaluated patients who had mild to moderate VaD. Mean ages of the subjects in the studies ranged from 69.7¹⁷¹ to 74.1 years.¹⁷³ The proportion of males in the trials varied from 34 - 69%, and only one trial¹⁷³ specified the proportion of Caucasians.

Intervention. All of the studies used the same dose of cerebrolysin, 30 ml per day, for 5 days per week. One trial¹⁷² was for 28 days, four studies^{169,170,171} lasted 4 weeks, one trial¹⁶⁸ lasted 16 weeks, and one trial¹⁷³ lasted 24 weeks.

Primary outcomes. Most studies evaluated general cognitive function and three trials^{168,169,173} used the ADAS-cog. Baseline MMSE was reported in a single trial¹⁷³ with a score of 21. All studies evaluated global function, and at least two studies evaluated one outcome in each of the remaining domains with the exception of caregiver burden.

Analysis. All but one¹⁷³ of the studies used ITT analysis.

Results and interpretation. See Summary Table 16. Four of the five studies that evaluated general cognitive function showed significant differences.^{169,168,170,171} Figure 19 displays the pooled estimate for those studies for which the appropriate data could be extracted for the ADAS-cog. Although a summary estimate was calculated, the test for heterogeneity was positive, suggesting the estimate should be interpreted with caution. Moreover, the overall estimate was not significant. One study¹⁷³ showed no significant difference in MMSE or ADAS-cog. This was the only study to report non-industry funding and coincidentally the only study to use OC population analysis. Three studies used specific cognitive measures, two of which^{172,171} found significant differences and one of which¹⁷⁰ showed mixed results.

All trials evaluated global assessment, and all except one trial¹⁷¹ reported a significant difference. Figure 20 shows the pooled estimate for the CGI. The pooled estimate was calculated, the test for heterogeneity was positive, suggesting the estimate should be interpreted with caution. However, the overall estimate is significant. Three trials reported results for a measure of behavior/mood, one showing significant effects¹⁶⁸ and the other two¹⁷¹ showing none. All trials carried out evaluations of quality of life/ADL measures; one did not report the effect,¹⁶⁸ one had mixed results¹⁷⁰ and the other four showed no significant difference.^{169,173,172,171} No study measured caregiver burden.

Figure 19. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing cerebrolysin versus placebo.





Figure 20. Odd Ratio (OR) from the Random Effects Model (Random) for the CGI comparing cerebrolysin versus placebo.



Two^{169,172} of the six trials scored 5 out of 5 on our quality scale for rating adverse events, yet they did not report any adverse events. Two studies^{173,168} scored 4, and the other two trials scored 3¹⁷¹ and 2.¹⁷⁰ All the studies with scores equals to 4 or less tested for statistical differences in adverse events between placebo and treatment groups. Withdrawals due to adverse events were not reported in one study,¹⁷⁰ and were 1% in two studies.^{173,168} Three studies^{169,172,171} reported no withdrawals. A significant difference between treatment and control group was reported in one study¹⁷³ for weight change, anxiety, and headache. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 10 - 24%, all doses = 3 - 21%), 2) dizziness (placebo = 0 - 12, all doses = 1 - 8%), and 3) agitation (placebo = 1%, all doses = 0%), and none reported diarrhea or eating disorder as an adverse event.

Estrogens. See Evidence Tables 169 through 175 at <http://www.ahrq.gov/clinic/epcindex.htm>. *Number of studies.* Five studies^{174,175,176,177,178} evaluated estrogens for dementia patients: one published in 1999,¹⁷⁸ three in 2000,^{176,177,175} and one in 2001.¹⁷⁴ None compared estrogens to another drug.

Design/methodology. The number of subjects included in the studies ranged from 15¹⁷⁸ to 120 subjects¹⁷⁷ with a total of 247 patients. Quality of the studies ranged from 5¹⁷⁴ to 8¹⁷⁸ points out of a possible 8 points. All studies were partially or fully funded by industry.

Populations. Four of the studies^{176,177,175,174} included patients with mild to moderate AD, and one study¹⁷⁸ included moderate to severe dementia patients who were all institutionalized. Only one of the studies¹⁷⁸ included male subjects. Mean age ranged from 71.8¹⁷⁵ to 80.0 years¹⁷⁴ in the AD studies, and it was 83.8 in the dementia study.¹⁷⁸

Intervention. One of the studies with AD patients used 0.10 mg per day¹⁷⁴ for 8 weeks, and the others used 1.25 mg per day for 12 weeks,¹⁷⁵ 16 weeks,¹⁷⁶ and 52 weeks.¹⁷⁷ The study¹⁷⁸ including subjects with severe disease used 2.5 mg per day for 4 weeks.

Primary outcomes. At least one study evaluated each of the included domains with the exception of caregiver burden.

Analysis. Two of the studies^{177,175} performed ITT analysis and the other three used OC analysis.

Results and interpretation. See Summary Table 17. Three^{176,177,175} trials evaluated general cognitive function and all showed non-significant findings; two trials^{176,177} lacked sufficient power (PW = 0.10, PW = 0.44) for the ADAS-cog. Attempts were made to combine the ADAS-cog, but the random-effects model was positive for heterogeneity and the overall effect was not significant. Two trials^{174,177} evaluated specific cognitive function, and only one of these, using the Stroop Color Word Interference Test (SCWIT) measure, showed significant differences.¹⁷⁴ Global assessment was undertaken in all trials and found to be not significant in any of these trials. For those trials where power could be estimated,^{176,177,175} there was insufficient power for the CGIC, CDR, and CIBIC+ outcomes. For the outcomes of behavior/mood and quality of life/ADL, none of the trials reported significant differences; power for the outcomes used in the trials could not be estimated. Overall, the evidence that estrogen affected general and specific cognitive function, global assessment, behavior/mood, and quality of life/ADL is inconclusive. There were not enough similar outcomes reported to complete a pooled analysis for estrogens.

One¹⁷⁸ of the five trials scored 5 out of 5 on our quality scale for rating adverse events, and surprisingly, this same trial did not report any adverse event. Two trials^{176,177} scored 3; one trial¹⁷⁵ scored 2, and one¹⁷⁴ scored 1. This latter study reported adverse events, but did not test for significant differences between groups. Withdrawal rates due to adverse events ranged from 0 - 5% for placebo and 0 -14% for the treatment group. The most frequently reported adverse event was vaginal bleeding,^{175,177,176} and a single trial¹⁷⁵ reported a significant difference between placebo and treatment group for vaginal bleeding. It was not clear from the descriptions provided in the study if they had ascertained whether vaginal bleeding was present prior to the trial commencement. Nausea was the single a priori symptom of interest that was reported and by a single trial; frequencies varied from 0% for the placebo group and 4% for the treatment group.

Ginkgo biloba. See Evidence Tables 176 through 180 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Three studies^{179,180,181} evaluating Ginkgo biloba were eligible to be included in this review. All of the studies compared the drug to placebo only. One of the studies was reported in 1996¹⁸¹ and two were reported in 1997.^{179,180}

Design/methodology. The studies included evaluated 20 subjects,¹⁸⁰ 216 subjects,¹⁸¹ and 327 subjects¹⁷⁹ (totaling 563 subjects). Two of the reports^{179,181} scored 8 quality points out of a possible 8 points, and the other¹⁸⁰ earned 6 points. One study did not indicate the funding source,¹⁸⁰ and the other two had industry funding.

Populations. All of the studies included a mix of dementia diagnoses as follows: 1) mild to moderately severe AD and MID,¹⁷⁹ 2) mild to moderate DAT and PDD¹⁸⁰, and 3) mild to moderate DAT and MID¹⁸¹ in community dwelling patients. Two of the studies reported subgroup analysis, one comparing diagnoses and comparing effects based on baseline MMSE score¹⁷⁹ and the other based on diagnosis.¹⁸¹ The patients in these trials had mean ages of 64.6,¹⁸⁰ 69.0,¹⁷⁹ and 69.6 years.¹⁸¹

Intervention. Two of the trials gave 240 mg per day for 3 months¹⁸⁰ and 6 months,¹⁸¹ and the other trial¹⁷⁹ gave 40 mg three times daily for 12 months.

Primary outcomes. None of the studies reported on quality of life/ADL or caregiver burden.

Analysis. All of the trials used an ITT analysis.

Results and interpretation. See Summary Table 18. Two of the three trials evaluated general cognitive function, and only one of these showed significant results.¹⁷⁹ Two studies^{181,180} showed positive results with specific cognitive function. The results for global assessment are inconsistent as only one trial had positive findings,¹⁸¹ one study had mixed results,¹⁷⁹ and one trial¹⁸⁰ showed non-significant results. This latter study had a very small sample size and lacked sufficient power for some outcomes. Only one trial¹⁸¹ reported behavior/mood outcomes and found no difference between groups. None of the studies evaluated quality of life/ADL and caregiver burden. There were not enough similar outcomes reported to complete a pooled analysis for ginkgo biloba.

One¹⁸⁰ of the three trials scored 5 out of 5 on our quality scale for rating adverse events. One study¹⁸¹ scored 4, and one trial¹⁷⁹ scored 3. Two studies^{181,180} had no withdrawals due to adverse events, and one trial¹⁷⁹ had a withdrawal rate of 6% for both placebo and treatment groups. Two studies^{179,180} did not report any adverse event. One study¹⁸¹ reported a statistically significant difference between the treatment and the placebo group for skin disorders. The same study reported gastrointestinal and headache adverse effects, but did not test for statistical differences between the placebo and the treatment group. None of the trials reported any of the a priori symptoms of interest.

Idebenone. See Evidence Tables 181 through 187 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Four studies^{182,183,184,185} were included in this review that evaluated idebenone versus placebo, and one study²⁶ compared idebenone to tacrine but not to placebo. The placebo trials were published in 1992,¹⁸⁴ 1994,¹⁸² 1997,¹⁸⁵ and 1998,¹⁸³ the tacrine trial was published in 2002 by the same author as a previous placebo trial.¹⁸³

Design/methodology. Sample sizes in the studies ranged from 92¹⁸² to 450 subjects¹⁸³ with a total of 950 patients in the placebo-controlled studies. The study comparing idebenone with tacrine included 203 subjects, but a large number withdrew, and only 44 completed the trial. One of the trials¹⁸⁵ earned 5 points out of a possible 8 points on the quality scale, two of the trials earned 6 points,^{182,183} and one earned 7 points.¹⁸⁴ None of the placebo studies reported their funding source. The tacrine study earned 7 points on the quality scale and was partially funded by industry.

Populations. The studies included patients with AD, MID, PDD, and DAT. Two of the trials^{182,183} reported that the subjects had mild to moderately severe disease and the remainder reported mild to moderate disease. Two of the studies reported subgroup analysis based on disease severity.^{185,183} Mean ages in the studies ranged from 69.9¹⁸³ to 73.6 years.¹⁸⁴

Intervention. Dosing schemes were 30 or 90 mg per day for 6 months,¹⁸⁵ 30 mg three times per day for 3 months,¹⁸² 45 mg twice daily for 4 months,¹⁸⁴ and 120 mg three times per day for 12 months.¹⁸³ The tacrine trial used 360 mg per day for 14 months.

Primary outcomes. Caregiver burden was the only domain in this review that was not evaluated in at least one of the studies.

Analysis. Two of the studies^{183,185} used ITT analysis while the other two used OC analysis. The tacrine trial used ITT analysis.

Results and interpretation. See Summary Table 19. Three trials^{183,184,185} found significant differences for general cognitive function. Two trials^{185,183} used the ADAS-cog and reported changes that varied from -4.5 to -4.9 for placebo versus -4.4 to -8.8 for the treatment group. The doses varied in these two trials from 90 to 360 mg per day. A single trial¹⁸² evaluated specific cognitive function and showed inconsistent findings. Three trials^{182,183,185} evaluated global assessment and all found significant differences relative to placebo. A single trial¹⁸⁵ evaluated behavior/mood and was statistically significant, even though it was a secondary outcome. Two trials^{183,184} evaluated quality of life/ADL and were both statistically significant. No study evaluated caregiver burden. These findings suggest some evidence of benefit for general cognitive function, global assessment, and quality of life/ADL. There were not enough similar outcomes reported to complete a pooled analysis for idebenone.

Quality scores for reporting adverse events varied from 1 to 5. Rates of withdrawal due to adverse events varied from 0 - 5% for the placebo group and 0 - 5% in the treatment group; a single trial¹⁸³ did not report withdrawal rates. Two trials^{183,185} tested for statistical differences between groups and found no differences. Although no clear pattern emerges, three studies identified at least one balance-related adverse event across studies. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 2%, all doses = 2 - 11%), 2) dizziness (placebo = not reported, all doses = 2%), and 3) not reported for diarrhea, agitation, or eating disorder as an adverse event.

Oxiracetam. See Evidence Tables 188 through 194 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Five trials^{186,187,188,189,190} included in this review evaluated oxiracetam versus placebo. The studies were published in 1988,¹⁸⁸ 1989,¹⁸⁷ and 1992.^{190,189,186}

Design/methodology. A total of 554 patients were included in the studies, ranging from 30 patients¹⁸⁸ to 289 patients.¹⁸⁷ Four of the studies earned 6 points out of a possible 8 points on the quality scale, and the other study¹⁸⁹ earned 4 points. Two of the studies^{189,187} did not report the source of their funding, and the other three trials had partial industry funding.

Populations. The trials included a mixture of diagnoses, including AD, PDD, mixed dementia, and MID, and none of the studies reported severe disease. One of the studies¹⁸⁷ performed subgroup analysis based on diagnosis, comparing MID to PDD. The mean age of the subjects included in the trials ranged from 62.0¹⁸⁸ to 73.8 years.¹⁸⁹

Intervention. All of the trials used a dose of 800 mg twice daily, for a duration of 12^{186,187} to 26 weeks.¹⁸⁹

Primary outcomes. At least one trial evaluating oxiracetam evaluated one of the outcome domains examined in this review with the exception of caregiver burden. A single trial¹⁸⁹ reported baseline MMSE at 22 for both placebo and treatment groups.

Analysis. None of the trials used ITT analysis.

Results and interpretation. See Summary Table 20. Three trials^{187,190,189} out of the five studies tested for outcomes on general cognitive function. Only two of these trials^{187,190} reported the findings, which were both significant, even though the NMIC and MMSE were used to measure this attribute. Three trials^{186,188,190} evaluated specific cognitive function and showed mixed results. A single large trial¹⁸⁷ evaluated global assessment and found significant differences between groups using the Blessed Dementia Scale (Italian version). Three trials^{187,188,190} evaluated behavior/mood with the IPSC-E, and of these, a single trial¹⁹⁰ did not show significant differences. One trial¹⁸⁹ reported on Beck Depression Inventory (BDI) but did not show statistical comparisons. Similarly, three trials^{186,189,190} evaluated quality of life/ADL, and a single trial¹⁸⁹ showed no significant findings. No study evaluated caregiver burden. There were not enough similar outcomes reported to complete a pooled analysis for oxiracetam.

The quality scores for reporting adverse events varied from 2 to 5. The proportion of withdrawals due to adverse events varied from 0 – 9% for the placebo group and 0 – 6% for the treatment group. No clear pattern for adverse events is evident, but three of the five studies reported gastrointestinal related problems, primarily associated with abdominal pain. Although, only single trials evaluated the range of frequencies of the a priori symptoms of interest are as follows: 1) dizziness (placebo = not reported, all doses = 11%), and 2) agitation (placebo = 1%, all doses = not reported); no trial reported nausea, eating disorder, or diarrhea as an adverse event.

Pentoxifylline. See Evidence Tables 195 through 200 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Three trials^{191,192,193} in this review evaluated pentoxifylline versus placebo. One trial,¹⁹⁴ published in 1997, compared pentoxifylline to sulodexide rather than placebo. The placebo trials were published in 1987,¹⁹³ 1992,¹⁹² and 1996.¹⁹¹

Design/methodology. The studies included 36 patients,¹⁹³ 64 patients¹⁹² and 289 patients.¹⁹¹ The sulodexide trial included 93 patients. All placebo trials had 6 points out of a possible 8 points on the quality scale and had partial or full industry funding. The sulodexide trial earned 5 points on the quality scale and did not report the source of funding.

Populations. The three placebo-controlled trials included patients with mild to moderate MID, and one trial¹⁹³ also included PDD patients. The sulodexide trial had only patients with mild to moderate VaD. Subgroup analysis was performed in two trials, looking at MID versus PDD

diagnosis¹⁹³ and grouping by vascular change versus discrete stroke.¹⁹² The mean age of the studies ranged from 69.7¹⁹¹ to 77.0 years.¹⁹³

Intervention. All of the studies gave 1200 mg per day of pentoxifylline; one study gave the drug once a day for 9 months,¹⁹¹ one study gave 400 mg three times per day for 9 months,¹⁹² and one gave 400 mg three times per day for 3 months.¹⁹³ The sulodexide study gave the drug once a day for 6 months.

Primary outcomes. At least one trial evaluated one of the outcome domains examined in this review with the exception of caregiver burden.

Analysis. A single¹⁹¹ trial used an ITT analysis.

Results and interpretation. See Summary Table 21. All three placebo trials showed non-significant findings for any primary outcome evaluated on all subjects in the study. It should be noted that two of these trials^{192,193} had very small sample sizes (n = 38, n = 28) that were evaluated in the OC analyses; this suggests that the trials lacked sufficient power to evaluate multiple outcomes. Knezevic et al.¹⁹¹ had a large sample size (n = 289) and employed an ITT analysis; all primary outcomes evaluated were not significant. The evidence for all outcomes considered in this review are inconclusive for pentoxifylline. There were not enough similar outcomes reported to complete a pooled analysis for pentoxifylline.

The quality scores for reporting adverse events were generally low, varying from 1 to 3. Withdrawal rates due to adverse events varied from 0 – 25% in the placebo group and 0 – 22% in the treatment group. The two studies that reported adverse events indicated the presence of gastrointestinal disturbances, including abdominal pain or nausea and vomiting (placebo = 7% and all doses = 14%). None of the trials reported dizziness, agitation, eating disorder or diarrhea.

Propentofylline. See Evidence Tables 201 through 206 at <http://www.ahrq.gov/clinic/epcindex.htm>.

Number of studies. Four studies^{195,196,197,198} in this review evaluated propentofylline versus placebo. The first trial was published in 1990.¹⁹⁸ Two, by the same author, were published in 1996¹⁹⁶ and 1998.¹⁹⁷ One was published in 1997.¹⁹⁵

Design/methodology. The number of subjects in the studies ranged from 30 subjects^{197,196} to 260 subjects,¹⁹⁵ with a total of 510 subjects. Three of the studies^{197,195,196} earned 5 points out of a possible 8 points on the quality scale, and the other study¹⁹⁸ earned 6 points. Only one study indicated the source of funding for the trial,¹⁹⁶ and it was industry-supported.

Populations. The trials included subjects with mild to moderate AD only,¹⁹⁷ VaD only,¹⁹⁶ mild dementia only,¹⁹⁸ and mild to moderate combined AD and VaD.¹⁹⁵ Two trials presented subgroup analysis: one for AD versus VaD,¹⁹⁵ and one based on MMSE baseline score. The mean age in the studies ranged from 64.8¹⁹⁷ to 72.4 years.¹⁹⁵

Intervention. All four studies gave 300 mg three times a day for 3 months with the exception of one trial¹⁹⁵ which had a duration of 12 months.

Primary outcomes. At least one trial evaluated one outcome in each of the domains examined in this review with the exception of caregiver burden. Baseline MMSE was reported in two trials^{197,198} and varied from 20 and 21 for both placebo and treatment groups.

Analysis. One of the studies¹⁹⁵ used an ITT analysis.

Results and interpretation. See Summary Table 22. All four trials evaluated general cognitive function and the pooled estimate can be seen in Figure 21. Two of the trials^{197,196} had small sample sizes and these trials had the widest confidence intervals. The test for heterogeneity did not exceed our threshold of 0.10 for significance; the overall summary effect was significant. Figure 22 shows the pooled estimate for the DSST, a measure of specific cognitive function; this pooled estimate should be interpreted with caution as the test for heterogeneity was significant and the overall effect was not significant. Thus, there is some evidence of benefit for general cognitive function, and inconclusive evidence for specific cognitive function as measured by the DSST. Similarly, there is inconclusive evidence for global assessment. Behavior/mood outcomes (using the NAB) were evaluated by a single trial¹⁹⁵ and shown to be significantly different; this same trial evaluated quality of life/ADL (using the NAA) and showed no significant difference.

The quality scores for reporting adverse events varied from 1 to 4. The percentage of withdrawals varied from 0 – 13% for the placebo group and 0 – 12% for the treatment group. None of the trials tested for differences between groups. Three of the trials^{195,197,198} reported gastrointestinal events that included abdominal pain, constipation, and nausea and vomiting (placebo = 2%, all doses = 7%). Dizziness (placebo = 3 - 5%, all doses = 1 - 6%) was the only other a priori symptom of interest.

Figure 21. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the MMSE change score comparing propentofylline versus placebo.





Figure 22. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the DSST change score comparing propentofylline versus placebo.



Various other agents. See Evidence Tables 207 through 249 at <http://www.ahrq.gov/clinic/epcindex.htm>. See Summary Table 23.

Interventions with two studies included for review. Six other agents were compared with placebo in only two included trials:

Choto-san (Evidence Tables 207, 213, 214, 249) Two studies compared Choto-San with placebo in patients with VaD. Both studies included all Asian subjects and co-morbid disorders were present in both studies. Each study lasted 12 week, with doses of drug at 7.5 g three times a day¹⁹⁹ and 2.5 g three times a day.²⁰⁰ The studies were published in 1994²⁰⁰ and 1997¹⁹⁹ and both had a quality score of 5 out of 8 points. Both studies measured global assessment and behavior/mood and disagreed on both results. One study¹⁹⁹ showed a significant change in global assessment and no significant difference in behavior/mood while the other²⁰⁰ showed mixed results for global assessment and a significant difference for behavior/mood.

Cyclandelate (Evidence Tables 207, 215, 216, 249). Cyclandelate was evaluated in two studies: one²⁰¹ in 139 AD patients and another²⁰² in 196 PDD, VaD, and mixed dementia patients. The mixed population study²⁰² had a subgroup analysis based on the MMSE, ADAS-cog, and treatment center. The AD patients received 400 mg four times per day for 16 weeks and the mixed population study used 800 mg twice a day for 24 weeks. Only caregiver burden was not evaluated by either study and only global assessment was evaluated by both studies. The study with AD patients²⁰¹ showed a significant change in global assessment and in behavior/mood and mixed results in the specific cognitive function measures. The study with the mixed population²⁰² showed no significant difference in global assessment or general cognitive measures or quality of life/ADL or function.

Ergokryptine (Evidence Tables 207, 212, 218, 249). Two trials, which were not similar, evaluated ergokryptine. One trial²⁰³ that did not indicate the source of funding included 125 PDD patients and treated them with a dose titrated up to 2 mg per day for 8 weeks. The other trial,²⁰⁴ which was industry funded, treated 215 AD patients with a dose titrated up to 20 mg twice a day for one year. Neither study reported caregiver burden, but both reported general cognitive function and global assessment. They differed on the results of both of those domains: one study²⁰⁴ showed significant change in cognition and mixed results in global assessment, and the other study²⁰³ showing no significant difference and significant change, respectively. Significant change was demonstrated in specific cognitive measures in the study with AD patients.²⁰⁴ Mixed

results were shown for behavior/mood outcomes, and no significant difference was seen for quality of life/ADL in the study with PDD patients.²⁰³

Hydroxychloroquine/Nimesulide (Evidence Tables 207, 226, 229, 249). No significant difference from placebo was seen in either of two studies^{205,206} for cognition, behavior/mood, or quality of life/ADL. Global assessment was evaluated in one of the studies²⁰⁶ and there was no significant difference found. One trial²⁰⁵ included minimal to mild AD patients and the other²⁰⁶ included mild to moderate AD patients. One study²⁰⁵ treated patients for 18 months with a dose of 400 or 200 mg per day based on weight. The other study treated for 3 months with 100 mg twice a day. Both studies had non-industry funding, and one study²⁰⁵ also had industry support.

Nimodipine (Evidence Tables 207, 230, 231, 249). Nimodipine was evaluated in one study²⁰⁷ with 259 mild to moderate MID patients receiving 30 mg twice a day for 26 weeks. The other study²⁰⁸ evaluated 178 patients with mild to moderately severe MID and PDD receiving 90 mg per day for 12 weeks. The trials received industry funding or support and were published ten years apart, in 1990²⁰⁸ and in 2000.²⁰⁷ The trial using ITT analysis²⁰⁷ showed no significant difference in the domains of general cognitive function measures, specific cognitive measures, global assessment, and quality of life/ADL. The trial using OC²⁰⁸ analysis found significant differences in the domains of general cognitive function measures, specific cognitive measures, global assessment, and behavior/mood.

ORG2766 (ACTH peptides) (Evidence Tables 207, 233, 234, 249) Org2766 versus placebo was reported for a total of 233 patients with AD or primary degenerative senile dementia (PDS) in reports from 1985²⁰⁹ and 1986.^{210,209} One study²⁰⁹ was industry-supported and used 20 mg twice daily for 6 months, and the other was non-industry-funded²¹⁰ and used 80 mg twice daily for 1 month. One study²¹⁰ found a statistical difference between drug and placebo in the domains of specific cognitive function measures and in global assessment, and the other study²⁰⁹ found no significant difference in the domains they evaluated: global assessment and behavior.

Interventions with only one trial included for review. Twenty-two drugs in this drug grouping were compared to placebo in only one included trial. Eleven of these trials showed a significant difference from placebo and are summarized briefly here. See Evidence Tables 207 to 240, 244, 248 and 249 for greater detail concerning the trials.

Drugs compared to placebo in one trial only (Evidence Tables 207, 208, 209, 210, 217, 220, 222, 224, 237, 238, 240, 248, 249) Aniracetam was better for cognition and global assessment in 109 community patients for 6 months, Ateroid²¹¹ in 155 PDD, MID or SDAT patients for 12 weeks was better for general cognition. Desferrioxamine²¹² in 48 probable AD patients for 2 years was better for behavior/mood. Glycosaminoglycan polysulfate²¹³ in 155 moderate to severe PDD or MID patients for 12 weeks was better for behavior/mood. Guanfacine²¹⁴ in 29 mild to moderate AD or PDD patients for 13 weeks was better for specific cognitive measures and global assessment. Nootropic agent BMY²¹⁵ in 69 mild to moderate AD patients for 12 weeks was better for general cognitive measures. Thiamine²¹⁶ in 15 mild to moderate AD patients for 12 months was better for general and specific cognitive function measures. Vincamine²¹⁷ for 12 weeks in 152 institutionalized patients with mild to moderate PDD or VaD was better for global assessment. Vitamin E¹³⁵ in 341 moderate AD patients for 2 years was better for delaying

institutionalization. Deamino-D-arginine-vasopressin²¹⁸ in 14 PDD patients was better for behavior and had mixed results for global assessment. Xantinolnicotinate²¹⁹ in 313 mild to moderate AD or MID patients for 12 weeks was better for specific cognitive function measures and global assessment.

Twelve trials (Evidence Tables 207, 211, 219, 221, 223, 225, 227, 228, 232, 235, 236, 239, 244, 249) found no significant differences from placebo or mixed results when evaluating buflomedil,²²⁰ citicoline,¹⁰¹ denbufylline,²²¹ diclofenac and misoprostol,²²² hydergine,²²³ indomethacin,²²⁴ monosialotetrahexosylgan,²²⁵ N-acetylcysteine,²²⁶ nizatidine,²²⁷ piracetam,²²⁸ prednisone,²²⁹ and simvastatin.²³⁰

Question 2: Does pharmacotherapy delay cognitive deterioration or delay disease onset of dementia syndromes?

Delay of Onset of Dementia

The concept of “delay onset” was operationalized to imply delay in conversion from a cognitive disturbance state, classified as MCI, CLOND or CIND, to a true dementia state. No studies with this population met the final eligibility criteria, although four trials^{231,232,233,234} advanced to the full text screening stage. The lack of studies eligible for evaluation in this systematic review points to a gap in the literature for pharmacological interventions (attempting to demonstrate a delay in disease onset) in MCI-type populations.

Delay of Progression

In general, very few studies evaluated patients who were classified as “severe”. Five studies^{126,208,129,178,132} had moderate to severe groups of dementia patients, and only one trial reported all three levels¹⁶³ of the disease spectrum. The interventions evaluated in these trials were estrogen, haloperidol, glycosaminoglycan polysulfate, memantine, and naftidrofuryl. This suggests that there is a bias in the trials eligible in this systematic review towards evaluating mild to moderate disease; this in turn reflects the underlying assumption that the less severe groups are most likely to benefit from drug trials. Since so few studies have evaluated the more severe groups, this assumption may require some empirical justification. Therefore, delay in progression has not been considered in severe patients.

The selected studies used two approaches for showing “delaying disease progression”. The first method for evaluating the potential for a drug to delay disease progression used longer-term follow-up; survival analyses (time to a relevant event) were then used to show differences between the two groups. The second design approach used withdrawal from treatment for a period and continued monitoring of the treatment and placebo groups (to demonstrate a deviation of the treatment group from the natural history as represented by the placebo group). Such designs have been termed withdrawal, active-extension, randomized withdrawal, randomized start, and staggered start.^{235,236,19,21} From our 186 included studies, we then further selected a subgroup of papers that had the potential to demonstrate delay in disease progression through the

use of one of these two designs. Therefore, any eligible trial that employed a survival analysis or a two-period approach, where the pharmacological agent was withdrawn during one of the periods, was selected for further evaluation to answer this question.

Survival Analyses

Two studies^{135,61} using survival analyses were identified. In a 2-year study¹³⁵ that compared placebo to three other groups (selegiline, selegiline with vitamin E, and vitamin E), time to the development of significant dementia milestones (death, institutionalization, loss of ability to perform ADL, or score on scale indicating severe dementia) was used as the time to event in the survival analysis. The results of this study showed that the vitamin E, selegiline, and combined groups were statistically different (i.e. declined less) from the placebo group in analyses that included baseline MMSE score as a covariate (not significant when excluded). The median survival was 230 days (vitamin E), 215 days (selegiline), and 145 days (combined group). Moreover, the vitamin E group showed a statistically significant difference for the endpoint of institutionalization, and the other treatment groups did not. There were no statistical differences between groups with respect to adverse events. Thus, the findings of this study suggest that selegiline and vitamin E may delay clinically important deterioration in patients with moderately severe AD; this delay varied from 20 to 32 weeks. The second study⁶¹ used survival analyses to evaluate the time to the development of severe functional impairments in a comparison of placebo and donepezil with a follow-up of 54 weeks. The results of the Kaplan-Meier analysis showed a mean number of days to significant functional decline of 252 days for placebo and 357 days for the donepezil group (mean difference of 100 days). The treatment group was 38% less likely to decline over a 1-year period. Both these studies demonstrated some delay in disease progress varying from 100 to 230 days for these three different pharmacological agents.

Staggered Withdrawal

Delay in disease progression can also be evaluated using a “time to return to baseline” following withdrawal of treatment. Similarly, staggering the start of the treatment parallels the staggered withdrawal and can be used to evaluate disease progression. In this design approach, the time to return to baseline is compared to the placebo group, which represents the natural course of the disease. Of the studies that were eligible for this research question used a classic withdrawal design (withdrawal in period II after the intervention was administered); none of these studies were able to maintain double blinding after the withdrawal of the intervention. Justification for the selection of the length of the washout or follow-up period was not consistently provided (which possibly reflects the lack of a priori aim to show delay in progression).

Tables 5 and 6 detail any study that attempted to withdraw the drug in the treatment group and then continue observations over time. All studies that reported outcomes after the drug trial endpoint subsequently interrupted protocol and switched to “open-label” circumstances. In open-label conditions, blinding was broken and greater proportions of patients withdrew from the study as the follow-up increased. From a methodological perspective, these data were considered to be biased and would not meet our review eligibility criteria. However, we

summarize in these Tables the same observations that were reported in all the studies eligible for this systematic review.

Table 5. Studies that withdrew the treatment agent but maintained at least single blinding.

Study	Drug	Schedule	Result
Ruether 2001	Cerebrolysin	4w drug + 8w washout + 4w drug + 12w washout	All patients: ADAS-noncog, maintained difference from placebo NAI returned to baseline Subgroup MMSE<20: ADAS Noncog, CGI, ADAS-cog, SKT maintained difference from placebo
Nyth 1990	Citalopram	4w drug + 8 w open drug + 4w new random drug	NR
Rogers 1996	Donepezil	12w drug + 2w SB PI washout	5 mg maintained effect, 3 mg no maintenance of effect for ADAS-cog (NS)
Rogers 1998b	Donepezil	24w drug + 6w SB and placebo washout	Return to placebo levels for ADAS-cog, MMSE, CIBIC (all NS)
Wilcock 2002	Memantine	2w SB and placebo + 28w drug + 2w SB placebo washout	NR
McKeith 2000	Rivastigmine	20w drug + 3w rest	Return to placebo levels for NPI and computerized cognitive assessment (NS)
Antuono 1995	Velnacrine	2w SB placebo + 24w drug + 6w SB placebo washout	Return to placebo levels for the ADAS-cog but SC for CGIC remained for washout
Bodick 1997	Xanomeline	24w drug + 4 w SB placebo	SC at week 24 with CNTB No differences vs. placebo at w4 of washout

In Table 5, single blinding was maintained in a placebo-controlled trial of cerebrolysin, which had an 8- and 12-week follow-up and showed continuing statistical differences after drug withdrawal. The remaining drug interventions listed in Table 5 suggest that the treatment provided predominately symptomatic relief lasting 2 to 6 weeks and then returning to placebo levels. Similarly, the pharmacological agents in Table 6 suggest that treatment provided only symptomatic relief.

Table 6. Studies that withdrew treatment and did not specify if blinding for washout or extension was maintained.

Study	Drug	Schedule	Result
Dehlin 1985	Alaproclate	2w placebo + 4w drug + 2w placebo	SC for GBS intellectual subscale at w4 of treatment No significant difference at w2 of washout
Cutler 1993	BMY 21,502	12w drug + 4w placebo washout	Treatment showed no significant change and follow-up showed no change
Amaducci 1988	Phosphatidylserine	3m drug + 21 m follow-up	SC remained for severe disease patients, not moderate
Raskind 1997	Metrifonate	26w drug + 8w follow-up	NR
Parnetti 1995	Posatirelin	90d IM + 30d follow-up placebo	"Maintained positive effect" but specific numbers not reported
Agid 1998	Rivastigmine	10w drug + 2w placebo washout	NR

Question 3: Are certain drugs, including alternative medicines (non-pharmaceutical), more effective than others?

From a methodological perspective, addressing the question of being “more effective” requires head to head comparisons of pharmacological interventions. If one intervention (Drug A) has been shown to be effective relative to placebo of a specified effect size, and a second intervention (Drug B) has been shown to be effective at a lower magnitude relative to placebo, it does not necessarily follow that Drug A is more effective than Drug B. Comparisons of the relative effectiveness of certain drugs can only be evaluated in the context of head to head evaluation within the same trial. Those studies undertaken as direct comparisons are summarized below.

Head to Head Comparisons

A total of 26^{125,152,129,237,130,238,239,145,150,124,114,131,127,240,241,135,242,101,243,70,194,128,244,26,92,245} studies compared efficacy of two or more pharmacological agents relative to each other. In general, few drugs showed statistically significant differences relative to each other. Those that did include the following (drug performing better is listed first):

- 1) Sulphomucopolysaccharides versus CDP-choline²³⁸ - Significant differences were seen in favor of sulphomucopolysaccharides in measures of behavior and global assessment in 30 institutionalized patients with mild to moderate MID.
- 2) Donepezil and vitamin E⁷⁰ - Significant differences were seen in favor of donepezil in general cognitive function in 54 patients with mild AD.
- 3) Antagonic stress versus nicergoline⁹² - Significant differences were seen in favor of antagonic stress in cognition as well as a global assessments in 62 subjects with mild to moderate AD.
- 4) Antagonic stress versus meclofenate²⁴² - Significant differences were seen in favor of antagonic stress in measures of cognition and global assessment in 63 patients with mild to moderate AD.
- 5) Posatirelin versus citicoline¹⁰¹ - Significant differences were seen in favor of posatirelin in general cognitive measure and mood in 222 community living patients with mild to moderate AD.
- 6) Pyritinol versus hydergine²⁴³ - A significant difference was found in favor of pyritinol in a global assessment measure in 102 Hispanic patients with mild to moderate AD.
- 7) Idebenone²⁶ versus tacrine-Mixed results were observed; the Efficacy Index Score showing a significant benefit over tacrine, while the global assessment showed no difference in 203 AD patients, 44 of whom completed the study.

Relative comparisons of FDA approved drugs for the treatment of dementia. Although no head to head trials compared drugs that are likely to be used in current practice in the United States, it was recognized that an assessment of the relative effectiveness of those drugs approved for the treatment of dementia would be of interest to clinicians. Four drug interventions that are currently approved for the treatment of dementia include donepezil, galantamine, rivastigmine,

and tacrine. We caution the reader that inferences drawn from the following figures are limited because these FDA-approved drugs were not compared within the same study. The evidence for benefits and harms has been previously discussed in this report. The pooled estimates (WMD and RR) of two outcomes (ADAS-cog, CIBIC) frequently used in clinical practice have been presented together to illustrate the relative benefit of these approved drugs (Figures 23 to 30). For the purposes of this relative comparison, the pooled estimate reflecting the largest effect size (i.e. the dose showing the greatest magnitude) was selected. Several relevant details should be noted before comparing these estimates as follows: 1) the 5 mg dose of donepezil was selected because the magnitude of the pooled estimate was largest, 2) the 32 mg dose of galantamine had the largest pooled estimate, 3) the rivastigmine pooled estimate for the ADAS-cog was significant for heterogeneity, so the pooled estimate should be considered with great caution, and 4) none of the studies that evaluated tacrine and measured the CIBIC reported sufficient data to estimate an effect size; hence the effect size of the CGIC was substituted for comparison.

Figure 23. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing donepezil versus placebo.




Figure 24. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing galantamine versus placebo.




Figure 25. Weighted Mean Difference (WMD) from the Random Effects Model (Random) for the ADAS-cog comparing rivastigmine versus placebo.




Figure 26. Weighted Mean Difference (WMD) from the Fixed Effects Model (Fixed) for the ADAS-cog comparing tacrine versus placebo.




Figure 27. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC comparing donepezil versus placebo.




Figure 28. Relative Risk (RR) from the Random Effects Model (Random) for the CIBIC comparing galantamine versus placebo.




Figure 29. Relative comparison of effect sizes for studies using the CIBIC rivastigmine versus placebo.





Figure 30. Relative comparison of effect sizes for studies using the CIBIC comparing tacrine versus placebo.



Question 4: Do certain patient populations benefit more from pharmacotherapy than others?

The following studies contained stratified analyses of outcomes for different clinical populations. A total of 22^{245,211,146,179,181,132,134,142,183,185,56,168,201,63,64,55,173,62,76,75,60,159} studies addressed this question. During data abstraction, these trials were identified if the methods sections (including analyses) stated that stratified analyses were undertaken. Eight different variables were identified for which stratified analyses were reported. These included: age, gender, APOE genotype, disease type, disease severity (as determined by MMSE/ ADAS-cog

threshold levels), treatment center, care dependence, and presence of depression. Of these 22 studies, seven trials^{245,211,146,179,181,132,134} evaluated disease type (AD, PDD, SDAT, MID, VaD). They will be discussed in Question 5 (see below).

Table 7 details the 15^{142,26,185,56,246,201,63,64,55,173,62,76,75,60,159} studies that provided stratified analyses other than for disease type. For disease severity, no clear pattern emerges. For APOE, no significant difference was noted between groups^{173,62,76,75} for the three interventions that included cerebrolysin, donepezil, and galantamine. For age, Thal et al.⁵⁵ conducted a post-hoc analysis to assess the effect of age on the rate of decline. Patients were categorized according to age (< 65 years, 65 years and older). The results of the study indicate that a subgroup of patients, aged 65 years or younger, may benefit more from carnitine as compared to older subjects. Specifically, in the younger population, the significant difference between the treatment and the placebo group was observed for ADAS-cog but not for CDR.

Table 7. Studies with stratified analyses.

CITATION	DRUG	SUBGROUP	DRUG EFFECT
Alvarez 2000	Anapso	Disease severity	SC in ADAS-cog in patients with mild cognitive deterioration and with AD NS in patients with VD
Gutzmann 1998	Idebenone	Disease severity	NR
Weyer 1997	Idebenone	Disease severity ADAS total score \geq 20	SC for ADAS Total
Sano 1992	Carnitine	MMSE	Low mMMSE group SC on the SRT and CSF levels of drug High mMMSE group NS neuropsychological test scores, CGI ratings and CSF levels of drug
Ruether 2001	Cerebrolysin	MMSE	Subgroup MMSE < 20: SC in CGI, ADAS-cog, NAI and ADAS-Noncog. Suggests it's because this group had reduced placebo response.
Schellenberg 1997	Cyclandelate	MMSE, ADAS-cog, Treatment center	NR
Feldman 2001	Donepezil	MMSE Psychoactive drug use	NR
Tariot 2001a	Donepezil	MMSE (10-26) Age	MMSE group: SC greater differences than for the whole group for MMSE, GDR Older patients group: SC for MMSE, CDR
Thal 1996a	Carnitine	Age	SC age-by-treatment interaction on the ADAS-cog ITT population Patients \leq 65 years significant difference in decline for ADAS-cog favoring Carnitine but not for CDR Patients > 65 years NS
Panisset 2002	Cerebrolysin	APOE genotype	NS association of the APOE e4 status and response to study drug
Winblad 2001b	Donepezil	APOE genotype Gender	NS difference for the subgroups

Table 7. Studies with stratified analyses (continued).

CITATION	DRUG	SUBGROUP	DRUG EFFECT
Raskind 2000	Galantamine	APOE genotype	NS
Wilcock 2000	Galantamine	APOE genotype MMSE	NS for APOE group SC for MMSE < 18
Prasher 2002	Donepezil	Down syndrome ONLY in trial	NR
Reifler 1989	Imipramine	Depression	Depressed patients SC higher HAM-D scale score. For MMSE patients with AD + depression had higher scores initially and improved significantly more over time

SC = Significant change

NS = Not statistically significant

NR = Not reported

In general, very few studies examined the efficacy of drugs with respect to dementia by population characteristics. Three additional studies attempted to evaluate unique populations or population characteristics. Prasher et al.⁶⁰ evaluated subjects who had Down's Syndrome with dementia and were treated with donepezil, and found none of the outcomes to be significant; this study had a sample size of 30 subjects and was underpowered. Ban et al.²¹³ conducted a multicenter, placebo-controlled, double-blind study with Hispanic and Italian populations. This study was not designed to specifically evaluate the efficacy of glycosaminoglycan polysulfate by ethnicity. However, the study included centers from Mexico, Panama, Naples, and Trieste. This study examined whether the changes encountered in the different outcome measures could be related to center effect, but no statistically significant center effect was found. While this study suggests that ethnicity may have minimal impact, future studies should specifically assess the impact of racial composition on the efficacy of drugs.

Question 5: What is the evidence for the treatment of VaD?

Summary Table 25 details the results of studies in which patients had VaD, or stratified data were presented with respect to VaD subgroups identified as VaD or MID. The trial details for all these studies are provided in evidence tables of key study characteristics, evidence tables of study results, and evidence tables of adverse events found in Appendix C; summary results of trials were also discussed in the results sections of Question 1.

A total of 20 pharmacological interventions in 29 studies^{211,220,238,171,200,199,146,68,181,184,133,134,132,161,89,91,93,247,187,191,192,194,193,100,98,196,195,245,217} were applied specifically to dementias classified as VaD. Sixteen studies evaluated populations entirely composed of patients with VaD (or MID), and the remaining 13 trials had VaD as a subgroup. The majority of these pharmacological interventions (n = 14) were represented by a single trial, limiting the extent of the evidence; these included ataroid, buflomedil, cerebrolisin, sulphomucopolysaccharides (CDP choline), citalopram, donepezil, Ginkgo biloba, idebenone, minaprine, nimodipine, nicergoline, oxiracetam, 5-THF (trazodone), vincamine, and xantinolnicotinate. Surprisingly, four of these trials did not report any results relative to placebo,

and these included buflomedil, Ginkgo biloba, oxiracetam, and 5-THF (trazodone); all but one of these trials²²⁰ evaluated subgroups of VaD patients and likely did not possess sufficient power to evaluate differences. Six interventions had more than a single trial, and these included Choto-san (n = 2), memantine (n = 3), nicergoline (n = 2), pentoxifylline (n = 4), posatirelin (n = 2), and propentofylline (n = 2).

Several of the trials with sample sizes greater than 100 subjects showed significant differences in general cognitive function: ateroid, cerebrolysin, donepezil, idebenone, and nicergoline. Similarly, these larger sample studies showed statistical differences for global assessment: Choto-san, donepezil, memantine, nicergoline, propentofylline, vincamine, and xantinolnicotinate. Findings for other outcome domains were inconclusive, as these were rarely evaluated (see Summary Table 25).

Table 8 below lists the studies that undertook comparisons between VaD populations and other dementia types. Although, not consistent across all trials, three of the studies suggests possible differences between 1) MID and AD for 5'-MTHF-trazodone,²⁴⁵ 2) AD/SDAT and VaD for citalopram,¹⁴⁶ and 3) DAT and MID for Ginkgo biloba.¹⁸¹

Table 8. Studies evaluating vascular dementia patients relative to other dementias.

CITATION	DRUG	SUBGROUP	DRUG EFFECT
Passeri 1993	5'-MTHF Trazodone (TRZ)	AD vs. MID	Equivalence study When patients with AD were analyzed separately the same pattern of response to MTHF and TRZ was found in the HDRS and RVM as when they were analyzed together with patients with MID. MID as separate group: HDRS was significantly reduced vs. baseline after 8 weeks of treatment in the TRZ group and only at the end of the follow-up period in the MTHF group. RVM remained unchanged in MID pts in both treatment groups.
Ban 1991b	Ateroid	PDD vs. MID	NR
Nyth 1990	Citalopram	AD/SDAT vs. VaD	A period: No improvement in the VaD group SC in the AD/SDAT group in emotional bluntness, confusion, irritability, anxiety, fear-panic, depressed mood, and restlessness. MADRS scores significantly reduced B period: AD/SDAT group SC in emotional bluntness at week 8. NS at week 4 and 12. NS for the VaD group.
LeBars 1997	Ginkgo biloba	AD vs. MID+AD MMSE	AD subgroup: SC for ADAS-cog and GERRI
Kanowski 1996	Ginkgo biloba	DAT vs. MID	Improvements at 24 weeks of treatment in comparison to baseline values were consistently slightly greater in the DAT group than in the MID group. Calculation of descriptive p-values seemed inappropriate due to the very small number of patients with MID in the sample.
Winblad 1999	Memantine	AD/VaD Care dependence	NR for differences between dementia types Care dependence: Patients with < 20 points on the CGI and BGP Care dependence subscore shows slightly higher response rates than those with >20 points in the memantine group.
Wilcock 2002	Minaprine	SDAT vs. MID	The largest treatment effect occurred in patients with baseline MMSE score < 15 (p = 0.04) and in those without cerebrovascular macro-lesions (p = 0.002)

SC = Significant change

NS = Not statistically significant

NR = Not reported

Table 9: Guide to Overall Summary Tables – Outcome Measures Classified by Domain

General cognitive function measure	Specific cognitive function measure		Global Assessment	Behavior/Mood	Quality of Life /ADL/ Function	Caregiver Burden	Other
ADAS-Cog (also ADAS-11) AMTS BCRS CamCOG CASI CETM IQCODE MCPT MMSE MMMSE SMMSE CMMSE MQ RMT RVM SIB SMQ SMST TP,TPAT WAIS	ACPT Babcock Story recall Barbizet Visuospatial BLM BNT BSRT BSV BVR CCASSS Category Fluency CDT CNTB Controlled Challenge Word Association COWAT CVLT Digit Span Test DSST EFR FCMT FIGT FOM GAGS Grooved Pegboard Test Letter Cancellation Letter Fluency LMT LNNB MAE MEMT MNLT NCT NDT NLT NMIC NST OLT OMDR	R-AVL RM RPM Rey Memory Test Set test Snodgrass Picture Naming Task SRT-DR SWFIT SWFT SKT TK TMT WMS (MQ) WMS-RR ZVT	ADAS ADCS-CGIC ADS AGS-E Bf-S BGP Blessed-D/ BDRS CAPE CDR-NH CDR-SB CGAE CGI CGIC CGRS CIBIC CICIC+ DBDS DMR DRS EIS FCCA FRS GERRI GBS GDS GIS GPI-E HDS HIS MAC-F NOSGER NOSIE NPI (NPI-4, NPI10) PDRS PGIR Plutchik CGS RAGS RGRS SCAG Stockton GRS TSI VRGI	ABID ABSR ACES ADAS-Non-cog AFBS BDI BEHAVE-AD BPRS BRMS BRDS CERAD-BRSD CMAI CS or CSDD DSCS DSS Facial Behavior GS HAM-A HAM-D HDRS HDS-R IPSC-E LPRS MAACL-R MADRS MOSES NAB NMS NOSGER-IADL NPI-NH NSL OAS PANSS-EC POMS RMBPC RPT SBI SHGRT SRT VHB	ABS ADCS-ADL ADFACS ADL ADL-C ADL-PDS BI Dependency Scale DAD FAST FIM IADL IDDD NAA NAI OARS-ADL PDS PSMS PSQI QoL QoL-P QoL-C SF-36 SIP Time to functional decline	CATS CSS CSI SCB	CAUST SAS AIMS BARS/ BAS ERP ESRS Finger Tapping Test SAS UPDR

Summary Evidence Tables

Summary Table 1. Carnitine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ADL	Caregiver Burden	Other
Livingston, 1991	NS*	MX*	NT	NT	NS*	NT	NT
Rai, 1989	NT	NS*	NS*	NT	NS*	NT	NT
Sano, 1992	NS*	NS*	NT	NT	NS*	NT	NT
Spagnoli, 1991	NT	MX	MX	NS	NT	NT	NT
Thal, 2000a	NS	NT	NS	2° NS	2° NS	NT	NT
Thal, 1996a Brooks, 1998	NS	NT	NS	2° NS	2° NS	SUBGROUP	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 2. Donepezil.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ADL	Caregiver Burden	Other
Burns, 1999	SC	NT	SC	NT	SC	NT	NT
Feldman, 2001 Gauthier, 2002	2° SC	NT	SC	NR	2° SC	NR	SUBGROUP
Mohs, 2001	SC	NT	NT	NT	SC	NT	SC Time to functional decline
Prasher, 2002	2° NS*	NT	NS*	2° MX*	NT	NT	NT
Rogers, 1996 Rogers, 2000 Neumann, 1999 Rogers, 1998	SC	NT	SC	NT	2°MX	NT	NT
Rogers, 1998b Doody, 2001 Sparano, 1998	SC	NT	SC	NT	2° NS	NT	NT
Rogers, 1998a Doody, 2001 Steele, 1999	SC	NT	SC	NT	2° SC	NT	NT
Tariot, 2001	2° NS	NT	2° SC	NS	2° NS	NT	SUBGROUP
Winblad, 2001	2° SC	NT	SC	2° NS	2° SC	NT	SUBGROUP
Pratt, 2002	SC	NT	SC	NT	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 3. Galantamine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ADL	Caregiver Burden	Other
Erkinjuntti, 2002	SC*	NT	SC*	2°SC*	2°SC*	NT	NT
Raskind, 2000	SC	NT	SC	NT	2°NS	NT	NT
Rockwood, 2001	SC	NT	SC	NS	2°SC	NT	NT
Tariot, 2000	SC	NT	SC	SC	SC	NT	NT
Wilcock, 2000 Wilcock, 2001	SC	NT	SC	NT	SC	NT	NT
Wilkinson, 2001	MX	NT	2°NS	NT	2°NS	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 4. Metrifonate.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL n	Caregiver Burden	Other
Becker, 1996	SC*	NT	2°MX*	2°NS*	2°NS*	NT	NT
Becker, 1998	SC*	NT	2°NS*	2°NS*	2°NS*	NT	NT
Cummings, 1997	SC*	NT	SC*	NT	NT	NT	NT
Cummings, 1998b Cummings, 1998a	SC	NT	SC	NT	2°NS	NT	NT
Dubois, 1999 McKeith, 1998	SC	NT	SC	2°SC	2°SC	NT	NT
Jann, 1999	SC	NT	2°MX	2°NS	NT	NT	NT
Morris, 1998	SC	NT	SC	2°NS	2°NS	NT	NT
Pettigrew, 1998	NR	NT	NR	NR	NT	NT	NT
Raskind, 1999	SC	NT	MX	MX	NS	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 5. Nicergoline.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Herrmann, 1997	SC	NT	SC	NT	NT	NT	NT
Nappi, 1997	SC*	NT	SC*	NT	NT	NT	NT
Saletu, 1995 Saletu, 1997	SC*	NT	MX*	NS*	NT	NT	SUBGROUP
Winblad, 2001a	SC	NT	NS	2°NS	2°NS	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 6. Physostigmine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Van Dyck, 2000	SC	NT	MX	NT	2°NS	NT	NT
Moller, 1999	NR	NT	NS*	NR	NT	NT	NT
Thal, 1996b	SC	NT	SC	NT	2°NS	NT	NT
Thal, 1999	SC	NT	SC	NT	2°NS	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 7. Posatirelin.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Ferrari, 1998	SC	NT	SC	NS	SC	NT	SUBGROUP
Gasbarrini, 1997	SC	NT	NT	SC	SC	NT	NT
Parnetti, 1995	NR	NT	NT	NR	NR	NT	NT
Parnetti, 1996	MX*	NT	NT	NS*	SC*	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2^o Secondary outcome
* OC analysis

Summary Table 8. Rivastigmine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Agid, 1998	NR	SC*	SC*	NS*	NS*	NT	NT
Corey-Bloom, 1998 Farlow, 2001 Farlow, 2000 Kumar, 2000 Del Ser, 2000 Doraiswamy, 2002	SC	NT	SC	NT	SC	NT	NT
Forette, 1999	SC*	NT	SC*	NS*	NS*	NT	NT
McKeith, 2000	SC	NT	NS	MX	NT	NT	NT
Potkin, 2001	NT	NS*	SC*	NT	NT	NT	NT
Rosler, 1999 Rosler, 2001 Farlow, 2000 Rosler, 1998 Doraiswamy, 2002	SC	NT	SC	NT	SC	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 9. Tacrine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Knapp, 1994b Farlow, 1998 Gracon, 1996 Henke, 1997 Knapp, 1994a Knopman, 1996 Raskind, 1997 Schneider, 1997 Schneider, 1996 Smith, 1996	SC	NT	SC	NS	NT	NT	NT
Maltby, 1994	NS*	NS*	NT	NS*	NS*	NS*	NT
Prentice, 1996	NS*	NT	NT	NS*	NT	NT	NT
Weinstein, 1991 Gool, 1991	NS*	NT	NT	NT	NS*	NS*	NT
Wong, 1999	MX	NT	NS	NT	NT	NT	NT
Wood, 1994	NS	NT	SC	NS	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 10. Velnacrine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Zemlan, 1996	SC*	NT	SC*	2°NS*	2°NS*	NT	NT
Antuono, 1995	SC	NT	SC	NT	2°SC	2°SC	NT
Huff, 1991	NT	NS*	MX*	NT	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 11. Various cholinergic neurotransmitter modifying agents.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
CHOLINERGIC NEUROTRANSMITTER MODIFYING AGENTS							
Eptastigmine							
Imbimbo, 1999	SC	NT	SC	NT	SC	NT	NT
Canal, 1996	NS*	NS*	MX*	NT	MX*	NT	NT
Huperzine							
Xu, 1995	SC*	NT	NT	SC*	SC*	NT	NT
Linopirdine							
Van Dyck, 1997	NS*	NT	NS*	NS*	NT	NT	NT
Rockwood, 1997	SC	2°NS	NS	2°NS	2°NS	NT	NT
Rockwood, 2000							
Sabeluzole							
Mohr, 1997	NS*	NS*	NT	NT	NT	NT	NT

MX
NS

Mixed results
Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 12. Haloperidol.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver/ Burden	Other
Allain, 2000	2°NS	NT	2° SC	SC	NT	NT	NT
Auchus, 1997	NT	NT	NT	NS*	NT	2° NS*	NT
De Deyn, 1999	NR	NT	NR	SC	NR	NT	NT
Petrie, 1982	NT	NT	NR	SC*	NT	NT	NT
Teri, 2000	NS*	NT	NS*	NS*	SC* favors Placebo	NS*	NT

MX Mixed results
 NS Not statistically significant

NT Not tested
 SC Significant change

2° Secondary outcome
 * OC analysis

Summary Table 13. Memantine.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Orgogozo, 2002	SC	NT	NS	NT	NT	NT	NT
Wilcock, 2002	SC	NT	NS	NT	NT	NT	NT
Winblad, 1999	NT	NT	SC	SC	SC	NT	NT

MX Mixed results
 NS Not statistically significant

NT Not tested
 SC Significant change

2° Secondary outcome
 * OC analysis

Summary Table 14. Selegeline.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Agnoli, 1992	MX*	NT	NT	NT	NT	NT	NT
Burke, 1993a Burke, 1993b	NS*	NT	NS*	NS*	NT	NT	NT
Filip, 1999	MX*	NT	MX*	NT	NT	NT	SUBGROUP
Freedman, 1998	2° NS	2° NS	2° NS	NS	NT	NT	NT
Mangoni, 1991 Smirne, 1993	NR	SC*	SC*	SC*	NT	NT	NT
Sano, 1997 Sano, 1996	NT	NT	NT	NT	NT	NT	NS Survival SUBGROUP

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 15. Various non-cholinergic neurotransmitter/neuropeptide modifying agents.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Perphenazine							
Pollock, 2002	NR	NT	NT	NS	NR	NR	NT
Thioridazine							
Barnes, 1982	NT	NT	NS	NS	NT	NT	NT
Alaproclate							
Dehlin, 1985	NS	NT	NT	NS	SC	NT	NT
Anapsos							
Alvarez, 2000	SC*	NT	NT	NT	NT	NT	SUBGROUPS
Cutler, 1993	NR*	NT	NS*	NT	NT	NT	NT
Citalopram							
Nyth, 1990	NT	NT	MX*	NS*	NT	NT	NT
Pollock, 2002	NT	NT	NT	SC	NT	NT	NT
Divalproex Sodium							
Tariot, 2001b	2° NS	NT	2° SC favors Placebo	NS	NT	NT	NT
Porsteinsson, 2001	2° NS*	NT	2° NS*	NS*	2° NS*	NT	NT
Fluvoxamine							
Olafsson, 1992	NS*	NS*	NS*	NT	NT	NT	NT
Fluoxetine							
Petracca, 2001	NS	NT	NR	NS	NS	NT	NT
Auchus, 1997	NT	NT	NT	NS	NT	2°NS	NT
Imipramine							
Reifler, 1989	NS*	NT	SC*	NS*	NS*	NT	SUBGROUPS
Lisuride							
Claus, 1998	SC*	NS*	NS*	NS*	NT	NT	NT
Lorazepam							
Meehan, 2002 Clark, 2001 Kennedy, 2001 Mintzer, 2001 Street, 2001	2° NS*	NT	2° NS*	NS*	NT	NT	NT
Loxapine							
Barnes, 1982	NT	NT	NS	NS	NT	NT	NT
Petrie, 1982	NT	NT	NR	SC*	NT	NT	NT

LU25

MX Mixed results
NS Not statistically significant
NT Not tested
SC Significant change
2° Secondary outcome
* OC analysis

Summary Table 15. Various non-cholinergic neurotransmitter/neuropeptide modifying agents.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Thal, 2000b	NS	NT	NS	2° NS	2° NS	NT	NT
Maprotiline							
Fuchs, 1993	2° NS*	NT	NS*	NT	NT	NT	NT
Minaprine							
Passeri, 1987	NT	NT	NT	MX	NT	NT	NT
Moclobemide							
Roth, 1996	SC	NT	MX	SC	NT	NT	NT
<u>Naftidrofuryl</u>							
Moller, 2001	SC	NT	SC	NT	NT	NT	NT
Olanzapine							
Meehan, 2002	2° NS*	NT	2° NS*	SC*	NT	NT	NT
Street, 2000	2° NS	NT	NT	SC	NT	NT	NT
Phosphatidylserine							
Amaducci, 1988 SMID Group, 1987 Amaducci, 1986	SC*	SC*	SC*	NT	NT	NT	SUBGROUP
Crook, 1992a	NT	NT	SC	NT	2° NS	NT	NT
Risperidone							
Katz, 1999 Jeste, 2000 Pryse-Phillips, 2000	NT	NT	2° SC	SC	NT	NT	NT
De Deyn, 1999	NS	NT	SC	MX	NS	NT	NT
Sertraline							
Lyketsos, 2000	2° NS	NT	SC	2° MX	2° SC	NT	NT
Magai, 2000	NS	NT	NT	NS	NT	NT	NT
Tiapride							
Allain, 2000	NR	NT	NR	2° SC	NT	NT	NT
Trazodone							
Teri, 2000	NS*	NT	NS*	NS*	SC*	NS*	NT
Xanomeline							
Bodick, 1997 Veroff, 1998 Satlin, 1997	SC	2° SC	SC	NT	2° SC	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 16. Cerebrolysin.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Bae, 2000	SC	NT	SC	NT	2° NS	NT	NT
Panisset, 2002	NS*	NT	SC*	NT	2° NS*	NT	NT
Ruther, 2001 Ruther, 2002	SC	NR	SC	2° SC	NR	NT	NT
Ruther, 1994 Ruther, 2000	NT	SC	SC	NR	2° NS	NT	NT
Xiao, 2000	SC	2° MX	SC	NT	2° MX	NT	NT
Xiao, 1999	SC	2° SC	NS	2° NS	2° NS	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 17. Estrogens.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Asthana, 2001	NT	SC	2°NS*	NR	2°NS*	NT	NT
Henderson, 2000	NS*	NT	2°NS*	2° NS*	2°NS*	NT	NT
Kyomen, 1999 Kyomen, 2002	NT	NR	NS*	MX*	NS*	NT	NT
Mulnard, 2000	2° NS	2° MX	NS	2° NS	2° NS	NT	NT
Wang, 2000	NS	NT	NS	2° NS	NT	NT	NT

MX
NS

Mixed results
Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 18. Ginkgo Biloba.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver/ Burden	Other
Kanowski, 1996	NT	SC	SC	NS	NT	NT	NT
Le Bars, 1997 Le Bars, 2000 Le Bars, 2002 Por, 1998	SC	NT	MX	NT	NT	NT	NT
Maurer, 1997	2° NS	SC	2° NS	NR	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 19. Idebenone.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Bergamasco, 1994	NT	MX*	SC*	NT	NT	NT	NT
Gutzmann, 1998 Weyer, 1996	2° SC	NT	SC	NT	2° SC	NT	NT
Marigliano, 1992	SC*	NT	NT	NT	SC*	NT	NT
Weyer, 1997	2° SC	NT	SC	2° SC	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 20. Oxiracetam.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Bottini, 1992	NT	MX*	NT	NT	SC*	NT	NT
Maina, 1989	SC*	NT	SC*	SC*	NT	NT	NT
Mangoni, 1988	NT	SC*	NT	SC*	NT	NT	NT
Rozzini, 1992 Rozzini, 1993	NR	NR	NT	NR	NS*	NT	NT
Villardita, 1992	SC*	MX*	NT	NS*	SC*	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 21. Pentoxifylline.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Black, 1992	2° NS*	NT	NS*	2° NS*	NT	NT	SUBGROUP
Ghose, 1987	MX	2° NS*	NS*	NT	NT	NT	SUBGROUP
Knezevic, 1996	2° NS	NT	NS	2° NS	2° NS	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 22. Propentofylline.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Marcusson, 1997	2° SC	SC	MX	2° SC	2° NS	NT	NT
Mielke, 1998	NS*	NS*	NT	NT	NT	NT	NT
Mielke, 1996	NS*	NS*	NT	NT	NT	NT	NT
Saletu, 1990 Moller, 1994	SC*	NS*	SC*	NT	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 23. Additional pharmacological agents.

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
ACTH Neuropeptides							
Soininen, 1985 Partanen, 1986	NT	NT	NS*	NT	NT	NT	NT
Kragh-Sorensen, 1986	NT	SC*	SC*	NT	NT	NT	NT
Aniracetam							
Senin, 1991	SC*	SC*	SC*	NT	NT	NT	NT
Ateriod							
Ban, 1991b	SC*	NT	MX*	NS*	NT	NT	NT
Buflomedil							
Cucinotta, 1992	NT	NR	NR	NR	NT	NT	NT
Choto-san							
Shimada, 1994	NT	NT	MX*	SC*	NT	NT	NT
Terasawa, 1997	NT	NT	SC*	NS*	NT	NT	NT
Citicoline							
Parnetti, 1995	NR	NT	NT	NR	NR	NT	NT
Cyclandelate							
Schellenberg, 1997	NT	MX	SC	SC	NT	NT	NT
Weyer, 2000	NS	NT	NS	NT	NS	NT	NT
DDAVP (Deamino-D-arginine-vasopressin)							
Peabody, 1986	NS*	NT	MX*	MX*	NT	NT	NT
Denbufylline							
Treves, 1999	NS*	NS*	NT	NT	NT	NT	NT
Desferrioxamine							
Crapper -McLachlan, 1991	NT	NT	NT	SC*	NT	NT	NT
Diclofenac/misoprostol							
Scharf, 1999	NS	NT	NS	2° NS	2° NS	NT	NT
Ergokryptine							
Cucinotta, 1996 Cucinotta, 1998	2° SC	2° SC	MX	NR	NT	NT	NT
Danielczyk, 1988	NS*	NR	SC*	MX*	NS*	NT	NT
Glycosaminoglycan polysulfate							
Ban, 1991a	MX*	NT	MX*	SC*	NS*	NT	NT
Guanfacine							
Crook, 1992b	NR	SC	SC	NT	NT	NT	NT
Hydergine							

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Thompson, 1990	MX*	NT	MX*	NS*	NT	NT	NT
Hydroxchloroquine							
Van Gool, 2001	2° NS	NT	NT	2° NS	NS	NT	NT
Aisen, 2002b	NS	NT	NS	NS	NS	NT	NT
Indomethacin							
Rogers, 1993	NS*	MX*	NS	NT	NT	NT	NT
Monosialotetrahexosylgan GM1							
Ala, 1990	NS*	NS*	NS*	NS*	NS*	NT	NT
NAC (N-Acetylcysteine)							
Adair, 2001	NS	2°NS	NT	NT	NS	NT	NT
Nimodipine							
Pantoni, 2000a	NS	NS	NS	NT	NS	NT	NT
Ban, 1990	SC*	SC*	SC*	SC*	NT	NT	NT
Nizatidine							
Carlson, 2002	NT	NS	NT	NT	NS	NT	NT
Breitner, 1999							
Nootropic agent - BMY							
Shrotriya, 1996	SC*	NT	NS*	NT	NT	NT	NT
Piracetam							
Croisile, 1993	NS*	NS*	NS*	NS*	NT	NT	NT
Prednisone							
Aisen, 2000b	NS	NT	2° NS	2° MX	NT	NT	NT
Aisen, 2000a							
Simvastatin							
Simons, 2002	MX*	NT	NT	NT	NT	NT	NT
Thiamine							
Nolan, 1991	SC*	SC*	NT	NT	NT	NT	NT
Vincamine							
Fischhof, 1996	NT	NR	SC*	NT	NR	NT	SUBGROUP
Vitamin E							
Sano, 1997	NT	NT	NT	NT	NT	NT	SC
Sano, 1996							Institutionalization
							SUBGROUP
Xantinolnicotinate							
Kanowski, 1990	NT	SC*	SC*	NT	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 24. Drug vs drug studies

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Haloperidol / Tradozone							
Teri, 2000	NS	NT	NS	NR	NR	NR	NT
Olanzapine / Lorazepam							
Meehan, 2002	NR	NT	NT	NT	NT	NT	NT
Haloperidol/Loxapine							
Carlyle, 1993	NT	NT	NT	NS*	NT	NT	NT
Alprazolam / Lorazepam							
Ancill, 1991	NT	NT	NS*	NT	NT	NT	NT
Haloperidol / Oxazepam / Diphenhydramine							
Coccaro, 1990	NT	NT	NS*	NS*	NS*	NT	NT
Sulphomucopolysaccharides / CDP-choline							
Cucinotta, 1987	NT	MX*	SC* favors sulphomucopolysaccharides	SC* favors sulphomucopolysaccharides	NT	NT	NT
Citalopram / Mianserin							
Karlsson, 2000	NT	NT	NT	NS*	NT	NT	NT
Citalopram/Perphenazine							
Pollock, 2002	NR	NT	NT	NR	NR	NR	NT
Thoridazine / Loxapine							
Barnes, 1982	NT	NT	NS	NR	NT	NT	NT
Tiapride / Haloperidol							
Allain, 2000	2° NS	NT	2° S	NS	NT	NT	NT
Tacrine / Silymarin							
Allain, 1998	NR	NR	NT	NT	NT	NT	NT
Risperidone / Haloperidol							
Chan, 2001	NR	NT	NT	NS*	NR	NT	NT
De Deyn, 1999	NR	NT	NR	NR	NR	NT	NT
Paroxetine / Imipramine							
Katona, 1998	NT	NT	NS	NS	NT	NT	NT
Fluoxetine / Amitriptyline							
Taragano, 1997	NS*	NT	NT	NS*	NT	NT	NT
Selegiline / Alpha-Tocopherol							
Sano, 1997	NT	NT	NT	NT	NT	NT	NR
Sano, 1996							
Meclofenoxate / Antagonic Stress							

MX Mixed results
 NS Not statistically significant

NT Not tested
 SC Significant change

2° Secondary outcome
 * OC analysis

Summary Table 24. Drug vs drug studies

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Popa, 1994	SC* favors Antagonic Stress	SC* favors Antagonic Stress	SC* favors Antagonic Stress	NT	NT	NT	NT
Posatirelin / Citicoline							
Parnetti, 1995	SC* favors Posatirelin	NT	NT	SC* favors Posatirelin	NR	NT	NT
Pyritinol / Hydergine							
Spilich, 1996	NR	NT	SC* favors Pyritinol	NT	NT	NT	NT
Donepezil / Vitamin E							
Thomas, 2001	SC* favors Donepezil	NT	NT	NR	NT	NT	NT
Sulodexide / Pentoxifylline							
Parnetti, 1997	NR	NT	NT	NR	NR	NT	NT
Haloperidol / Fluoxetine							
Auchus, 1997	NT	NT	NT	NS*	NT	2° NS*	NT
Melperone / Tiapride							
Gutzmann, 1997	NT	NT	NS	NR	NR	NT	NT
Idebenone/Tacrine							
Gutzmann, 2002	NS	NT	SC favors Idebenone	NR	NT	NT	NT
Nicergoline/Antagonic Stress							
Schneider, 1994	SC* favors Antagonic Stress	NT	SC* favors Antagonic Stress	NT	NT	NT	NT
Tradozone / 5'-MTHF Folate							
Passeri, 1993	NR	NT	NT	NR	NT	NT	NT

MX Mixed results
 NS Not statistically significant

NT Not tested
 SC Significant change

2° Secondary outcome
 * OC analysis

Summary Table 25. VaD/MID Studies

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Ban, 1991b Ateroid Subgroup MID	SC*	NT	MX*	NS*	NT	NT	NT
Cucinotta, 1992 Buflomedil VaD	NT	NR	NR	NR	NT	NT	NT
Cucinotta, 1987 sulphomucopolysaccari des vs CDP-choline MID	NS	SC*	SC* favors sulphomucopoly saccharides	SC* favors sulphomucopoly saccharides	NT	NT	NT
Xiao, 1999 Cerebrolysin VaD	SC	2° SC	NS	2° NS	2° NS	NT	NT
Shimada, 1994 Choto-san VaD	NT	NT	MX*	SC*	NT	NT	NT
Terasawa, 1997 Choto-san VaD	NT	NT	SC*	NS*	NT	NT	NT
Nyth, 1990 Citalopram Subgroup VaD	NT	NT	NS*	NS*	NT	NT	NT
Pratt, 2002 Donepezil VaD	SC	NT	SC	NT	NT	NT	NT
Kanowski, 1996 Ginkgo Biloba Subgroup MID	NT	NR	NR	NR	NT	NT	NT
Marigliano, 1992 Idebenone MID	SC*	NT	NT	NT	SC*	NT	NT
Orgogozo, 2002 Memantine VaD	SC	NT	NS	NT	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 25. VaD/MID Studies

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Wilcock, 2002 Memantine VaD	SC	NT	NS	NT	NT	NT	NT
Winblad, 1999 Memantine Subgroup HIS \geq 5	NT	NT	2° SC*	NT	NT	NT	NT
Passeri 1987 Minaprine Subgroup MID	NT	NT	NT	MX	NT	NT	NT
Herrmann, 1997 Nicergoline MID	SC	NT	SC	NT	NT	NT	NT
Saletu1995 Saletu1997 Nicergoline Subgroup MID	SC*	NT	MX*	NS*	NT	NT	NT
Pantoni, 2000a Nimodipine MID	NS	NS	NS	NT	NS	NT	NT
Maina, 1989 Oxiracetam Subgroup MID	NT	NR	NR	NR	NT	NT	NT
Knezevic, 1996 Pentoxifylline MID	2° NS	NT	NS	2° NS	2° NS	NT	NT
Black, 1992 Pentoxifylline Vascular damage or strokes	2° NS*	NT	NS*	2° NS*	NT	NT	SUBGROUP
Parnetti, 1997 Pentoxifylline vs Sulodexide VaD	NR	NT	NT	NR	NR	NT	NT
Ghose 1987 Pentoxifylline Subgroup MID	SC*	NS*	NS*	NT	NT	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

2° Secondary outcome
* OC analysis

Summary Table 25. VaD/MID Studies

Author, Year	General Cognitive Function Measure	Specific Cognitive Function Measure	Global Assessment	Behavior/ Mood	Quality of Life/ ADL	Caregiver Burden	Other
Parnetti, 1996 Posatirelin VaD	MX*	NT	NT	NS*	SC*	NT	NT
Ferrari 1998 Posatirelin Subgroup VaD	NT	NT	NR	NT	NT	NT	NT
Mielke, 1996 Propentofylline VaD	NS*	NS*	NT	NT	NT	NT	NT
Marcusson 1997 Propentofylline Subgroup VaD	NT	SC	SC	NR	NR	NT	NT
Passeri 1993 5'-MTHF vs Tradozone Subgroup MID	NT	NR	NT	NR	NT	NT	NT
Fischhof, 1996 Vincamine Subgroup MID	NT	NR	SC*	NT	NR	NT	NT

MX Mixed results
NS Not statistically significant

NT Not tested
SC Significant change

^{2°} Secondary outcome
* OC analysis

Chapter 4. Discussion

This systematic review was undertaken primarily to evaluate the efficacy of pharmacological agents in the treatment of dementia. The studies were limited to parallel design RCTs with quality scores greater than 3 on the Jadad scale. The interventions were not limited to those currently on label by the FDA; it was of interest to cast a wide net and capture reports of pharmacological agents that are used off-label for the treatment of dementia. Since a variety of agents with different therapeutic effects were evaluated, the outcomes were not restricted to a specific subset of all available outcomes used in the dementia literature. The psychometric properties of some of the most commonly used outcomes have been critically appraised and found to be limited. Moreover, there is no current consensus as to which domains, and the outcomes within these, that best reflect clinically important change.

Strength of the Evidence

The studies eligible for review in this dementia report represent the highest form of evidence. This strongly suggests that these trials are more likely to be “well-designed, well conducted studies in representative populations that assess the effects of health outcomes”.⁵³ The high quality scores also indicate that the studies evaluated in this systematic review have a relatively high level of internal validity. The characteristics of the population and the interventions were detailed to assist the reader in evaluating the degree of external validity. Similarly, attempts were made to highlight “consistency” in the evidence as well as the quantity of evidence and the magnitude of the reported changes.

Although, there is greater understanding on evaluating the evidence for the “benefits” of therapies, there is less clarity on determining the potential for harms from pharmacological interventions for treating dementia. With respect to adverse events and the potential for serious harms, greater variability in systematic collection and reporting of these were observed in the dementia pharmacological literature. Evaluation of the potential for harm is considered with three main points: 1) the most frequently reported adverse events across studies for a specific drug, 2) the overall withdrawal rate due to adverse events for both the control and treatment groups, and 3) the range of frequencies reported for a subset of symptoms (nausea, diarrhea, dizziness, agitation, eating disorder) selected **a priori** and evaluated for all pharmacological interventions.

At present there is no coherent framework that captures the disease processes present in dementia patients for the range of outcomes evaluated in this systematic review. This report details the highest evidence from both a design and internal validity perspective. It is our view that determining the clinical relevance (external validity) of such high-quality evidence must ultimately be reached by consensus amongst multidisciplinary experts within the decision-making body that will use this evidence for such purposes as developing practice guidelines.

Question 1: Does pharmacotherapy for dementia syndromes improve cognitive symptoms and outcomes?

Summary of the Systematic Review Results

A total of 97 interventions in 186 studies were eligible for evaluation in this systematic review and were distributed as follows:

- A total of 16 different cholinergic neurotransmitter modifying pharmacological agents in 72 studies.
- A total of 35 non-cholinergic neurotransmitter/neuropeptide modifying agents in 61 studies.
- A total of 46 other pharmacological agents in 76 studies*.

* there are more than 186 studies here because some studies compared a drug from one class with a drug from another, so that study would be in both categories and therefore counted twice.

- two studies compared two NCNMA with an OTHER.
- two studies compared a CNMA with an OTHER.
- one study compared a CNMA with two OTHERS.
- two studies compared an NCNMA with an OTHER.

The evidence for all these pharmacological agents was presented in great detail in Chapter 3 and in Evidence Tables of Key Study Characteristics, Tables of Study Results, and Tables of Study Adverse Events contained in Appendix C. Conclusions regarding those pharmacological agents that had a minimum of three trials are summarized here. The summary of the pharmacological agents that had fewer than three trials can be found in Chapter 3.

Summary of Cholinergic Neurotransmitter Modifying Agents

Carnitine. Six trials evaluated carnitine in 925 subjects with mild to moderate severity, recruited predominately from the community at doses of 2 to 3 g for either 24 or 52 weeks. Evidence of benefit is conflicting for the domains of cognition. Most studies were not statistically significant and the lack of sufficient power may have been an important factor. Similarly, no significant differences were found in the domains of global assessment, behavior/mood, and quality of life/ADL; power could not be evaluated for the majority of these outcomes.

Four of the six studies scored 3 for quality on reporting adverse events. Withdrawal rates due to adverse events varied from 0 - 3% (excluding results from one outlier trial²⁴⁸), and gastrointestinal symptoms were the most frequently reported types of adverse events. The percent of subjects reporting the a priori symptoms of interest across all studies were as follows: 1) nausea (placebo = 6 - 14%, all doses carnitine = 28%), and 2) agitation (placebo = 6%, all doses carnitine = 7%). Dizziness, diarrhea, or eating disorder were not reported by any study. No serious adverse events requiring hospitalization and associated with carnitine were reported.

Donepezil. Ten trials in 3239 subjects evaluated the efficacy of donepezil compared to placebo, and one trial compared donepezil to a group given vitamin E. The majority of studies (n = 8) evaluated AD patients, for which half were recruited from the community (other studies did not specify). The subjects had predominately mild to moderate disease and doses of 5 or 10 mg were used with varying duration from 12 to 56 weeks.

There is consistent evidence of benefit in the domains of general cognitive function and global assessment. The combined effect sizes for the ADAS-cog and the CIBIC were estimated. Evidence is inconsistent for a dose response in these domains based on the three studies that evaluated two different doses (5 and 10 mg); the benefit was of similar magnitude for both dose groups for global assessment outcomes. Similarly, two of the three studies that evaluated behavior/mood outcomes (NPI) showed no statistically significant changes relative to placebo; these trials lacked sufficient power to detect a difference. There is some evidence of benefit in ADL outcomes, although this outcome domain was evaluated with a variety of instruments. Caregiver burden outcomes was evaluated in a single study that did not report the findings for this domain.

Adverse events quality scores were 3 or greater for the majority of studies (n=7). Four trials provided evidence of a dose response for adverse events. One study showed a statistical difference for balance-related problems and asthenia (neurological fatigue) between placebo and treatment groups. Withdrawal due to adverse events ranged from 0 - 18% for treatment groups and 0 - 11% for placebo. Four out of six studies testing differences between groups were statistically significant for diarrhea (placebo = 3 - 21%, all doses donepezil = 0 - 38%), nausea and vomiting (placebo = 4 - 9%, all doses donepezil = 4 - 25%). The other a priori symptom reported was agitation and frequencies for placebo varied from 0 - 8% and for all doses from 3 - 19%; but these were not shown to be statistically different.

Galantamine. Six trials in 3530 subjects evaluated the efficacy of galantamine compared to placebo. Doses of 24 and 32 mg were evaluated in half of these studies. Five studies evaluated AD patients and there was limited information regarding whether the subjects were from the general community or institutional settings. All studies recruited subjects with mild to moderate disease and the drug was administered with varying duration of 3 or 6 months.

Evidence of benefit is consistent in the domains of general cognitive function, global assessment, quality of life/ADL. Two of the three studies that evaluated, behavior/mood found statistically significant differences. A small dose effect was evident in the ADL domain when comparing the pooled estimates of the DAD; no dose effect was observed for outcomes in the global assessment domain, and dose effect could not be evaluated for the general cognition domain. The caregiver burden domain was not evaluated in any trials.

Five of the six trials scored 3 out of 5 on our quality scale for rating adverse events. Withdrawal rates due to adverse events ranged from 4 - 9% for placebo and 8 - 27% for the treatment group. One study showed a dose response for adverse events. Although, most trials did not report testing for differences between groups, two trials reported a statistically significant

difference in weight loss with the treatment group having more than the placebo group. The most common types of adverse events reported were gastrointestinal symptoms (nausea and vomiting, diarrhea), eating disorders/weight loss, and dizziness. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea and vomiting (placebo = 3 - 13%, all doses = 6 - 44%), 2) dizziness (placebo = 3 - 11%, all doses = 4 - 19%), 3) diarrhea (placebo = 2 - 10%, all doses = 4 - 19%), 4) agitation (placebo = 1 - 9%, all doses = 6 - 15%), and 5) eating disorder (placebo = 0 - 6%, all doses = 4 - 20%).

Metrifonate. Nine studies compared metrifonate to placebo in 2759 subjects with mild to moderate AD (likely from community settings as the majority of studies did not specify this). Metrifonate dosages evaluated varied from 50 to 80 mg, and study duration ranged from 21 days to 26 weeks duration.

All but one study showed metrifonate to have a consistent positive effect on measures of general cognitive function; none of the studies evaluated any specific cognitive function measures. The effects on global assessment were less consistent, but suggested a positive effect in four of the eight studies that reported this outcome. Evidence for effect in the domains of behavior/mood and quality of life/ADL were not significant in the majority of studies that evaluated these domains, however these were primarily evaluated as secondary outcomes and likely lacked sufficient power.

With the exception of a single study, quality scores for reporting adverse events were greater than 3. However, only one trial⁸³ tested for differences between groups and found nausea and vomiting, diarrhea, and muscle and joint disorder to be significantly different. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea and vomiting (placebo = 3 - 14%, all doses = 2 - 50%), 2) dizziness (placebo = 1%, all doses = 3 - 4%), 3) diarrhea (placebo = 4 - 14%, all doses = 11 - 19%), 4) agitation (placebo = 2 - 14%, all doses = 8 - 33%), and none reported eating disorder as an adverse event. Withdrawal due to adverse events varied from 0 - 9% for placebo and 0 - 12% for the treatment group. Overall, it was difficult to determine which types of adverse events reported had the potential to cause serious harm. This is noteworthy as metrifonate has been withdrawn from use in North America, and Bayer has suspended Phase III trials,⁸⁷ because some patients in clinical trials have experienced serious muscle weakness. This decision was based on the results of an experimental study showing risk of respiratory paralysis with the use of metrifonate. Other adverse events of concern included severe leg cramps, dyspepsia, and bradycardia. None of the studies we reviewed indicated that if present, these events differed significantly between groups. It is not clear if this inconsistency is a function of the methods used to collect and report adverse events or a limitation of RCTs as a source of detecting serious adverse events when incidence is low.

Nicergoline. Four trials in 705 subjects compared nicergoline to placebo and one trial compared it to antagonistic-stress in mixed populations that included AD, MID, PDD, VaD, mixed dementia, and SDAT, which were classified as mild to moderate in severity.

All placebo-controlled trials found a positive effect for general cognitive outcomes, but half the results were based on OC analyses. The evidence was mixed for benefit in the domain of global assessments. No significant differences were found for behavior/mood, and quality of life/ADL outcomes, but these were evaluated in few studies and as secondary outcomes (suggesting that sufficient power was an issue).

Quality scores for reporting adverse events varied from 2 to 5 for these four trials, and none tested for differences between groups. Withdrawal due to adverse events varied from 0 - 8% for placebo and 0 - 9% for the treatment group. With the exception of headache, which was reported in all four trials, it was difficult to determine which types of adverse events most characterized exposure to this pharmacological agent. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = 3%), 2) dizziness (placebo = 1-2%, all doses = 0% or not reported), 3) diarrhea (placebo = 2 - 6%, all doses = 2 - 4%), 4) agitation (placebo = 5%, all doses = not reported), and none reported eating disorder as an adverse event.

Physostigmine. Four studies of 1198 subjects with mild to moderate AD evaluated physostigmine administered in patch and oral form (30 to 60 mg dose) for study duration varying from 6 to 24 weeks. All subjects were recruited from the community.

There is evidence that physostigmine has a statistically significant effect on general cognitive function, as three of the four studies showed improvement. Evidence for an effect on global function was mixed with no consistent benefit. Similarly, for quality of life/ADL outcomes, all three studies that evaluated this domain were not statistically significant but these were secondary outcomes and may reflect a lack of power. Behavior/mood and caregiver burden were not tested in these trials.

The quality scores for reporting adverse events were generally low, scoring 1 or 2 out of 5. Withdrawal rates due to adverse events varied from 1 - 5% for placebo and 12 - 55% in the treatment group, with one study not reporting rates. The high withdrawal rates were in studies with sample sizes that varied from 181 to 475 subjects. A single study tested for differences between groups, and found that dizziness, tremor, weight loss, asthenia, confusion, delirium, and respiratory problems were significantly different. The cluster of reported types of adverse events suggests that gastrointestinal problems (abdominal pain, diarrhea) (placebo = 1 - 9%, all doses = 13 - 28%), nausea and vomiting (placebo = 1 - 9%, all doses = 9 - 75%) and eating disorder (placebo = 2 - 6%, all doses = 5 - 16%) were most frequently reported. Dizziness (placebo = 4 - 13%, all doses = 11 - 38%) and agitation (placebo = 6 - 16%, all doses = 4 - 8%) were also reported.

Posatirelin. Four trials evaluated posatirelin in 931 subjects in a variety of mild to moderate dementia populations (AD, PDD, VaD) using a dose of 10 mg per day over 3 months duration.

Three of the four trials showed statistical significance for general cognitive function and quality of life/ADL (as measured by GBS subscales for these domains). The evidence remains

inconsistent for benefit in global assessment (evaluated in only one trial) and behavior/mood (mixed results). Caregiver burden and specific cognitive function were not evaluated in any trial.

Quality scores for reporting adverse events varied from 2 to 4. Withdrawal rates due to adverse events ranged from 0 - 3% in placebo and 0 - 4% in the treatment group. None of the studies tested for significant differences between groups. All studies reported the presence of agitation, and at least three studies reported arrhythmia, nausea/vomiting, headache, rash/skin disorder, and sleep disorder. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = 1 - 4%), 2) dizziness (placebo = not reported, all doses = 1%), 3) diarrhea (placebo = 2%, all doses = 2%), 4) agitation (placebo = 1 - 5%, all doses = 1 - 5%), and none reported eating disorder as an adverse event.

Rivastigmine. Six studies evaluated 2071 subjects and three of these studies were limited to AD patients only. Doses for rivastigmine varied from 1 to 12 mg, and treatment ranged from 14 to 26 weeks and only one study specified a community sample.

The evidence shows that general cognitive function improves with rivastigmine at a dose of 12 mg, but there is mixed results for efficacy at lower doses. Two trials also evaluated specific cognitive function but the results were not consistent within studies (between general and specific measures) and between studies for these domains. There is consistent evidence of benefit for the outcome of global assessment but the dosage at which this is significant varies highly between studies. In the domains of behavior/mood and quality of life/ADL, the findings were not statistically significant nor consistent; most of these analyses were not based on intention to treat analysis and lack of sufficient power cannot be ruled out. Caregiver burden outcomes were not evaluated by any trial.

Quality scores for reporting adverse events varied from 2 to 5. Withdrawal rates due to adverse events ranged from 4 - 11% in the placebo and 11 - 27% in the treatment group. Two trials demonstrated a dose response; however, one of these trials showed significant differences for nausea and vomiting only, and the other trial showed significant difference for all the adverse events reported. The majority of studies reported dizziness, nausea and vomiting, eating disorder/weight loss, and headache. It should be noted that one study allowed intentional prescribed anti-emetic drugs to increase the tolerance of subjects taking rivastigmine. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3 - 10%, all doses = 8 - 58%), 2) dizziness (placebo = 0 - 7%, all doses = 6 - 20 %), 3) diarrhea (placebo = 2 - 9%, all doses = 7 - 17%), 4) eating disorder (placebo = 4 - 8%, all doses = 4 - 19%), and 5) agitation was not reported.

Tacrine. Six studies^{108,109,110,111,112,113} evaluated tacrine in 994 subjects predominately with mild to moderate AD at doses of 80 to 160 mg lasting either 12 - 13 or 3 - 36 weeks in duration. Two other studies^{114,26} involving 425 patients were non-placebo controlled studies.

A single trial¹⁰⁸ was found to show benefit for general cognitive function with a small effect and this was based on a series of related publications. The five trials showing no benefit for general cognitive function comprised small sample sizes and much shorter study duration. Overall, the evidence for benefit for general cognitive function is limited to this single trial. There is evidence for benefit in global function from two of the three trials that evaluated this domain. Changes in behavior/mood, quality of life/ADL domains, specific cognitive function, and caregiver burden were all not significant, but lack of sufficient power cannot be ruled out.

The quality scores for reporting adverse events varied from 1 to 3. The proportion of subjects withdrawing due to adverse events ranged from 0 - 12% for placebo and 0 - 55% in the treatment group. The higher rates of withdrawal were associated with higher doses. Elevated alanine transaminase (ALT) or hepatic abnormality (placebo = 4 - 13%, all doses tacrine = 7 - 67%) was reported in six studies, raising concerns for the potential for serious liver damage. None of these trials tested for differences between treatment and placebo with respect to adverse events. Five of the studies reported nausea and vomiting (placebo = 0 - 9%, all doses = 9 - 37%); gastrointestinal problems; dizziness (placebo = 0 - 16%, all doses = 4 - 14%) was also noted in several studies. Frequencies of other a priori symptoms of interest are as follows: 1) agitation (placebo = 5 - 12%, all doses = 5 - 9%), and 2) diarrhea (placebo = 0 - 13%, all doses = 4 - 18%). There is evidence for the potential for serious adverse events associated with liver function in six trials.

Velnacrine. Three studies evaluated the effects of velnacrine in 774 AD patients with a diagnosis of AD. The doses that were shown to effect significant changes were 75 mg twice daily and 225 mg daily in studies with a 15 and 24 week duration. Location of recruitment was not specified.

Statistically significant effects were observed for general cognitive function, and global assessment in the two studies with sample sizes over 300 subjects. Behavior/mood and caregiver burden showed some benefit in one trial¹¹⁶ at the highest dose only. Quality of life/ADL was tested as a secondary outcome and showed mixed findings.

Quality scores for reporting adverse events were 3 for all studies. Withdrawal rates varied from 0 - 22% for the placebo group and 5 - 33% for the treatment group. None of the studies reported a dose response. None of the studies tested for statistical differences between the placebo and treatment groups. Two studies reported aberrant hematology and hepatic abnormality;^{116,117} for these two studies the rates of occurrence were 2 - 21% for placebo, and 32 - 40% for all doses. All studies reported diarrhea and nausea and vomiting. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 0 - 4%, all doses = 3 - 8%), 2) dizziness (placebo = 3%, all doses = 0 - 8%), 3) diarrhea (placebo = 3%, all doses = 2 - 33%), 4) agitation (placebo = 4%, all doses = 1 - 4%), and 5) eating disorder (placebo = 1%, all doses 2 - 4%). The potential for serious liver effects was not well specified in these trials.

Summary of Non-cholinergic Neurotransmitter/Neuropeptide Modifying Agents

Haloperidol. Five studies evaluated the effect of haloperidol relative to placebo in a total of 622 subjects with mild to moderate disease and included AD patients^{124,125,128} and mixed populations (MID/VaD/ PDD).^{126,127} One trial¹²⁸ had only 15 patients, and one trial¹²⁴ lasted only three weeks. Two studies recruited subjects from institutions; one from the community and two did not specify.

Mixed results were observed for improvement in global assessment. In three of the trials there was benefit in the domain of behavior/mood which reached statistical significance. Two trials evaluated caregiver burden and found no statistically significant differences but lack of sufficient power cannot be ruled out. Few studies evaluated outcomes in quality of life/ADL. Haloperidol did not affect general cognitive function in two trials and was not evaluated in the other studies.

The quality scores for reporting adverse events varied from 1 to 5 and only three of five studies reported withdrawal rates; the proportion of subjects withdrawing due to adverse events ranged from 5% to 17% for placebo and 17 – 33% in the treatment group. One trial¹²⁴ showed a dose-response effect but the study only lasted for three weeks. Three trials tested for differences between treatment and placebo with respect to extra pyramidal symptoms (placebo = 17 - 32%, all doses = 34 - 97%), and two found statistically significant differences.^{124,125} One study¹²⁵ found significant differences between groups for balance-related problems. Although reported by only two trials, the range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = not reported, and 2) dizziness (placebo = 24%, all doses = 21%). No frequencies were reported for agitation, diarrhea, or eating disorder.

Memantine. Three trials evaluated memantine in 1066 patients, primarily with VaD, with 10 or 20 mg doses lasting 12 or 28 weeks. Disease severity was moderate to severe in a single study¹³² and mild to moderate in the remaining two studies^{133,134}. One study included patients that were institutionalized, one from the community and the third did not specify.

There is consistent evidence of benefit for general cognitive function in the two studies that evaluated this domain. The findings for global assessment are mixed. The sole trial that evaluated mixed dementia populations (including some VaD) with moderate to severe dementia found significant differences for global function, behavior/mood, and quality of life/ADL outcomes, but did not evaluate general cognitive function. It should be noted that this trial with mixed populations used half the dose of memantine for half the study duration in patients with greater disease severity, and had approximately half the sample size of the other two trials evaluated in this systematic review. Despite a lower dose, a smaller number of more severely affected patients, and a shorter duration, a statistically significant difference was found.

The quality scores for reporting adverse events varied from 3 to 4. Only two of three studies reported withdrawal rates; the proportion of subjects withdrawing due to adverse events ranged from 7% to 13% for placebo and 9 - 12% in the treatment group. A single trial tested for differences between treatment and placebo, and none of the comparisons were significant. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 3%, all doses = 5%), 2) dizziness (placebo = 3 - 8%, all doses = 6 - 11%), 3) diarrhea (placebo = 4%, all doses = 4%), 4) agitation (placebo = 7 - 8%, all doses = 4 - 5%), and none reported eating disorder as an adverse event.

Selegiline. Six trials^{135,136,249,138,139,140} evaluated selegiline in 733 patients with AD, PDD, and DAT with 10 mg per day and a study duration of 60 days or 2 years. Only one study reported that patients were from institutional settings.

All but one trial that evaluated general cognition showed no statistically significant changes. A single trial found statistical improvements in specific cognitive tests (Sternberg Memory tests); this trial also showed statistically significant improvements in global assessment and behavior/mood. Only this trial, which had the highest quality score (7), showed consistently positive findings across domains tested. Three of the five trials that evaluated part or all of these domains had very small sample sizes and were likely underpowered, possibly accounting for the inconsistent findings. There is some evidence that selegiline and selegiline combined with vitamin E, increases the time to important functional decline milestones; this is based on a single study.

The quality scores for reporting adverse events varied from 0 to 3. The proportion of subjects withdrawing due to adverse events ranged from 0 - 4% for placebo and 0 - 9% in the treatment group. Only one trial¹³⁵ tested for differences between the treatment and placebo groups and showed that balance and falls were significantly different (worse) between groups (22% for the group with selegiline combined with vitamin E versus 5% in the placebo). When adjusted for multiple comparisons, these were no longer significant. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 2%, all doses = 0%), 2) dizziness (placebo = 2 - 20%, all doses = 0 - 30%), and 3) agitation (placebo = 4 - 16%, all doses = 4 - 23%); no trial reported diarrhea or eating disorder as an adverse event.

Summary of Other Pharmacological Agents

Cerebrolysin. Six studies evaluated the effect of cerebrolysin in a total of 819 subjects. All but one of the trials¹⁷¹ included only AD patients with mild to moderate disease. All of the studies used the same dose of cerebrolysin, 30 ml per day for 5 days per week for 4 to 24 weeks. Location of recruitment was not specified.

Cerebrolysin showed a statistically significant effect on cognition in four out of five studies. Although, a pooled estimate for the ADAS-cog was calculated, the model was positive for heterogeneity and the overall estimate was not significant. The results for specific cognitive tests for the three trials that evaluated this domain were inconsistent. Global assessment measures

showed a significant effect in five of the trials. This model was also positive for heterogeneity but significant for an overall effect. Two out of three studies showed an effect for behavior/mood, and none of the six studies showed an effect on quality of life/ADL. No study measured caregiver burden.

Two of the six trials scored 5 out of 5 on our quality scale for rating adverse events, yet they did not report any adverse events. Two studies scored 4, and the other two trials scored 3 and 2. All the studies with scores equals to 4 or less tested for statistical differences in adverse events between placebo and treatment groups. Withdrawals due to adverse events were not reported in one study¹⁷⁰ and were 1% in two studies^{173,168} and none withdrew in three studies.^{169,172,173} One study reported significant differences between treatment and control group¹⁷³ for weight change, anxiety, and headache. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 10 - 24%, all doses = 3 - 21%), 2) dizziness (placebo = 0 - 12, all doses = 1 - 8%), and 3) agitation (placebo = 1%, all doses = 0%), and none reported diarrhea or eating disorder as an adverse event.

Estrogen. Five studies evaluated estrogens for dementia in 247 patients with primarily mild to moderate AD, with the exception of one study¹⁷⁸ that included moderate to severe dementia patients who were all institutionalized. One of the studies with AD patients provided 0.10 mg per day¹⁷⁴ by skin patch for 8 weeks and the others used 1.25 mg per day for 12 to 52 weeks.¹⁷⁷ The study including severe subjects used 2.5 mg per day for 4 weeks.¹⁷⁸

Three trials evaluated general cognitive function and all showed non-significant findings; two of these trials lacked sufficient power for the ADAS-cog. Similarly, two trials evaluated specific cognitive function but results were mixed. Most of the outcomes evaluated in the domains of global assessment, behavior/mood, and quality of life/ADL were secondary outcomes and none showed significance; lack of power could be a factor in these trials.

One of the five trials scored 5 out of 5 on our quality scale for rating adverse events, and surprisingly, this same trial did not report any adverse event. Withdrawal rates due to adverse events ranged from 0 - 5% for placebo and 0 -14% for the treatment group. The most frequently reported adverse event was vaginal bleeding and a single trial reported a significant difference between placebo and treatment group for this symptom. It was not clear from the descriptions provided in the study if they had ascertained whether vaginal bleeding was present prior to the trial commencement. Nausea was the single a priori symptom of interest that was reported and by a single trial; frequencies varied from 0% for the placebo group and 4% for the treatment group.

Ginkgo biloba. Three trials evaluated Ginkgo biloba, 120 to 240 mg per day for 3 to 12 months, in a total of 563 subjects with mixed dementias of mild to moderate severity.

The largest trial¹⁷⁹ had the longest treatment interval but the lowest daily dosage and reported a significant effect for general cognitive function but had mixed findings for global assessment.

A second large trial¹⁸¹ found positive changes for neuropsychological tests, global assessment, and behavior outcomes with double the dosage of the previously described trial and half the treatment interval. In this same RCT, clinical efficacy was assessed by using a responder analysis, with therapy response being defined as response in at least two of the three variables: CGI (global function), SKT (special cognitive function), and NAB (ADL). A single trial evaluated behavior/mood and was not significant. No trial evaluated caregiver burden or quality of life/ADL.

All three trials scored 3 or greater on the quality scale for rating adverse events. Two studies had no withdrawals due to adverse events, and one trial had a withdrawal rate of 6% for both placebo and treatment groups. Two studies did not report any adverse event. One study reported a statistically significant difference between the treatment and the placebo group for skin disorders. The same study reported gastrointestinal and headache adverse effects, but did not test for statistical differences between the placebo and the treatment group. None of the trials reported the presence of the a priori symptoms of interest.

Idebenone. Four studies^{185,183,182,184} evaluated the drug idebenone in 1153 subjects of mixed dementia populations of mild to moderate severity; one of these trials²⁶ evaluated idebenone relative to tacrine. Doses varied from 30 mg per day to 360 mg per day, and the treatment interval ranged from 90 days to 60 weeks.

There was evidence of benefit for general cognitive function and global assessment. Several studies evaluated behavior/mood and quality of life/ADL and these outcomes were found to be significantly different. None of the trials evaluated caregiver burden.

Quality scores for reporting adverse events varied from 1 to 5. Rates of withdrawal due to adverse events varied from 0 - 5% for the placebo group and 0 - 5% in the treatment group; a single trial¹⁸³ did not report withdrawal rates. Two trials^{183,185} tested for statistical differences between groups and found no differences. Although no clear pattern emerges, three studies identified at least one balance-related adverse event most consistently reported across studies. The range of frequencies of the a priori symptoms of interest are as follows: 1) nausea (placebo = 2%, all doses = 2 - 11%), 2) dizziness (placebo = not reported, all doses = 2%), and 3) not reported for diarrhea, agitation, or eating disorder as an adverse event.

Oxiracetam. Five studies^{186,187,188,189,190} evaluated oxiracetam in 554 subjects with different dementia syndromes of mild to moderate severity. All analyses were observed cases and not ITT. All studies used 1600 mg daily, with one exception¹⁸⁹ where the dose ranged between 1600 - 2400 mg per day. The treatment interval ranged from 90 days to 26 weeks.

All outcomes shown to be positive for this drug were based on observed case evaluation. The two trials that evaluate general cognitive function showed benefit. The findings for specific cognitive function were mixed. A single trial evaluated global assessment and showed statistically significant change. Behavior/mood, and quality of life/ADL outcomes showed mixed results. No study evaluated caregiver burden.

The quality scores for reporting adverse events varied from 2 to 5. The proportion of withdrawals due to adverse events varied from 0 - 9% for the placebo group and 0 - 6% for the treatment group. No clear pattern for adverse events is evident, but three of the five studies reported gastrointestinal related problems, primarily associated with abdominal pain. Although, only single trials evaluated the range of frequencies of the a priori symptoms of interest are as follows: 1) dizziness (placebo = not reported, all doses = 11%), and 2) agitation (placebo = 1%, all doses = not reported); no trial reported nausea, eating disorder, or diarrhea as an adverse event.

Pentoxifylline. Three placebo-controlled studies^{193,192,191} evaluated pentoxifylline and one study¹⁹⁴ compared pentoxifylline to sulodexide, with a total of 482 subjects with predominately MID. The total dose administered in all studies was 1200 mg per day but varied from 400 mg three times per day to 1200 mg once per day. The treatment intervals ranged from 12 to 36 weeks.

All three placebo trials showed non-significant findings for any primary outcome evaluated on all subjects in the study. It should be noted that two of these trials had very small sample sizes (n = 38, n = 28) that were evaluated in the OC analyses; this suggests that the trials lacked sufficient power to evaluate multiple outcomes. The remaining trial had a large sample size (n = 289) and employed an ITT analysis; all primary outcomes evaluated were not significant.

The quality scores for reporting adverse events were generally low, varying from 1 to 3. Withdrawal rates due to adverse events varied from 0 - 25% in the placebo group and 0 - 22% in the treatment group. The two studies that reported adverse events indicated the presence of gastrointestinal disturbances, including abdominal pain or nausea and vomiting (placebo = 7% and all doses = 14%). None of the trials reported dizziness, agitation, eating disorder or diarrhea.

Propentofylline. Four trials^{197,196,250,198} using propentofylline in 510 patients with AD and VaD were included. A dose of 900 mg per day was consistent across all studies, and the treatment duration ranged from 3 to 12 months.

The two studies with small sample sizes (n = 30) showed no significant results for any outcome evaluated but lack of power cannot be ruled out. There were two trials that found benefit in general cognitive function based on the MMSE. The results for specific cognitive function as measured by the DSST were mixed, as were those for global assessment. Behavior/mood outcomes were evaluated by a single trial¹⁹⁵ and shown to be significant; this same trial evaluated quality of life/ADL and showed no significant difference. No trial evaluated caregiver burden.

The quality scores for reporting adverse events varied from 1 to 4. The percentage of withdrawals varied from 0 - 13% for the placebo group and 0 - 12% for the treatment group. None of the trials tested for differences between groups. Three of the trials^{195,197,198} reported gastrointestinal events that included abdominal pain, constipation, and nausea and vomiting (placebo = 2%, all doses = 7%). Dizziness (placebo = 3 - 5%, all doses = 1 - 6%) was the only other a priori symptom of interest.

Methodological Issues and Limitations in Assessing Efficacy of Dementia Agents

Definition of clinically significant or meaningful difference. The stance undertaken in this review has been cautious with regards to interpreting “clinically significant” differences within and across studies. This systematic review has highlighted some of the concerns expressed in the literature on pharmacological efficacy research in dementia. Ultimately, clinical significance is a complex issue, and its definition can vary across individuals and groups of individuals. Wherever possible, attempts were made to identify the magnitude of differences in the studies and the limitations of the data from some of these primary studies.

In drug development programs, an ordered series of trials are undertaken: dose tolerance (phase I), dose finding (phase II), dose efficacy (Phase III), and post-marketing (phase IV). However, due to the pressures on pharmaceutical companies to develop drugs quickly and cost-efficiently, a drug may move into the next phase of development before evidence of the previous phase is known.²⁵¹ Even when phase III trials are carried out in an adequate manner, the interpretation of the efficacy results is hampered by multiple p-values, disagreement over the need for multiplicity corrections, and the potential for conflicting evidence from trials of different sizes.²⁵¹ Some of these difficulties can be minimized by measuring a single primary efficacy variable at one point in time and using a p-value of less than 0.025 (one-tailed, as the aim is for the statistical test to determine if the drug performs better than the placebo or low dose).²⁵² This presumes that good dose-response data exist, identifying a single dose level as the best candidate for further evaluation. Lastly, interpreting differences on the basis of statistical significance has long been recognized as problematic. Clinically meaningful change reflects a different level of “significance” and often requires consensus among experts within the field for these criteria.

Issues of diagnosis and severity. Three methodological issues related to population classifications have limited the inferences that can be garnered from this systematic review. The first issue concerns the classification models used for diagnosing dementia; they are not interchangeable among the various types of dementia and the “pre-clinical” forms of slight cognitive impairment. Moreover, there are still concerns about the accuracy of these criteria. For example, in the American Academy of Neurology’s (AAN) recent evidence-based review of dementia case definitions, none met the AAN’s highest evidence standard. A clinical diagnosis of AD is only 28% specific after age 79 years. Similarly, no dementia screening measure is accurate enough to be recommended by the American Society of Internal Medicine.²⁵³ The AAN specifically faulted the emphasis on memory function in dementia case definitions.²⁵⁴ Yet tests like the ADAS-cog emphasize memory loss at the expense of other cognitive domains, especially executive control function,³¹ and many anti-dementia treatment strategies target neurotransmitters and structures (like acetylcholine and the hippocampus), which mediate memory test performance.

A second consideration in defining populations of dementia patients concerns determination of severity level. The MMSE, although frequently used, may not best capture severity. Many studies were observed to define the severity (mild, moderate, severe) of dementia populations based on the MMSE. For example, a range from 10 to 26 has been used to define a mild to moderate severity level.⁵⁰ Given that the maximum and minimum instrument scores are 0 and 30, this suggests that the extreme ends of the spectrum, particularly the “severe” end (i.e. <10), represent a very narrow proportion of patients. These two broad categories (mild to moderate and severe) may not actually reflect the cognitive and functional differences in a clinically meaningful manner. The MMSE does not address issues of executive control function (as required by the DSM-IV dementia case definition), which is known to be a good predictor of functional status. From a research perspective, a better classification reflecting disease severity may be an important factor for stratification and determining the efficacy of pharmacological interventions.

Outcome issues. The studies evaluated in our review used 181 different outcomes across seven domains. This raises the issue of which of these outcomes are considered by clinicians to be most “clinically relevant”. Let us assume that the most clinically relevant outcomes for all the drug interventions for dementia are the ADAS-cog and the MMSE because they are very commonly reported in studies.

In this dementia review, numerous studies did not measure outcomes evaluating cognition, as the intended effect of the drug was not always in the domain of cognition (e.g. neuroleptics for behavior control). Moreover, a large number of the studies that used important clinical cognition outcomes, such as the MMSE, did so only to establish baseline severity, or they used it as a secondary outcome. This presents us with some difficulty in the consistency of reporting on this limited set of “clinically relevant” outcomes. There is also the issue of which domain (i.e. cognitive function versus ADL versus behavior) is the most clinically relevant. The FDA guidelines suggest cognition and global assessment; the EMEA guidelines suggest the addition of an ADL or quality of life/ADL measure as being most clinically relevant. Thus, some consensus work needs to be done among experts in the field to determine the most clinically relevant outcomes and domains. For example, the choice of most clinically relevant outcome may depend upon type and stage of dementia (e.g. for mild AD, neuropsychological outcomes may be the most important domain while for severe AD, behavior may be the most relevant outcome), which may challenge the achievement of consensus.

To our knowledge, no specific set of outcomes that define “clinical relevance” applies to all the drug interventions we evaluated. The FDA has recommended that “dual efficacy” of dementia drug interventions be established by significant change in both a psychological measure and a global change measure. The outcomes measuring these attributes within these two domains were not specified. However, there was a general trend for using the outcomes ADAS-cog and CIBIC+ to capture these two attributes when evaluating drugs for AD populations.

Ideally, all outcomes should have demonstrated acceptable psychometric properties, such as reliability, validity (construct), and responsiveness. We did not a priori evaluate the properties of outcomes reported in the eligible studies. In some cases, these outcomes were developed in non-

English languages but the original study was reported in English. In considering the psychometric properties of some of the outcome instruments used, the attribute of responsiveness is critical, and some have suggested that this has not been adequately evaluated in many outcome measures.^{33,30,255}

We might envision a clinically relevant pharmacological treatment as one that has made a real difference, where the change is both relevant and important to the patient or to clinicians. This fundamentally shows the difference between clinically significant (relevant and important) versus statistically significant (associated with probabilities), where the latter determines that the results are not due to chance. Moreover, a clinically important change will vary depending on whether importance is defined from the patient or clinician perspective.

Five different levels of responsiveness (ability to detect change) of outcome measures have been defined:²⁵⁶ 1) Minimal change potentially detectable (essentially an attribute of the scoring method of the outcome), 2) Minimal change actually detectable beyond measurement error of the instrument (also defined as Minimum Detectable Change (MDI) or Reliability Change Index (RCI), which includes the Standard Error of the Measurement (SEM)), 3) Observed change (often reported as the standardized response mean (SRM) or effect size (ES). 4) Observed change in those estimated to have improved; the key to understanding change in this instance is that an external standard is used to determine whom has improved (often reported as comparison between groups that have improved versus those who have not; the improved group can be defined by either patient and/or clinician or a combination), and 5) Observed change in those estimated to have important improvement (often reported as the minimal clinically important difference and can be determined by the patient or clinician, or a combination of both).

Consider the ADAS-cog and the CIBIC+: The minimal change detectable is $1/70 = 0.0143$ for the ADAS-cog and $1/7 = 0.143$ for the CIBIC+, suggesting that the ADAS-cog can detect smaller increments of change relative to the CIBIC+. Thus different instruments have differing sensitivities to detecting change. There is scant literature on the responsiveness of outcome measures as defined in number 4 above, observed change in those that have improved, or as in number 5 above, observed change in those estimated to have important improvement. Thus, we have identified a significant gap in the literature with regard to estimating clinically important changes. Much greater consideration of issues of responsiveness should be given in future research in efficacy trials of pharmacological agents. Greater understanding of clinically important change suggests that some of our current judgments of efficacy are limited as these important differences need to be established.

Analysis issues. The inability to estimate the power of a study to detect a difference presented significant limitations in interpreting those studies that showed no significant differences. Similarly, the lack of sufficient data for estimating effect size limited the ability to show the magnitude of the change. It is recommended that future trials evaluating the efficacy of pharmacological agents adhere to the CONSORT guidelines in order to provide sufficient data to estimate power and effect size for all relevant outcomes.

Although the difficulty of maintaining adherence to long-term drug interventions among dementia patients is acknowledged, the ITT analysis should continue to be the analysis of choice in trials. Ideally, both ITT and OC analyses should be presented. If both suggested the same conclusion, confidence in the study results would be increased.

Problems with funding/ sponsorship exclusively from drug companies. The sponsorship of studies by for-profit organizations has led to bias towards the publishing of positive results.²⁵⁷ These findings suggest that there are powerful disincentives for pharmaceutical companies to publish negative trials. This is contrary to what academic-based, non-industry funded trials show, where the publication of negative trials are more likely.

A recent evaluation of FDA databases for antidepressant drugs²⁵⁸ in the US, suggested that less than half of antidepressant trials were negative, which does not correspond to the published literature. In this systematic review, no attempts were made to contact industry for unpublished trials, which introduces the possibility of a bias associated with not reporting negative trials. Additionally, we did not contact authors who did not specify funding sources for their studies. Future research on the efficacy of pharmacological agents to treat dementia should indicate all sources of funding and who undertook the study analyses.

Adverse events. In this systematic review, the type and frequency of adverse events associated with the use of a drug intervention were scrutinized and reported to a greater extent than previous reviews of anti-dementia drugs. Attempts were made to weigh the potential for harm against the benefits when determining the efficacy of pharmacological interventions. Empirical evidence across diverse medical fields indicates that reporting of safety information (including milder adverse events) receives much less attention than the positive efficacy outcomes.³⁵ Thus, it was recognized that an evaluation of the benefits of anti-dementia pharmacological agents alone may present a biased view of the efficacy of the intervention.

The ability to capture and evaluate adverse events proved to be difficult for several reasons. For example, although metrifonate had good evidence of positive effects on cognitive function, it was banned from use due to the risk of respiratory paralysis. The description of serious adverse events in the trials we evaluated did not capture this type of event, nor did different studies identify “serious events” in a consistent manner. This points to several fundamental limitations. The first of these relates to the limitation associated with the RCT design itself, which is less likely than the longitudinal cohort study designs to capture serious adverse events that are rare. Secondly, many trials were of relatively short duration and captured “idealized” dementia populations. Many of these trials were from pre-marketing studies contracted by pharmaceutical companies in carefully controlled research settings. Dementia patients seen in practice may have more complex medical illnesses and are at greater risk for potential side effects. In addition, drugs used in “polypharmacy” have even greater potential for pharmacological interactions. Furthermore, practitioners may prescribe these pharmacological agents for wider indications than originally intended, or may not refrain from withholding the drug from certain high-risk subgroups, leading to increased risk of adverse events. Thus, published rates of adverse events in well-controlled trials may underestimate true rates seen in practice.

Thirdly, by their nature, some adverse events are not easily anticipated, and therefore are not screened for in some trials. Adverse events may be hard to predict or anticipate but can be captured only if a trial protocol was designed to measure these events. This problem is compounded by the lack of consistency in what constitutes “serious” events or how the severity of the typical events is rated. A limited number of standardized instruments exist to capture these events reliably, but the overwhelming majority of studies in this systematic review did not use these instruments. Furthermore, capturing information from individuals with cognitive decline can create problems; the validity of the self-report instrument, even if completed by the caregiver, can be problematic. More research on the reliable collection of adverse events in dementia populations (with compromised cognition) may be required.

A fourth consideration concerns the issue of off-label use of pharmacological agents. Given that only four drugs are currently approved by the FDA for the treatment of dementia, the other 97 interventions evaluated in this review are classified as “off label use” but many are not approved by the FDA and not, therefore, available. For some of these off-label medications the potential mechanism of action on the disease process has not been fully established (if even considered), yet they have been applied to dementia populations. This off-label use of pharmacological agents may present further difficulties in evaluating adverse events.

Question 2: Does pharmacotherapy delay cognitive deterioration or delay disease onset of dementia syndromes?

Summary of Systematic Review Results

Few studies evaluated delay of onset or delay in disease progression. A definite gap for evaluating disease onset (as defined by the selection of populations at risk such as MCI populations) has been identified in this review.

Conversely, the need for good evaluation of disease progression in trials was also identified. In general, few studies evaluated subjects in more severe state of the disease. This suggests that a bias exists towards evaluating mild to moderate disease in the trials eligible in this systematic review. This in turn reflects the underlying assumption that the less severe groups are most likely to benefit from drug trials. Since so few studies have evaluated the more severe groups, this assumption may require some empirical justification in future research. Those studies that evaluated severe patients showed some potential for benefit. Future research in this area may require some consensus regarding the classification of severity levels.

Three studies evaluating cerebrolysin,¹⁶⁸ selegiline and vitamin E,¹³⁵ and donepezil⁶¹ have shown significant effects in delaying disease progress in mild to moderate^{61,168} and moderately severe disease in patients with AD. This delay in progress was expressed in terms of delay in days to primary event^{135,61} or statistical differences between placebo at a specified time interval.¹⁶⁸ Although these two trials coincidentally evaluated dementia patients over the longest time interval, it did not withdraw the drug at the end of the study. Theoretically, conclusive evidence of disease delay would be demonstrated if the treatment groups did not return to the level of the placebo. Thus, distinguishing between symptomatic and disease modifying effects is

not possible unless the drug is withdrawn and the treatment group(s) are observed for these changes.

When studies attempted to evaluate disease progression, long-term (1 year or greater) trials continued in an "open-label fashion", where blinding was no longer maintained. This limits the confidence that bias did not affect the subsequent changes in the outcomes. It was observed that increasing levels of dropout (for a variety of reasons) also plagued these open-label phases of evaluation. From a practical perspective, maintaining adherence in longer-term trials in dementia patients are challenging,¹⁹ particularly for those in the placebo arm or for those interventions that have a high proportion of adverse events.

A number of trial designs have been proposed to capture delay in disease progression versus symptomatic treatment. Some of these trial designs include withdrawal of treatment, active-extension, randomized withdrawal, randomized start, and staggered start designs.^{235,236,19,21} One important aspect of these designs is the selection of an adequate washout period or an adequate follow-up period. In addition, longer evaluation with survival analyses may be a good strategy to evaluate delay in disease progress for some drugs. One advantage of this design is the selection of clinically relevant milestones (functional changes over time), which was utilized in two studies^{63,136}; the selection of such events may merit greater consideration in future trials evaluating delay. A more critical analysis of the staggered/start/stagger withdrawal design in comparison to the survival analysis would be helpful. Also, one could provide a more extensive analysis of the data on propentofylline and vitamin E,¹³⁶ which represent the most extensive efforts to use the stagger/start/stagger withdrawal and survivor analysis approaches, respectively. Future research seeking to establish efficacy should clearly specify if symptomatic treatment or delay in progression is the therapeutic aim. This is important for determining specifically if efficacy is considered with respect to these two aims. Accordingly, a design that can establish this aim should be selected.


Methodological Issues

Determining symptomatic treatment versus affecting delay in disease progress. Figure 31 depicts hypothetical responses of dementia patients to two similar pharmacological interventions relative to placebo. In this example, the placebo group changes over time were modeled according to the natural history of AD as described by Stern et al. (1994)²⁹; the progressive decline of the AD subjects may not be representative of all dementia types. For simplicity's sake, the decline is assumed to be linear, although the literature has suggested the rate of decline varies between the different types of dementia and within each of these groups as a function of the disease severity.^{21,259} The two drugs depicted in Figure 31 are similar in that they have the identical titration (approximately 8 weeks) and washout periods. In this hypothetical scenario, the drugs are both withdrawn at 6 months (DW) and the washout periods have ended at 8 months. Within the active treatment period (first 6 months), the response to Drug I depicts the maintenance or stabilization of cognition function relative to the placebo, whereas the response to Drug II suggests improvement (or restoration) of cognition for a short period. However, the rapid decline of cognition scores within the two treatment groups to the level of placebo at 8 months (end of the washout period (EW)) suggests that the treatment effect was symptomatic relief. Upon withdrawal for subjects exposed to either Drug I (maintenance or stabilization) or

Drug II (improvement relative to baseline and placebo), the cognition scores declined to the same rate of placebo, and thus no delays in disease progression were demonstrated.


In contrast, Figure 32 shows a delayed rate of decline relative to placebo after the withdrawal of the pharmacological interventions. The response depicted for the treatment group exposed to Drug I shows that cognitive function is maintained until the drug is withdrawn (DW) and then the rate of decline is slower relative to the placebo (different slope of change) following the washout period. The response of the treatment group exposed to Drug II would suggest that cognition is improved for an interval (relative to baseline and placebo); when the drug is withdrawn, the rate of decline in cognitive function approximates that of the placebo group but is offset by approximately 6 months. Comparison of the slopes of the decline of cognition (Figure 32) would indicate a greater rate of decline for Drug II relative to Drug I, but both exemplify delay in progression of the disease effects. Theoretically, the treatment group rates of decline will never meet the decline rate of the placebo group when true disease modification has been effected by the pharmacological agent.

Figure 31. Delay of symptomatic treatment effects.



DW = Drug Withdrawn; EW = End of Washout

Figure 32. Delay in disease progression treatment effects.



DW = Drug Withdrawn; EW = End of Washout

In these two Figures (31, 32), the responses depicted have been idealized to clearly show the differences between symptomatic treatments versus disease modifying treatments. Additionally, two examples of the rate of decline as a characteristic of delay in disease progress have been explicated. However, in practical terms the most effective time interval for bringing about meaningful change in cognition (or other important outcomes), and the best time period to observe whether or not the effect is maintained (or lost), is not known. The difficulty in estimating these ideal time intervals is further compounded when the uncertainty of the rate of cognitive change (or decline) is considered.²¹ It is likely that treatment effects may not be equal across all stages of the disease (mild, moderate, severe) or between the various types of dementia diagnoses.

The evidence provided in dementia trials to demonstrate the three broad therapeutic aims of pharmacological interventions has been expressed in a variety of comparisons. Ideally the changes due to the pharmacological intervention would be expressed in terms of differences between the treatment and placebo groups. Surprisingly, many trials have reported statistical significance between baseline and endpoint of the treatment group(s) as evidence of a therapeutic effect. Change has been described as “improvement” relative to the baseline for either of the treatment or control groups. Although, it is unlikely that AD subjects would ever improve spontaneously relative to baseline, it may be possible in some dementias. Additionally, the magnitude of the “improvement” is dependent on the time interval for which the differences

were estimated. Consider Drug II in Figure 31 at the 4- and 6-month intervals; clearly, the magnitude of the difference is greatest at 4 months. Similarly, the evidence for “stabilization” or estimates for “delay in progression” was dependent on the interval used for evaluation.

Question 3: Are certain drugs, including alternative medicines (including non-pharmaceutical) more effective than others?

Summary of Systematic Review Results

What may be most relevant to clinicians are head to head comparisons of the cholinergic modifying neurotransmitter pharmacological agents, particularly those currently approved for the treatment of dementia (tacrine, rivastigmine, galantamine, donepezil) in the United States. The evidence for each of these drugs has been extensively detailed, and the relative merits and handicaps of each were outlined in chapter 3. Relative effectiveness as demonstrated by effect sizes for the ADAS-cog and the CIBIC were also shown in chapter 3. Although, the psychometric properties of these two outcomes are well accepted, comparison across the populations in these pooled estimates may not lend themselves to direct comparison across these four different specific drugs. Thus, inferences about the relative effectiveness of these four medications specific for the treatment of dementia should be made cautiously as head to head comparisons were not undertaken.

Relative efficacy must be evaluated in direct comparison trials

From a methodological perspective, addressing the question of being “more effective” requires head to head comparisons of pharmacological interventions.

An evaluation of the trials that undertook direct head to head comparison of two distinct pharmacological agents was limited because only seven trials were identified. Although, these trials may have shown some relative benefit of one drug versus another, the clinical relevance of these particular agents is limited as none of the drugs currently approved by the FDA specifically for the treatment of dementia is represented in these eligible studies. Moreover, these studies are essentially limited to single trials and are not sufficiently strong to base recommendations on the relative effectiveness of drugs. Head to head comparison studies are beginning to appear in abstract form only and a significant gap in the literature has been identified.

Question 4: Do certain patient populations benefit more from pharmacotherapy than others?

Summary of Systematic Review Results

In general, very few trials examined the efficacy of dementia drugs across different populations or population characteristics. From the 13 studies that reported stratified analyses, eight different variables were identified, which included age, gender, APOE genotype, disease type, disease severity (as determined by MMSE/ ADAS-cog threshold levels), treatment center, care dependence, and presence of depression. Additionally, three trials were identified that evaluated efficacy in 1) patients with Down's syndrome and dementia, 2) different ethnicities as a function of treatment center in a multicenter trial, and 3) depressed patients. Given the relatively small number of trials evaluating these variables within different populations and different pharmacological interventions, the findings of this review are limited with respect to these patient variables. These reflect merely what has been reported in the literature rather than variables of importance with respect to efficacy of pharmacological therapies. A significant gap in the literature has been identified.

Representativeness of populations in the drug trials

The study population characteristics were detailed for the trials evaluated. A recent study,²² suggests that many "real world" dementia patients in Ontario would not have met the eligibility criteria for participation in several of the cholinesterase inhibitor studies. This study highlights an important limitation of the pharmacological literature in that dementia patients recruited are not representative of the general dementia population. Additionally, clinicians and researchers should note that when a drug is approved for use, it is for a specific indication and a specific patient population. Evidence for one type of patient population may not necessarily be applied to another population. This is critical information to have when establishing clinical practice guidelines.

Question 5: What is the evidence-base for the treatment of vascular dementia?

Summary of Systematic Review Results

A total of 20 pharmacological interventions in 29 studies^{211,220,238,171,200,199,146,68,181,184,133,134,132,161,89,91,93,247,187,191,192,194,193,100,98,196,195,245,217} were applied specifically to VaD classified dementias. The majority of these pharmacological interventions (n = 14) were represented by single trials, these interventions included ateroid, buflomedil, cerebrolysin, sulphomucopolysaccharides (CDP choline), citalopram, donepezil, Ginkgo biloba, idebenone, minaprine, nimodipine, oxiracetam, 5-THF (trazodone), vincamine, and xantinolnicotinate. Six interventions had more than a single trial, and these included Choto-

san (n = 2), memantine (n = 3), nicergoline(n = 2), pentoxifylline (n = 4), posatirelin (n = 2), and propentofylline (n = 2). In general, when the drug interventions were shown to be effective, it was in the domains of cognitive function (both general and specific) and global assessment. Other domains were less frequently evaluated. Several trials attempted to test for differences between VaD groups and other dementia types.

Diagnosis Classification of VaD

Erkinjuntti et al (1997)¹ compared six commonly used classification schemes (DSM-III, DSM-III-R, DSM-IV, ICD-9, ICD-10 and the CAMDEX) and demonstrated that the prevalence of dementia can differ by a factor of 10 depending on the diagnostic criteria used. Two other studies have demonstrated that the prevalence of VaD varies with the classification system; therefore these criteria for diagnosis are not interchangeable.^{10,11}

There is controversy about the validity of the clinical classification of VaD, as autopsy confirmation often does not substantiate the clinical diagnosis.^{12,13} The majority of dementias were actually AD with co-existing VaD and PDD lesions.¹⁴ In contrast, the clinical accuracy of AD diagnosis is relatively high.⁷ Future research in vascular dementia should attempt to better distinguish this subgroup.

Determining Clinical Relevance

With rare exceptions, dementias are inevitably progressive and eventually lead to severe cognitive deficits, functional impairment, and often behavioral problems, unless death supervenes from intercurrent disease. The trajectories, sequence of clinical features, and burden on caregivers vary depending upon the type of dementia. For example, cognitive decline typically precedes functional impairment and behavioral disturbances in AD, while behavior and/or language problems typically announce the onset of frontotemporal degeneration.

Physicians and other health care practitioners have numerous roles in the management of individuals with dementia. These include identification, assessment and staging, classification, and prognostication, in addition to treatment of the individual and caregiver and planning for future disabilities (e.g. arranging alternatives to driving, assigning power of attorney and compiling living wills/advance directives).

Given these multiple tasks, how is the treating physician to interpret the results of therapeutic trials, which mostly deal with the pharmacological treatment of individuals with predominantly one type of dementia (AD) in the mild to moderate stages?

The traditional view of most physicians is that treatment success is measured by reversal of a disease, which is not a realistic goal in dementia. (While the older literature suggested that as many as 15 to 30% of dementias were “reversible,” more recent studies indicate that at most a few percent of dementias presenting to physicians are potentially reversible.)

Thus, the treating practitioner must begin by setting a realistic goal for therapeutic intervention. Symptom relief, alleviation of caregiver burden, prevention of complications (such as injury prevention or avoidance of aspiration pneumonia), and delay in progression of disease might be potential treatment targets. From this list, only symptom relief and delay in progression could be inferred from the studies examined in this systematic evidence review.

Outside the specialty clinic or clinical trial setting, most physicians have limited time and resources to expend on their patients with dementia. Few will have access to psychometrists or other individuals capable of administering extensive assessment instruments such as those used in clinical trials (e.g. ADAS-cog). Thus the typical practitioner must be able to complete a brief assessment, which provides sufficient information to determine whether a treatment is 1) indicated and 2) effective.

Deciding if a treatment is indicated depends upon the correct diagnosis (does this person have a dementia, and if so what type?), potential contraindications to the treatment (e.g. active peptic ulcer or heart block in the case of cholinesterase inhibitors), and the severity of disease. Determination of severity of dementia has given rise to several global scores such as the Global Deterioration Scale (GDS)²⁶⁰ and the Clinical Dementia Rating (CDR)²⁶¹. In practice, the Mini-Mental State Examination (MMSE)⁵⁰ (a short, 30-item, cognitive screening test) is frequently used as a measure of severity. Not only is it part of the usual diagnostic protocol for suspected dementia, but it also has the advantage of being included in the entry criteria of many of the RCTs of anti-dementia medications. It is therefore useful for determining whether a patient fulfils the appropriate severity criterion for therapeutic intervention.

With regard to deciding whether a treatment is effective, much has been written about the relative importance of statistically significant and clinically significant changes in measures of cognition, function, and behavior in dementia. A distinction must be drawn between clinically detectable change and clinically meaningful change. While psychometric measures (standardized instruments, which are highly reliable and relatively free from the influence of judgment) may detect changes too small to be appreciated by the clinician, clinometric tools (measures that are based on a clinical judgment about an individual patient²⁶²) may be considered more relevant to practice. Results expressed as a change from baseline measured by clinometric instruments such as the Clinicians Interview Based Impression of Change (CIBIC) or its derivative the CIBIC plus, which incorporates observations of the caregiver, mimic clinical practice more closely than most psychometric tools. The CIBIC aims to cover multiple domains relevant to the clinician (i.e. cognitive, functional, and behavioral). Clinicians may therefore interpret statistically significant changes on the CIBIC or similar scales with more confidence than changes on the many psychometric scales used in therapeutic trials. However, if an effect size of ~0.5 or greater is included in the analysis of psychometric outcomes, one can be reasonably confident of a robust response to the treatment under investigation.

Another measure of efficacy is the response rate—the percentage of study participants who experience an improvement (defined as a change of a specific magnitude on one or more scales.) This figure is useful for the clinician who may then indicate to the individual with dementia the chances of a positive outcome from the planned treatment.

Clinicians are faced with a bewildering array of results from clinical trials. Convergence of results (different studies of the same medication showing similar results) or studies of drugs in the same class showing similar results may help to reassure clinicians that the results are genuine. Conversely, when trials show differing results, clinicians should be especially vigilant in accepting only the results of the more positive trials.

As always, the translation of clinical trial evidence into practice demands careful scrutiny by the practitioner. Attention to external validity (is my patient sufficiently similar to those in the clinical trial that I can expect the same result from treatment?), interpretation of the outcome measures (clinically as well as statistically significant benefit), and weighing potential risks against potential benefits remain the responsibility of the treating practitioner.

Limitations of the McMaster AHRQ Review

A systematic review that has evaluated 91 pharmacological interventions in 186 RCTs with high internal validity has several limitations. The studies selected for this review are English-language trials. Based on our search results, we estimate that we could have potentially retrieved 1385 foreign-language articles (after de-duplication 1213) distributed among databases as follows: 346 from Cochrane Central, 444 from EMBASE, 559 from MEDLINE/PreMedline® 36 from other databases before review for title and abstract. If we assumed the same rate of potentially eligible studies for these non-English studies, an additional 16 non-English studies may have been eligible for review. It is possible that agents, such as Ginkgo biloba, may have had important trials published in non-English languages. The budget and timelines available, however, were a limiting factor to obtaining, translating, and abstracting non-English trials.

Secondly, no contact with authors of the eligible trials was undertaken to collect additional unpublished studies or provide results/data that were not presented in the published article. Although contact with the original authors of the trials (to supplement the missing information from the included studies) could have compensated for many of the reporting challenges we encountered, this strategy was not feasible given the timeline of this systematic review. Our experience at the McMaster EPC suggests that the majority of authors do not respond in a timely fashion if at all. Additionally, efforts were not made to contact industry for unpublished trials. It is likely that industry sponsors of trials that are not published in the public domain are under no obligation to share trials (particularly negative trials). Not contacting authors of eligible trials for additional data and not attempting to locate unpublished trials (either by other authors/ experts or by industry) may introduce publication bias in this systematic review.

Thirdly, we employed two eligibility criteria that may account for some differences in acceptance of well-known studies. The first of these was a minimum threshold for quality score as determined by the modified Jadad scale. Despite the fact that this scale has excellent reliability and content validity, some may argue that the threshold score of 3 is arbitrary and may have unnecessarily eliminated studies of historical importance. It is our view that given the amount of literature available, all efforts should be aimed at selecting only the trials with the highest internal validity rather than selecting the largest number of eligible trials.

The second eligibility criteria concerned the exclusion of crossover trials. Although crossover trials are suitable for chronic diseases, they may be prone to period effects or period-by-treatment interactions. Period effects are systematic changes in the outcome that apply to all patients due to temporal changes in the disease or to the measurement instrument. Period-by-treatment interactions occur when the efficacy of the intervention varies by period. This is a significant concern for studies that attempt to show disease modification and are carried out over a longer period of time. Additionally, a carry-over effect may occur if the washout period is not adequate. In addition to the weaknesses of this design, some limitations arise when considering the potential for meta-analytic analyses. Traditionally, first period data from a crossover trial are abstracted and can be potentially combined with parallel trials for analyses of a pooled estimate; the reporting of the study results (positive or negative) would also be based on this first period data. In a preliminary phase of the review, several crossover trials were examined. It was noted that many did not report first period data, which precludes any potential for combining with parallel trials; many trials also did not undertake statistical tests during the first experience, thus making it difficult to report the direction of the findings, even if the trial could be combined. Finally, the TEP considered the fact that this systematic review was evaluating a variety of drug interventions administered over differing time intervals, and so period effects might be an important source of bias. For all these reasons, the TEP made the decision to exclude crossover trials from this systematic review. Thus, this review is limited to evidence based on high-quality parallel trials only.

A final limitation to our study was the use of a checklist developed to address the issue of quality of reporting adverse events. The Jadad scale was not designed to evaluate the quality of reporting adverse events. Thus, when determining the “harms” or risks associated with an intervention, the quality or “internal validity” of collecting and reporting these adverse events needed to be evaluated. Although our checklist has face validity, it has not undergone formal psychometric testing.

Future Research Recommendations

The findings of this report suggest several important areas for future research on pharmacological treatments for dementia. These include:

Analytic Framework of the intended aim of the therapy on the disease

- Better conceptualization and research design to capture “delay in progression”.
- Clearer consensus on defining efficacy (benefits and clinically important change).
- Longer term studies (> 12 months).

Potential for bias

- Clarification of the role of industry sponsorship; one recommendation should be that all studies are required to disclose such information in future, including who analyzed the results.
- More concerted effort to incorporate unpublished studies and negative trials in future reviews.

Population

- Inclusion of the spectrum of severity in the patient populations (there is nothing to suggest that severe patients may not benefit from pharmacotherapy aimed at cognitive function improvement).
- The need for validation of trials and testing processes within cultures other than the traditional white population.
- Examining the efficacy of interventions in different sub-populations (age, disease severity levels, etc.).
- Better measurement and reporting of important patient characteristics (including baseline cognition scores, co-morbid conditions, the use of other medications, etc.).
- Inclusion of MCI type groups of subjects to evaluate “delay of onset”.

Outcomes

- Expansion of outcomes collected to include more than just cognitive function, and especially include caregiver burden and quality of life/ADL.
- Clear operational definitions for determining critical outcomes (delay to onset, delay to progression, important effect size, etc).
- Better understanding of how outcomes perform cross-culturally.
- Production of other diagnostic instruments to detect both onset and responses to therapies across varied cultural groups.
- Improvement in the reporting of adverse events to evaluate harm.

Analysis

- Appropriate analytical strategies that take into account intention to treat (ITT)/ last observation carried forward (LOCF) analyses; where possible both observed case and ITT/LOCF analyses should be presented.
- Sufficient data to estimate effect size, taking into account variability in both treated and control populations on the primary measures.
- Reporting the power of the study when findings are non-significant.

Intervention

- Undertake more studies with direct comparison of drugs to determine the relative efficacy of agents.
- Improved description of the titration process.
- Improved collection of adverse events undertaken in a systematic fashion with standardized instruments.

References

1. Erkinjuntti T, Ostbye T, Steenhuis R, et al. The effect of different diagnostic criteria on the prevalence of dementia. *N Engl J Med* 1997 Dec 4; 337(23):1667-74.
2. Jorm AF, Korten AE, Henderson AS. The prevalence of dementia: a quantitative integration of the literature. *Acta Psychiatr Scand* 1987 Nov; 76(5):465-79.
3. Brookmeyer R, Gray S, Kawas C. Projections of Alzheimer's disease in the United States and the public health impact of delaying disease onset. *Am J Public Health* 1998 Sep; 88(9):1337-42.
4. Canadian Study of Health and Aging Working Group. Canadian study of health and aging: study methods and prevalence of dementia. *CMAJ* 1994 Mar 15; 150(6):899-913.
5. Baumgarten M, Hanley JA, Infante-Rivard C, et al. Health of family members caring for elderly persons with dementia. A longitudinal study. *Ann Intern Med* 1994 Jan 15; 120(2):126-32.
6. The Canadian Study of Health and Aging Working Group. Patterns of caring for people with dementia in Canada: The Canadian Study of Health and Aging. *Can J Aging* 1994; 13:470-87.
7. Morris JC, McKeel DW, Jr., Fulling K, et al. Validation of clinical diagnostic criteria for Alzheimer's disease. *Ann Neurol* 1988 Jul; 24(1):17-22.
8. Tierney MC, Fisher RH, Lewis AJ, et al. The NINCDS-ADRDA Work Group criteria for the clinical diagnosis of probable Alzheimer's disease: a clinicopathologic study of 57 cases. *Neurology* 1988 Mar; 38(3):359-64.
9. McKhann G, Drachman D, Folstein M, et al. Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984 Jul; 34(7):939-44.
10. Pohjasvaara T, Mantyla R, Ylikoski R, et al. Comparison of different clinical criteria (DSM-III, ADDTC, ICD-10, NINDS-AIREN, DSM-IV) for the diagnosis of vascular dementia. National Institute of Neurological Disorders and Stroke-Association Internationale pour la Recherche et l'Enseignement en Neurosciences. *Stroke* 2000 Dec; 31(12):2952-7.
11. Chui HC, Mack W, Jackson JE, et al. Clinical criteria for the diagnosis of vascular dementia: A multicenter study of comparability and interrater reliability. *Arch Neurol* 2000 Feb; 57(2):191-6.
12. Galasko D, Hansen LA, Katzman R, et al. Clinical-neuropathological correlations in Alzheimer's disease and related dementias. *Arch Neurol* 1994 Sep; 51(9):888-95.
13. Nolan KA, Lino MM, Seligmann AW, et al. Absence of vascular dementia in an autopsy series from a dementia clinic. *J Am Geriatr Soc* 1998 May; 46(5):597-604.
14. Lim A, Tsuang D, Kukull W, et al. Clinico-neuropathological correlation of Alzheimer's disease in a community-based case series. *J Am Geriatr Soc* 1999 May; 47(5):564-9.
15. Schroder J, Kratz B, Pantel J, et al. Prevalence of mild cognitive impairment in an elderly community sample. *J Neural Transm Suppl* 1998; 54:51-9.
16. Petersen RC, Smith GE, Waring SC, et al. Aging, memory, and mild cognitive impairment. *Int Psychogeriatr* 1997; 9 Suppl 1:65-9.
17. Petersen RC. Mild cognitive impairment: transition between aging and Alzheimer's disease. *Neurologia* 2000 Mar; 15(3):93-101.
18. Ritchie K, Artero S, Touchon J. Classification criteria for mild cognitive impairment: a population-based validation study. *Neurology* 2001 Jan 9; 56(1):37-42.
19. Whitehouse PJ, Kittner B, Roessner M, et al. Clinical trial designs for demonstrating disease-course-altering effects in dementia. *Alzheimer Dis Assoc Disord* 1998 Dec; 12(4):281-94.
20. Rockwood K, MacKnight C. Assessing the clinical importance of statistically significant improvement in anti-dementia drug trials. *Neuroepidemiology* 2001 May; 20(2):51-6.

21. Leber P. Slowing the progression of Alzheimer disease: methodologic issues. *Alzheimer Dis Assoc Disord* 1997; 11 Suppl 5:S10-21; discussion S37-9.:S10-S21
22. Gill SS. Representation and eligibility of real-world subjects with dementia in clinical trials of donepezil. *Geriatrics Today: Journal of the Canadian Geriatrics Society* 2003; 6(2):67
23. Leber P. Guidelines for the clinical evaluation of anti-dementia drugs. 1st draft. Rockville,MD: US Food and Drug Administration; 1990
24. Rockwood K, Joffres C. Improving clinical descriptions to understand the effects of dementia treatment: Consensus recommendations. *Int J Geriatr Psychiatry* 2002; 17(11):1006-11.
25. European Medicine Evaluation Agency (EMA). Note for guidance on medicinal products in the treatment of Alzheimer's disease. London: EMA; 1997
26. Gutzmann H, Kuhl KP, Hadler D, et al. Safety and efficacy of idebenone versus tacrine in patients with Alzheimer's disease: Results of a randomized, double-blind, parallel-group multicenter study. *Pharmacopsychiatry* 2002 Jan; 35(1):12-8.
27. Mayeux R, Sano M. Drug therapy: Treatment of Alzheimer's disease. *N Engl J Med* 1999; 341(22):1670-9.
28. Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. *Am J Psychiatry* 1984 Nov; 141(11):1356-64.
29. Stern RG, Mohs RC, Davidson M, et al. A longitudinal study of Alzheimer's disease: measurement, rate, and predictors of cognitive deterioration. *Am J Psychiatry* 1994 Mar; 151(3):390-6.
30. Demers L, Oremus M, Perrault A, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Psychometric properties of functional and quality of life scales. *J Geriatr Psychiatry Neurol* 2000; 13(4):170-80.
31. Royall DR, Lauterbach EC, Cummings JL, et al. Executive control function: a review of its promise and challenges for clinical research. A report from the Committee on Research of the American Neuropsychiatric Association. *J Neuropsychiatry Clin Neurosci* 2002; 14(4):377-405.
32. Oremus M, Perrault A, Demers L, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Psychometric properties of global scales. *J Geriatr Psychiatry Neurol* 2000; 13(4):197-205.
33. Demers L, Oremus M, Perrault A, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Introduction. *J Geriatr Psychiatry Neurol* 2000; 13(4):161-9.
34. Perrault A, Oremus M, Demers L, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Psychometric properties of behavior and mood scales. *J Geriatr Psychiatry Neurol* 2000; 13(4):181-96.
35. Ioannidis JP, Lau J. Improving safety reporting from randomised trials. *Drug Saf* 2002; 25(2):77-84.
36. Lasagna L. Balancing risks versus benefits in drug therapy decisions. *Clin Ther* 1998; 20 Suppl C:C72-C79
37. The Cochrane Non Randomized Studies Method Group: Adverse effects subgroup. Proposed draft addition to Cochrane Handbook: Including adverse effects in Cochrane reviews. Internet. <http://www.dsru.org/wwwboard/latestdraft.pdf>
38. Fergusson D, Aaron SD, Guyatt G, et al. Post-randomisation exclusions: the intention to treat principle and excluding patients from analysis. *BMJ* 2002 Sep 21; 325(7365):652-4.
39. Pocock SJ, Abdalla M. The hope and the hazards of using compliance data in randomized controlled trials. *Stat Med* 1998 Feb 15; 17(3):303-17.
40. Lefebvre C, Clarke MJ. Identifying randomised trials. In: Egger M, Smith DG, Altman DG, editors. *Systematic Reviews in Health Care: Meta-analysis in context*, 2nd Edition London: BMJ Books; 2001. Chapter 4 p. 69-86.
41. Anonymous. The International classification of diseases, 9th revision, clinical modification: ICD-9-CM. 3rd edition. Washington, DC: US Department of Health and Human Services; 1989

42. World Health Organization. The tenth revision of the International Classification of Diseases and relative health problems (ICD-10). Geneva: WHO; 1992
43. American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 3rd ed. Washington, DC: American Psychiatric Association; 1980
44. American Psychiatric Association. Diagnostic and statistical manual of mental disorders (3rd ed. rev.): DSM-III-R. Washington, DC: American Psychiatric Association; 1987
45. American Psychiatric Association. Diagnostic criteria from DSM-IV. 1994
46. Roman GC, Tatemichi TK, Erkinjuntti T, et al. Vascular dementia: diagnostic criteria for research studies. Report of the NINDS-AIREN International Workshop. *Neurology* 1993 Feb; 43(2):250-60.
47. Petersen RC, Stevens JC, Ganguli M, et al. Practice parameter: Early detection of dementia: Mild cognitive impairment (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001 May; 56(9):1133-42.
48. Graham JE, Rockwood K, Beattie BL, et al. Prevalence and severity of cognitive impairment with and without dementia in an elderly population. *Lancet* 1997 Jun 21; 349(9068):1793-6.
49. Graham JE, Rockwood K, Beattie BL, et al. Standardization of the diagnosis of dementia in the Canadian Study of Health and Aging. *Neuroepidemiology* 1996; 15(5):246-56.
50. Folstein MF, Folstein SE, McHugh PR. "Minimal state". A practical method for grading the cognitive state of patients for the clinician. *J Psychiatr Res* 1975 Nov; 12(3):189-98.
51. Hachinski VC, Iliff LD, Zilhka E, et al. Cerebral blood flow in dementia. *Arch Neurol* 1975 Sep; 32(9):632-7.
52. Oremus M, Wolfson C, Perrault A, et al. Interrater reliability of the modified Jadad quality scale for systematic reviews of Alzheimer's disease drug trials. *Dement Geriatr Cogn Disord* 2001 May; 12(3):232-6.
53. Harris RP, Helfand M, Woolf SH, et al. Current methods of the US Preventive Services Task Force: a review of the process. *Am J Prev Med* 2001 Apr; 20(3 Suppl):21-35.
54. Thal LJ, Calvani M, Amato A, et al. A 1-year controlled trial of acetyl-L-carnitine in early-onset AD. *Neurology* 2000a; 55(6):805-10.
55. Thal LJ, Carta A, Clarke WR, et al. A 1-year multicenter placebo-controlled study of acetyl-L-carnitine in patients with Alzheimer's disease. *Neurology* 1996a; 47(3):705-11.
56. Sano M, Bell K, Cote L, et al. Double-blind parallel design pilot study of acetyl levocarnitine in patients with Alzheimer's disease. *Arch Neurol* 1992 Nov; 49(11):1137-41.
57. Livingston GA, Sax KB, McClenahan Z, et al. Acetyl-L-carnitine in dementia. *Int J Geriatr Psychiatry* 1991; 6(12):853-60.
58. Spagnoli A, Lucca U, Menasce G, et al. Long-term acetyl-L-carnitine treatment in Alzheimer's disease. *Neurology* 1991 Nov; 41(11):1726-32.
59. Rai G, Wright G, Scott L, et al. Double-blind, placebo-controlled study of acetyl-L-carnitine in patients with Alzheimer's disease. *Curr Med Res Opin* 1990; 11(10):638-47.
60. Prasher VP, Huxley A, Haque MS, et al. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Down syndrome and Alzheimer's disease: Pilot study. *Int J Geriatr Psychiatry* 2002 Mar; 17(3):270-8.
61. Mohs RC, Doody RS, Morris JC, et al. A 1-year, placebo-controlled preservation of function survival study of donepezil in AD patients. *Neurology* 2001 Aug 14; 57(3):481-8.
62. Winblad B, Engedal K, Soininen H, et al. A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology* 2001b; 57(3):489-95.
63. Feldman H, Gauthier S, Hecker J, et al. A 24-week, randomized, double-blind study of donepezil in moderate to severe Alzheimer's disease. *Neurology* 2001 Aug 28; 57(4):613-20.
64. Tariot PN, Cummings JL, Katz IR, et al. A randomised, double-blind, placebo-controlled study of the efficacy and safety of Donepezil in patients with Alzheimer's disease in the

- nursing home setting. *J Am Geriatr Soc* 2001a; 49(12):1590-9.
65. Burns A, Rossor M, Hecker J, et al. The effects of donepezil in Alzheimer's disease - results from a multinational trial. *Dement Geriatr Cogn Disord* 1999 May; 10(3):237-44.
 66. Rogers SL, Farlow MR, Doody RS, et al. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. *Neurology* 1998b; 50(1):136-45.
 67. Rogers SL, Doody RS, Mohs RC, et al. Donepezil improves cognition and global function in Alzheimer disease: A 15-week, double-blind, placebo-controlled study. *Arch Intern Med* 1998a; 158(9):1021-31.
 68. Pratt RD, Perdomo CA. Donepezil-treated patients with probable vascular dementia demonstrate cognitive benefits. *Ann N Y Acad Sci* 2002 Nov; 977:513-22.:513-22.
 69. Rogers SL, Friedhoff LT. The efficacy and safety of donepezil in patients with Alzheimer's disease: Results of a US Multicentre, Randomized, double-blind, placebo-controlled trial. *The Donepezil Study Group. Dementia* 1996 Nov; 7(6):293-303.
 70. Thomas A, Iacono D, Bonanni L, et al. Donepezil, rivastigmine, and vitamin E in Alzheimer disease: A combined P300 event-related potentials/neuropsychologic evaluation over 6 months. *Clin Neuropharmacol* 2001 Jan; 24(1):31-42.
 71. Erkinjuntti T, Kurz A, Gauthier S, et al. Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: A randomised trial. *Lancet* 2002 Apr 13; 359(9314):1283-90.
 72. Rockwood K, Mintzer J, Truyen L, et al. Effects of a flexible galantamine dose in Alzheimer's disease: a randomised, controlled trial. *J Neurol Neurosurg Psychiatry* 2001; 71(5):589-95.
 73. Wilkinson D, Murray J. Galantamine: A randomized, double-blind, dose comparison in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 2001 Sep; 16(9):852-7.
 74. Tariot PN, Solomon PR, Morris JC, et al. A 5-month, randomized, placebo-controlled trial of galantamine in AD. *The Galantamine USA-10 Study Group. Neurology* 2000 Jun 27; 54(12):2269-76.
 75. Wilcock GK, Lilienfeld S, Gaens E. Efficacy and safety of galantamine in patients with mild to moderate Alzheimer's disease: Multicentre randomised controlled trial. *Galantamine International-1 Study Group. BMJ* 2000 Dec 9; 321(7274):1445-9.
 76. Raskind MA, Peskind ER, Wessel T, et al. Galantamine in AD: A 6-month randomized, placebo-controlled trial with a 6-month extension. *The Galantamine USA-1 Study Group. Neurology* 2000 Jun 27; 54(12):2261-8.
 77. Gelinas I, Gauthier L, McIntyre M, et al. Development of a functional measure for persons with Alzheimer's disease: the disability assessment for dementia. *Am J Occup Ther* 1999 Sep; 53(5):471-81.
 78. Becker RE, Colliver JA, Markwell SJ, et al. Double-blind, placebo-controlled study of metrifonate, an acetylcholinesterase inhibitor, for Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1996; 10(3):124-31.
 79. Becker RE, Colliver JA, Markwell SJ, et al. Effects of metrifonate on cognitive decline in Alzheimer's disease: A double-blind, placebo-controlled, 6-month study. *Alzheimer Dis Assoc Disord* 1998 Mar; 12(1):54-7.
 80. Cummings J, Bieber F, Mas J, et al. Metrifonate in Alzheimer's disease results of a dose finding study. *Alzheimers Dis Biol Diagn Ther* 1997:665-9.
 81. Cummings JL, Cyrus PA, Bieber F, et al. Metrifonate treatment of the cognitive deficits of Alzheimer's disease. *Neurology* 1998b; 50(5):1214-21.
 82. Dubois B, McKeith I, Orgogozo JM, et al. A multicentre, randomized, double-blind, placebo-controlled study to evaluate the efficacy, tolerability and safety of two doses of metrifonate in patients with mild-to-moderate Alzheimer's disease: The MALT study. *Int J Geriatr Psychiatry* 1999 Nov; 14(11):973-82.
 83. Jann MW, Cyrus PA, Eisner LS, et al. Efficacy and safety of a loading-dose regimen versus a no-loading-dose regimen of metrifonate in the symptomatic treatment of Alzheimer's disease: A randomized, double-masked, placebo-controlled trial. *Clin Ther* 1999; 21(1):88-102.
 84. Morris JC, Cyrus PA, Orazem J, et al. Metrifonate benefits cognitive, behavioral, and

- global function in patients with Alzheimer's disease. *Neurology* 1998 May; 50(5):1222-30.
85. Pettigrew LC, Bieber F, Lettieri J, et al. Pharmacokinetics, pharmacodynamics, and safety of metrifonate in patients with Alzheimer's disease. *J Clin Pharmacol* 1998 Mar; 38(3):236-45.
 86. Raskind MA, Cyrus PA, Ruzicka BB, et al. The effects of metrifonate on the cognitive, behavioral, and functional performance of Alzheimer's disease patients. *J Clin Psychiatry* 1999; 60(5):318-25.
 87. Nordberg A, Svensson AL. Cholinesterase inhibitors in the treatment of Alzheimer's disease: a comparison of tolerability and pharmacology. *Drug Saf* 1998 Dec; 19(6):465-80.
 88. Winblad B, Bonura ML, Rossini BM, et al. Nicergoline in the treatment of mild-to-moderate Alzheimer's disease: A European multicentre trial. *Clin Drug Investig* 2001a(9):621-32.
 89. Herrmann WM, Stephan K, Gaede K, et al. A multicenter randomized double-blind study on the efficacy and safety of nicergoline in patients with multi-infarct dementia. *Dement Geriatr Cogn Disord* 1997 Jan; 8(1):9-17.
 90. Nappi G, Bono G, Merlo P, et al. Long-term nicergoline treatment of mild to moderate senile dementia. Results of a multicentre, double-blind, placebo-controlled study. *Clin Drug Investig* 1997(6):308-16.
 91. Saletu B, Paulus E, Linzmayer L, et al. Nicergoline in senile dementia of Alzheimer type and multi-infarct dementia: A double-blind, placebo-controlled, clinical and EEG/ERP mapping study. *Psychopharmacologia* 1995 Feb; 117(4):385-95.
 92. Schneider F, Popa R, Mihalas G, et al. Superiority of antagonistic-stress composition versus nicergoline in gerontopsychiatry. *Ann N Y Acad Sci* 1994 Jun; 717:332-42.
 93. Saletu B, Anderer P, Semlitsch HV. Relations between symptomatology and brain function in dementias: Double-blind, placebo-controlled, clinical and EEG/ERP mapping studies with nicergoline. *Dement Geriatr Cogn Disord* 1997; Vol 8(Suppl 1):12-21.
 94. Moller HJ, Hampel H, Hegerl U, et al. Double-blind, randomized, placebo-controlled clinical trial on the efficacy and tolerability of a physostigmine patch in patients with senile dementia of the Alzheimer type. *Pharmacopsychiatry* 1999 May; 32(3):99-106.
 95. Thal LJ, Schwartz G, Sano M, et al. A multicenter double-blind study of controlled-release physostigmine for the treatment of symptoms secondary to Alzheimer's disease. *Neurology* 1996b; 47(6):1389-95.
 96. Thal LJ, Ferguson JM, Mintzer J, et al. A 24-week randomized trial of controlled-release physostigmine in patients with Alzheimer's disease. *Neurology* 1999 Apr 12; 52(6):1146-52.
 97. Van Dyck CH, Newhouse P. Extended-release physostigmine in Alzheimer's disease: A multicenter, double-blind, 12-week study with dose enrichment. *Arch Gen Psychiatry* 2000; 57(2):157-64.
 98. Ferrari E, Cucinotta D, Albizzati MG, et al. Effectiveness and safety of posatirelin in the treatment of senile dementia: A multicenter, double-blind, placebo-controlled study. *Arch Gerontol Geriatr* 1998; 27(Suppl 6):163-74.
 99. Gasbarrini G, Stefanini G, Addolorato G, et al. Posatirelin for the treatment of degenerative and vascular dementia: Results of explanatory and pragmatic efficacy analyses. *Arch Gerontol Geriatr* 1997; 26(1):33-47.
 100. Parnetti L, Ambrosoli L, Agliati G, et al. Posatirelin in the treatment of vascular dementia: A double-blind multicentre study vs placebo. *Acta Neurol Scand* 1996 Jun; 93(6):456-63.
 101. Parnetti L, Ambrosoli L, Abate G, et al. Posatirelin for the treatment of late-onset Alzheimer's disease: A double-blind multicentre study vs citicoline and ascorbic acid. *Acta Neurol Scand* 1995 Aug; 92(2):135-40.
 102. Agid Y, Dubois B, Anand R, et al. Efficacy and tolerability of rivastigmine in patients with dementia of the Alzheimer type. *Curr Ther Res Clin Exp* 1998; 59(12):837-45.
 103. Corey-Bloom JR, Anand JV, Veach J, et al. A randomized trial evaluating the efficacy and safety of ENA 713 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe Alzheimer's disease. *Int J Geriatr Psychopharmacol* 1998; 1:55-65.

104. Rosler M, Anand R, Cicin-Sain A, et al. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: International randomised controlled trial. *BMJ* 1999 Mar 6; 318(7184):633-8.
105. Forette F, Anand R, Gharabawi G. A phase II study in patients with Alzheimer's disease to assess the preliminary efficacy and maximum tolerated dose of rivastigmine (Exelon registered). *Eur J Neurol* 1999; 6(4):423-9.
106. McKeith I, Del Ser T, Spano P, et al. Efficacy of rivastigmine in dementia with Lewy bodies: A randomised, double-blind, placebo-controlled international study. *Lancet* 2000 Dec 16; 356(9247):2031-6.
107. Potkin SG, Anand R, Fleming K, et al. Brain metabolic and clinical effects of rivastigmine in Alzheimer's disease. *Int J Neuropsychopharmacol* 2001 Sep; 4(3):223-30.
108. Knapp MJ, Knopman DS, Solomon PR, et al. A 30-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. The Tacrine Study Group. *JAMA* 1994b; 271(13):985-91.
109. Maltby N, Broe GA, Creasey H, et al. Efficacy of tacrine and lecithin in mild to moderate Alzheimer's disease: Double blind trial. *BMJ* 1994 Apr 2; 308(6933):879-83.
110. Prentice N, Van BM, Dougall NJ, et al. A double-blind, placebo-controlled study of tacrine in patients with Alzheimer's disease using SPET. *J Psychopharmacol (Oxf)* 1996; 10(3):175-81.
111. Weinstein HC, Teunisse S, Van Gool WA. Tetrahydroaminoacridine and lecithin in the treatment of Alzheimer's disease. Effect on cognition, functioning in daily life, behavioural disturbances and burden experienced by the carers. *J Neurol* 1991 Feb; 238(1):34-8.
112. Wong WJ, Liu HC, Fuh JL, et al. A double-blind, placebo-controlled study of tacrine in Chinese patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 1999 Jul; 10(4):289-94.
113. Wood PC, Castleden CM. A double-blind, placebo controlled, multicentre study of tacrine for Alzheimer's disease. *Int J Geriatr Psychiatry* 1994; 9(8):649-54.
114. Allain H, Schuck S, Lebreton S, et al. Aminotransferase levels and silymarin in de novo tacrine-treated patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 1999 May; 10(3):181-5.
115. Huff FJ, Antuono P, Murphy M, et al. Potential clinical use of an adrenergic/cholinergic agent (HP 128) in the treatment of Alzheimer's disease. *Ann N Y Acad Sci* 1991; 640:263-7.
116. Antuono PG. Effectiveness and safety of velnacrine for the treatment of Alzheimer's disease. A double-blind, placebo-controlled study. Mentane Study Group. *Arch Intern Med* 1995 Sep 11; 155(16):1766-72.
117. Zemlan FP, Folks DG, Goldstein BJ, et al. Velnacrine for the treatment of Alzheimer's disease: A double-blind, placebo-controlled trial. *J Neural Transm Gen Sect* 1996; 103(8-9):1105-16.
118. Imbimbo BP, Martelli P, Troetel WM, et al. Efficacy and safety of eptastigmine for the treatment of patients with Alzheimer's disease. *Neurology* 1999 Mar 10; 52(4):700-8.
119. Canal N, Imbimbo BP. Relationship between pharmacodynamic activity and cognitive effects of eptastigmine in patients with Alzheimer's disease. *Clin Pharmacol Ther* 1996; 60(2):218-28.
120. Rockwood K, Beattie BL, Eastwood MR, et al. A randomized, controlled trial of linopirdine in the treatment of Alzheimer's disease. *Can J Neurol Sci* 1997 May; 24(2):140-5.
121. Van Dyck CH, Lin CH, Robinson R, et al. The acetylcholine releaser linopirdine increases parietal regional cerebral blood flow in Alzheimer's disease. *Psychopharmacologia* 1997 Aug; 132(3):217-26.
122. Xu SS, Gao ZX, Weng Z, et al. Efficacy of tablet huperzine-A on memory, cognition, and behavior in Alzheimer's disease. *Zhongguo Yao Li Xue Bao/Acta Pharmacologica Sinica* 1995 Sep; 16(5):391-5.
123. Mohr E, Nair NP, Sampson M, et al. Treatment of Alzheimer's disease with sabeluzole: Functional and structural correlates. *Clin Neuropharmacol* 1997 Aug; 20(4):338-45.
124. Allain H, Dautzenberg PH, Maurer K, et al. Double blind study of tiapride versus haloperidol and placebo in agitation and aggressiveness in elderly patients with

- cognitive impairment. *Psychopharmacologia* 2000 Mar; 148(4):361-6.
125. Teri L, Logsdon RG, Peskind E, et al. Treatment of agitation in AD: A randomized, placebo-controlled clinical trial. *Neurology* 2000 Nov 14; 55(9):1271-8.
 126. Petrie WM, Ban TA, Berney S, et al. Loxapine in psychogeriatrics: A placebo- and standard-controlled clinical investigation. *J Clin Psychopharmacol* 1982 Apr; 2(2):122-6.
 127. De Deyn PP, Rabheru K. A randomized trial of risperidone, placebo, and haloperidol for behavioral symptoms of dementia. *Neurology* 1999 Sep 22; 53(5):946-55.
 128. Auchus AP, Bissey-Black C. Pilot study of haloperidol, fluoxetine, and placebo for agitation in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1997; 9(4):591-3.
 129. Carlyle W, Ancill RJ, Sheldon L. Aggression in the demented patient: a double-blind study of loxapine versus haloperidol. *Int Clin Psychopharmacol* 1993; 8(2):103-8.
 130. Coccaro EF, Kramer E, Zemishlany Z, et al. Pharmacologic treatment of noncognitive behavioral disturbances in elderly demented patients. *Am J Psychiatry* 1990 Dec; 147(12):1640-5.
 131. Chan WC, Lam LC, Choy CN, et al. A double-blind randomised comparison of risperidone and haloperidol in the treatment of behavioural and psychological symptoms in Chinese dementia patients. *Int J Geriatr Psychiatry* 2001 Dec; 16(12):1156-62.
 132. Winblad B, Poritis N. Memantine in severe dementia: Results of the 9M-Best Study (Benefit and efficacy in severely demented patients during treatment with memantine). *Int J Geriatr Psychiatry* 1999 Feb; 14(2):135-46.
 133. Orgogozo J, Rigaud AS, Stoffler A, et al. Efficacy and safety of memantine in patients with mild to moderate vascular dementia. *Stroke* 2002; 33:1834-9.
 134. Wilcock G, Mobius HJ, Stoffler A. A double-blind, placebo-controlled multicentre study of memantine in mild to moderate vascular dementia (MMM500). *Int Clin Psychopharmacol* 2002; 17(6):297-305.
 135. Sano M, Ernesto C, Thomas RG, et al. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *N Engl J Med* 1997; 336(17):1216-22.
 136. Mangoni A, Grassi MP, Frattola L, et al. Effects of a MAO-B inhibitor in the treatment of Alzheimer disease. *Eur Neurol* 1991; 31(2):100-7.
 137. Agnoli A, Fabbri G, Fioravanti M, et al. CBF and cognitive evaluation of Alzheimer type patients before and after IMAO-B treatment: a pilot study. *Eur Neuropsychopharmacol* 1992 Mar; 2(1):31-5.
 138. Burke WJ, Roccaforte WH, Wengel SP, et al. L-deprenyl in the treatment of mild dementia of the Alzheimer type: Results of a 15-month trial. *J Am Geriatr Soc* 1993a; 41(11):1219-25.
 139. Filip V, Kolibas E. Selegiline in the treatment of Alzheimer's disease: A long-term randomized placebo-controlled trial. Czech and Slovak Senile Dementia of Alzheimer Type Study Group. *J Psychiatry Neurosci* 1999 May; 24(3):234-43.
 140. Freedman M, Rewilak D, Xerri T, et al. L-deprenyl in Alzheimer's disease: Cognitive and behavioral effects. *Neurology* 1998; 50(3):660-8.
 141. Cutler NR, Shrotriya RC, Sramek JJ, et al. The use of the Computerized Neuropsychological Test Battery (CNTB) in an efficacy and safety trial of BMY 21,502 in Alzheimer's disease. *Ann N Y Acad Sci* 1993; 695(Sep 24):332-6.
 142. Alvarez XA, Pichel V, Perez P, et al. Double-blind, randomized, placebo-controlled pilot study with anapsos in senile dementia: Effects on cognition, brain bioelectrical activity and cerebral hemodynamics. *Methods & Findings in Experimental & Clinical Pharmacology* 2000 Sep; 22(7):585-94.
 143. Olin JT, Fox LS, Pawluczyk S, et al. A pilot randomized trial of carbamazepine for behavioral symptoms in treatment-resistant outpatients with Alzheimer's disease. *Am J Geriatr Psychiatry* 2001; 9(4):400-5.
 144. Tariot PN, Erb R, Podgorski CA, et al. Efficacy and tolerability of carbamazepine for agitation and aggression in dementia. *Am J Psychiatry* 1998 Jan; 155(1):54-61.

145. Pollock BG, Mulsant BH, Rosen J, et al. Comparison of citalopram, perphenazine, and placebo for the acute treatment of psychosis and behavioral disturbances in hospitalized, demented patients. *Am J Psychiatry* 2002; 159(3):460-5.
146. Nyth AL, Gottfries CG. The clinical efficacy of citalopram in treatment of emotional disturbances in dementia disorders. A Nordic multicentre study. *Br J Psychiatry* 1990 Dec; 157:894-901.
147. Tariot PN, Schneider LS, Mintzer JE, et al. Safety and tolerability of divalproex sodium in the treatment of signs and symptoms of mania in elderly patients with dementia: Results of a double-blind, placebo-controlled trial. *Curr Ther Res Clin Exp* 2001b; 62(1):51-67.
148. Porsteinsson AP, Tariot PN, Erb R, et al. Placebo-controlled study of divalproex sodium for agitation in dementia. *Am J Geriatr Psychiatry* 2001; 9(1):58-66.
149. Petracca GM, Chmerinski E, Starkstein SE. A double-blind, placebo-controlled study of fluoxetine in depressed patients with Alzheimer's disease. *Int Psychogeriatr* 2001 Jun; 13(2):233-40.
150. Barnes R, Veith R, Okimoto J. Efficacy of antipsychotic medications in behaviorally disturbed dementia patients. *Am J Psychiatry* 1982; 139(9):1170-4.
151. Street JS, Clark WS, Gannon KS, et al. Olanzapine treatment of psychotic and behavioral symptoms in patients with Alzheimer disease in nursing care facilities: A double-blind, randomized, placebo-controlled trial. The HGEU Study Group. *Arch Gen Psychiatry* 2000 Oct; 57(10):968-76.
152. Meehan KM, Wang H, David SR, et al. Comparison of rapidly acting intramuscular olanzapine, lorazepam, and placebo: A double-blind, randomized study in acutely agitated patients with dementia. *Neuropsychopharmacology* 2002 Apr; 26(4):494-504.
153. Amaducci L. Phosphatidylserine in the treatment of Alzheimer's disease: Results of a multicenter study. *Psychopharmacol Bull* 1988; 24(1):130-4.
154. Crook T, Petrie W, Wells C, et al. Effects of phosphatidylserine in Alzheimer's disease. *Psychopharmacol Bull* 1992a; 28(1):61-6.
155. Katz IR, Jeste DV, Mintzer JE, et al. Comparison of risperidone and placebo for psychosis and behavioral disturbances associated with dementia: A randomized, double-blind trial. *J Clin Psychiatry* 1999; 60(2):107-15.
156. Magai C, Kennedy G, Cohen CI, et al. A controlled clinical trial of sertraline in the treatment of depression in nursing home patients with late-stage Alzheimer's disease. *Am J Geriatr Psychiatry* 2000; 8(1):66-74.
157. Lyketsos CG, Sheppard JM, Steele CD, et al. Randomized, placebo-controlled, double-blind clinical trial of sertraline in the treatment of depression complicating Alzheimer's disease: Initial results from the Depression in Alzheimer's Disease study. *Am J Psychiatry* 2000 Oct; 157(10):1686-9.
158. Dehlin O, Hedenrud B, Jansson P, et al. A double-blind comparison of alaproclate and placebo in the treatment of patients with senile dementia. *Acta Psychiatr Scand* 1985 Feb; 71(2):190-6.
159. Reifler BV, Teri L, Raskind M, et al. Double-blind trial of imipramine in Alzheimer's disease patients with and without depression. *Am J Psychiatry* 1989 Jan; 146(1):45-9.
160. Claus JJ, de K, I, van Harskamp F, et al. Lisuride treatment of Alzheimer's disease. A preliminary placebo-controlled clinical trial of safety and therapeutic efficacy. *Clin Neuropharmacol* 1998 May; 21(3):190-5.
161. Passeri M, Cucinotta D, de Mello M, et al. Comparison of minaprine and placebo in the treatment of Alzheimer's disease and multi-infarct dementia. *Int J Geriatr Psychiatry* 1987; 2(2):97-103.
162. Roth M, Mountjoy CQ, Amrein R. Moclobemide in elderly patients with cognitive decline and depression: An international double-blind, placebo-controlled trial. *Br J Psychiatry* 1996 Feb; 168(2):149-57.
163. Moller HJ, Hartmann A, Kessler C, et al. Naftidrofuryl in the treatment of vascular dementia. *Eur Arch Psychiatry Clin Neurosci* 2001; 251(6):247-54.
164. Bodick NC, Offen WW, Levey AI, et al. Effects of xanomeline, a selective muscarinic receptor agonist, on cognitive function and behavioral symptoms in Alzheimer's disease. *Arch Neurol* 1997 Apr; 54(4):465-73.

165. Olafsson K, Jorgensen S, Jensen HV, et al. Fluvoxamine in the treatment of demented elderly patients: A double-blind, placebo-controlled study. *Acta Psychiatr Scand* 1992 Jun; 85(6):453-6.
166. Thal LJ, Forrest M, Loft H, et al. Lu 25-109, a muscarinic agonist, fails to improve cognition in Alzheimer's disease. Lu25-109 Study Group. *Neurology* 2000b; 54(2):421-6.
167. Fuchs A, Hehnke U, Erhart C, et al. Video rating analysis of effect of maprotiline in patients with dementia and depression. *Pharmacopsychiatry* 1993 Mar; 26(2):37-41.
168. Ruether E, Husmann R, Kinzler E, et al. A 28-week, double-blind, placebo-controlled study with Cerebrolysin in patients with mild to moderate Alzheimer's disease. *Int Clin Psychopharmacol* 2001 Sep; 16(5):253-63.
169. Bae CY, Cho CY, Cho K, et al. A double-blind, placebo-controlled, multicenter study of Cerebrolysin for Alzheimer's disease. *J Am Geriatr Soc* 2000 Dec; 48(12):1566-71.
170. Xiao S, Yan H, Yao P, et al. Efficacy of FPF 1070 (cerebrolysin) in patients with Alzheimer's disease: A multicentre, randomised, double-blind, placebo-controlled trial. *Clin Drug Investig* 2000(1):43-53.
171. Xiao S, Yan H, Yao P, et al. The efficacy of cerebrolysin in patients with vascular dementia: Results of a Chinese multicentre, randomised, double-blind, placebo-controlled trial. *Hong Kong Journal of Psychiatry* 1999(2):13-9.
172. Ruether E, Ritter R, Apecechea M, et al. Efficacy of the peptidergic nootropic drug cerebrolysin in patients with senile dementia of the Alzheimer type (SDAT). *Pharmacopsychiatry* 1994 Jan; 27(1):32-40.
173. Panisset M, Gauthier S, Moessler H, et al. Cerebrolysin in Alzheimer's disease: A randomized, double-blind, placebo-controlled trial with a neurotrophic agent. *J Neural Transm Gen Sect* 2002; 109(7-8):1089-104.
174. Asthana S, Baker LD, Craft S, et al. High-dose estradiol improves cognition for women with AD: Results of a randomized study. *Neurology* 2001 Aug 28; 57(4):605-12.
175. Wang PN, Liao SQ, Liu RS, et al. Effects of estrogen on cognition, mood, and cerebral blood flow in AD: A controlled study. *Neurology* 2000 Jun 13; 54(11):2061-6.
176. Henderson VW, Paganini-Hill A, Miller BL, et al. Estrogen for Alzheimer's disease in women: Randomized, double-blind, placebo-controlled trial. *Neurology* 2000 Jan 25; 54(2):295-301.
177. Mulnard RA, Cotman CW, Kawas C, et al. Estrogen replacement therapy for treatment of mild to moderate Alzheimer disease: A randomized controlled trial. *JAMA* 2000(8):1007-15.
178. Kyomen HH, Satlin A, Hennen J, et al. Estrogen therapy and aggressive behavior in elderly patients with moderate-to-severe dementia: Results from a short-term, randomized, double-blind trial. *Am J Geriatr Psychiatry* 1999; 7(4):339-48.
179. Le Bars PL, Katz MM, Berman N, et al. A placebo-controlled, double-blind, randomized trial of an extract of Ginkgo biloba for dementia. *JAMA* 1997(16):1327-32.
180. Maurer K, Ihl R, Dierks T, et al. Clinical efficacy of Ginkgo biloba special extract EGb 761 in dementia of the Alzheimer type. *J Psychiatr Res* 1997 Nov; 31(6):645-55.
181. Kanowski S, Herrmann WM, Stephan K, et al. Proof of efficacy of the Ginkgo biloba special extract EGb 761 in outpatients suffering from mild to moderate primary degenerative dementia of the Alzheimer type or multi-infarct dementia. *Pharmacopsychiatry* 1996 Mar; 29(2):47-56.
182. Bergamasco B, Scarzella L, La Commare P. Idebenone, a new drug in the treatment of cognitive impairment in patients with dementia of the Alzheimer type. *Funct Neurol* 1994 May; 9(3):161-8.
183. Gutzmann H, Hadler D. Sustained efficacy and safety of idebenone in the treatment of Alzheimer's disease: Update on a 2-year double-blind multicentre study. *J Neural Transm Suppl* 1998; 54:301-10.
184. Marigliano V, Abate G, Barbagallo-Sangiorgi G, et al. Randomized, double-blind, placebo controlled, multicentre study of idebenone in patients suffering from multi-infarct dementia. *Arch Gerontol Geriatr* 1992; 15(3):239-48.
185. Weyer G, Babej-Dolle RM, Hadler D, et al. A controlled study of 2 doses of idebenone in the treatment of Alzheimer's disease. *Neuropsychobiology* 1997; 36(2):73-82.

186. Bottini G, Vallar G, Cappa S, et al. Oxiracetam in dementia: A double-blind, placebo-controlled study. *Acta Neurol Scand* 1992 Sep; 86(3):237-41.
187. Maina G, Fiori L, Torta R, et al. Oxiracetam in the treatment of primary degenerative and multi-infarct dementia: A double-blind, placebo-controlled study. *Neuropsychobiology* 1989; 21(3):141-5.
188. Mangoni A, Perin C, Smirne S, et al. A double-blind, placebo-controlled study with oxiracetam in demented patients administered the Luria-Nebraska Neuropsychological Battery. *Drug Dev Res* 1988; 14(3-4):217-4.
189. Rozzini R, Zanetti O, Bianchetti A. Effectiveness of oxiracetam therapy in the treatment of cognitive deficiencies secondary to primary degenerative dementia. *Acta Neurol (Napoli)* 1992 Apr; 14(2):117-26.
190. Villardita C, Grioli S, Lomeo C, et al. Clinical studies with oxiracetam in patients with dementia of Alzheimer type and multi-infarct dementia of mild to moderate degree. *Neuropsychobiology* 1992; 25(1):24-8.
191. Knezevic S. European Pentoxifylline Multi-Infarct Dementia Study. *Eur Neurol* 1996; 36(5):315-21.
192. Black RS, Barclay LL, Nolan KA, et al. Pentoxifylline in cerebrovascular dementia. *J Am Geriatr Soc* 1992 Mar; 40(3):237-44.
193. Ghose K. Oxpentifylline in dementia: A controlled study. *Arch Gerontol Geriatr* 1987 Apr; 6(1):19-26.
194. Parnetti L, Mari D, Abate G, et al. Vascular dementia Italian sulodexide study (VA.D.I.S.S.). Clinical and biological results. *Thromb Res* 1997 Jul 15; 87(2):225-33.
195. Marcusson J, Rother M, Kittner B, et al. A 12-month, randomized placebo-controlled trial of propentofylline (HWA 285) in patients with dementia according to DSM III-R. *Dement Geriatr Cogn Disord* 1997; 8(5):320-8.
196. Mielke R, Kittner B, Ghaemi M, et al. Propentofylline improves regional cerebral glucose metabolism and neuropsychologic performance in vascular dementia. *JNS* 1996 Sep 15; 141(1-2):59-2.
197. Mielke R, Ghaemi M, Kessler J, et al. Propentofylline enhances cerebral metabolic response to auditory memory stimulation in Alzheimer's disease. *JNS* 1998 Jan 21; 154(1):76-82.
198. Saletu B, Moller HJ, Grunberger J, et al. Propentofylline in adult-onset cognitive disorders: double-blind, placebo-controlled, clinical, psychometric and brain mapping studies. *Neuropsychobiology* 1990 Sep; 24(4):173-84.
199. Terasawa K, Shimada Y, Kita T, et al. Choto-san in the treatment of vascular dementia: A double-blind, placebo-controlled study. *Phytomedicine* 1997; 4(1):15-22.
200. Shimada Y, Terasawa K, Yamamoto T, et al. A well-controlled study of Choto-san and placebo in the treatment of vascular dementia. *J Tradit Med* 1994; 11:246-55.
201. Schellenberg R, Todorova A, Wedekind W, et al. Pathophysiology and psychopharmacology of dementia: A new study design. *Neuropsychobiology* 1997; 35(3):132-42.
202. Weyer G, Eul A, Milde K, et al. Cyclandelate in the treatment of patients with mild to moderate primary degenerative dementia of the Alzheimer type or vascular dementia: Experience from a placebo controlled multicenter study. *Pharmacopsychiatry* 2000 May; 33(3):89-97.
203. Danielczyk W, Simanyi B, Forette F, et al. CBM 36-733 (2-methyl-alpha-ergokryptine) in primary degenerative dementia: Results of a European multicentre trial. *Int J Geriatr Psychiatry* 1988; 3(2):107-14.
204. Cucinotta D, De LD, Frattola L, et al. Dihydroergokryptine vs. placebo in dementia of Alzheimer type: Interim results of a randomized multicenter study after a 1-year follow-up. *Arch Gerontol Geriatr* 1996; 22(2):169-80.
205. Van Gool WA, Weinstein HC, Scheltens PK, et al. Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: An 18-month randomised, double-blind, placebo-controlled study. *Lancet* 2001 Aug 11; 358(9280):455-60.
206. Aisen PS, Schmeidler J, Pasinetti GM. Randomized pilot study of nimesulide treatment in Alzheimer's disease. *Neurology* 2002 Apr 9; 58(7):1050-4.
207. Pantoni L, Bianchi C, Beneke M, et al. The Scandinavian Multi-Infarct Dementia Trial: A

- double-blind, placebo-controlled trial on nimodipine in multi-infarct dementia. *JNS* 2000a; 175(2):116-23.
208. Ban TA, Morey L, Aguglia E, et al. Nimodipine in the treatment of old age dementias. *Prog Neuropsychopharmacol Biol Psychiatry* 1990; 14(4):525-51.
209. Soininen H, Koskinen T, Helkala EL, et al. Treatment of Alzheimer's disease with a synthetic ACTH 4-9 analog. *Neurology* 1985 Sep; 35(9):1348-51.
210. Kragh-Sorensen P, Olsen RB, Lund S, et al. Neuropeptides: ACTH-peptides in dementia. *Prog Neuropsychopharmacol Biol Psychiatry* 1986; 10(3-5):479-92.
211. Ban TA, Morey LC, Santini V. Clinical investigations with ateroid in old-age dementias. *Semin Thromb Hemost* 1991b; 17(Suppl 2):161-3.
212. Crapper McLachlan DR, Dalton AJ, Kruck TP, et al. Intramuscular desferrioxamine in patients with Alzheimer's disease. *Lancet* 1991 Jun 1; 337(8753):1304-8.
213. Ban TA, Morey LC, Aguglia E, et al. Glycosaminoglycan polysulfate in the treatment of old age dementias. *Prog Neuropsychopharmacol Biol Psychiatry* 1991a; 15(3):323-42.
214. Crook T, Wilner E, Rothwell A, et al. Noradrenergic intervention in Alzheimer's disease. *Psychopharmacol Bull* 1992b; 28(1):67-70.
215. Shrotriya RC, Cutler NR, Sramek JJ, et al. Efficacy and safety of BMY 21,502 in Alzheimer's disease. *Ann Pharmacother* 1996 Dec; 30(12):1376-80.
216. Nolan KA, Black RS, Sheu KF, et al. A trial of thiamine in Alzheimer's disease. *Arch Neurol* 1991 Jan; 48(1):81-3.
217. Fischhof PK, Moslinger-Gehmayr R, Herrmann WM, et al. Therapeutic efficacy of vincamine in dementia. *Neuropsychobiology* 1996; 34(1):29-35.
218. Peabody CA, Davies H, Berger PA, et al. Desamino-D-arginine-vasopressin (DDAVP) in Alzheimer's disease. *Neurobiol Aging* 1986 Jul; 7(4):301-3.
219. Kanowski S, Fischhof PK, Grobe-Einsler R, et al. Efficacy of xantinolnicotinate in patients with dementia. *Pharmacopsychiatry* 1990 May; 23(3):118-24.
220. Cucinotta D, Aveni Casucci MA, Pedrazzi F, et al. Multicentre clinical placebo-controlled study with buflomedil in the treatment of mild dementia of vascular origin. *J Int Med Res* 1992 Apr; 20(2):136-49.
221. Treves TA, Korczyn AD. Denbufylline in dementia: a double-blind controlled study. *Dement Geriatr Cogn Disord* 1999 Nov; 10(6):505-10.
222. Scharf S, Mander A, Ugoni A, et al. A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology* 1999 Jul 13; 53(1):197-201.
223. Thompson TL, Filley CM, Mitchell WD, et al. Lack of efficacy of hydergine in patients with Alzheimer's disease. *N Engl J Med* 1990 Aug 16; 323(7):445-8.
224. Rogers J, Kirby LC, Hempelman SR, et al. Clinical trial of indomethacin in Alzheimer's disease. *Neurology* 1993 Aug; 43(8):1609-11.
225. Ala T, Romero S, Knight F, et al. GM-1 treatment of Alzheimer's disease. A pilot study of safety and efficacy. *Arch Neurol* 1990 Oct; 47(10):1126-30.
226. Adair JC, Knoefel JE, Morgan N. Controlled trial of N-acetylcysteine for patients with probable Alzheimer's disease. *Neurology* 2001 Oct 23; 57(8):1515-7.
227. Carlson MC, Tschanz JT, Norton MC, et al. H2 histamine receptor blockade in the treatment of Alzheimer disease: A randomized, double-blind, placebo-controlled trial of nizatidine. *Alzheimer Dis Assoc Disord* 2002 Jan; 16(1):24-30.
228. Croisile B, Trillet M, Fondarai J, et al. Long-term and high-dose piracetam treatment of Alzheimer's disease. *Neurology* 1993 Feb; 43(2):301-5.
229. Aisen PS, Davis KL, Berg JD, et al. A randomized controlled trial of prednisone in Alzheimer's disease. *Alzheimer's Disease Cooperative Study*. *Neurology* 2000b; 54(3):588-93.
230. Simons M, Schwarzler F, Lutjohann D, et al. Treatment with simvastatin in

- normocholesterolemic patients with Alzheimer's disease: A 26-week randomized, placebo-controlled, double-blind trial. *Ann Neurol* 2002; 52(3):346-50.
231. Convit A, de Asis J, de Leon MJ, et al. Atrophy of the medial occipitotemporal, inferior, and middle temporal gyri in non-demented elderly predict decline to Alzheimer's disease. *Neurobiol Aging* 2000 Jan; 21(1):19-26.
 232. Rapp S, Brenes G, Marsh AP. Memory enhancement training for older adults with mild cognitive impairment: A preliminary study. *Aging Ment Health* 2002 Feb; 6(1):5-11.
 233. Johnson SA, Simmon VF. Randomized, double-blind, placebo-controlled international clinical trial of the AMPAKINE CX516 in elderly participants with mild cognitive impairment. A progress report. *J Mol Neurosci* 2002; 19(1-2):197-200.
 234. Sherwin BB. Estrogen and cognitive functioning in men with Mild Cognitive Impairment. *J Mol Neurosci* 2002; 19(1-2):219-23.
 235. McDermott MP, Hall WJ, Oakes D, et al. Design and analysis of two-period studies of potentially disease-modifying treatments. *Control Clin Trials* 2002 Dec; 23(6):635-49.
 236. Bodick N, Forette F, Hadler D, et al. Protocols to demonstrate slowing of Alzheimer's disease progression: Position paper from the International Working Group on Harmonization of Dementia Drug Guidelines. *Alzheimer Dis Assoc Disord* 1997; 11(Suppl 3):50-3.
 237. Ancill RJ, Carlyle WW, Liang RA, et al. Agitation in the demented elderly: A role for benzodiazepines? *Int Clin Psychopharmacol* 1991; 6(3):141-6.
 238. Cucinotta D, Romagnoli S, Godoli G, et al. Comparison of sulfomucopolysaccharides and cytidine diphosphocholine in the treatment of multi-infarct dementia. A randomized double-blind test. *Curr Ther Res Clin Exp* 1988; 43(1):12-20.
 239. Karlsson I, Godderis J, Augusto De Mendonca LC, et al. A randomised, double-blind comparison of the efficacy and safety of citalopram compared to mianserin in elderly, depressed patients with or without mild to moderate dementia. *Int J Geriatr Psychiatry* 2000 Apr; 15(4):295-305.
 240. Katona CL, Hunter BN, Bray J. A double-blind comparison of the efficacy and safety of paroxetine and imipramine in the treatment of depression with dementia. *Int J Geriatr Psychiatry* 1998 Feb; 13(2):100-8.
 241. Taragano FE, Lyketsos CG, Mangone CA, et al. A double-blind, randomized, fixed-dose trial of fluoxetine vs. amitriptyline in the treatment of major depression complicating Alzheimer's disease. *Psychosomatics* 1997 May; 38(3):246-52.
 242. Popa R, Schneider F, Mihalas G, et al. Antagonic-Stress superiority versus meclofenoxate in gerontopsychiatry. *Arch Gerontol Geriatr* 1994; 18(Suppl 4):197-206.
 243. Spilich GJ, Wannenmacher W, Duarte A, et al. Efficacy of pyritinol versus hydergine upon cognitive performance in patients with senile dementia of the Alzheimer's type: A double-blind multi-center trial. *Alzheimers Res* 1996(3):79-84.
 244. Gutzmann H, Kuhl KP, Kanowski S, et al. Measuring the efficacy of psychopharmacological treatment of psychomotoric restlessness in dementia: clinical evaluation of tiapride. *Pharmacopsychiatry* 1997 Jan; 30(1):6-11.
 245. Passeri M, Cucinotta D, Abate G, et al. Oral 5'-methyltetrahydrofolic acid in senile organic mental disorders with depression: Results of a double-blind multicenter study. *Aging (Milano)* 1993 Feb; 5(Milano):63-71.
 246. Ruether E, Alvarez XA, Rainer M, et al. Sustained improvement of cognition and global function in patients with moderately severe Alzheimer's disease: A double-blind, placebo-controlled study with the neurotrophic agent Cerebrolysin. *J Neural Transm Suppl* 2002(62):265-75.
 247. Pantoni L, Rossi R, Inzitari D, et al. Efficacy and safety of nimodipine in subcortical vascular dementia: A subgroup analysis of the Scandinavian Multi-Infarct Dementia Trial. *JNS* 2000b; 175(2):124-34.
 248. Dunne MP, Hartley LR. Scopolamine and the control of attention in humans. *Psychopharmacologia* 1986; 89(1):94-7.
 249. Black DW, Monahan P, Wesner R, et al. The effect of fluvoxamine, cognitive therapy, and

- placebo on abnormal personality traits in 44 patients with panic disorder. *J Personal Disord* 1996; 10(2):185-94.
250. Dingemans J, Bury M, Roncari G, et al. Pharmacokinetics and pharmacodynamics of Ro 41-3696, a novel nonbenzodiazepine hypnotic. *J Clin Pharmacol* 1995 Aug; 35(8):821-9.
 251. Huster WJ, Enas GG. A framework establishing clear decision criteria for the assessment of drug efficacy. *Stat Med* 1998 Aug 15; 17(15-16):1829-38.
 252. Ruberg S, Cairns V. Providing evidence of efficacy for a new drug. *Stat Med* 1998 Aug 15; 17(15-16):1813-23.
 253. Boustani M, Peterson B, Hanson L, et al. Screening for dementia in primary care: a summary of the evidence for the U.S. Preventive Services Task Force. *Ann Intern Med* 2003 Jun 3; 138(11):927-37.
 254. Knopman DS, DeKosky ST, Cummings JL, et al. Practice parameter: diagnosis of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001 May 8; 56(9):1143-53.
 255. Rockwood K, Stolee P. Responsiveness of outcome measures used in an antidementia drug trial. *Alzheimer Dis Assoc Disord* 2000 Jul; 14(3):182-5.
 256. Beaton DE, Bombardier C, Katz JN, et al. A taxonomy for responsiveness. *J Clin Epidemiol* 2001 Dec; 54(12):1204-17.
 257. Als-Nielsen B, Chen W, Gluud C, et al. Association of funding and conclusions in randomized drug trials: a reflection of treatment effect or adverse events? *JAMA* 2003 Aug 20; 290(7):921-8.
 258. Khan A. Are placebo controls necessary to test new antidepressants and anxiolytics? *Int J Neuropsychopharmacol* 2002; 5(3):193-7.
 259. Bowler JV, Eliasziw M, Steenhuis R, et al. Comparative evolution of Alzheimer disease, vascular dementia, and mixed dementia. *Arch Neurol* 1997 Jun; 54(6):697-703.
 260. Reisberg B, Ferris SH, de Leon MJ, et al. The Global Deterioration Scale for assessment of primary degenerative dementia. *Am J Psychiatry* 1982 Sep; 139(9):1136-9.
 261. Hughes CP, Berg L, Danziger WL, et al. A new clinical scale for the staging of dementia. *Br J Psychiatry* 1982 Jun; 140:566-72.
 262. Feinstein AR. "Clinical Judgment" revisited: the distraction of quantitative models. *Ann Intern Med* 1994 May 1; 120(9):799-805.

Acronyms and Abbreviations

Acronyms and Abbreviations

AAMI	Age-Associated Memory Impairment
ABID	Agitated Behavior Inventory for Dementia
ABS	Adaptive Behavior Scale
ACES	Agitation-Calmness Evaluation Scale
ACFP	American College of Family Physicians
AChE	Acetylcholinesterase
ACP-ASIM	American College of Physicians – American Society of Internal Medicine
ACPT	Auditory Continuous Performance Test
ACTH	Adrenocorticotrophic hormone
AD	Alzheimer's Disease
ADAS	Alzheimer's Disease Assessment Scale
ADAS-11; ADAS-13	Alzheimer's Disease Assessment Scale (11 and 13 items)
ADAS-Cog ADAS-NonCog	Alzheimer's Disease Assessment Scale-Cognitive and Non-Cognitive Sections
ADCS-ADL	Alzheimer's Disease Cooperative Study – Activities of Daily Living
ADCS-CGIC	Alzheimer's Disease Cooperative Study – Clinical Global Impression of Change
ADFACS	Alzheimer's Disease Functional Assessment and Change Scale
ADL; ADLC	Activities of Daily Living (Checklist)
ADL-BDRS	Activities of Daily Living-Blessed Dementia Rating Scale
ADL-PDS	Activities of Daily Living- Progressive Deterioration Scale
ADS	Alzheimer's Deficit Scale
ADSS	Alzheimer's Disease Symptomatology Scale
AFBS	Aversive Feeding Behavior Scale
AGGR	Aggressiveness subscale of the Personality Psychopathology Five (PSY-5) Scales
AGS-E	Assessment of Global Symptomatology-Elderly
AHRQ	Agency for Healthcare Research and Quality
AIDS	Acquired Immune Deficiency Syndrome
AIMS	Abnormal Involuntary Movement Scale
ALCAR	Carnitine
AMED	Allied and Complementary Medicine Database
AMI	Attention Matrices
AMPA	Alpha-amino-3-hydroxy-5-methyl-4-isoxazole propionic acid
AMTS	Abbreviated Mental Test Score
APOE	Apolipoprotein E gene
BADL	Basic Activities of Daily Living
BARS, BAS	Barnes Akathisia Rating Scale
BCRS	Brief Cognitive Rating Scale
BDI	Beck Depression Inventory
BDRS	Blessed Dementia Rating Scale
BEHAVE – AD	Behavioral Pathology in Alzheimer's Disease Rating Scale
BePU	Berlin rating scale for psychomotoric restlessness
Bf-S	Zerssen Adjective Mood Scale (German test: Befindlichkeitsskala)
BGP	Behavioural Rating Scale for Geriatric Patients
BGP	Behavioural Rating Scale for Geriatric Patients
BI	Barthel Index
bid	Twice a day
BL-A	Blessed A scale
Blessed-D BDRS	Blessed Dementia Rating Scale
BMI	Body Mass Index
BMICT	Blessed Memory Information and Concentration Test
BMV	Nootropic agent; Bristol-Myers Squibb
BNT	Boston Naming Test
BPRS	Brief Psychiatric Rating Scale

Acronyms and Abbreviations cont'd.

BRMS	Bech-Rafaelsen Mania Scale
BRSD	Behavioral Rating Scale for Dementia
BSRT	Babcock Story Recall Test
BSRT	Buschke Selective Reminding Test
BSS	Behavioral Syndromes Scale for Dementia
BTT	Block Tapping Test
CADASIL	Cerebral Autosomal Dominant Ischemia with Subcortical Leukoencephalopathy
CAMCOG	Cognitive section of the Cambridge Examination for Mental Disorders in the Elderly
CAMDEX	Cambridge Examination for Mental Disorders in the Elderly
CAMTOT	CAMCOG Total Score
CANTAB	Cambridge Automated Neuropsychological Test Assessment Battery
CAPE	Clifton Assessment Procedures for the Elderly
CASE	Clifton Assessment Scale for the Elderly
CASI	Cognitive Abilities Screening Instrument
CATS	Caregiver's Activity Time Survey
CAUST	Canadian Utilization of Service Tracking questionnaire
CBC	Complete Blood Count
CBM 36-733	2-methyl-alpha-ergokryptine
CCASSS	Computerized Cognitive Assessment System Speed Score
CCT	Controlled Clinical Trial
CDR, CDRS	Clinical Dementia Rating; Clinical Dementia Rating Scale
CDR-NH	Clinical Dementia Rating – Nursing Home Version
CDR-SB	Clinical Dementia Rating – Sum of Boxes
CDT	Clock-Drawing Test
CEB	Clinical Epidemiology and Biostatistics
CERAD	Consortium to Establish a Registry for Alzheimer's Disease
CERE	Cerebrolysin
CETM	Dynamic measure of comprehension process (Spilich)
CGAE	Clinical Global Assessment of Efficacy
CGC+	Clinical Global Change-Plus
CGI	Clinical Global Impression
cGIC	Caregiver-rated Global Impression of Change
CGIC	Clinical Global Impression of Change
CGI-GI	Global Improvement
CGI-CGC	Clinical Global Impression-Clinical Global Change
CGI-S; CGI-S/C	Clinical Global Impression-Severity/Change
CGRS	Clinicians' Global Rating Score
chisq	Chi-Square Test
chisq _{M-H}	Mantel-Haenszel Chi-Square Test
CI	Confidence interval
CIBI	Clinician's Interview-Based Impression
CIBIC	Clinician's Interview-Based Impression of Change
CIBIC+	Clinician's Interview Based Impression of Change plus Caregiver
CIBIS+	Clinician's Interview-Based Impression of Severity with Caregiver Input
CINAHL	Cumulative Index to Nursing & Allied Health Literature ®
CIND	Cognitive Impairment Not yet Diagnosed
CLEX	Clinical Examination
CloND	Cognitive Loss No Dementia
CMAI	Cohen Mansfield Agitation Inventory
CNTB	Computerized Neuropsychological Test Battery
COSTART	Coding Symbols for a Thesaurus of Adverse Reaction Terms
COWAT	Controlled Oral Word Association Test
CPRS	Comprehensive Psychopathological Rating Scale
CPT	Cognitive Performance Test
CSDD	Cornell Scale for Depression in Dementia
CSGDS	Collateral Source Geriatric Depression Scale
CSI	Caregiver Stress Inventory

Acronyms and Abbreviations cont'd.

CSS	Caregiver Stress Scale
CT	Computerized Tomography
CVD	Cerebrovascular Disease
CVLT	California Verbal Learning Test
d	day
d	Effect Size Value – (d) is the average amount of change in standard deviation units achieved by individuals in a treated group versus the change achieved by members of a control/comparison group for a particular study
DAD	Disability Assessment for Dementia
DAT	Dementia Alzheimer's Type
D-B	Delay relative to Baseline
DBDS	Dementia Behavior Disturbance Scale
DCT	Digit Copying Test
DDAVP	Deamino-D-Arginine-Vasopressin
DEK	Dihydroergokryptine
Df	Degrees of Freedom
DMR	Dementia Questionnaire for Mentally Retarded Persons
DMSE	Delayed Matching-to-Sample Exam
D-P	Delay relative to Placebo
DPZ	Donepezil
DRS	Dementia Rating Scale
DSCS	Depressive Symptoms Collateral Source
DSM	Diagnostic and Statistical Manual of Mental Disorders (Edition III, III-R, IV)
DSPT	Digit Span Test
DSS	Depressive Signs Scale
DST; DSST	Digit Symbol (Substitution) Test
DTIC	Discovering Things in Common
e.g.,	example
ECG	Electrocardiogram
EEG	Electroencephalography
EFR	Emotional Face Recognition
EIS	Efficacy Index Score
EMBASE	Excerpta Medica Database
EPS	Extrapyramidal Symptoms
ERP	Event-Related Potential
ESRS	Extrapyramidal Symptom Rating Scale
FAST	Functional Assessment Staging
FCCA	Final Comprehensive Consensus Assessment
FCMT	Figure Copy/ Memory Test
FDA	Food and Drug Administration
FDG-PET	Positron Emission Tomography with 18-fluorodeoxyglucoseis
FIGT	Figure Detection Test
FIM	Functional Independence Measure
FRS	Functional Rating Scale test
g	gram
GABA	Gamma-aminobutyric acid
GBS	Gottfries-Bråne-Steen
GBS-SDS	Gottfries-Bråne-Steen – Scale for Dementia Syndromes
GDS	Global Deterioration Scale
GERRI	Geriatric Evaluation by Relative's Rating Instrument
GIS	Global Improvement Scale
GM-1	Monosialoganglioside
GMS-A	Geriatric Mental State questionnaire
GPI-E	General Psychiatric Impression-Elderly
GS	Gestalt Scale
h	hour
HAM-A; HARS	Hamilton Anxiety Rating Scale

Acronyms and Abbreviations cont'd.

HAM-D; HDRS	Hamilton Depression Rating Scale
HDS-R	Hasegawa Dementia Scale-Revised
HIS	Hachinski Ischemic Score
HIV	Human Immunodeficiency Virus
HMI	Hachinski-Marshall Ischaemic Index
HVLT	Hopkins Verbal Learning Test
IADL	Instrumental Activities of Daily Living
ICC	Item Characteristic Curve analysis
ICD	International Classification of Diseases (Version 9 or 10)
IDDD	Interview for Deterioration in Daily Living Activities in Dementia-complex task
IF	Industry Funded
IM	Intramuscular
I-P	Improvement relative to Placebo
IPSC-E	Raskin's and Crook's Inventory of Psychic and Somatic Complaints for the Elderly
IQCODE	Informant Questionnaire on Cognitive Decline in the Elderly
IS	Industry provided Supplies
ITT	Intention-to-treat
IU	International Units
kg	kilogram
KOLT	Kendrick Object Learning Test
LAS	Luria Alternating Series
lbs	pounds
LFT	Liver Function Test
LMT	Logical Memory Test
LNNB	Luria-Nebraska Neuropsychological Battery
LOCF	Last Observation Carried Forward
LPRS	London Psychogeriatric Rating Scale
LRU	Lipaseic Releasing Units
m	month
M	male
MAACL-R	Multiple Affect Adjective Checklist-Revised
MACF	Microtubule Actin Crosslinking Factor
MADR-S	Montgomery-Asberg Depression Rating Scale
MCI	Mild Cognitive Impairment
MCPT	Modified Continuous Performance Test
MDB	Mental Deterioration Battery
MeSH	Medical Subject Heading
µg	microgram
mg	milligram
MID	Multi Infarct Dementia
Min	Minimal
MITT	Modified Intention-to-treat
ml	milliliter
MMSE (MMSE-CE)	Mini-Mental Status Exam (estimated score)
CMMSE	Cantonese MMSE
MMMSE	Modified MMSE
SMMSE	Standardized MMSE
MNLT	Modified Names Learning Test
Mod	Moderate
Modly Sev	Moderately Severe
MQ	Memory Quotient
MRI	Magnetic Resonance Imaging
MRS	Magnetic Resonance Spectroscopy
MU-EPC	McMaster University Evidence-based Practice Center
MWF	Mattis Word Fluency
MX	Mixed results
n	number included in study
N	No

Acronyms and Abbreviations cont'd.

NA	Not available
NAA	Nuremberg gerontopsychological inventory for Assessing Activities of daily living
NAB	Nurnberger-Alters-Beobachtungs-Skala
NAC	N-Acetylcysteine
NAI	Nuremberg Age Inventory
NART	Nelson Adult Reading Test
NCT	Number Connection Test
NDT	New Dot Test
NI	Non-Industry funding source
NIMCS	Newcastle Memory, Information and Concentration Scale
NINCDS	National Institute of Neurological and Communicative Disorders and Stroke
NINCDS-ADRDA	National Institute of Neurological and Communicative Disorders and Stroke – Alzheimer's Disease and Related Disorders Association
NINDS-AIREN	National Institute of Neurological Disorders and Stroke – Association Internationale pour la Recherche et l'Enseignement en Neurosciences
NLT	Names Learning Test
NMDA	N-methyl-D-aspartate
NMICS	Newcastle Memory, Information and Concentration Scale
NMS	Nowlis Mood Scale
NNI	Number Needed to Intervene
NOSGER	Nurses Observation Scale for Geriatric Patients
NOSGER-IADL	Nurses Observation Scale for Geriatric Patients – Instrumental Activities of Daily Living subscale
NOSIE	Nurses Observation Scale for Inpatients
NPI (NPI-4, NPI-10)	Neuropsychiatric Inventory Subscores 4,10
NPI-NH	Neuropsychiatric Inventory – Nursing Home Version
NR	Not Reported
NRSMG	Non-Randomised Studies Methods Group
NS	Not significant
NSL	Neuropsychological Aging Self-Evaluation – List for Age Symptoms
NST	Non-Stress Test
NT	Not tested
OARS – ADL	Older Americans Resource Scale
OAS	Overt Aggression Scale
OC	Observed Cases
OLT	Object Learning Test
OMDR	Oculomotor Delayed Response
OR	Odds Ratio
ORG 2766	Adrenocorticotrophic hormone derivative
OXIR	Oxiracetam
oz	ounce
p	p value
P300	Electrophysiological potential that is indicator of associative and cognitive processes and latency in decision making processes
PAD	Presenile Alzheimer's Disease
PADL	Performance of Activities of Daily Living
PANSS-EC	Positive and Negative Syndrome Scale-Excited Component
PD	Parkinson's Disease
PDD	Progressive Degenerative Dementia
PDS	Progressive Deterioration Scale
PDSD	Primary Degenerative Senile Dementia
PET	Positron Emission Tomography
PGIR	Patient's Global Improvement Rating
PI	Partially funded by Industry
PICD	Presenile Idiopathic Cognitive Decline
POMS	Profile of Mood States
PRL	Prolactin

Acronyms and Abbreviations cont'd.

PSMS	Physical Self-Maintenance Scale
PSP	Progressive Supranuclear Palsy
PSQI	Pittsburgh Sleep Quality Index
qid	Four times daily
QoL	Quality of Life
R	Correlation Coefficient
RA	Research Assistant
RAGS; RAGS-E	Relative's Assessment of Global Symptomatology (Elderly)
RAPSU	Scale for psychomotoric agitation
R-AVL	Rey auditory-verbal-learning test
RCT	Randomized Controlled Trial
RDS	Rapid Disability Scale
RefMan	Reference Manager Version 10®
RGRS	Relatives' Global Rating Score
RMBPC	Revised Memory and Behavior Problems Checklist
RMT	Rey Memory Test; Randt Memory Test
RMT-A&R	Randt Memory Test – Acquisition and Recall
RMT-DR	Randt Memory Test – Delayed Recall
RMT-MI	Randt Memory Test – Memory Index
RPM	Raven's Progressive Matrices
RPT	Rivermead Behavioural Memory Test-Profile Score
RR	Relative Risk
RT	Reaction Time
RTI	Research Triangle Institute
SADS	Schedule for Affective Disorders and Schizophrenia
SAS	Simpson-Angus Scale
SAS-G	Self Assessment Scale – Geriatric
S-B	Stabilization relative to Baseline
SBI	Spontaneous Behavior Interview
SC	Significant change
SCAG	Sandoz Clinical Assessment – Geriatric
SCB	Screen for Caregiver Burden
SCWIT	Stroop Color Word Interference Test
SD	Standard Deviation
SDAT	Senile Dementia of the Alzheimer's Type
SEM	Standard Error
Sev	Severe
SF-36	Medical Outcomes Study Short-Form 36-Item Health Survey
SGRS	Stockton Geriatric Rating Scale
SHGRS	Stuard Hospital Geriatric Rating Scale
SIB	Severe Impairment Battery
SIP	Sickness Impact Profile
SKT	Syndrome Kurz test; Syndrome Short Test
SMQ	Squire's Memory Questionnaire
SMST	Sternberg's Memory Scanning Test
SPECT-TcHMPAO	Single Photon Emission Computed Tomography with hexamethylpropyleneamineoxime
SPET	Single Photon Emission Tomography
SRT	Selective Reminding Procedure
SRT-DR	Selective Reminding Procedure-Delayed Recall
SWFT	Semantic Word Fluency Test
TEP	Technical Expert Panel
TESS	Treatment Emergent Symptom Scale
TESS-DOTES	Dosage Record and Treatment Emergent Symptom Scales
tid	Three times daily
TK	Token Test
TOO	Task Order Officer
TP	Toulouse Piéron
TPAT	Toulouse-Pieron Attention Test

Acronyms and Abbreviations cont'd.

TSI	Test for Severe Impairment
UK	United Kingdom
UKU	Side effect rating scale
UPDRS	Unified Parkinson's Disease Rating Scale
US	United States
VaD	Vascular Dementia
VAMS	Visual Analog Mood Scale
VAS	Visual Analogue Scales
VHB	Videorecorder Home-Behavioral assessment
vs.	versus
w	week
WAIS	Wechsler Adult Intelligence Scale
WAIS-DI	Deterioration Index
WAIS-DSPT	Wechsler Adult Intelligence Scale – Digit Span Test
WAIS-DSST	Wechsler Adult Intelligence Scale –Digit Symbol Substitution Test
WAIS-DTIC	Wechsler Adult Intelligence Scale – Discovering Things in Common
WAIS-VOC	Wechsler Adult Intelligence Scale-Vocabulary Subset
WHO	World Health Organization
WLM	Word List Memory test
WMS-MQ	Wechsler Memory Scale-Memory Learning Restoration
WMS-R	Wechsler Memory Scale-Revised
χ^2	chi-square
y	year
Y	yes
ZVT	Zahlen-Verbindungs Test -Trail Making Test

Bibliography

Aarsland D, Larsen JP, Lim NG, et al. Olanzapine for psychosis in patients with Parkinson's disease with and without dementia. *J Neuropsychiatry Clin Neurosci* 1999; 11(3):392-4.

Status: Not included because dementia population not randomized to treatment

Aarsland D, Laake K, Larsen JP, et al. Donepezil for cognitive impairment in Parkinson's disease: A randomised controlled study. *J Neurol Neurosurg Psychiatry* 2002; 72(6):708-12.

Status: Cross-over trial

Aarsland D. Erratum: Donepezil for cognitive impairment in Parkinson's disease. A randomised controlled study. *J Neurol Neurosurg Psychiatry* 2002; 73(3):354.

Status: Not included because not a full article

Abalan F, Manciet G, Dartigues JF, et al. Nutrition and SDAT. *Biol Psychiatry* 1992 Jan 1; 31(1):103-5.

Status: Not included because not a full article

Abuzzahab FS, Sr., Merwin GE, Zimmermann RL, et al. A double-blind investigation of piracetam (nootropil) versus placebo in the memory of geriatric inpatients. *Psychopharmacol Bull* 1978 Jan; 14(1):23-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Abyad A. Prevalence of vitamin B12 deficiency among demented patients and cognitive recovery with cobalamin replacement. *J Nutr Health Aging* 2002; 6(4):254-60.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Adair JC, Knoefel JE, Morgan N. Controlled trial of N-acetylcysteine for patients with probable Alzheimer's disease. *Neurology* 2001 Oct 23; 57(8):1515-7.

Status: Included

Adler LA, Peselow E, Rosenthal M, et al. A controlled comparison of the effects of propranolol, benzotropine, and placebo on akathisia: An interim analysis. *Psychopharmacol Bull* 1993; 29(2):283-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Aerssens J, Raeymaekers P, Lilienfeld S, et al. APOE genotype: no influence on galantamine treatment efficacy nor on rate of decline in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2001 Mar; 12(2):69-77.

Status: Not included because does not meet criteria for treatment for dementia patients

Agid Y, Dubois B, Anand R, et al. Efficacy and tolerability of rivastigmine in patients with dementia of the Alzheimer type. *Curr Ther Res Clin Exp* 1998; 59(12):837-45.

Status: Included

Agnoli A, Martucci N, Manna V, et al. Effect of cholinergic and anticholinergic drugs on short-term memory in Alzheimer's dementia: a neuropsychological and computerized electroencephalographic study. *Clin Neuropharmacol* 1983; 6(4):311-23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Agnoli A, Martucci N, Manna V. Quantitative EEG as a tool in neuropharmacological studies: The effect of naftidrofuryl in chronic cerebrovascular diseases (C.C.V.D.). *Curr Ther Res Clin Exp* 1985; 37(3):387-97.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Agnoli A, Manna V, Martucci N, et al. Randomized double-blind study of flunarizine versus placebo in patients with chronic cerebrovascular disorders. *Int J Clin Pharmacol Res* 1988; 8(3):189-97.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Agnoli A, Martucci N, Fabbrini G, et al. Monoamine oxidase and dementia: Treatment with an inhibitor of MAO-B activity. *Dementia* 1990; 1(2):109-14.

Status: Article not retrievable

Agnoli A, Fabbrini G, Fioravanti M, et al. CBF and cognitive evaluation of Alzheimer type patients before and after IMAO-B treatment: a pilot study. *Eur Neuropsychopharmacol* 1992 Mar; 2(1):31-5.

Status: Included

Aguglia E, Caraceni T, Genitrini S, et al. Comparison of teniloxazine and piracetam in

Alzheimer-type or vascular dementia. *Curr Ther Res Clin Exp* 1995; 56(3):250-7.
Status: Not included because Jadad Quality Scale score less than three

Ahlin A, Nyback H, Junthe T, et al. THA in Alzheimer's dementia clinical biochemical and pharmacokinetic findings. *Alzheimer's disease basic mechanisms diagnosis and therapeutic strategies* 1990; 621-5.
Status: Not included because not a full article

Ahlin A, Nyback H, Junthe T, et al. Tetrahydroaminoacridine in Alzheimer's dementia: Clinical and biochemical results of a double-blind crossover trial. *Hum Psychopharmacol* 1991; (2):109-18.
Status: Cross-over trial

Ahlin A, Hassan M, Junthe T, et al. Tacrine in Alzheimer's disease: Pharmacokinetic and clinical comparison of oral and rectal administration. *Int Clin Psychopharmacol* 1994; 9(4):263-70.
Status: Cross-over trial

Aisen PS, Marin D, Davis KL. Anti-inflammatory drug studies in Alzheimer's disease. *Biol Psychiatry* 1996; 39(7):563
Status: Not included because not a full article

Aisen PS, Davis KL, Berg JD, et al. A randomized controlled trial of prednisone in Alzheimer's disease. *Alzheimer's Disease Cooperative Study. Neurology* 2000b; 54(3):588-93.
Status: Included

Aisen PS. Anti-inflammatory therapy for Alzheimer's disease: Implications of the prednisone trial. *Acta Neurol Scand Suppl* 2000a; 176:85-9.
Status: Companion of an included article

Aisen PS, Marin DB, Brickman AM, et al. Pilot tolerability studies of hydroxychloroquine and colchicine in Alzheimer disease. *Alzheimer Dis Assoc Disord* 2001 Apr; 15(2):96-101.
Status: Not included because dementia population not randomized to treatment

Aisen PS, Schmeidler J, Pasinetti GM. Randomized pilot study of nimesulide treatment in Alzheimer's disease. *Neurology* 2002 Apr 9; 58(7):1050-4.
Status: Included

Aisen PS, Berg JD, Craft S, et al. Steroid-induced elevation of glucose in Alzheimer's disease: Relationship to gender, apolipoprotein E genotype and cognition. *Psychoneuroendocrinology* 2003; 28(1):113-20.
Status: Not included because dementia population not randomized to treatment

Ala T, Romero S, Knight F, et al. GM-1 treatment of Alzheimer's disease. A pilot study of safety and efficacy. *Arch Neurol* 1990 Oct; 47(10):1126-30.
Status: Included

Alafuzoff I, Helisalmi S, Heinonen EH, et al. Selegiline treatment and the extent of degenerative changes in brain tissue of patients with Alzheimer's disease. *Eur J Clin Pharmacol* 2000 Feb; 55(11-12):815-12.
Status: Not included because no extractable data relevant to review

Albizzati MG, Bassi S, Calloni E, et al. Cyclandelate versus flunarizine. A double-blind study in a selected group of patients with dementia. *Drugs* 1987; 33(Suppl 2):90-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Aldenkamp AP, van Wieringen A, Alpherts WC, et al. Double-blind placebo-controlled, neuropsychological and neurophysiological investigations with oxiracetam (CGP 21690E) in memory-impaired patients with epilepsy. *Neuropsychobiology* 1990 Sep; 24(2):90-101.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Alexopoulos GS, Meyers BS, Young RC, et al. Executive dysfunction and long-term outcomes of geriatric depression. *Arch Gen Psychiatry* 2000 Mar; 57(3):285-90.
Status: Not included because does not meet criteria for treatment for dementia patients

Allain H, Denmat J, Bentue-Ferrer D, et al. Randomized, double-blind trial of exifone versus cognitive problems in Parkinson's disease. *Fundam Clin Pharmacol* 1988; 2(1):1-12.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Allain H, Raoul P, Lieury A, et al. Effect of two doses of Gingko biloba extract (EGb 761) on the dual-coding test in elderly subjects. *Clin Ther* 1993; 15(3):549-58.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Allain H, Neuman E, Malbezin M, et al. Bridging study of S12024 in 53 in-patients with Alzheimer's disease. *J Am Geriatr Soc* 1997; 45(1):125-6.
Status: Not included because not a full article

Allain H, Schuck S, Lebreton S, et al. Aminotransferase levels and silymarin in de novo tacrine-treated patients with Alzheimer's disease. *Dement Geriatr Cogn Disord* 1999 May; 10(3):181-5.
Status: Included

Allain H, Dautzenberg PH, Maurer K, et al. Double blind study of tiapride versus haloperidol and placebo in agitation and aggressiveness in elderly patients with cognitive impairment. *Psychopharmacologia* 2000 Mar; 148(4):361-6.
Status: Included

Almkvist O, Jelic V, Amberla K, et al. Responder characteristics to a single oral dose of cholinesterase inhibitor: A double-blind placebo-controlled study with tacrine in Alzheimer patients. *Dement Geriatr Cogn Disord* 2001 Jan; 12(1):22-32.
Status: Cross-over trial;
Cross-over trial

Als-Nielsen B, Chen W, Gluud C, et al. Association of funding and conclusions in randomized drug trials: a reflection of treatment effect or adverse events? *JAMA* 2003 Aug 20; 290(7):921-8.
Status: Background article;

Altman H, Mehta D, Evenson RC, et al. Behavioral effects of drug therapy on psychogeriatric inpatients. II. Multivitamin supplement. *J Am Geriatr Soc* 1973 Jun; 21(6):249-52.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Altman H, Mehta D, Evenson RC, et al. Behavioral effects of drug therapy on psychogeriatric inpatients. I. Chlorpromazine and thioridazine. *J Am Geriatr Soc* 1973 Jun; 21(6):241-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Alvarez XA, Laredo M, Corzo D, et al. Citicoline improves memory performance in elderly

subjects. *Methods & Findings in Experimental & Clinical Pharmacology* 1997 Apr; 19(3):201-10.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Alvarez XA, Mouzo R, Pichel V, et al. Double-blind placebo-controlled study with citicoline in APOE genotyped Alzheimer's disease patients. Effects on cognitive performance, brain bioelectrical activity and cerebral perfusion. *Methods & Findings in Experimental & Clinical Pharmacology* 1999 Nov; 21(9):633-44.
Status: Not included because Jadad Quality Scale score less than three

Alvarez XA, Pichel V, Perez P, et al. Double-blind, randomized, placebo-controlled pilot study with anapso in senile dementia: Effects on cognition, brain bioelectrical activity and cerebral hemodynamics. *Methods & Findings in Experimental & Clinical Pharmacology* 2000 Sep; 22(7):585-94.
Status: Included

Amaducci L. Phosphatidylserine in the treatment of Alzheimer's disease: Results of a multicenter study. *Psychopharmacol Bull* 1988; 24(1):130-4.
Status: Included

Amaducci L, Maurer K, Winblad B, et al. A long-term, double-blind, placebo-controlled efficacy and safety study of nicergoline in patients with mild to moderate Alzheimer's disease. *J Eur Coll Neuropsychopharmacol* 1999; (Suppl 5):S323.
Status: Not included because not a full article

Amaducci LA, Fratiglioni L, Rocca WA, et al. Risk factors for clinically diagnosed Alzheimer's disease: a case-control study of an Italian population. *Neurology* 1986 Jul; 36(7):922-31.
Status: Background article

Amar K, Wilcock GK, Scot M, et al. The presence of leuko-araiosis in patients with Alzheimer's disease predicts poor tolerance to tacrine, but does not discriminate responders from non-responders. *Age Ageing* 1997 Jan; 26(1):25-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ambrozi L, Danielczyk W. Treatment of impaired cerebral function in psychogeriatric patients with memantine: Results of a phase II double-blind study. *Pharmacopsychiatry* 1988 May; 21(3):144-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

American Psychiatric Association. Diagnostic and statistical manual of mental disorders. 3rd ed. Washington, DC: American Psychiatric Association; 1980.

Status: Background article

American Psychiatric Association. Diagnostic and statistical manual of mental disorders (3rd ed. rev.): DSM-III-R. Washington, DC: American Psychiatric Association; 1987.

Status: Background article

American Psychiatric Association. Diagnostic criteria from DSM-IV. 1994.

Status: Background article

Anand R, Gharabawi G, Enz A. Efficacy and safety results of the early phase studies with Exelon(tm) (ENA-713) in Alzheimer's disease: An overview. J Drug Dev Clin Pract 1996; 1-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ananth JV, Deutsch M, Ban TA. Senilex in the treatment of geriatric patients. Curr Ther Res Clin Exp ; 13(5):316-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ancill RJ, Carlyle WW, Liang RA, et al. Agitation in the demented elderly: A role for benzodiazepines? Int Clin Psychopharmacol 1991; 6(3):141-6.

Status: Included

Ancoli-Israel S, Martin JL, Kripke DF, et al. Effect of light treatment on sleep and circadian rhythms in demented nursing home patients. J Am Geriatr Soc 2002; 50(2):282-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anderer P, Barbanoj MJ, Saletu B, et al. Restriction to a limited set of EEG-target variables may lead to misinterpretation of pharmaco-EEG results. Neuropsychobiology 1993; 27(2):112-6.

Status: Not included because no extractable data relevant to review

Anderson J, Arens K, Johnson R, et al. Spaced retrieval vs. memory tape therapy in memory rehabilitation for dementia of the Alzheimer's type.

Clin Gerontol 2001; (1-2):123-39.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Darvon and Darvon-N. Med Lett Drugs Ther 1972 May; 14(11):37-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Severely demented patients beyond help of drugs. Modern Geriatrics 1976; (10):36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Double-blind, placebo-controlled evaluation of cinromide in patients with the Lennox-Gastaut Syndrome. The Group for the Evaluation of Cinromide in the Lennox-Gastaut Syndrome. Epilepsia 1989 Jul; 30(4):422-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. The International classification of diseases, 9th revision, clinical modification: ICD-9-CM. 3rd edition. Washington, DC: US Department of Health and Human Services; 1989.

Status: Background article

Anonymous. Searle clinical research report. A two year, multicenter, randomized, double-blind controlled parallel group study of cycloserine in the treatment of Alzheimer's disease with six months placebo control. Report No. NC6-93-06-009. Skokie, IL: GD Searle. 1993.

Status: Article not retrievable

Anonymous. Safety and tolerability of the antioxidant OPC-14117 in HIV-associated cognitive impairment. The Dana Consortium on the Therapy of HIV Dementia and Related Cognitive Disorders. Neurology 1997 Jul; 49(1):142-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Study suggests antioxidants slow decline in Alzheimer's disease. Am J Health Syst Pharm 1997; 54(13):1478.

Status: Not included because not a full article

Anonymous. Benefits of new Alzheimer disease therapies. J Pharm Technol 1998; 14(3):125.

Status: Not included because not a full article

Anonymous. Eptastigmine tartrate: Cognition enhancer acetylcholinesterase inhibitor. *Drugs of the Future* 1998; 23(2):217-8.

Status: Not included because not a full article

Anonymous. Selegiline hydrochloride: Antiparkinsonian cognition enhancer. *Drugs of the Future* 1998; 23(2):240-1.

Status: Not included because not a full article

Anonymous. Tacrine and Alzheimer disease. *WHO Drug Information* 1999; 13(1):7-8.

Status: Not included because not a full article

Anonymous. The alternative to tube-feeding patients with advanced dementia. *Volunt Leader* 1999; 40(4):13-4.

Status: Not included because not a full article

Anonymous. Phosphatidylserine. *Altern Med Rev* 1999; 4(2):115-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Rivastigmine for Alzheimer's disease. *Drug Ther Bull* 2000; 38(2):15-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Erratum: Estrogen replacement therapy for treatment of mild to moderate Alzheimer disease: A randomized controlled trial (*Journal of the American Medical Association* (February 23,2000) 283 (1007-1015)). *JAMA* 2000; 284(20):2597

Status: Not included because not a full article

Anonymous. New hope for early Alzheimer's disease. *Harv Womens Health Watch* 2000 Apr; 7(8):7

Status: Not included because not a full article

Anonymous. Lead success for Nuerogen. *Manuf Chem* 2001; 72(4):11

Status: Not included because not a full article

Anonymous. Caregiver experience relates to clinical trial involvement: research looks at how caregivers of Alzheimer's patients make decisions regarding care. *Caremanagement* 2001 Jun; 7(3):55.

Status: Not included because not a full article

Anonymous. Idebenone. *Altern Med Rev* 2001; 6(1):83-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Galantamine (Reminyl) for Alzheimer's disease. *Med Lett Drugs Ther* 2001; 43(1107):53-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Galantamine: New preparation. The fourth cholinesterase inhibitor for Alzheimer's disease. *Prescrire Int* 2001 Dec; 10(56):180-1.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Aromatherapy trial. *Journal of Dementia Care* 2001; 9(6):38.

Status: Not included because not a full article

Anonymous. Erratum: A 24-week, randomized, double-blind study of donepezil in moderate to severe alzheimer's disease (*Neurology* (2001) 57 (613-620)). *Neurology* 2001; 57(11):2153.

Status: Not included because not a full article

Anonymous. New Alzheimer's drug is first therapy to show efficacy in vascular dementia. *Hosp Formul* 2001; 36(8):569.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Double-blind trial will compare two anti-Alzheimer's drugs. *Journal of Dementia Care* 2001; 9(5):6

Status: Not included because not a full article

Anonymous. Colostrinin. *Journal of Dementia Care* 2001; 9(6):37.

Status: Not included because not a full article

Anonymous. Galantamine effective in treating dementia in patients with cerebrovascular disease. *Pharm J* 2001; 266(7153):842.

Status: Not included because not a full article

Anonymous. Drug that modulates glutamate levels promising for Alzheimer's disease. *Pharm J* 2002; 269(7209):152.

Status : Not included because not a full article

Anonymous. NSAIDs: Protection against Alzheimer's? *Med Today* 2002; 3(2):9

Status: Not included because not a full article

Anonymous. Drugs to treat dementia and psychosis: Management of Parkinson's disease. *Mov Disord* 2002; 17(Suppl 4):S120-S127.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Trial of new immunotherapeutic agent for Alzheimer's suspended. *Pharm J* 2002; 268(7187):279
Status: Not included because not a full article

Anonymous. Reminyl of benefit in vascular dementia. *Pharm J* 2002; 268(7194):526.
Status: Not included because not a full article

Anonymous. Perindopril protects against dementia. *Pharm J* 2002; 268(7204):899.
Status: Not included because not a full article

Anonymous. Memantine launched for treatment of Alzheimer's. *Pharm J* 2002; 269(7219):516.
Status: Not included because not a full article

Anonymous. Greater satisfaction, ease of use reported with donepezil versus galantamine. *Hosp Formul* 2002; 37(8):383-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. NSAID use could reduce Alzheimer's risk. *Pharm J* 2002; 269(7217):428.
Status: Not included because not a full article

Anonymous. Risperdal (risperidone) and cerebrovascular adverse events in placebo-controlled dementia trials. http://www.hc-sc.gc.ca/hpb-dgps/therapeut/zfiles/english/advisory/industry/risperdal1_e.pdf. *Background article*

Anonymous. USP DI: Drug information for the health care professional. 23rd Edition. Taunton, Massachusetts: Thomsom Micromedex; 2003.
Status: Background article

Anonymous. CPS: Compendium of pharmaceuticals and specialties: The Canadian drug reference for health professionals. Mississauga, ON: Canadian Pharmacists Association; 2003.
Status: Background article

Antuono PG. Effectiveness and safety of velnacrine for the treatment of Alzheimer's disease. A double-blind, placebo-controlled study.

Mentane Study Group. *Arch Intern Med* 1995 Sep 11; 155(16):1766-72.
Status: Included

Antuono P, Doody R, Gilman S, et al. Diagnostic criteria for dementia in clinical trials: Position paper from the International Working Group on Harmonization of Dementia Drug Guidelines. *Alzheimer Dis Assoc Disord* 1997; 11(Suppl 3):22-5.
Status: Background article

Aranda B, Dumoulin P, Groothold G. Controlled study of the effect of dihydroergocristine on organic brain syndrome. *Arzneimittelforschung* 1992; 42(11 A):1406-9.
Status: Background article

Arendt G, von Giesen HJ, Hefter H, et al. Therapeutic effects of nucleoside analogues on psychomotor slowing in HIV infection. *AIDS* 2001 Mar 9; 15(4):493-500.
Status: Not included because dementia population not randomized to treatment

Areosa Sastre A, Sherriff F. Memantine for dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software
Status: Background article

Areosa Sastre A, Sherriff F. Memantine for dementia. In: *The Cochrane Library*, 2003. Issue 1. Oxford: Update Software
Status: Background article

Arkin SM. Alzheimer memory training: Quizzes beat repetition, especially with more impaired. *Am J Alzheimers Dis* 1997; (4):147-58.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Arkin SM. Alzheimer memory training: Students replicate learning successes. *Am J Alzheimers Dis* 2000 May; 15(3):152-62.
Status: Not included because dementia population not randomized to treatment

Arrigo A, Moglia A, Borsotti L. A double-blind, placebo-controlled, crossover trial with nicergoline in patients with senile dementia. *Int J Clin Pharmacol Res* 1982; 2(4 Suppl 1):33-41.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Arrigo A, Casale R, Giorgi I, et al. Effects of intravenous high dose c-dergocrine mesylate ('Hydergine' (R)) in elderly patients with severe multi-infarct dementia: A double-blind, placebo-controlled trial. *Curr Med Res Opin* 1989; 11(8):491-500.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ashford JW, Soldinger S, Schaeffer J, et al. Physostigmine and its effect on six patients with dementia. *Am J Psychiatry* 1981; 138(6):829-30.
Status: Not included because dementia population not randomized to treatment

Asthana S, Raffaele KC, Berardi A, et al. Treatment of Alzheimer's disease by continuous intravenous infusion of physostigmine. *Alzheimer Dis Assoc Disord* 1995; 9(4):223-32.
Status: Cross-over trial;
Cross-over trial

Asthana S, Greig NH, Holloway HW, et al. Clinical pharmacokinetics of arecoline in subjects with Alzheimer's disease. *Clin Pharmacol Ther* 1996 Sep; 60(3):276-82.
Status: Not included because no extractable data relevant to review

Asthana S. Estrogen Patch Boosts Memory in Alzheimer's? *Fam Pract News* 1998;
Status: Article not retrievable

Asthana S, Raffaele KC, Greig NH, et al. Neuroendocrine responses to intravenous infusion of physostigmine in patients with Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1999 Apr; 13(2):102-8.
Status: Cross-over trial;
Cross-over trial

Asthana S, Craft S, Baker LD, et al. Cognitive and neuroendocrine response to transdermal estrogen in postmenopausal women with Alzheimer's disease: Results of a placebo-controlled, double-blind, pilot study. *Psychoneuroendocrinology* 1999 Aug; 24(6):657-77.
Status: Not included because Jadad Quality Scale score less than three

Asthana S, Baker LD, Craft S, et al. High-dose estradiol improves cognition for women with AD: Results of a randomized study. *Neurology* 2001 Aug 28; 57(4):605-12.
Status: Included

Ather SA, Shaw SH, Stoker MJ. A comparison of chlormethiazole and thioridazine in agitated confusional states of the elderly. *Acta Psychiatr Scand* 1986; 73(Suppl 329):81-91.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Auchus AP, Bissey-Black C. Pilot study of haloperidol, fluoxetine, and placebo for agitation in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1997; 9(4):591-3.
Status: Included

Auer SR, Monteiro IM, Reisberg B. Behavioral symptoms in dementia: community-based research. *Int Psychogeriatr* 1996; 8(Suppl 3):363-6, 381, 382.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Avorn J, Soumerai SB, Everitt DE, et al. A randomized trial of a program to reduce the use of psychoactive drugs in nursing homes. *N Engl J Med* 1992 Jul 16; 327(3):168-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Avorn J, Benner J, Ford I, et al. Measuring the cost-effectiveness of lipid-lowering drugs in the elderly: The outcomes research and economic analysis components of the PROSPER trial. *Control Clin Trials* 2002; 23(6):757-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Azuma T, Nagai Y, Saito T, et al. The effect of dehydroepiandrosterone sulfate administration to patients with multi-infarct dementia. *JNS* 1999 Jan 1; 162(1):69-73.
Status: Not included because dementia population not randomized to treatment

Bach D, Bach M, Bohmer F, et al. Reactivating occupational therapy: A method to improve cognitive performance in geriatric patients. *Age Ageing* 1995 May; 24(3):222-6.
Status: Not included because does not meet criteria for treatment for dementia patients

Bachynsky J, McCracken P, Lier D, et al. Propentofylline treatment for Alzheimer disease and vascular dementia: An economic evaluation based on functional abilities. *Alzheimer Dis Assoc Disord* 2000 Apr; 14(2):102-11.
Status: Not included because Jadad Quality Scale score less than three

Backonja M, Beydoun A, Edwards KR, et al. Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus: A randomized controlled trial. *JAMA* 1998 Dec 2; 280(21):1831-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Badia X, Herdman M. The importance of health-related quality-of-life data in determining the value of drug therapy. *Clin Ther* 2001 Jan; 23(1):168-75.

Status: Background article

Bae CY, Cho CY, Cho K, et al. A double-blind, placebo-controlled, multicenter study of Cerebrolysin for Alzheimer's disease. *J Am Geriatr Soc* 2000 Dec; 48(12):1566-71.

Status: Included

Baines S, Saxby P, Ehler K. Reality orientation and reminiscence therapy. A controlled cross-over study of elderly confused people. *Br J Psychiatry* 1987; Vol 151:222-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baker R, Dowling Z, Wareing LA, et al. Snoezelen: Its long-term and short-term effects on older people with dementia. *Br J Occup Ther* 1997; (5):213-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baker R, Bell S, Baker E, et al. A randomized controlled trial of the effects of multi-sensory stimulation (MSS) for people with dementia. *Br J Clin Psychol* 2001 Mar; 40(Pt 1):1-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baladi JF, Bailey PA, Black S, et al. Rivastigmine for Alzheimer's disease: Canadian interpretation of intermediate outcome measures and cost implications. *Clin Ther* 2000 Dec; 22(12):1549-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baldereschi M, Di Carlo A, Lepore V, et al. Estrogen-replacement therapy and Alzheimer's disease in the Italian Longitudinal Study on Aging. *Neurology* 1998 Apr; 50(4):996-1002.

Status: Not included because dementia population not randomized to treatment

Balestreri R, Bompani R, Cerrato G. Comparative study of suloctidil and dihydroergotoxine in chronic cerebrovascular insufficiency. Results of a double blind double dummy multicentric trial. *Acta Ther* 1984; 10(2):163-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Balestreri R, Fontana L, Astengo F. A double-blind placebo controlled evaluation of the safety and efficacy of vinpocetine in the treatment of patients with chronic vascular senile cerebral dysfunction. *J Am Geriatr Soc* 1987 May; 35(5):425-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ball JA, Taylor AR. Effect of cyclandelate on mental function and cerebral blood flow in elderly patients. *BMJ* 1967 Aug 26; 3(564):525-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ballard C, O'Brien J, James I, et al. Quality of life for people with dementia living in residential and nursing home care: The impact of performance on activities of daily living, behavioral and psychological symptoms, language skills, and psychotropic drugs. *Int Psychogeriatr* 2001 Mar; 13(1):93-106.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ballard C, Powell I, James I, et al. Can psychiatric liaison reduce neuroleptic use and reduce health service utilization for dementia patients residing in care facilities? *Int J Geriatr Psychiatry* 2002 Feb; 17(2):140-5.

Status: Not included because dementia population not randomized to treatment

Ballard CG, O'Brien JT, Reichelt K, et al. Aromatherapy as a safe and effective treatment for the management of agitation in severe dementia: The results of a double-blind, placebo-controlled trial with Melissa. *J Clin Psychiatry* 2002; 63(7):553-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Balldin J, Gottries CG, Karlson I, et al. Relationship between DST and the serotonergic system. Results from treatments with two 5-HT reuptake blockers in dementia disorders. *Int J Geriatr Psychiatry* 1988; 3(1):17-26.

Status: Not included because no extractable data relevant to review

Bambasova E, Bilkova J, Budinska K. Papaverin in the treatment of geriatric patients. *Act Nerv Super (Praha)* 1974 Aug; 16(3):192-3.

Status: Not included because dementia population not randomized to treatment

Ban TA, Modafferi A, Morey L. Global changes with glycosaminoglycan polysulfate in primary degenerative and multi-infarct dementia. *Curr Ther Res Clin Exp* 1987; 41(5):631-6.

Status: Not included because Jadad Quality Scale score less than three

Ban TA, Morey L, Aguglia E, et al. Nimodipine in the treatment of old age dementias. *Prog Neuropsychopharmacol Biol Psychiatry* 1990; 14(4):525-51.

Status: Included

Ban TA, Morey LC, Aguglia E, et al. Glycosaminoglycan polysulfate in the treatment of old age dementias. *Prog Neuropsychopharmacol Biol Psychiatry* 1991a; 15(3):323-42.

Status: Included

Ban TA, Morey LC, Santini V. Clinical investigations with ateroid in old-age dementias. *Semin Thromb Hemost* 1991b; 17(Suppl 2):161-3.

Status: Included

Ban TA, Morey LC, Fjetland OK, et al. Early manifestations of dementing illness: Treatment with glycosaminoglycan polysulfate. *Prog Neuropsychopharmacol Biol Psychiatry* 1992 Sep; 16(5):661-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Banerjee S. Randomized controlled trials. *Int Rev Psychiatry* 1998; 10(4):291-303.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Barak Y, Levine J, Glasman A, et al. Inositol treatment of Alzheimer's disease: A double blind, cross-over placebo controlled trial. *Prog Neuropsychopharmacol Biol Psychiatry* 1996 May; 20(4):729-35.

*Status: Cross-over trial;
Cross-over trial*

Barnes R, Veith R, Okimoto J. Efficacy of antipsychotic medications in behaviorally disturbed dementia patients. *Am J Psychiatry* 1982; 139(9):1170-4.

Status: Included

Baro F, Malfroid M, Waegemans T, et al. Double-blind trial of suloctidil versus placebo in moderate to severe mental deterioration. *Pharmatherapeutica* 1985; 4(6):399-404.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bass DM, McClendon MJ, Deimling GT, et al. The influence of a diagnosed mental impairment on family caregiver strain. *J Gerontol* 1994 May; 49(3):S146-S155

Status: Background article

Bass DM, McClendon MJ, Brennan PF, et al. The buffering effect of a computer support network on caregiver strain. *J Aging Health* 1998; 10(1):20-43.

Status: Not included because does not meet criteria for treatment for dementia patients

Bassi S, Albizzati MG, Corsini GU, et al. Therapeutic experience with transdihydrolisuride in Huntington's disease. *Neurology* 1986; 36(7):984-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Battaglia A, Bruni G, Ardia A, et al. Nicergoline in mild to moderate dementia. A multicenter, double-blind, placebo-controlled study. *J Am Geriatr Soc* 1989 Apr; 37(4):295-302.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Battaglia A, Bruni G, Sacchetti G, et al. A double-blind randomized study of two ergot derivatives in mild to moderate dementia. *Curr Ther Res Clin Exp* 1990; 48(4):597-612.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Battistin L, Pizzolato G, Dam M, et al. Effects of acetyl-L-carnitine (ALC) treatment in dementia: A multicentric, randomized, double-blind study. *New Trends in Clinical Neuropharmacology* 1989; (2):131-2.

Status: Not included because Jadad Quality Scale score less than three

Baumgarten M. The health of persons giving care to the demented elderly: a critical review of the literature. *J Clin Epidemiol* 1989; 42(12):1137-48.
Status: Background article

Baumgarten M, Hanley JA, Infante-Rivard C, et al. Health of family members caring for elderly persons with dementia. A longitudinal study. *Ann Intern Med* 1994 Jan 15; 120(2):126-32.
Status: Background article

Bavazzano A, Guarducci R, Gestri G, et al. Clinical trial with amantadine and hydergine in elderly patients. *J Clin Exp Gerontol* 1980; (4):289-99.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bazo AJ. An ergot alkaloid preparation (Hydergine) versus papaverine in treating common complaints of the aged: Double-blind study. *J Am Geriatr Soc* 1973 Feb; 21(2):63-71.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beck C, Heacock P, Mercer SO, et al. Improving dressing behavior in cognitively impaired nursing home residents. *Nurs Res* 1997 May; 46(3):126-32.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beck C, Cody M, Souder E, et al. Dementia diagnostic guidelines: Methodologies, results, and implementation costs. *J Am Geriatr Soc* 2000; 48(10):1195-203.
Status: Background article

Beck CK, Vogelpohl TS, Rasin JH, et al. Effects of behavioral interventions on disruptive behavior and affect in demented nursing home residents. *Nurs Res* 2002; 51(4):219-28.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beck C, Heacock P, Mercer S, et al. The impact of cognitive skills remediation training on persons with Alzheimer's disease or mixed dementia. *J Geriatr Psychiatry* 1988; 21(1):73-88.
Status: Not included because dementia population not randomized to treatment

Becker RE, Colliver JA, Markwell SJ, et al. Double-blind, placebo-controlled study of metrifonate, an acetylcholinesterase inhibitor, for Alzheimer's disease. *Alzheimer Dis Assoc Disord*

1996; 10(3):124-31.
Status: Included

Becker RE, Colliver JA, Markwell SJ, et al. Effects of metrifonate on cognitive decline in Alzheimer's disease: A double-blind, placebo-controlled, 6-month study. *Alzheimer Dis Assoc Disord* 1998 Mar; 12(1):54-7.
Status: Included

Beckers T, Wagemans J, Boucart M, et al. Different effects of lorazepam and diazepam on perceptual integration. *Vision Res* 2001 Aug; 41(17):2297-303.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beckett LA. Community-based studies of Alzheimer's disease: statistical challenges in design and analysis. *Stat Med* 2000 Jun 15; 19(11-12):1469-80.
Status: Background article

Beckmann J. Basic aspects of risk-benefit analysis. *Semin Thromb Hemost* 1999; 25(1):89-95.
Status: Background article

Bedard M, Molloy DW, Standish T, et al. Clinical trials in cognitively impaired older adults: Home versus clinic assessments. *J Am Geriatr Soc* 1995 Oct; 43(10):1127-30.
Status: Not included because does not meet criteria for treatment for dementia patients

Bedard MA, Pillon B, Dubois B, et al. Acute and long-term administration of anticholinergics in Parkinson's disease: Specific effects on the subcortico-frontal syndrome. *Brain Cogn* 1999 Jul; 40(2):289-313.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beiser A, D'Agostino RB, Sr., Seshadri S, et al. Computing estimates of incidence, including lifetime risk: Alzheimer's disease in the Framingham Study. The Practical Incidence Estimators (PIE) macro. *Stat Med* 2000 Jun 15; 19(11-12):1495-522.
Status: Background article

Belanoff JK, Jurik J, Schatzberg LD, et al. Slowing the progression of cognitive decline in Alzheimer's disease using mifepristone. *J Mol Neurosci* 2002; 19(1-2):201-6.

Status: Not included because no extractable data relevant to review

Belfiore G, Di Maio L, Napolitano G, et al. Long-term effect of a single dose of flunarizine in Huntington's disease. *Eur J Neurol* 1998; 5(3):249-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bell IR, Edman JS, Morrow FD, et al. Brief communication. Vitamin B1, B2, and B6 augmentation of tricyclic antidepressant treatment in geriatric depression with cognitive dysfunction. *J Am Coll Nutr* 1992 Apr; 11(2):159-63.

Status: Not included because does not meet criteria for treatment for dementia patients

Beller SA, Overall JE, Swann AC. Efficacy of oral physostigmine in primary degenerative dementia. A double-blind study of response to different dose level. *Psychopharmacologia* 1985; 87(2):147-51.

*Status: Cross-over trial;
Cross-over trial*

Bellus SB, Vergo JG, Kost PP, et al. Behavioral rehabilitation and the reduction of aggressive and self-injurious behaviors with cognitively impaired, chronic psychiatric inpatients. *Psychiatr Q* 1999; 70(1):27-37.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ben Shlomo Y, Churchyard A, Head J, et al. Investigation by Parkinson's Disease Research Group of United Kingdom into excess mortality seen with combined levodopa and selegiline treatment in patients with early, mild Parkinson's disease: Further results of randomised trial and confidential inquiry. *BMJ* 1998 Apr 18; 316(7139):1191-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Benedict RH, Shapiro A, Priore R, et al. Neuropsychological counseling improves social behavior in cognitively-impaired multiple sclerosis patients. *Mult Scler* 2000 Dec; 6(6):391-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bentham PW. A double-blind placebo-controlled trial of L-tryptophan to assess the degree of cognitive and behavioural improvement in patients with Alzheimer-type dementia and to compare differential response in clinical sub-groups. *Int*

Clin Psychopharmacol 1990; 5(4):261-72.

Status: Not included because Jadad Quality Scale score less than three

Bergamasco B, Villardita C, Coppi R. Effects of idebenone in elderly subjects with cognitive decline. Results of a multicentre clinical trial. *Arch Gerontol Geriatr* 1992; 15(3):279-86.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bergamasco B, Villardita C, Coppi R. Idebenone in the treatment of multi-infarct dementia: A randomised, double-blind, placebo controlled multicentre trial. *Arch Gerontol Geriatr* 1992; 15(3):271-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bergamasco B, Scarzella L, La Commare P. Idebenone, a new drug in the treatment of cognitive impairment in patients with dementia of the Alzheimer type. *Funct Neurol* 1994 May; 9(3):161-8.

Status: Included

Bergman I, Brane G, Gottfries CG, et al. Alaproclate: A pharmacokinetic and biochemical study in patients with dementia of Alzheimer type. *Psychopharmacology (Berl)* 1983; 80(3):279-83.

Status: Background article

Bernardi F, Lanzone A, Cento RM, et al. Allopregnanolone and dehydroepiandrosterone response to corticotropin-releasing factor in patients suffering from Alzheimer's disease and vascular dementia. *Eur J Endocrinol* 2000; 142(5):466-71.

Status: Not included because does not meet criteria for treatment for dementia patients

Besson JAO, Palin AN, Ebmeier KP, et al. Calcium antagonists and multi-infarct dementia: A trial involving sequential NMR and psychometric assessment. *Int J Geriatr Psychiatry* 1988; 3(2):99-105.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Biegel DE, Sales E, Schultz R. Family caregiving in chronic illness. Newbury Park: Sage; 1991.

Status: Background article

Bierer LM, Aisen PS, Davidson M, et al. A pilot study of oral physostigmine plus yohimbine in patients with Alzheimer disease. *Alzheimer Dis*

Assoc Disord 1993; 7(2):98-104.

Status: Not included because dementia population not randomized to treatment

Bierer LM, Aisen PS, Davidson M, et al. A pilot study of clonidine plus physostigmine in Alzheimer's disease. Dementia 1994 Sep; 5(5):243-6.

Status: Not included because Jadad Quality Scale score less than three

Binder EF, Schechtman KB, Birge SJ, et al. Effects of hormone replacement therapy on cognitive performance in elderly women. Maturitas 2001 Apr; 38(2):137-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Birge SJ. The role of estrogen in the treatment of Alzheimer's disease. Neurology 1997; 48(5 Suppl 7):S36-S41.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Birkett DP, Boltuch B. Chlorpromazine in geriatric psychiatry. J Am Geriatr Soc 1972 Aug; 20(8):403-6.

Status: Not included because no extractable data relevant to review

Birks J, Grimley EG, Van Dongen M. Ginkgo biloba for dementia and cognitive impairment. In: The Cochrane Library, 2002. Issue 3. Oxford:Update Software.

Status: Background article

Birks J, Grimley EJ, Iakovidou V, et al. Rivastigmine for Alzheimer's disease. In: The Cochrane Library, 2000. Issue 4. Oxford: Update Software

Status: Background article

Birks J, Flicker L. Selegiline for Alzheimer's disease. In: The Cochrane Library, 2000. Issue 2. Oxford: Update Software.

Status: Background article

Birks JS, Melzer D, Beppu H. Donepezil for mild and moderate Alzheimer's disease. In: The Cochrane Library, 2000. Issue 4. Oxford: Update Software.

Status: Background article

Bischkopf J, Busse A, Angermeyer MC. Mild cognitive impairment: A review of prevalence,

incidence and outcome according to current approaches. Acta Psychiatr Scand 2002 Dec; 106(6):403-14.

Status: Background article

Bjorkman T, Hansson L, Sandlund M. Outcome of case management based on the strengths model compared to standard care. A randomised controlled trial. Soc Psychiatry Psychiatr Epidemiol 2002 Apr; 37(4):147-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Black RS, Barclay LL, Nolan KA, et al. Pentoxifylline in cerebrovascular dementia. J Am Geriatr Soc 1992 Mar; 40(3):237-44.

Status: Included

Black SE, Patterson C, Feightner J. Preventing dementia. Can J Neurol Sci 2001 Feb; 28(Suppl 1):S56-S66

Status: Background article

Blaha L, Erzigkeit H, Adamczyk A, et al. Clinical evidence of the effectiveness of vinpocetine in the treatment of organic psychosyndrome. Hum Psychopharmacol 1989; (2):103-11.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blass JP, Gleason P, Brush D, et al. Thiamine and Alzheimer's disease. A pilot study. Arch Neurol 1988 Aug; 45(8):833-5.

Status: Cross-over trial;

Cross-over trial

Blass JP, Cyrus PA, Bieber F, et al. Randomized, double-blind, placebo-controlled, multicenter study to evaluate the safety and tolerability of metrifonate in patients with probable Alzheimer's disease. Alzheimer Dis Assoc Disord 2000; 14(1):39-45.

Status: Not included because no extractable data relevant to review

Blazer DG, Landerman LR, Hays JC, et al. Symptoms of depression among community-dwelling elderly African-American and white older adults. Psychol Med 1998 Nov; 28(6):1311-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blin J, Piercey MF, Giuffra ME, et al. Metabolic effects of scopolamine and physostigmine in human brain measured by positron emission tomography. JNS 1994 May; 123(1-2):44-2.

Status: Not included because does not meet criteria for treatment for dementia patients

Blin J, Ivanoiu A, Coppens A, et al. Cholinergic neurotransmission has different effects on cerebral glucose consumption and blood flow in young normals, aged normals, and Alzheimer's disease patients. *Neuroimage* 1997 Nov; 6(4):335-43.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blin J, Ivanoiu A, De Volder A, et al. Physostigmine results in an increased decrement in brain glucose consumption in Alzheimer's disease. *Psychopharmacologia* 1998 Apr; 136(3):256-63.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Block RI, DeVoe M, Stanley B, et al. Memory performance in individuals with primary degenerative dementia: Its similarity to diazepam-induced impairments. *Exp Aging Res* 1985; 11(3-4):151-4.

Status: Not included because does not meet criteria for treatment for dementia patients

Blume J, Ruhlmann KU, de la Haye R, et al. Treatment of chronic cerebrovascular disease in elderly patients with pentoxifylline. *J Med* 1992; 23(6):417-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bodick NC, Offen WW, Levey AI, et al. Effects of xanomeline, a selective muscarinic receptor agonist, on cognitive function and behavioral symptoms in Alzheimer's disease. *Arch Neurol* 1997 Apr; 54(4):465-73.

Status: Included

Bodick NC, Offen WW, Shannon HE, et al. The selective muscarinic agonist xanomeline improves both the cognitive deficits and behavioral symptoms of Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1997; 11(Suppl 4):S16-S22.

Status: Not included because Jadad Quality Scale score less than three

Bodick N, Forette F, Hadler D, et al. Protocols to demonstrate slowing of Alzheimer's disease progression: Position paper from the International Working Group on Harmonization of Dementia Drug Guidelines. *Alzheimer Dis Assoc Disord*

1997; 11(Suppl 3):50-3.

Status: Background article

Boelhouwer C, Henry CE, Glueck BC, Jr. Positive spiking: A double-blind control study on its significance in behavior disorders, both diagnostically and therapeutically. *Am J Psychiatry* 1968 Oct; 125(4):473-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bollen ELEM, Gaw A, Buckley BM, et al. Statin therapy and the prevention of dementia. *Arch Neurol* 2001; 58(6):1023-4.

Status: Not included because not a full article

Boller F, Barba GD. Neuropsychological tests in Alzheimer's disease. *Aging (Milano)* 2001 Jun; 13(3):210-20.

Status: Background article

Bompani R, Scali G. Fipexide, an effective cognition activator in the elderly: A placebo-controlled, double-blind clinical trial. *Curr Med Res Opin* 1986; 10(2):99-106.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bonavita E. Study of the efficacy and tolerability of L-acetylcarnitine therapy in the senile brain. *Int J Clin Pharmacol Ther Toxicol* ; 24(9):511-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bonura ML, Grigoletto F. A 6-month, multicentre, double-blind trial of nicergoline in the treatment of mild to moderate Alzheimer's disease and its 12-month follow-up: Preliminary results. *J Neural Transm Gen Sect* 2000; XVIII.

Status: Not included because not a full article

Boon AJ, Tans JT, Delwel EJ, et al. Does CSF outflow resistance predict the response to shunting in patients with normal pressure hydrocephalus? *Acta Neurochir Suppl* 1998; 71:331-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Borjesson A, Karlsson T, Adolfsson R, et al. Linopirdine (DUP 996): Cholinergic treatment of older adults using successive and non-successive tests. *Neuropsychobiology* 1999; 40(2):78-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Borromei A, Gaggi R, Giancola LC. Involutional dementias: New perspectives. Ital J Neurol Sci 1985; 6(2):167-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Borroni B, Colciaghi F, Pastorino L, et al. Amyloid precursor protein in platelets of patients with Alzheimer's disease: Effect of acetylcholinesterase inhibitor treatment. Arch Neurol 2001; 58(3):442-6.

Status: Not included because dementia population not randomized to treatment

Bottiglieri T, Godfrey P, Flynn T, et al. Cerebrospinal fluid S-adenosylmethionine in depression and dementia: Effects of treatment with parenteral and oral S-adenosylmethionine. J Neurol Neurosurg Psychiatry 1990; 53(12):1096-8.

Status: Not included because dementia population not randomized to treatment

Bottini G, Vallar G, Cappa S, et al. Oxiracetam in dementia: A double-blind, placebo-controlled study. Acta Neurol Scand 1992 Sep; 86(3):237-41.

Status: Included

Bourgeois MS, Burgio LD, Schulz R, et al. Modifying repetitive verbalizations of community-dwelling patients with AD. Gerontologist 1997 Feb; 37(1):30-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Bourgeois MS, Dijkstra K, Burgio L, et al. Memory aids as an augmentative and alternative communication strategy for nursing home residents with dementia. AAC: Augmentative & Alternative Communication 2001 Sep; 17(3):196-210.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Boustani M, Peterson B, Harris R et al. Screening for Dementia. Technical Support for the U.S. Preventive Services Task Force, No. 3. Rockville, MD: U.S. Department of Health and Human Services, Agency for Healthcare Research and Quality. 2002.

Status: Background article

Bower HM, McDonald C. A controlled trial of A.N.P. 235 ("Lucidril") in senile dementia. Med J Aust 1966 Aug 6; 2(6):270-1.

Status: Not included because dementia population not randomized to treatment

Bowles EJ, Griffiths DM, Quirk L, et al. Effects of essential oils and touch on resistance to nursing care procedures and other dementia-related behaviours in a residential care facility. Int J Aromather 2002; 12(1):22-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bowling A, Formby J, Grant K, et al. A randomized controlled trial of nursing home and long-stay geriatric ward care for elderly people. Age Ageing 1991 Sep; 20(5):316-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO. A memory assessment technique for use in geriatric psychopharmacology: Drug efficacy trial with naftidrofuryl. J Am Geriatr Soc 1977 Apr; 25(4):186-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO. The impairment index as a symptom-independent parameter of drug efficacy in geriatric psychopharmacology. J Gerontol 1978 Mar; 33(2):217-23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO, Gardos G. ACTH 4-10 in the amelioration of neuropsychological symptomatology associated with senile organic brain syndrome. Psychopharmacologia 1979 Mar; 61(2):161-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO, Ghazvinian S, et al. Treating the depressed elderly patient: The comparative behavioral pharmacology of mianserin and amitriptyline. Adv Biochem Psychopharmacol 1982; Vol 32:195-212.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO, Dessain EC, et al. The therapeutic efficacy of pramiracetam in Alzheimer's disease: Preliminary observations. Psychopharmacol Bull 1983; 19(4):726-30.

Status: Not included because dementia population not randomized to treatment

Branconnier RJ, Harto NE, Dessain EC, et al. Speech blockage, memory impairment, and age: a prospective comparison of amitriptyline and maprotiline. *Psychopharmacol Bull* 1987; 23(1):230-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Harto NE, Dessain EC, et al. Palliation of the progressive memory impairment of Alzheimer's disease by nimodipine. *Psychopharmacologia* 1988; 96(Suppl):242.

Status: Not included because not a full article

Branconnier RJ, Cole JO. Effects of chronic papaverine administration on mild senile organic brain syndrome. *J Am Geriatr Soc* 1977; 25(10):458-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brandimonte MA, Passolunghi MC. The effect of cue-familiarity, cue-distinctiveness, and retention interval on prospective remembering. *Q J Exp Psychol A* 1994 Aug; 47(3):565-87.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brass EP, Polinsky R, Sramek JJ, et al. Effects of the cholinomimetic SDZ ENS-163 on scopolamine-induced cognitive impairment in humans. *J Clin Psychopharmacol* 1995 Feb; 15(1):58-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brautigam MRH, Blommaert FA, Verleye G, et al. Treatment of age-related memory complaints with ginkgo biloba extract: A randomized double blind placebo-controlled study. *Phytomedicine* 1998; 5(6):425-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brauzer B, Goldstein BJ. The differential response to parenteral chlorpromazine and mesoridazine in psychotic patients. *J Clin Pharmacol New Drugs* 1970 Mar; 10(2):126-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Braverman AM, Naylor CD. Vasoactive substances in the management of elderly patients suffering from dementia. *Modern Geriatrics* 1975; (5):20-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brazzelli M, Capitani E, Della SS, et al. A neuropsychological instrument adding to the description of patients with suspected cortical dementia: The Milan overall dementia assessment. *J Neurol Neurosurg Psychiatry* 1994; 57(12):1510-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Breeze RW, Cox S, Rodgers CJ. Changes in P-300 latency as a result of co-dergocrine mesylate therapy in patients with senile dementia. *Int J Geriatr Psychiatry* 1988; 3(4):263-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Breitbart W, Marotta R, Platt MM, et al. A double-blind trial of haloperidol, chlorpromazine, and lorazepam in the treatment of delirium in hospitalized AIDS patients. *Am J Psychiatry* 1996 Feb; 153(2):231-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Breitner JC, Gau BA, Welsh KA, et al. Inverse association of anti-inflammatory treatments and Alzheimer's disease: Initial results of a co-twin control study. *Neurology* 1994 Feb; 44(2):227-32.

Status: Not included because dementia population not randomized to treatment

Breitner JC. The role of anti-inflammatory drugs in the prevention and treatment of Alzheimer's disease. *Annu Rev Med* 1996; Vol 47:401-11.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Breitner JC, Wyse BW, Anthony JC, et al. APOE-epsilon4 count predicts age when prevalence of AD increases, then declines: The Cache County Study. *Neurology* 1999 Jul 22; 53(2):321-31.

Status: Companion of an included article

Breitner JCS, Zandi PP, In't Veld BA. Long-term use of non-steroidal anti-inflammatory drugs (NSAIDs) reduce the relative risk of Alzheimer's disease to 0.2. *J Neurol* 2002; 249(3):355-6.

Status: Not included because not a full article

Breteler MM, van Duijn CM, Chandra V, et al. Medical history and the risk of Alzheimer's disease: a collaborative re-analysis of case-control studies. *EURODEM Risk Factors*

Research Group. *Int J Epidemiol* 1991; 20 Suppl 2:S36-S42

Status: Background article

Bretz F, Hothorn LA, Hsu JC. Identifying effective and/or safe doses by stepwise confidence intervals for ratios. *Stat Med* 2003 Mar 30; 22(6):847-58.

Status: Background article

Breuil V, de Rotrou J, Forette F, et al. Cognitive stimulation of patients with dementia: Preliminary results. *Int J Geriatr Psychiatry* 1994; 9(3):211-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Bridges-Parlet S, Knopman D, Steffes S. Withdrawal of neuroleptic medications from institutionalized dementia patients: Results of a double-blind, baseline-treatment-controlled pilot study. *J Geriatr Psychiatry Neurol* 1997 Jul; 10(3):119-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brinkman SD, Smith RC, Meyer JS. Lecithin and memory training in suspected Alzheimer's disease. *J Gerontol* 1982; 37(1):4-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brinkman SD, Pomara N, Goodnick PJ, et al. A dose-ranging study of lecithin in the treatment of primary degenerative dementia (Alzheimer's disease). *J Clin Psychopharmacol* 1982 Aug; 2(4):281-5.

*Status: Cross-over trial;
Cross-over trial*

Brody H, Roberts K, Peters K. Quasi-experimental evaluation of an educational model for dementia caregivers. *Int J Geriatr Psychiatry* 1994; 9(3):195-204.

Status: Not included because dementia population not randomized to treatment

Brody H, Gresham M, Luscombe G. The Prince Henry Hospital dementia caregivers' training programme. *Int J Geriatr Psychiatry* 1997 Feb; 12(2):183-92.

Status: Not included because does not meet criteria for treatment for dementia patients

Brody H, Dresser R, Eisner M, et al. Alzheimer's Disease International and International Working Group for Harmonization of

Dementia Drug Guidelines for research involving human subjects with dementia. *Alzheimer Dis Assoc Disord* 1999 Apr; 13(2):71-9.

Status: Background article

Broderick JP, Gaskill M, Dhawan A, et al. Temporal changes in brain volume and cognition in a randomized treatment trial of vascular dementia. *J Neuroimaging* 2001 Jan; 11(1):6-12.
Status: Not included because Jadad Quality Scale score less than three

Brodersen P, Philbert A, Gulliksen G, et al. The effect of L-Deprenyl on on-off phenomena in Parkinson's disease. *Acta Neurol Scand* 1985 Jun; 71(6):494-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brodie NH. A double-blind trial of naftidrofuryl in treating confused elderly patients in general practice. *Practitioner* 1977 Feb; 218(1304):274-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brody EM, Kleban MH, Lawton MP, et al. A longitudinal look at excess disabilities in the mentally impaired aged. *J Gerontol* 1974; 29(1):79-84.

Status: Not included because dementia population not randomized to treatment

Brooker D, Duce L. Wellbeing and activity in dementia: A comparison of group reminiscence therapy, structured goal-directed group activity and unstructured time. *Aging Ment Health* 2000; 4(4):354-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brooker DJ, Snape M, Johnson E, et al. Single case evaluation of the effects of aromatherapy and massage on disturbed behaviour in severe dementia. *Br J Clin Psychol* 1997 May; 36(Pt 2):287-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brookmeyer R, Gray S, Kawas C. Projections of Alzheimer's disease in the United States and the public health impact of delaying disease onset. *Am J Public Health* 1998 Sep; 88(9):1337-42.

Status: Background article

Brookmeyer R, Gray S. Methods for projecting the incidence and prevalence of chronic diseases in

aging populations: application to Alzheimer's disease. *Stat Med* 2000 Jun 15; 19(11-12):1481-93.

Status: Background article

Brooks JO, III, Yesavage JA, Carta A, et al. Acetyl L-carnitine slows decline in younger patients with Alzheimer's disease: A reanalysis of a double-blind, placebo-controlled study using the trilinear approach. *Int Psychogeriatr* 1998 Jun; 10(2):193-203.

Status: Companion of an included article

Brouwers P, Hendricks M, Lietzau JA, et al. Effect of combination therapy with zidovudine and didanosine on neuropsychological functioning in patients with symptomatic HIV disease: A comparison of simultaneous and alternating regimens. *AIDS* 1997 Jan; 11(1):59-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brown S, Gotell E, Ekman SL. Singing as a therapeutic intervention in dementia care. *Journal of Dementia Care* 2001; (4):33-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Brun A, Gustafson L. The lund longitudinal dementia study a 25 year prespective on neuropathology differential diagnosis and treatment. In: Corian B, Iqbal K, Nicolini M, Winblad B, Wisniewski HM, Zatta PF, editors. *Alzheimer's Disease: Advances in Clinical and Basic Research*, Chichester: John Wiley & Sons; 1993. Chapter 1. p. 3-18.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bruno G, Mohr E, Gillespie M, et al. Muscarinic agonist therapy of Alzheimer's disease. A clinical trial of RS-86. *Arch Neurol* 1986 Jul; 43(7):659-61.

*Status: Cross-over trial;
Cross-over trial*

Buettner LL, Lundegren H, Lago D, et al. Therapeutic recreation as an intervention for persons with dementia and agitation: An efficacy study. *Am J Alzheimers Dis* 1996; (5):4-10.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Buettner LL, Fitzsimmons S. AD-venture program: Therapeutic biking for the treatment of depression in long-term care residents with dementia. *Am J*

Alzheimers Dis Other Demen 2002 Mar; 17(2):121-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Buettner LL. Focus on caregiving. Falls prevention in dementia populations: Following a trial program of recreation therapy, falls were reduced by 164 percent. *Provider* 2002 Feb; 28(2):41-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bukatina EE, Grigor'eva IV, Sokol'chik EI. The effectiveness of amiridin in senile dementia of the Alzheimer's type. *Neurosci Behav Physiol* 1993 Jan; 23(1):83-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Burback D, Molnar FJ, St John P, et al. Key methodological features of randomized controlled trials of Alzheimer's disease therapy. Minimal clinically important difference, sample size and trial duration. *Dement Geriatr Cogn Disord* 1999 Nov; 10(6):534-40.

Status: Background article;

Burgener SC, Bakas T, Murray C, et al. Effective caregiving approaches for patients with Alzheimer's disease. *Geriatr Nurs (Minneap)* 1998 May; 19(3):121-6.

Status: Not included because does not meet criteria for treatment for dementia patients

Burgio LD, Reynolds CFI, Janosky JE, et al. A behavioral microanalysis of the effects of haloperidol and oxazepam in demented psychogeriatric inpatients. *Int J Geriatr Psychiatry* 1992; 7(4):253-62.

Status: Not included because Jadad Quality Scale score less than three

Burke WJ, Roccaforte WH, Wengel SP, et al. L-deprenyl in the treatment of mild dementia of the Alzheimer type: Results of a 15-month trial. *J Am Geriatr Soc* 1993a; 41(11):1219-25.

Status: Included

Burke WJ, Ranno AE, Roccaforte WH, et al. L-deprenyl in the treatment of mild dementia of the Alzheimer type: Preliminary results. *J Am Geriatr Soc* 1993b; 41(4):367-70.

Status: Companion of an included article

Burns A, Marsh A, Bender DA. A trial of vitamin supplementation in senile dementia. *Int J Geriatr*

Psychiatry 1989; 4(6):333-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Burns A, Rossor M, Hecker J, et al. The effects of donepezil in Alzheimer's disease - results from a multinational trial. Dement Geriatr Cogn Disord 1999 May; 10(3):237-44.

Status: Included

Cahn LA, Diesfeldt HF. The use of neuroleptics in the treatment of dementia in old age. A critical analysis with reference to an experiment with a long-acting oral neuroleptic (penfluridol Janssen). Psychiatr Neurol Neurochir 1973; 76(6):411-20.
Status: Not included because dementia population not randomized to treatment

Caligiuri MP, Lacro JP, Jeste DV. Incidence and predictors of drug-induced parkinsonism in older psychiatric patients treated with very low doses of neuroleptics. J Clin Psychopharmacol 1999 Aug; 19(4):322-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Camberg L, Woods P, Ooi WL, et al. Evaluation of Simulated Presence: A personalized approach to enhance well-being in persons with Alzheimer's disease. J Am Geriatr Soc 1999 Apr; 47(4):446-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Campi N, Todeschini GP, Scarzella L. Selegiline versus L-acetylcarnitine in the treatment of Alzheimer-type dementia. Clin Ther 1990 Jul; 12(4):306-14.

Status: Not included because Jadad Quality Scale score less than three

Campion D, Brice A, Hannequin D, et al. A large pedigree with early-onset Alzheimer's disease: Clinical, neuropathologic, and genetic characterization. Neurology 1995 Jan; 45(1):80-5.

Status: Not included because does not meet criteria for treatment for dementia patients

Canadian Study of Health and Aging Working Group. Canadian study of health and aging: study methods and prevalence of dementia. CMAJ 1994 Mar 15; 150(6):899-913.

Status: Background article

Canadian Study of Health and Aging Working Group. The Canadian Study of Health and Aging:

risk factors for Alzheimer's disease in Canada. Neurology 1994 Nov; 44(11):2073-80.

Status: Background article

Canadian Task Force on the Periodic Health Examination. The periodic health examination. Can Med Assoc J 1979 Nov 3; 121(9):1193-254.

Status: Background article

Canal N, Imbimbo N. A 25-week double-blind randomized placebo-controlled trial of Eptastigmine in patients with diagnosis of probable Alzheimer's disease. J Neural Transm Gen Sect 1996; 103:XXIV.

Status: Not included because not a full article

Canal N, Imbimbo BP. Relationship between pharmacodynamic activity and cognitive effects of eptastigmine in patients with Alzheimer's disease. Clin Pharmacol Ther 1996; 60(2):218-28.

Status: Included

Cantillon M, Brunswick R, Molina D, et al. Buspirone vs. haloperidol: A double-blind trial for agitation in a nursing home population with Alzheimer's disease. Am J Geriatr Psychiatry 1996; 4(3):263-7.

Status: Not included because Jadad Quality Scale score less than three

Cantor MH. Family and community: changing roles in an aging society. Gerontologist 1991 Jun; 31(3):337-46.

Status: Background article

Capote B, Parikh N. Cyclandelate in the treatment of senility: a controlled study. J Am Geriatr Soc 1978 Aug; 26(8):360-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Capurso A, Capurso S, Panza F, et al. Efficacy of cytidine diphosphate choline in patients affected by chronic cerebrovascular disease. Clin Drug Investig 1996; (1):26-38.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Caraceni TA, Girotti F, Celano I, et al. 2-dimethylaminoethanol (Deanol) in Huntington's chorea. J Neurol Neurosurg Psychiatry 1978; 41(12):1114-8.

Status: Not included because no extractable data relevant to review

Carbonin PU, Greco A, Pisanti P, et al. Efficacy of almitrine-raubasine in cognitive disorders of aging: A double-blind, placebo-controlled, clinical and psychometric study. Clin Neuropharmacol 1990; Vol 13(Suppl 3):S92-S99

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cardebat D, Demonet J-F, Puel M, et al. Brain correlates of memory processes in patients with dementia of Alzheimer's type: A SPECT activation study. J Cereb Blood Flow Metab 1998; 18(4):457-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Carlson MC, Tschanz JT, Norton MC, et al. H2 histamine receptor blockade in the treatment of Alzheimer disease: A randomized, double-blind, placebo-controlled trial of nizatidine. Alzheimer Dis Assoc Disord 2002 Jan; 16(1):24-30.

Status: Included

Carlyle W, Ancill RJ, Sheldon L. Aggression in the demented patient: a double-blind study of loxapine versus haloperidol. Int Clin Psychopharmacol 1993; 8(2):103-8.

Status: Included

Carman JS, Shoulson I, Chase TN. Huntington's chorea treated with lithium carbonate. Lancet 1974; 1(7861):811.

Status: Not included because not a full article

Carmel R. Mild cobalamin deficiency. West J Med 1998 Jun; 168(6):522-3.

Status: Not included because not a full article

Caro AJ, Caro S. Vitamin E in treatment of Huntington's chorea. BMJ 1978 Jan 21; 1(6106):153.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Caro JJ, Salas M, Ward A, et al. Economic analysis of galantamine, a cholinesterase inhibitor, in the treatment of patients with mild to moderate Alzheimer's disease in the Netherlands. Dement Geriatr Cogn Disord 2002; 14(2):84-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Carter LT, Howard BE, O'Neil WA. Effectiveness of cognitive skill remediation in acute stroke patients. Am J Occup Ther 1983 May; 37(5):320-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Carver A, Dobson AM. Effects of dietary supplementation on demented elderly hospital residents. Age Ageing 1993; 22(Suppl 3):37.

Status: Not included because not a full article

Carver AD, Dobson AM. Effects of dietary supplementation of elderly demented hospital residents. J Hum Nutr Diet 1995; 8(6):389-94.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Casale R, Giorgi I, Guarnaschelli C. Evaluation of the effect of vincamine teprosilate on behavioural performances of patients affected with chronic cerebrovascular disease. Int J Clin Pharmacol Res 1984; 4(4):313-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cebul RD. Aspirin and MID Notes of caution. J Am Geriatr Soc 1989; 37(6):573-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cenacchi T, Bertoldin T, Farina C, et al. Cognitive decline in the elderly: A double-blind, placebo-controlled multicenter study on efficacy of phosphatidylserine administration. Aging (Milano) 1993 Apr; 5(Milano):123-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chabriat H, Pappata S, Ostergaard L, et al. Cerebral hemodynamics in CADASIL before and after acetazolamide challenge assessed with MRI bolus tracking. Stroke 2000 Aug; 31(8):1904-12.

Status: Not included because dementia population not randomized to treatment

Challis D, Von Abendorff R, Brown P, et al. Care management, dementia care and specialist mental health services: An evaluation. Int J Geriatr Psychiatry 2002; 17(4):315-25.

Status: Not included because dementia population not randomized to treatment

Chambers CA, Bain J, Rosbottom R. Carbamazepine in senile dementia and overactivity: A placebo controlled double blind trial. IRCS Med Sci 1982; (6):505-6.

Status: Article not retrievable

Chan WC, Lam LC, Choy CN, et al. A double-blind randomised comparison of risperidone and haloperidol in the treatment of behavioural and psychological symptoms in Chinese dementia patients. *Int J Geriatr Psychiatry* 2001 Dec; 16(12):1156-62.

Status: Included

Chandra B. Treatment of multi-infarct dementia with citicholine. *Journal of Stroke and Cerebrovascular Diseases* 1992; 232-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chandra B. Treatment of vascular dementia with CDP choline. *Journal of Stroke and Cerebrovascular Diseases* 2000; 9(Suppl 2):128-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chang BL. Cognitive-behavioral intervention for homebound caregivers of persons with dementia. *Nurs Res* 1999 May; 48(3):173-82.
Status: Not included because does not meet criteria for treatment for dementia patients

Chatellier G, Lacomblez L, and Group. Tacrine (tetrahydroaminoacridine; THA) and lecithin in senile dementia of the Alzheimer type: A multicentre trial. *BMJ* 1990; 300(6723):495-9.
Status: Cross-over trial;
Cross-over trial

Christe C, Janssens JP, Armenian B, et al. Midazolam sedation for upper gastrointestinal endoscopy in older persons: A randomized, double-blind, placebo-controlled study. *J Am Geriatr Soc* 2000 Nov; 48(11):1398-403.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Christensen DB, Benfield WR. Alprazolam as an alternative to low-dose haloperidol in older, cognitively impaired nursing facility patients. *J Am Geriatr Soc* 1998 May; 46(5):620-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Christie JE, Shering A, Ferguson J, et al. Physostigmine and arecoline: Effects of intravenous infusions in Alzheimer presenile dementia. *Br J Psychiatry* 1981; Vol 138:46-50.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chui HC, Mack W, Jackson JE, et al. Clinical criteria for the diagnosis of vascular dementia: A multicenter study of comparability and interrater reliability. *Arch Neurol* 2000 Feb; 57(2):191-6.
Status: Background article

Chung JCC, Lai CKM, Chung PMB, et al. Snoezelen for dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford:Update Software.
Status: Background article

Churchill M, Safaoui J, McCabe BW, et al. Using a therapy dog to alleviate the agitation and desocialization of people with Alzheimer's disease. *J Psychosoc Nurs Ment Health Serv* 1999 Apr; 37(4):16-22.
Status: Not included because dementia population not randomized to treatment

Citrin RS, Dixon DN. Reality Orientation: A Milieu Therapy Used in an Institution for the Aged. *Gerontologist* 1977; 17(1):39-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clair AA, Bernstein B. Effect of no music, stimulative background music and sedative background music on agitated behaviors in persons with severe dementia. *Activities Adaptation Aging* 1994; (1):61-70.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clair AA. Effect of singing on alert responses in persons with late stage dementia. *J Music Ther* 1996; (No. 4):234-47.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clare L, Woods RT, Moniz Cook EL, et al. Cognitive rehabilitation interventions to improve memory functioning in early-stage Alzheimer's disease and vascular dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software
Status: Background article

Clark LR, Fraaza V, Schroeder S, et al. Alternative nursing environments: Do they affect hospital outcomes? *J Gerontol Nurs* 1995 Nov; 21(11):32-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clark ME, Lipe AW, Bilbrey M. Use of music to decrease aggressive behaviors in people with dementia. J Gerontol Nurs 1998 Jul; 24(7):10-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clark WS, Street JS, Feldman PD, et al. The effects of olanzapine in reducing the emergence of psychosis among nursing home patients with Alzheimer's disease. J Clin Psychiatry 2001 Jan; 62(1):34-40.
Status: Companion of an included article

Claus JJ, Mohr E, Chase TN. Clinical trials in dementia: Learning effects with repeated testing. J Psychiatry Neurosci 1991 Mar; 16(1):1-4.
Status: Not included because does not meet criteria for treatment for dementia patients

Claus JJ, Ludwig C, Mohr E, et al. Nootropic drugs in Alzheimer's disease: Symptomatic treatment with pramiracetam. Neurology 1991 Apr; 41(4):570-4.
Status: Cross-over trial;
Cross-over trial

Claus JJ, van Harskamp F, de K, I, et al. Serotonergic mechanisms in Alzheimer's disease: Preliminary results of a controlled clinical trial with lisuride. Can J Neurol Sci 1993; Vol 20:126.
Status : Not included because not a full article

Claus JJ, de K, I, van Harskamp F, et al. Lisuride treatment of Alzheimer's disease. A preliminary placebo-controlled clinical trial of safety and therapeutic efficacy. Clin Neuropharmacol 1998 May; 21(3):190-5.
Status: Included

Clifford DB, McArthur JC, Schifitto G, et al. A randomized clinical trial of CPI-1189 for HIV-associated cognitive-motor impairment. Neurology 2002 Nov 26; 59(10):1568-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clipp EC, Moore MJ. Caregiver time use: an outcome measure in clinical trial research on Alzheimer's disease. Clin Pharmacol Ther 1995 Aug; 58(2):228-36.
Status: Not included because Jadad Quality Scale score less than three

Coccaro EF, Kramer E, Zemishlany Z, et al. Pharmacologic treatment of noncognitive behavioral disturbances in elderly demented

patients. Am J Psychiatry 1990 Dec; 147(12):1640-5.
Status: Included

Coelho F, Birks J. Physostigmine for Alzheimer's disease. In: The Cochrane Library, 2001. Issue 2. Oxford: Update Software
Status: Background article

Cole MG, McCusker J, Bellavance F, et al. Systematic detection and multidisciplinary care of delirium in older medical inpatients: a randomized trial. CMAJ 2002 Oct 1; 167(7):753-9.
Status: Not included because does not meet criteria for treatment for dementia patients

Colling KB, Buettner LL. Simple pleasures. Interventions from the need-driven Dementia-Compromised Behavior model. J Gerontol Nurs 2002 Oct; 28(10):16-20.
Status: Not included because does not meet criteria for treatment for dementia patients

Comelli M, Lucca U, Spagnoli A. Statistical analysis of the clinical trial of a therapy for Alzheimer's disease. Univariate tests and logistic regression. Acta Neurol (Napoli) 1990 Jun; 12(3):222-30.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Commissaris K, Verhey FR, Jolles J. A controlled study into the effects of psychoeducation for patients with cognitive disturbances. J Neuropsychiatry Clin Neurosci 1996; 8(4):429-35.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Conti L, Re F, Lazzerini F, et al. Glycosaminoglycan polysulfate (Ateroid) in old-age dementias: Effects upon depressive symptomatology in geriatric patients. Prog Neuropsychopharmacol Biol Psychiatry 1989; 13(6):977.
Status: Not included because Jadad Quality Scale score less than three

Convit A, de Asis J, de Leon MJ, et al. Atrophy of the medial occipitotemporal, inferior, and middle temporal gyri in non-demented elderly predict decline to Alzheimer's disease. Neurobiol Aging 2000 Jan; 21(1):19-26.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Conway EL. A review of the randomized controlled trials of tacrine in the treatment of Alzheimer's disease: methodologic considerations. Clin Neuropharmacol 1998 Jan; 21(1):8-17.

Status: Background article

Cook WA. Methylperiodol: Clinical trials of a new tranquilizer. Med J Aust 1966 Jul 16; 2(3):117-9.

Status: Not included because dementia population not randomized to treatment

Cools R, Barker RA, Sahakian BJ, et al. Enhanced or impaired cognitive function in Parkinson's disease as a function of dopaminergic medication and task demands. Cereb Cortex 2001 Dec; 11(12):1136-43.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cooney C, Mortimer A, Smith A, et al. Carbamazepine use in aggressive behaviour associated with senile dementia. Int J Geriatr Psychiatry 1996; 11(10):901-5.

*Status: Cross-over trial;
Cross-over trial*

Cooper AJ, Wong YT, Packer H. A controlled trial of cosaldon in arteriosclerotic dementia: Penicillin versus penicillin-malaria in the treatment of dementia paralytica. British Journal Of Psychiatry: British Journal of Venereal Diseases 1964; 25(1):415-8, 439.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cooper AJ, Magnus RV. A placebo-controlled study of pyritinol ('Encephabol') in dementia. Pharmatherapeutica 1980; 2(5):317-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cooper JA, Sagar HJ, Doherty SM, et al. Different effects of dopaminergic and anticholinergic therapies on cognitive and motor function in Parkinson's disease. Brain 1992; 115(6):1701-25.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Corbeil RR, Quayhagen MP, Quayhagen M. Intervention effects on dementia caregiving interaction: A stress-adaptation modeling approach. J Aging Health 1999 Feb; 11(1):79-95.

Status: Not included because does not meet criteria for treatment for dementia patients

Corcoran MA, Gitlin LN. Family caregiver acceptance and use of environmental strategies provided in an occupational therapy intervention. Phys Occup Ther Geriatr 2001; 19(1):1-20.

Status: Not included because does not meet criteria for treatment for dementia patients

Corey-Bloom JR, Anand JV, Veach J, et al. A randomized trial evaluating the efficacy and safety of ENA 713 (rivastigmine tartrate), a new acetylcholinesterase inhibitor, in patients with mild to moderately severe Alzheimer's disease. Int J Geriatr Psychopharmacol 1998; 1:55-65.

Status: Included

Corkin S. Lecithin precursor treatment in Alzheimers-disease and amnesic syndromes. Int J Neurosci 1981; Vol 12:178.

Status: Article not retrievable

Corona GL, Cucchi ML, Frattini P, et al. Clinical and biochemical responses to therapy in Alzheimer's disease and multi-infarct dementia. Eur Arch Psychiatry Neurol Sci 1989; 239(2):79-86.

Status: Not included because Jadad Quality Scale score less than three

Corrigan FM, Van Rhijn A, Horrobin DF. Essential fatty acids in Alzheimer's disease. Ann N Y Acad Sci 1991; Vol 640:250-2.

Status: Not included because no extractable data relevant to review

Cott CA, Dawson P, Sidani S, et al. The effects of a walking/talking program on communication, ambulation, and functional status in residents with Alzheimer's disease. Alzheimer Dis Assoc Disord 2002; 16(2):81-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Coull JT, Sahakian BJ, Hodges JR. The alpha(2) antagonist idazoxan remediates certain attentional and executive dysfunction in patients with dementia of frontal type. Psychopharmacologia 1996 Feb; 123(3):239-49.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Covington JS. Alleviating agitation, apprehension, and related symptoms in geriatric patients: A double-blind comparison of a phenothiazine and a benzodiazepien. South Med Assoc J 1975 Jun; 68(6):719-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cowan P. "Special care" for dementia patients. *Can Nurse* 1999 Jun; 95(6):49-50.
Status: Not included because dementia population not randomized to treatment

Coward RT, Dwyer JW. The association of gender, sibling network composition, and patterns of parent care by adult children. *Research on Aging* 1990 Jun; 12(2):158-81.
Status: Background article

Cowley LM, Glen RS. Double-blind study of thioridazine and haloperidol in geriatric patients with a psychosis associated with organic brain syndrome. *J Clin Psychiatry* 1979 Oct; 40(10):411-9.
Status : Not included because dementia population not defined by DSM, NINCDS or ICD

Cox JR. Double-blind evaluation of naftidrofuryl in treating elderly confused hospitalised patients. *Gerontol Clin (Basel)* 1975; 17(3):160-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cox JR, Shaw AM. Controlled trial of naftidrofuryl in dementia in old age. *J Clin Exp Gerontol* 1981; (4):339-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Coyle J, Kershaw P. Galantamine, a cholinesterase inhibitor that allosterically modulates nicotinic receptors: Effects on the course of Alzheimer's disease. *Biol Psychiatry* 2001; 49(3):289-99.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Coyne AC, Potenza M, Broken-Nose MA. Caregiving and dementia: The impact of telephone helpline services. *Am J Alzheimers Dis* 1995 Jul; (4):27-32.
Status: Not included because does not meet criteria for treatment for dementia patients

Coyne ML, Hoskins L. Improving eating behaviors in dementia using behavioral strategies. *Clin Nurs Res* 1997 Aug; 6(3):275-90.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Craft S, Asthana S, Newcomer JW, et al. Enhancement of memory in Alzheimer disease with insulin and somatostatin, but not glucose. *Arch Gen Psychiatry* 1999 Dec; 56(12):1135-40.
Status: Not included because dementia population not randomized to treatment

Crapper McLachlan DR, Dalton AJ, Kruck TP, et al. Intramuscular desferrioxamine in patients with Alzheimer's disease. *Lancet* 1991 Jun 1; 337(8753):1304-8.
Status: Included

Croisile B, Trillet M, Fondarai J, et al. Long-term and high-dose piracetam treatment of Alzheimer's disease. *Neurology* 1993 Feb; 43(2):301-5.
Status: Included

Cronin-Stubbs D, DeKosky ST, Morris JC, et al. Promoting interactions with basic scientists and clinicians: the NIA Alzheimer's Disease Data Coordinating Center. *Stat Med* 2000 Jun 15; 19(11-12):1453-61.
Status: Background article

Crook, T.H. A 6-month, double-blind, placebo-controlled trial of nicergoline in patients with mild to moderate probable Alzheimer's disease. *J Neural Transm Gen Sect* 2000; 107:XVIII.
Status: Not included because not a full article

Crook T, Ferris S, Sathananthan G, et al. The effect of methylphenidate on test performance in the cognitively impaired aged. *Psychopharmacologia* 1977 May 9; 52(3):251-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Crook T, Wilner E, Rothwell A, et al. Noradrenergic intervention in Alzheimer's disease. *Psychopharmacol Bull* 1992b; 28(1):67-70.
Status: Included

Crook T, Petrie W, Wells C, et al. Effects of phosphatidylserine in Alzheimer's disease. *Psychopharmacol Bull* 1992a; 28(1):61-6.
Status: Included

Crook TH, Tinklenberg J, Yesavage J, et al. Effects of phosphatidylserine in age-associated memory impairment. *Neurology* 1991 May; 41(5):644-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Crook TH. Nicergoline in the treatment of probable Alzheimer's disease: Preliminary results of a double-blind, randomized, placebo-controlled study. JNS 1997; 150(Suppl 1):S18.

Status: Not included because not a full article

Cucinotta D, Passeri M, Ventura S, et al. Multicenter clinical placebo-controlled study with acetyl-L-carnitine (LAC) in the treatment of mildly demented elderly patients. Drug Dev Res 1988; 14(3-4):213-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cucinotta D, Romagnoli S, Godoli G, et al. Comparison of sulfomucopolysaccharides and cytidine diphosphocholine in the treatment of multi-infarct dementia. A randomized double-blind test. Curr Ther Res Clin Exp 1988; 43(1):12-20.

Status: Included

Cucinotta D, Aveni Casucci MA, Pedrazzi F, et al. Multicentre clinical placebo-controlled study with buflomedil in the treatment of mild dementia of vascular origin. J Int Med Res 1992 Apr; 20(2):136-49.

Status: Included

Cucinotta D, De LD, Frattola L, et al. Dihydroergokryptine vs. placebo in dementia of Alzheimer type: Interim results of a randomized multicenter study after a 1-year follow-up. Arch Gerontol Geriatr 1996; 22(2):169-80.

Status: Included

Cucinotta D, De Leo D, Frattola L, et al. Dihydroergokryptine as long-term treatment of Alzheimer type dementia: A multicenter two-year follow-up. Arch Gerontol Geriatr 1998; 27(suppl 6):103-10.

Status: Companion of an included article

Cui Y. Anti-senility potential of Zusanli. Int J Clin Acupunct 1995; 6(1):1-4.

Status: Not included because dementia population not randomized to treatment

Culebras A. Effect of papaverine on cerebral electrogenesis. Neurology 1976 Jul; 26(7):673-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Cummings J, Bieber F, Mas J, et al. Metrifonate in Alzheimer's disease results of a dose finding study. Alzheimers Dis Biol Diagn Ther 1997; 665-

9.

Status: Included

Cummings JL, Gorman DG, Shapira J. Physostigmine ameliorates the delusions of Alzheimer's disease. Biol Psychiatry 1993 Apr 1; 33(7):536-41.

Status: Not included because dementia population not randomized to treatment

Cummings JL, Cyrus PA, Bieber F, et al. Metrifonate treatment of the cognitive deficits of Alzheimer's disease. Neurology 1998a; 51(1):332.

Status: Companion of an included article

Cummings JL, Cyrus PA, Bieber F, et al. Metrifonate treatment of the cognitive deficits of Alzheimer's disease. Neurology 1998b; 50(5):1214-21.

Status: Included

Cummings JL, Knopman D. Advances in the treatment of behavioral disturbances in Alzheimer's disease. Neurology 2000; 53(5):899

Status: Not included because not a full article

Cummings JL, Nadel A, Masterman D, et al. Efficacy of metrifonate in improving the psychiatric and behavioral disturbances of patients with Alzheimer's disease. J Geriatr Psychiatry Neurol 2001; 14(2):101-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cummings JL, Street J, Masterman D, et al. Efficacy of olanzapine in the treatment of psychosis in dementia with lewy bodies. Dement Geriatr Cogn Disord 2002; 13(2):67-73.

Status: Not included because Jadad Quality Scale score less than three

Curless R, James OFW, McKeith I, et al. Effects of propranolol on aggressive behaviour in elderly patients with dementia. Age Ageing 1994; 23:P19.

Status: Not included because not a full article

Curtin F, Elbourne D, Altman DG. Meta-analysis combining parallel and cross-over clinical trials. II: Binary outcomes. Stat Med 2002 Aug 15; 21(15):2145-59.

Status: Background article

Curtin F, Elbourne D, Altman DG. Meta-analysis combining parallel and cross-over clinical trials. III: The issue of carry-over. *Stat Med* 2002 Aug 15; 21(15):2161-73.

Status: Background article

Curtin F, Altman DG, Elbourne D. Meta-analysis combining parallel and cross-over clinical trials. I: Continuous outcomes. *Stat Med* 2002 Aug 15; 21(15):2131-44.

Status: Background article

Curtis-Prior P, Vere D, Fray P. Therapeutic value of Ginkgo biloba in reducing symptoms of decline in mental function. *J Pharm Pharmacol* 1999 May; 51(5):535-41.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cutler NR, Haxby J, Kay AD, et al. Evaluation of zimeldine in Alzheimer's disease. Cognitive and biochemical measures. *Arch Neurol* 1985 Aug; 42(8):744-8.

*Status: Cross-over trial;
Cross-over trial*

Cutler NR, Haxby JV, Narang PK, et al. Evaluation of an analogue of somatostatin (L363,586) in Alzheimer's disease. *N Engl J Med* 1985 Mar; 312(11):725.

Status: Not included because not a full article

Cutler NR, Murphy MF, Nash RJ, et al. Clinical safety, tolerance, and plasma levels of the oral anticholinesterase 1,2,3,4-tetrahydro-9-aminoacridin-1- α -maleate (HP 029) in Alzheimer's disease: Preliminary findings. *J Clin Pharmacol* 1990 Jun; 30(6):556-61.

Status: Not included because no extractable data relevant to review

Cutler NR, Sramek JJ, Murphy MF, et al. Alzheimer's patients should be included in phase I clinical trials to evaluate compounds for Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1992 Oct; 5(4):192-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cutler NR, Sramek JJ, Murphy MF, et al. Implications of the study population in the early evaluation of anticholinesterase inhibitors for Alzheimer's disease. *Ann Pharmacother* 1992 Sep; 26(9):1118-22.

Status: Not included because no extractable data relevant to review

Cutler NR, Shrotriya RC, Sramek JJ, et al. The use of the Computerized Neuropsychological Test Battery (CNTB) in an efficacy and safety trial of BMY 21,502 in Alzheimer's disease. *Ann N Y Acad Sci* 1993; 695(Sep 24):332-6.

Status: Included

Cutler NR, Fakouhi TD, Smith WT, et al. Evaluation of multiple doses of milacemide in the treatment of senile dementia of the Alzheimer's type. *J Geriatr Psychiatry Neurol* 1993 Apr; 6(2):115-9.

Status: Not included because Jadad Quality Scale score less than three

Cutler NR, Sramek JJ, Viereck C, et al. Tolerability and pharmacodynamics of besipirdine in Alzheimer's disease. *Biol Psychiatry* 1995; 37(9):643.

Status: Not included because not a full article

Cutler NR, Sramek JJ, Anand R. Safety and tolerance of ENA 713 in patients with Alzheimer's disease. *Biol Psychiatry* 1995; 37(9):643.

Status: Not included because not a full article

Cutler NR, Jhee SS, Cyrus P, et al. Safety and tolerability of metrifonate in patients with Alzheimer's disease: Results of a maximum tolerated dose study. *Life Sci* 1998; 62(16):1433-41.

Status: Not included because dementia population not randomized to treatment

Cutler NR, Polinsky RJ, Sramek JJ, et al. Dose-dependent CSF acetylcholinesterase inhibition by SDZ ENA 713 in Alzheimer's disease. *Acta Neurol Scand* 1998 Apr; 97(4):244-50.

Status: Not included because dementia population not randomized to treatment

Czerwinski AW, Clark ML, Serafetinides EA, et al. Safety and efficacy of zinc sulfate in geriatric patients. *Clin Pharmacol Ther* 1974 Apr; 15(4):436-41.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Danielczyk W, Simanyi B, Forette F, et al. CBM 36-733 (2-methyl- α -ergokryptine) in primary degenerative dementia: Results of a European multicentre trial. *Int J Geriatr Psychiatry* 1988; 3(2):107-14.

Status: Included

Daniele A, Moro E, Bentivoglio AR. Zolpidem in progressive supranuclear palsy. *N Engl J Med* 1999 Aug 12; 341(7):543-4.

Status: Not included because not a full article

Daniels L. A group cognitive-behavioural and process-oriented approach to treating the social impairment and negative symptoms associated with chronic mental illness. *J Psychother Pract Res* 1998; 7:167-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dansky KH, Dellasega C, Shellenbarger T, et al. After hospitalization: Home health care for elderly persons. *Clin Nurs Res* 1996 May; 5(2):185-98.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Danyasz W. CX-516 Cortex Pharmaceuticals Inc. *Idrugs* 1999; 2(8):814-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Darreh-Shori T, Almkvist O, Guan ZZ, et al. Sustained cholinesterase inhibition in AD patients receiving rivastigmine for 12 months. *Neurology* 2002 Aug 27; 59(4):563-72.

Status: Not included because dementia population not randomized to treatment

Davidson M, Mohs RC, Hollander E, et al. Lecithin and piracetam in Alzheimer's disease. *Biol Psychiatry* 1987; 22(1):112-4.

Status: Not included because not a full article

Davidson M, Zemishlany Z, Mohs RC, et al. 4-Aminopyridine in the treatment of Alzheimer's disease. *Biol Psychiatry* 1988 Mar 1; 23(5):485-90.

*Status: Cross-over trial;
Cross-over trial*

Davies AE. A pilot study to measure aluminium levels in hair samples of patients with dementia and the influence of aluminium 30c compared with placebo. *Communications of the British Homoeopathy Research Group Issue* 18, 1988; 42-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davies B, Andrewes D, Stargatt R, et al. Tacrine in Alzheimer's disease. *Lancet* 1989 Jul 15; 2(8655):163-4.

Status: Not included because not a full article

Davies B, Andrewes D, Stargatt R, et al. Tetrahydroaminoacridine in Alzheimer's disease. *Int J Geriatr Psychiatry* 1990; 5(5):317-21.

*Status: Cross-over trial;
Cross-over trial*

Davies G, Hamilton S, Hendrickson E, et al. The effect of cyclandelate in depressed and demented patients: A controlled study in psychogeriatric patients. *Age Ageing* 1977 Aug; 6(3):156-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davies G. Drugs for dementia. *Modern Geriatrics* 1978; 8(1):56

Status: Not included because not a full article

Davis KJ, Sloane PD, Mitchell CM, et al. Specialized dementia programs in residential care settings. *Gerontologist* 2000 Feb; 40(1):32-42.

Status: Not included because does not meet criteria for treatment for dementia patients

Davis KL, Mohs RC. Enhancement of memory processes in Alzheimer's disease with multiple-dose intravenous physostigmine. *Am J Psychiatry* 1982; 139(11):1421-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davis KL, Mohs RC, Davis BM, et al. Oral physostigmine in Alzheimer's disease. *Psychopharmacol Bull* 1983; 19(3):451-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davis KL, Thal LJ, Gamzu ER, et al. A double-blind, placebo-controlled multicenter study of tacrine for Alzheimer's disease. *N Engl J Med* 1992; 327(18):1253-9.

Status: Not included because Jadad Quality Scale score less than three

Davis KL, Yang RK, Davidson M, et al. Alzheimer's disease: Tacrine and tacrine metabolite concentrations in plasma and cognitive change. *Drug Dev Res* 1995; 34(1):55-65.

Status: Not included because Jadad Quality Scale score less than three

Davis RN, Massman PJ, Doody RS. Cognitive intervention in Alzheimer disease: A randomized placebo-controlled study. *Alzheimer Dis Assoc Disord* 2001 Jan; 15(1):1-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Dawson P, Kontos P. Nursing assistants reduce aggressive behaviour during bathing cognitively impaired nursing home residents. *Perspectives (Montclair)* 1998; 22(2):20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dawson P. Bright light treatment for people with Alzheimer's disease. *Perspectives (Montclair)* 1999; 23(1):25-6.

Status: Not included because dementia population not randomized to treatment

Day JJ, Grant I, Atkinson JH, et al. Incidence of AIDS dementia in a two-year follow-up of AIDS and ARC patients on an initial phase II AZT placebo-controlled study: San Diego cohort. *J Neuropsychiatry Clin Neurosci* 1992; 4(1):15-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

de Boer JB, van Dam FS, Sprangers MA, et al. Longitudinal study on the Quality of Life of symptomatic HIV-infected patients in a trial of zidovudine versus zidovudine and interferon-alpha. *AIDS* 1993 Jul; 7(7):947-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

De Deyn PP, Rabheru K. A randomized trial of risperidone, placebo, and haloperidol for behavioral symptoms of dementia. *Neurology* 1999 Sep 22; 53(5):946-55.

Status: Included

De Deyn PP, Scheltens P, Kittner B. A double-blind placebo-controlled trial assessing the effects of propentofylline in patient's with Alzheimer's disease and vascular dementia: safety, efficacy, and impact on disease progression. *Alzheimers Rep* 1999; (2):51.

Status: Not included because not a full article

De Deyn PP, Rabheru K, Rasmussen A. Neuroleptic for behavioral symptoms of dementia. *J Fam Pract* 2000; 49(1):28-9.

Status: Not included because not a full article

Dehlin O, Hedenrud B, Jansson P, et al. A double-blind comparison of alaproclate and placebo in the treatment of patients with senile dementia. *Acta Psychiatr Scand* 1985 Feb; 71(2):190-6.

Status: Included

Deimling GT, Bass DM. Symptoms of mental impairment among elderly adults and their effects on family caregivers. *J Gerontol* 1986 Nov; 41(6):778-84.

Status: Background article

Del Ser T, McKeith I, Anand R, et al. Dementia with lewy bodies: Findings from an international multicentre study. *Int J Geriatr Psychiatry* 2000 Nov; 15(11):1034-45.

Status: Companion of an included article

DeLuca J, Johnson SK, Ellis SP, et al. Cognitive functioning is impaired in patients with chronic fatigue syndrome devoid of psychiatric disease. *J Neurol Neurosurg Psychiatry* 1997; 62(2):151-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Delwaide PJ, Devoitille JM, Ylief M. Acute effect of drugs upon memory of patients with senile dementia. *Acta Psychiatr Belg* 1980; 748-54.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Delwaide PJ, Hurler A. Bromocriptine and buccolingual dyskinesias in patients with senile dementia. A quantitative study. *Arch Neurol* 1980 Jul; 37(7):441-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Delwaide PJ, Gyselynck-Mambourg AM, Hurler A, et al. Double-blind randomized controlled study of phosphatidylserine in senile demented patients. *Acta Neurol Scand* 1986 Feb; 73(2):136-40.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Demers L, Oremus M, Perrault A, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Psychometric properties of functional and quality of life scales. *J Geriatr Psychiatry Neurol* 2000; 13(4):170-80.

Status: Background article

Demers L, Oremus M, Perrault A, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Introduction. *J Geriatr Psychiatry Neurol* 2000; 13(4):161-9.

Status: Background article

Dencker SJ, Lindberg D. A controlled double blind study of piracetam in the treatment of senile dementia. *Nord J Psychiatry* 1977; (1):48-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Denney A. Quiet music. An intervention for mealtime agitation? J Gerontol Nurs 1997 Jul; 23(7):16-23.

Status: Not included because dementia population not randomized to treatment

Denolle T, Sassano P, Allain H, et al. Effects of nicardipine and clonidine on cognitive functions and electroencephalography in hypertensive patients. Fundam Clin Pharmacol 2002; 16(6):527-35.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Derouesne C, Renault B, Gueguen B, et al. Neuropsychophysiological evaluation of three doses of S 12024-2 in mild-to-moderate Alzheimer's disease. Clin Drug Investig 1997; (4):301-6.

*Status: Cross-over trial;
Cross-over trial*

Desai A, Grossberg G. Review of rivastigmine and its clinical applications in Alzheimer's disease and related disorders. Expert Opin Pharmacother 2001; 2(4):653-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

desRosiers G, Hodges JR, Berrios G. The neuropsychological differentiation of patients with very mild Alzheimer's disease and/or major depression. J Am Geriatr Soc 1995 Nov; 43(11):1256-63.

Status: Not included because does not meet criteria for treatment for dementia patients

Devanand DP, Sackeim HA, Brown RP, et al. A pilot study of haloperidol treatment of psychosis and behavioral disturbance in Alzheimer's disease. Arch Neurol 1989; 46(8):854-7.

Status: Not included because dementia population not randomized to treatment

Devanand DP, Cooper T, Sackeim HA, et al. Low dose oral haloperidol and blood levels in Alzheimer's disease: A preliminary study. Psychopharmacol Bull 1992; 28(2):169-73.

Status: Not included because Jadad Quality Scale score less than three

Devanand DP, Marder K, Michaels KS, et al. A randomized, placebo-controlled dose-comparison

trial of haloperidol for psychosis and disruptive behaviors in Alzheimer's disease. Am J Psychiatry 1998 Nov; 155(11):1512-20.

*Status: Cross-over trial;
Cross-over trial*

di Bari M, Pahor M, Franse LV, et al. Dementia and disability outcomes in large hypertension trials: Lessons learned from the systolic hypertension in the elderly program (SHEP) trial. Am J Epidemiol 2001; 153(1):72-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Di Perri R, Coppola G, Ambrosio LA, et al. A multicentre trial to evaluate the efficacy and tolerability of alpha-glycerylphosphorylcholine versus cytosine diphosphocholine in patients with vascular dementia. J Int Med Res 1991 Jul; 19(4):330-41.

Status: Not included because Jadad Quality Scale score less than three

Dick MB, Nielson KA, Beth RE, et al. Acquisition and long-term retention of a fine motor skill in Alzheimer's disease. Brain Cogn 1995 Dec; 29(3):294-306.

Status: Not included because does not meet criteria for treatment for dementia patients

Dierks T, Maurer K, Ihl R. Influence of tenilsetam on AEP-P300 in Alzheimer's disease. J Neural Transm Gen Sect 1989; 1:49.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ditzler K. Efficacy and tolerability of memantine in patients with dementia syndrome. A double-blind, placebo controlled trial. Arzneimittelforschung 1991 Aug; 41(8):773-418.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Diwan S, Phillips VL. Agitation and dementia-related problem behaviors and case management in long-term care. Int Psychogeriatr 2001; 13(1):5-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Djernes JK, Gulmann NC, Abelskov KE, et al. Psychopathologic and functional outcome in the treatment of elderly inpatients with depressive disorders, dementia, delirium and psychoses. Int Psychogeriatr 1998 Mar; 10(1):71-83.

Status: Not included because does not meet criteria for treatment for dementia patients

Doble SE, Fisk JD, Rockwood K. Assessing the ADL functioning of persons with Alzheimer's disease: Comparison of family informants' ratings and performance-based assessment findings. *Int Psychogeriatr* 1999 Dec; 11(4):399-409.
Status: Background article

Dominguez D, De CCL, Gomensoro J, et al. Modification of psychometric, practical and intellectual parameters in patients with diffuse cerebrovascular insufficiency during prolonged treatment with pentoxifylline: A double blind, placebo controlled trial. *Pharmatherapeutica* 1977; 1(8):498-506.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Doody RS, Geldmacher DS, Gordon B, et al. Open-label, multicenter, phase 3 extension study of the safety and efficacy of donepezil in patients with Alzheimer's disease. *Arch Neurol* 2001 Mar; 58(3):427-33.
Status: Companion of an included article

Doody RS, Dunn JK, Clark CM, et al. Chronic donepezil treatment is associated with slowed cognitive decline in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2001 Jul; 12(4):295-300.
Status: Not included because dementia population not randomized to treatment

Doody RS, Stevens JC, Beck C, et al. Practice parameter: Management of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001 May 8; 56(9):1154-66.
Status: Background article

Doraiswamy PM, Kaiser L, Bieber F, et al. The Alzheimer's Disease Assessment Scale: Evaluation of psychometric properties and patterns of cognitive decline in multicenter clinical trials of mild to moderate Alzheimer's disease. *Alzheimer Dis Assoc Disord* 2001 Oct; 15(4):174-83.
Status: Not included because does not meet criteria for treatment for dementia patients

Doraiswamy PM, Krishnan KRR, Anand R, et al. Long-term effects of rivastigmine in moderately severe Alzheimer's disease: Does early initiation of therapy offer sustained benefits? *Prog*

Neuropsychopharmacol Biol Psychiatry 2002; 26(4):705-12.
Status: Companion of an included article

Doraiswamy PM, Krishen A, Stallone F, et al. NSAIDs and cognition in Alzheimer's disease. *Neurology* 1996; 46(4):1194.
Status: Not included because not a full article

Droes RM, Breebaart E, Ettema TP, et al. Effect of integrated family support versus day care only on behavior and mood of patients with dementia. *Int Psychogeriatr* 2000; 12(1):99-115.
Status: Not included because does not meet criteria for treatment for dementia patients

Dubois B, McKeith I, Orgogozo JM, et al. A multicentre, randomized, double-blind, placebo-controlled study to evaluate the efficacy, tolerability and safety of two doses of metrifonate in patients with mild-to-moderate Alzheimer's disease: The MALT study. *Int J Geriatr Psychiatry* 1999 Nov; 14(11):973-82.
Status: Included

Duffy FH, McAnulty G, Albert M, et al. Lecithin: Absence of neurophysiologic effect in Alzheimer's disease by EEG topography. *Neurology* 1987 Jun; 37(6):1015-9.
Status: Not included because no extractable data relevant to review

Dunn JC, Thiru-Chelvam B, Beck CH. Bathing. Pleasure or pain? *J Gerontol Nurs* 2002 Nov; 28(11):6-13.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Duret M, Goldman S, Messina D, et al. Effect of L-dopa on dementia-related rigidity. *Acta Neurol Scand* 1989 Jul; 80(1):64-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Durso R, Fedio P, Brouwers P. Lysine vasopressin in Alzheimer's disease. *Neurology* 1982; 32(6):674-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dykieriek P, Stadtmuller G, Schramm P, et al. The value of REM sleep parameters in differentiating Alzheimer's disease from old-age depression and normal aging. *J Psychiatr Res* 1998 Jan; 32(1):1-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Dysken M, Kuskowski M, Love S, et al. Ondansetron in the treatment of cognitive decline in Alzheimer's dementia. *Am J Geriatr Psychiatry* 2002 Mar; 10(2):212-5.

Status: Not included because Jadad Quality Scale score less than three

Dysken MW, Anton JS, Klein L, et al. CI-911: A placebo-controlled study in patients with primary degenerative dementia. *Drug Dev Res* 1988; 12(3-4):267-4.

*Status: Cross-over trial;
Cross-over trial*

Dysken MW, Mendels J, Lewitt P, et al. Milacemide: a placebo-controlled study in senile dementia of the Alzheimer type. *J Am Geriatr Soc* 1992 May; 40(5):503-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dysken MW, Johnson SB, Holden L, et al. Haloperidol concentrations in patients with Alzheimer's dementia. *Am J Geriatr Psychiatry* 1994; 2(2):124-33.

Status: Not included because Jadad Quality Scale score less than three

Eagger S, Levi I. Serum levels of tacrine in relation to clinical response in Alzheimer's disease. *Int J Geriatr Psychiatry* 1992a; 7(2):115-9.

*Status: Cross-over trial;
Cross-over trial*

Eagger S, Morant N, Levy R, et al. Tacrine in Alzheimer's disease. Time course of changes in cognitive function and practice effects. *Br J Psychiatry* 1992b; 160:36-40.

*Status: Cross-over trial;
Cross-over trial*

Eagger S. Searching for a treatment for Alzheimer's disease - tales from the cutting-room floor. *Int J Geriatr Psychiatry* 1996; 11(4):337-42.

Status: Not included because dementia population not randomized to treatment

Eagger SA, Levy R, Sahakian BJ. Tacrine in Alzheimer's disease. *Lancet* 1991 Apr 27; 337(8748):989-92.

*Status: Cross-over trial;
Cross-over trial*

Eagger SA, Morant NJ, Levy R. Parallel group analysis of the effects of tacrine versus placebo in Alzheimer's disease. *Dementia* 1991; 2(4):207-11.

Status: Article not retrievable

Eagger SA, Levy R, Sahakian BJ. Tacrine in Alzheimer's disease. *Acta Neurol Scand Suppl* 1992; 139:75-80.

*Status: Cross-over trial;
Cross-over trial*

Eagger SA, Richards M, Levy R. Long-term effects of tacrine in Alzheimer's disease: An open study. *Int J Geriatr Psychiatry* 1994; 9(8):643-7.

Status: Companion of an included article

Eaton M, Mitchell-Bonair IL, Friedmann E. The effect of touch on nutritional intake of chronic organic brain syndrome patients. *J Gerontol* 1986 Sep; 41(5):611-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ebmeier KP, Hunter R, Curran SM, et al. Effects of a single dose of the acetylcholinesterase inhibitor velnacrine on recognition memory and regional cerebral blood flow in Alzheimer's disease. *Psychopharmacologia* 1992; 108(1-2):103-2.

*Status: Cross-over trial;
Cross-over trial*

Eccles M, Clarke J, Livingstone M, et al. North of England evidence based guidelines development project: Guideline for the primary care management of dementia. *BMJ* 1998 Sep 19; 317(7161):802-8.

Status: Background article

Edberg A, Hallberg IR. Actions seen as demanding in patients with severe dementia during one year of intervention. Comparison with controls. *Int J Nurs Stud* 2001 Jun; 38(3):271-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Edberg AK, Hallberg IR. Effects of clinical supervision on nurse-patient cooperation quality: A controlled study in dementia care. *Clin Nurs Res* 1996 May; 5(2):127-46.

Status: Not included because does not meet criteria for treatment for dementia patients

Edberg AK, Norberg A, Hallberg I. Mood and general behavior of patients with severe dementia

during one year of supervised, individualized planned care and systematic clinical supervision: Comparison with a similar control group. *Aging Clin Exp Res* 1999; (6):395-403.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Edland SD. Blomqvist revisited: how and when to test the relationship between level and longitudinal rate of change. *Stat Med* 2000 Jun 15; 19(11-12):1441-52.
Status: Background article

Edwards NE, Beck AM. Animal-assisted therapy and nutrition in Alzheimer's disease. *West J Nurs Res* 2002 Oct; 24(6):697-712.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Egger M, Juni P, Bartlett C et al. How important are comprehensive literature searches and the assessment of trial quality in systematic reviews. *Health Technology Assessment* No. 7. 2003. 1.
Status: Background article

Ellingrod VL, Schultz SK, Ekstam-Smith K, et al. Comparison of risperidone with olanzapine in elderly patients with dementia and psychosis. *Pharmacotherapy* 2002 Jan; 22(1):1-5.
Status: Not included because dementia population not randomized to treatment

Eloniemi-Sulkava U, Notkola IL, Hentinen M, et al. Effects of supporting community-living demented patients and their caregivers: A randomized trial. *J Am Geriatr Soc* 2001 Oct; 49(10):1282-7.
Status: Not included because does not meet criteria for treatment for dementia patients

Emeriau JP, Leher P, Mosnier M. Efficacy of naftidrofuryl in patients with vascular or mixed dementia: Results of a multicenter, double-blind trial. *Clin Ther* 2000 Jul; 22(7):834-44.
Status: Article not retrievable

Emsley CL, Gao S, Hall KS, et al. Estimating odds ratios adjusting for misclassification in Alzheimer's disease risk factor assessment. *Stat Med* 2000 Jun 15; 19(11-12):1523-30.
Status: Background article

Engel RR, Satzger W, Gunther W, et al. Double-blind cross-over study of phosphatidylserine vs. placebo in patients with early dementia of the Alzheimer type. *Eur Neuropsychopharmacol* 1992 Jun; 2(2):149-55.

*Status: Cross-over trial;
 Cross-over trial*

Engelberts NH, Klein M, Ader HJ, et al. The effectiveness of cognitive rehabilitation for attention deficits in focal seizures: A randomized controlled study. *Epilepsia* 2002; 43(6):587-95.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Erkinjuntti T, Ostbye T, Steenhuis R, et al. The effect of different diagnostic criteria on the prevalence of dementia. *N Engl J Med* 1997 Dec 4; 337(23):1667-74.
Status: Background article

Erkinjuntti T, Kurz A, Gauthier S, et al. Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: A randomised trial. *Lancet* 2002 Apr 13; 359(9314):1283-90.
Status: Included

Etienne P, Dastoor D, Gauthier S. Alzheimer's disease: Lack of effect of lecithin treatment for 3 months. *Neurology* 1981; 31(12):1552-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

European Medicine Evaluation Agency (EMA). Note for guidance on medicinal products in the treatment of Alzheimer's disease. London: EMA; 1997.
Status: Background article

European Propentofylline Study Group. Propentofylline in dementia (vascular dementia and Alzheimer's disease). *Cardiovasc Dis* 1994; 4:258.
Status: Not included because not a full article

Evans M, Hammond M, Wilson K, et al. Treatment of depression in the elderly: Effect of physical illness on response. *Int J Geriatr Psychiatry* 1997 Dec; 12(12):1189-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Evans M, Ellis A, Watson D, et al. Sustained cognitive improvement following treatment of Alzheimer's disease with donepezil. *Int J Geriatr Psychiatry* 2000 Jan; 15(1):50-3.
Status: Not included because dementia population not randomized to treatment

Evers S, Grotemeyer KH, Reichelt D, et al. Impact of antiretroviral treatment on AIDS dementia: A longitudinal prospective event-related potential study. *J Acquir Immune Defic Syndr Hum Retrovirol* 1998 Feb 1; 17(2):143-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Exton Smith AN, Piper ME, Phillips MJ, et al. Management of elderly patients with dementia: A clinical trial using high doses of Hydergine. *Br J Clin Pract* 1982; Vol 16:55-8.

Status: Article not retrievable

Fakouhi TD, Jhee SS, Sramek JJ, et al. Evaluation of cycloserine in the treatment of Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1995 Oct; 8(4):226-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Falsaperla A, Monici Preti PA, Oliani C. Selegiline versus oxiracetam in patients with Alzheimer-type dementia. *Clin Ther* 1990 Sep; 12(5):376-84.

Status: Not included because Jadad Quality Scale score less than three

Farina E, Fioravanti R, Chiavari L, et al. Comparing two programs of cognitive training in Alzheimer's disease: A pilot study. *Acta Neurol Scand* 2002 May; 105(5):365-71.

Status: Not included because does not meet criteria for treatment for dementia patients

Farlow M, Gracon SI, Hershey LA, et al. A controlled trial of tacrine in Alzheimer's disease. The Tacrine Study Group. *JAMA* 1992 Nov 11; 268(18):2523-9.

Status: Not included because Jadad Quality Scale score less than three

Farlow M, Brashear A, Hiu S, et al. The effects of tacrine in patients with mild versus moderate stage Alzheimer's disease. In: Iqbal K, editors. *Research Advances in Alzheimer's disease and related disorders*, Chichester: John Wiley & Sons; 1995. p. 283-92.

Status: Not included because not a full article

Farlow M, Anand R, Messina J, Jr., et al. A 52-week study of the efficacy of rivastigmine in patients with mild to moderately severe Alzheimer's disease. *Eur Neurol* 2000; 44(4):236-41.

Status: Companion of an included article

Farlow MR, Lahiri DK, Poirier J, et al. Apolipoprotein E genotype and gender influence response to tacrine therapy. *Ann N Y Acad Sci* 1996; 802(Dec 16):101-10.

Status: Not included because Jadad Quality Scale score less than three

Farlow MR, Lahiri DK, Poirier J, et al. Treatment outcome of tacrine therapy depends on apolipoprotein genotype and gender of the subjects with Alzheimer's disease. *Neurology* 1998 Mar; 50(3):669-77.

Status: Companion of an included article

Farlow MR, Cyrus PA, Nadel A, et al. Metrifonate treatment of AD: Influence of APOE genotype. *Neurology* 1999 Dec 10; 53(9):2010-6.

Status: Not included because dementia population not randomized to treatment

Farlow MR, Cyrus PA. Metrifonate therapy in Alzheimer's disease: A pooled analysis of four randomized, double-blind, placebo-controlled trials. *Dement Geriatr Cogn Disord* 2000 Jul; 11(4):202-11.

Status: Not included because dementia population not randomized to treatment

Farlow MR, Hake A, Messina J, et al. Response of patients with Alzheimer's disease to rivastigmine treatment is predicted by the rate of disease progression. *Arch Neurol* 2001 Mar; 58(3):417-22.

Status: Companion of an included article

Faxen-Irving G, Andren-Olsson B, af GA, et al. The effect of nutritional intervention in elderly subjects residing in group-living for the demented. *Eur J Clin Nutr* 2002 Mar; 56(3):221-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Feldman H, Gauthier S, Hecker J, et al. Benefits of Donepezil on global function, behavior, cognition and ADLs in patients with moderate to severe Alzheimer's disease. *Neurology* 2000; 54(Suppl 3):A469.

Status: Not included because not a full article

Feldman H, Gauthier S, Hecker J, et al. A 24-week, randomized, double-blind study of donepezil in moderate to severe Alzheimer's disease. *Neurology* 2001 Aug 28; 57(4):613-20.

Status: Included

Feldman H, Sauter A, Donald A, et al. The disability assessment for dementia scale: A 12-month study of functional ability in mild to moderate severity Alzheimer's disease. *Alzheimer Dis Assoc Disord* 2001 Apr; 15(2):89-95.

Status: Not included because does not meet criteria for treatment for dementia patients

Fenn P, Gray A. Estimating long-term cost savings from treatment of Alzheimer's disease. A modelling approach. *Pharmacoeconomics* 1999 Aug; 16(2):165-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fergusson D, Aaron SD, Guyatt G, et al. Post-randomisation exclusions: the intention to treat principle and excluding patients from analysis. *BMJ* 2002 Sep 21; 325(7365):652-4.

Status: Background article

Fernandez HH, Trieschmann ME, Burke MA, et al. Quetiapine for psychosis in Parkinson's disease versus dementia with Lewy bodies. *J Clin Psychiatry* 2002 Jun; 63(6):513-5.

Status: Not included because dementia population not randomized to treatment

Ferrari E, Cucinotta D, Albizzati MG, et al. Effectiveness and safety of posatirelin in the treatment of senile dementia: A multicenter, double-blind, placebo-controlled study. *Arch Gerontol Geriatr* 1998; 27(Suppl 6):163-74.

Status: Included

Ferris SH, Sathananthan G, Gershon S, et al. Cognitive effects of ACTH 4-10 in the elderly. *Pharmacol Biochem Behav* 1976; 5(Suppl 1):73-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ferris SH, Mittelman MS. Behavioral treatment of Alzheimer's disease. *Int Psychogeriatr* 1996; 8(Suppl 1):87-90.

Status: Not included because no extractable data relevant to review

Fichter MM, Bruce ML, Schroppel H, et al. Cognitive impairment and depression in the oldest old in a German and in U.S. communities. *Eur Arch Psychiatry Clin Neurosci* 1995; 245(6):319-25.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Filip V, David I, Klatschka J, et al. Selegiline slows down the decline of cognitive and electrophysiological functions in Alzheimer's disease. *Basic and clinical science of mental and additive disorders. Bibl Psychiatr* 1997; 238-40.

Status: Not included because not a full article

Filip V, Kolibas E. Selegiline in the treatment of Alzheimer's disease: A long-term randomized placebo-controlled trial. *Czech and Slovak Senile Dementia of Alzheimer Type Study Group. J Psychiatry Neurosci* 1999 May; 24(3):234-43.

Status: Included

Finali G, Piccirilli M, Oliani C, et al. L-deprenyl therapy improves verbal memory in amnesic Alzheimer patients. *Clin Neuropharmacol* 1991 Dec; 14(6):523-36.

Status: Cross-over trial; Cross-over trial

Finali G, Piccirilli M, Oliani C, et al. Alzheimer-type dementia and verbal memory performances: Influence of selegiline therapy. *Ital J Neurol Sci* 1992 Mar; 13(2):141-8.

Status: Cross-over trial; Cross-over trial

Findlay DJ, Sharma J, McEwen J, et al. Double-blind controlled withdrawal of thioridazine treatment in elderly female inpatients with senile dementia. *Int J Geriatr Psychiatry* 1989; 4(2):115-20.

Status: Not included because Jadad Quality Scale score less than three

Finkel SI, Lyons JS, Anderson RL, et al. A randomized, placebo-controlled trial of thiothixene in agitated, demented nursing home patients. *Int J Geriatr Psychiatry* 1995; 10(2):129-36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Finkel SI, Lyons J. Nursing home research from investigators' perspective. *Int Psychogeriatr* 1996; 8(Suppl 3):371-3, 381-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Finnema E, Droes RM, Ribbe M, et al. The effects of emotion-oriented approaches in the care for persons suffering from dementia: a review of the literature. *Int J Geriatr Psychiatry* 2000 Feb; 15(2):141-61.

Status: Background article

Fioravanti M, Bergamasco B, Bocola V, et al. A multicentre, double-blind, controlled study of piracetam vs placebo in geriatric patients with nonvascular mild-moderate impairment in cognition. *New Trends in Clinical Neuropharmacology* 1991; (1):27-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fioravanti M, Di Cesare F. Memory improvements and pharmacological treatment: A method to distinguish direct effects on memory from secondary effects due to attention improvement. *Int Psychogeriatr* 1992; 4(1):119-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fioravanti M, Ferrario E, Massaia M, et al. Low folate levels in the cognitive decline of elderly patients and the efficacy of folate as a treatment for improving memory deficits. *Arch Gerontol Geriatr* 1997; 26(1):1-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fioravanti M, Yanagi M. Cytidinediphosphocholine (CDP choline) for cognitive and behavioural disturbances associated with chronic cerebral disorders in the elderly. In: *The Cochrane Library*, 2000. Issue 4. Oxford: Update Software.

Status: Background article

Fioravanti M, Flicker L. Efficacy of nicergoline in dementia and other age associated forms of cognitive impairment. In: *The Cochrane Library*, 2001. Issue 4. Oxford: Update Software.

Status: Background article

Fioravanti M, Birks J. Idebenone for Alzheimer's disease (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software.

Status: Background article

Fischer, Gotz P. Blood transferrin and ferritin in Alzheimer's disease. *Life Sci* 1997; 60(25):2273-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Fischhof PK, Saletu B, Ruther E, et al. Therapeutic efficacy of pyritinol in patients with senile dementia of the Alzheimer type (SDAT) and multi-infarct dementia (MID). *Neuropsychobiology* 1992; 26(1-2):65-2.

Status: Not included because Jadad Quality Scale score less than three

Fischhof PK. Divergent neuroprotective effects of nimodipine in PDD and MID provide indirect evidence of disturbances in Ca²⁺ homeostasis in dementia. *Methods & Findings in Experimental & Clinical Pharmacology* 1993 Oct; 15(8):549-55.

Status: Not included because Jadad Quality Scale score less than three

Fischhof PK, Moslinger-Gehmayr R, Herrmann WM, et al. Therapeutic efficacy of vincamine in dementia. *Neuropsychobiology* 1996; 34(1):29-35.

Status: Included

Fisman M, Merksey H, Helmes E. Double blind study of lecithin in patients with Alzheimer's disease. *Can J Psychiatry* 1981; (6):426-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fisman M, Merskey H, Helmes E. Double-blind trial of 2-dimethylaminoethanol in Alzheimer's disease. *Am J Psychiatry* 1981; 138(7):970-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fitten LJ, Perryman KM, Gross PL, et al. Treatment of Alzheimer's disease with short- and long-term oral THA and lecithin: A double-blind study. *Am J Psychiatry* 1990 Feb; 147(2):239-42.

Status: Cross-over trial;

Cross-over trial

Fitten LJ, Ganzell S. Spouses' assessments of Alzheimer patients' response to THA and lecithin. *Am J Psychiatry* 1992; 149(4):575.

Status: Not included because not a full article

Fitzsimmons S, Buettner LL. Therapeutic recreation interventions for need-driven dementia-compromised behaviors in community-dwelling elders. *Am J Alzheimers Dis Other Dement* 2002; 17(6):367-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fleischhacker WW, Buchgeher A, Schubert H. Memantine in the treatment of senile dementia of the Alzheimer type. *Prog Neuropsychopharmacol Biol Psychiatry* 1986; 10(1):87-93.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fleming J. Dementia with Lewy bodies. J Am Geriatr Soc 1999 Jan; 47(1):121-2.

Status: Background article

Flicker C, Ferris SH, Kalkstein D, et al. A double-blind, placebo-controlled crossover study of ganglioside GM1 treatment for Alzheimer's disease. Am J Psychiatry 1994 Jan; 151(1):126-9.

Status: Cross-over trial;

Cross-over trial

Flicker L, Grimley EG. Piracetam for dementia or cognitive impairment. In: The Cochrane Library, 2001. Issue 2. Oxford: Update Software.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Folnegovic Smalc V, Knezevic S, Bokonjic R, et al. European pentoxifylline multi-infarct dementia trial: The empid study. J Neurol 1994; 2(41):158.

Status: Not included because not a full article

Folstein MF, Folstein SE, McHugh PR. "Mini-mental state". A practical method for grading the cognitive state of patients for the clinician. J Psychiatr Res 1975 Nov; 12(3):189-98.

Status: Background article;

Fontana RJ, Turgeon DK, Woolf TF, et al. The caffeine breath test does not identify patients susceptible to tacrine hepatotoxicity. Hepatology 1996 Jun; 23(6):1429-35.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fontana RJ, deVries TM, Woolf TF, et al. Caffeine based measures of CYP1A2 activity correlate with oral clearance of tacrine in patients with Alzheimer's disease. Br J Clin Pharmacol 1998 Sep; 46(3):221-8.

Status: Cross-over trial;

Cross-over trial

Food and Drug Administration. Tacrine as a treatment for Alzheimer's disease: Editor's note. An interim report from the FDA. A response from Summers et al. N Engl J Med 1991 Jan 31; 324(5):349-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ford JM, Truman CA, Wilcock GK, et al. Serum concentrations of tacrine hydrochloride predict its adverse effects in Alzheimer's disease. Clin Pharmacol Ther 1993; 53(6):691-5.

Status: Not included because dementia population not randomized to treatment

Forette F, Amery A, Staessen J, et al. Is prevention of vascular dementia possible? The Syst-Eur Vascular Dementia Project. Aging (Milano) 1991; 3(4):373-82.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Forette F, Hoover T, Gracon S, et al. A double-blind, placebo-controlled, enriched population study of tacrine in patients with Alzheimer's disease. Eur J Neurol 1995; 2:229-38.

Status: Not included because Jadad Quality Scale score less than three

Forette F, Seux ML, Staessen JA, et al. Prevention of dementia in randomised double-blind placebo-controlled Systolic Hypertension in Europe (Syst-Eur) trial. Lancet 1998 Oct 24; 352(9137):1347-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Forette F, Anand R, Gharabawi G. A phase II study in patients with Alzheimer's disease to assess the preliminary efficacy and maximum tolerated dose of rivastigmine (Exelon registered). Eur J Neurol 1999; 6(4):423-9.

Status: Included

Forssell LG, Sjobqvist B, Winblad B. Early stages of late onset Alzheimer's disease. III. Double blind treatment with choline chloride and lecithin with and without L-dopa and L-tryptophan, alternatively placebo. Acta Neurol Scand Suppl 1989; 79(121):43-66.

Status: Not included because dementia population not randomized to treatment

Forster DP, Newens AJ, Kay DW, et al. Risk factors in clinically diagnosed presenile dementia of the Alzheimer type: A case-control study in northern England. J Epidemiol Commun Health 1995 Jun; 49(3):253-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Foster HG, Hillbrand M, Chi CC. Efficacy of carbamazepine in assaultive patients with frontal lobe dysfunction. Prog Neuropsychopharmacol Biol Psychiatry 1989; 13(6):865

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Foster NL, Aldrich MS, Bluemlein L, et al. Failure of cholinergic agonist RS-86 to improve cognition and movement in PSP despite effects on sleep. *Neurology* 1989 Feb; 39(2 Pt 1):257-61.

*Status: Cross-over trial;
Cross-over trial*

Foster NL, Petersen RC, Gracon SI, et al. An enriched-population, double-blind, placebo-controlled, crossover study of tacrine and lecithin in Alzheimer's disease. *Dementia* 1996; 7(5):260-6.

*Status : Cross-over trial;
Cross-over trial*

Foster NL, Gombosi E, Teboe C, et al. Balanced centralized and distributed database design in a clinical research environment. *Stat Med* 2000 Jun 15; 19(11-12):1531-44.

Status: Background article

Foster NA, Valentine ER. Effect of auditory stimulation on autobiographical recall in dementia. *Exp Aging Res* 2001 Jul; (3):215-28.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fovall P, Dysken MW, Lazarus LW. Choline bitartrate treatment of Alzheimer-type dementias. *Commun Psychopharmacol* 1980; 4(2):141-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Frampton M, Harvey R, Kirchner V. Propentofylline for Dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software

Status: Background article

Francese T. Research corner. The effects of regular exercise on muscle strength and functional abilities of late stage Alzheimer's residents. *Va Nurse Today* 1995; (4):25-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Francese T, Sorrell J, Butler FR. The effects of regular exercise on muscle strength and functional abilities of late stage Alzheimer's residents. *Am J Alzheimers Dis* 1997; (3):122-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Franciosi A, Zavattini G. Dihydroergocristine in the treatment of elderly patients with cognitive deterioration: A double-blind, placebo-controlled,

dose- response study. *Curr Ther Res Clin Exp* 1994; 55(11):1391-401.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Frattola L, Piolti R, Bassi S, et al. Multicenter clinical comparison of the effects of choline alfoscerate and cytidine diphosphocholine in the treatment of multi-infarct dementia. *Curr Ther Res Clin Exp* 1991; 49(4):683-93.

Status: Not included because Jadad Quality Scale score less than three

Frattola L, Trabucchi H, Cucinotta D, et al. Dopamine agonist and free-radicals scavenger activities of dihydroergokryptine in dementia of alzheimer type (DAT): Multicentre, long-term double-blind clinical study versus placebo. *J Neurol* 1994; 241:159.

Status: Not included because not a full article

Frederick B, Satlin A, Wald LL, et al. Brain proton magnetic resonance spectroscopy in Alzheimer's disease: Changes after treatment with xanomeline. *Am J Geriatr Psychiatry* 2002 Jan; 10(1):81-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Freedman M, Rewilak D, Xerri T, et al. L-deprenyl in Alzheimer's disease: Cognitive and behavioral effects. *Neurology* 1998; 50(3):660-8.

Status: Included

Fric M, Horn R, Hasse SI, et al. Effects of nimodipine-treatment in primary degenerative dementia. Results of a clinical and psychometric study. *Pharmacopsychiatry* 1995; 28:177.

Status: Not included because not a full article

Friedman JI, Adler DN, Temporini HD, et al. Guanfacine treatment of cognitive impairment in schizophrenia. *Neuropsychopharmacology* 2001 Sep; 25(3):402-9.

Status: Background article

Friedman JI, Adler DN, Howanitz E, et al. Erratum: A double blind placebo controlled trial of donepezil adjunctive treatment to risperidone for the cognitive impairment of schizophrenia. *Biol Psychiatry* 2002; 51(12):1014.

Status: Not included because not a full article

Friedman JI, Adler DN, Howanitz E, et al. A double blind placebo controlled trial of donepezil adjunctive treatment to risperidone for the

cognitive impairment of schizophrenia. *Biol Psychiatry* 2002 Mar 1; 51(5):349-57.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Friedman R, Tappen RM. The effect of planned walking on communication in Alzheimer's disease. *J Am Geriatr Soc* 1991 Jul; 39(7):650-4.
Status: Not included because does not meet criteria for treatment for dementia patients

Frisoni GB, Gozzetti A, Bignamini V, et al. Special care units for dementia in nursing homes: A controlled study of effectiveness. *Arch Gerontol Geriatr* 1998; 27(Suppl 6):215-24.
Status: Not included because dementia population not randomized to treatment

Frith CD, Stevens M, Johnstone EC, et al. Effects of ECT and depression on various aspects of memory. *Br J Psychiatry* 1983; Vol 142(Jun):610-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fuchs A, Hehnke U, Erhart C, et al. Video rating analysis of effect of maprotiline in patients with dementia and depression. *Pharmacopsychiatry* 1993 Mar; 26(2):37-41.
Status: Included

Fuglum E, Schillinger A, Andersen JB, et al. Zuclopenthixol and haloperidol/levomepromazine in the treatment of elderly patients with symptoms of aggressiveness and agitation: A double-blind, multi-centre study. *Pharmatherapeutica* 1989; 5(5):285-91.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fulop TJ, Worum I, Csongor J, et al. Effects of centrophenoxine on body composition and some biochemical parameters of demented elderly people as revealed in a double-blind clinical trial. *Arch Gerontol Geriatr* 1990; 10(3):239-51.
Status: Not included because no extractable data relevant to review

Funfgeld EW, Baggen M, Nedwidek P, et al. Double blind study with phosphatidylserine (PS) in parkinsonian patients with senile dementia of Alzheimer's type (SDAT). *Alzheimers Dis Relat Disord* 1989; 1235-46.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fuschillo C, La Pia S, Campana F, et al. Cognitive deficits in Alzheimer's disease: Treatment with acetylcholinesterase inhibitor agents. *Arch Gerontol Geriatr* 2001; 33(suppl 1):151-8.
Status: Not included because Jadad Quality Scale score less than three

Fusgen I, Bressel H-U, De Mey C. A randomised, placebo-controlled, double-blind study of the efficacy and tolerability of dimenhydrinate in multi-morbid patients with senile dizziness. *Eur J Geriatr* 2002; 4(2):92-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gaber S, Ronzoni S, Bruni A, et al. Sertaline versus small doses of haloperidol in the treatment of agitated behaviour in pateints with dementia. *Arch Gerontol Geriatr Psychiatry* 2001; (Suppl 1):159-62.
Status: Not included because Jadad Quality Scale score less than three

Gabrynowicz JW, Dumbrell M. A clinical trial of leptazole with nicotinic acid in the management of psycho-geriatric patients. *Med J Aust* 1968 May 11; 1(19):799-802.
Status: Not included because no extractable data relevant to review

Gainotti G, Nocentini U, Sena E. Can the pattern of neuropsychological improvement obtained with cholinergic drugs be used to infer a cholinergic mechanism in other nootropic drugs? *Prog Neuropsychopharmacol Biol Psychiatry* 1989; 13(Suppl):S47-S59
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gaitz CM, Varner RV, Overall JE. Pharmacotherapy for organic brain syndrome in late life. Evaluation of an ergot derivative vs placebo. *Arch Gen Psychiatry* 1977 Jul; 34(7):839-45.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Galasko D, Hansen LA, Katzman R, et al. Clinical-neuropathological correlations in Alzheimer's disease and related dementias. *Arch Neurol* 1994 Sep; 51(9):888-95.
Status: Background article

Galasko DR, Gould RL, Abramson IS, et al. Measuring cognitive change in a cohort of patients

with Alzheimer's disease. *Stat Med* 2000 Jun 15; 19(11-12):1421-32.

Status: Background article

Gallai V, Mazzotta G, Firenze C, et al. Study of the P300 and cerebral maps in subjects with multi-infarct dementia treated with cytidine. *Psychopharmacologia* 1991; 103(1):1-5.

Status: Not included because no extractable data relevant to review

Gallai V, Mazzotta G, Del Gatto F, et al. A clinical and neurophysiological trial on nootropic drugs in patients with mental decline. *Acta Neurol (Napoli)* 1991 Feb; 13(1):1-12.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Galluzzi S, Cimaschi L, Ferrucci L, et al. Mild cognitive impairment: Clinical features and review of screening instruments. *Aging Clin Exp Res* 2001; 13(3):183-202.

Status: Background article

Gandy S. Will "cerebral proteopathy" be a useful construct for discovering one drug that shows efficacy against multiple neurodegenerative diseases? *Neurobiol Aging* 2000 Jul; 21(4):565

Status: Background article

Gao H, Yan L, Liu B, et al. Clinical study on treatment of senile vascular dementia by acupuncture. *J Tradit Chin Med* 2001 Jun; 21(2):103-9.

Status: Not included because Jadad Quality Scale score less than three

Gao S, Hui SL, Hall KS, et al. Estimating disease prevalence from two-phase surveys with non-response at the second phase. *Stat Med* 2000 Aug 30; 19(16):2101-14.

Status: Background article

Garetz FK, Baron JJ, Barron PB, et al. Efficacy of nylidrin hydrochloride in the treatment of cognitive impairment in the elderly. *J Am Geriatr Soc* 1979 May; 27(5):235-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Garrett AD. Therapeutic equivalence: fallacies and falsification. *Stat Med* 2003 Mar 15; 22(5):741-62.

Status: Background article

Garzya G, Corallo D, Fiore A, et al. Evaluation of the effects of L-acetylcarnitine on senile patients suffering from depression. *Drugs Exp Clin Res* 1990; 16(2):101-6.

Status: Background article

Gasbarrini G, Stefanini G, Addolorato G, et al. Posatirelin for the treatment of degenerative and vascular dementia: Results of explanatory and pragmatic efficacy analyses. *Arch Gerontol Geriatr* 1997; 26(1):33-47.

Status: Included

Gasnault J, Gueguen B, Bourdel MC, et al. Oral tacrine effects on computerised EEG activity during a double blind cross-over study in dementia of the Alzheimer type. *J Neurol* 1990; 237:33.

Status: Not included because not a full article

Gauthier S, Leblanc R, Robitaille Y, et al. Transmitter-replacement therapy in Alzheimer's disease using intracerebroventricular infusions of receptor agonists. *Can J Neurol Sci* 1986 Nov; 13(4 Suppl):394-402.

Status: Not included because dementia population not randomized to treatment

Gauthier S, Etienne P, Dastoor D, et al. Lack of effect of a 3-month treatment with lecithin in Alzheimer's disease. *Neurology* 1989; 31:89.

Status: Article not retrievable

Gauthier S, Bouchard R, Lamontagne A, et al. Tetrahydroaminoacridine-lecithin combination treatment in patients with intermediate-stage Alzheimer's disease. *N Engl J Med* 1990; 322(18):1272-6.

Status: Cross-over trial;

Cross-over trial

Gauthier S. Update on diagnostic methods, natural history and outcome variables in Alzheimer's disease. *Dement Geriatr Cogn Disord* 1998; 9(Suppl 3):2-7.

Status: Background article

Gauthier S, Feldman H, Hecker J, et al. Functional, cognitive and behavioral effects of donepezil in patients with moderate Alzheimer's disease. *Curr Med Res Opin* 2002; 18(6):347-54.

Status: Companion of an included article

Gedye JL, Exton-Smith AN, Wedgwood J. A method for measuring mental performance in the elderly and its use in a pilot clinical trial of

meclofenoxate in organic dementia (preliminary communication). *Age Ageing* 1972 May; 1(2):74-80.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gedye JL, Ibrahimi GS, McDonald C. A double blind controlled trial of piracetam (2-pyrrolidone acetamide) on two groups of psychogeriatric patients. *IRCS Med Sci Clin Med* 1978; 6(5):202.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gelb DJ. Measurement of progression in Alzheimer's disease: a clinician's perspective. *Stat Med* 2000 Jun 15; 19(11-12):1393-400.

Status: Background article

Gelinas I, Gauthier L, McIntyre M, et al. Development of a functional measure for persons with Alzheimer's disease: the disability assessment for dementia. *Am J Occup Ther* 1999 Sep; 53(5):471-81.

Status: Background article

Gelinas I, Gauthier S, Cyrus PA. Metrifonate enhances the ability of Alzheimer's disease patients to initiate, organize, and execute instrumental and basic activities of daily living. *J Geriatr Psychiatry Neurol* 2000; 13(1):9-16.

Status: Not included because dementia population not randomized to treatment

Geng J. Treatment of 50 cases of senile dementia by acupuncture combined with inhalation of herbal drugs and oxygen. *J Tradit Chin Med* 1999 Dec; 19(4):287-9.

Status: Not included because Jadad Quality Scale score less than three

George TP, Vessicchio JC, Termine A, et al. Effects of smoking abstinence on visuospatial working memory function in schizophrenia. *Neuropsychopharmacology* 2002; 26(1):75-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gerber GJ, Prince PN, Snider HG, et al. Group activity and cognitive improvement among patients with Alzheimer's disease. *Hospital & Community Psychiatry* 1991; 42(8):843-5.

Status: Not included because does not meet criteria for treatment for dementia patients

Gerdner LA. Effects of individualized versus classical "relaxation" music on the frequency of

agitation in elderly persons with Alzheimer's disease and related disorders. *Int Psychogeriatr* 2000 Mar; 12(1):49-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gerdner LA, Buckwalter KC, Reed D. Impact of a psychoeducational intervention on caregiver response to behavioral problems. *Nurs Res* 2002 Nov; 51(6):363-74.

Status: Not included because does not meet criteria for treatment for dementia patients

Gessner B, Voelp A, Klasser M. Study of the long-term action of a Ginkgo biloba extract on vigilance and mental performance as determined by means of quantitative pharmaco-EEG and psychometric measurements. *Arzneimittelforschung* 1985; 35(9):1459-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Getsios D, Caro JJ, Caro G, et al. Assessment of health economics in Alzheimer's disease (AHEAD): Galantamine treatment in Canada. *Neurology* 2001 Sep 25; 57(6):972-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ghatak R. Effects of an intervention program on dementia patients and their caregivers. *Caring* 1994 Aug; (No. 8):34-9.

Status: Not included because dementia population not randomized to treatment

Ghose K. Oxpentifylline in dementia: A controlled study. *Arch Gerontol Geriatr* 1987 Apr; 6(1):19-26.

Status: Included

Ghose K. Plasma drug levels of oxpentifylline in patients with dementia. An assessment of compliance. *Brain Dysfunct* 1989; (2):105-10.

Status: Not included because no extractable data relevant to review

Giacobini E, Spiegel R, Enz A, et al. Inhibition of acetyl- and butyryl-cholinesterase in the cerebrospinal fluid of patients with Alzheimer's disease by rivastigmine: Correlation with cognitive benefit. *J Neural Transm Gen Sect* 2002 Jul; 109(7-8):1053-65.

Status: Not included because dementia population not randomized to treatment

Ginsburg R, Weintraub M. Caffeine in the "sundown syndrome." Report of negative results. J Gerontol 1976 Jul; 31(4):419-20.
Status: Not included because dementia population not randomized to treatment

Gitlin LN, Corcoran M, Winter L, et al. A randomized, controlled trial of a home environmental intervention: Effect on efficacy and upset in caregivers and on daily function of persons with dementia. Gerontologist 2001 Feb; 41(1):4-14.
Status: Not included because does not meet criteria for treatment for dementia patients

Giuffra M, Mouradian MM, Bammert J, et al. Prolonged intravenous infusion of physostigmine in Alzheimer's disease. Neurology 1990; 40(Suppl 1):229, 1990.
Status: Not included because not a full article

Gleason RP, Schneider LS. Carbamazepine treatment of agitation in Alzheimer's outpatients refractory to neuroleptics. J Clin Psychiatry 1990; 51(3):115-8.
Status: Not included because dementia population not randomized to treatment

Goad DL, Davis CM, Liem P, et al. The use of selegiline in Alzheimer's patients with behavior problems. J Clin Psychiatry 1991; 52(8):342-5.
Status: Not included because dementia population not randomized to treatment

Gobburu JV, Tammara V, Lesko L, et al. Pharmacokinetic-pharmacodynamic modeling of rivastigmine, a cholinesterase inhibitor, in patients with Alzheimer's disease. J Clin Pharmacol 2001 Oct; 41(10):1082-90.
Status: Not included because dementia population not randomized to treatment

Goety CG, Tanner CM, Cohen JA, et al. L-acetylcarnitine in Huntington's disease: Double-blind placebo controlled crossover study of drug effects on movement disorder and dementia. Mov Disord 1990; 5(3):263-5.
Status: Not included because not a full article

Goldstein SE, Birnbaum F. Nylidrin HCL in the treatment of symptoms of the aged: A double-blind placebo controlled study. J Clin Psychiatry 1979 Dec; 40(12):520-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Goldwasser AN, Auerbach SM, Harkins SW. Cognitive, affective, and behavioral effects of reminiscence group therapy on demented elderly. Int J Aging Hum Dev 1987; 25(3):209-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Golomb BA. Statins and dementia. Arch Neurol 2001; 58(7):1169-70.
Status: Not included because not a full article

Goodwin GM, Conway SC, Peyro-Saint-Paul H, et al. Executive function and uptake of 99mTc-exametazime shown by single photon emission tomography after oral idazoxan in probable Alzheimer-type dementia. Psychopharmacologia 1997 Jun; 131(4):371-8.
Status: Cross-over trial; Cross-over trial

Gool WA, Waardenburg J. The effect of tetrahydroaminoacridine (THA) on P300 in Alzheimer's disease. Biol Psychiatry 1991; 30(9):953-7.
Status: Companion of an included article

Gori G, Pientini S, Vespa A. The selection of meaningful activities as a treatment for day-care in dementia. Arch Gerontol Geriatr 2001; 33(Suppl):207-12.
Status: Not included because dementia population not randomized to treatment

Gorman DG, Read S, Cummings JL. Cholinergic therapy of behavioral disturbances in Alzheimer's disease. Neuropsychiatry Neuropsychol Behav Neurol ; 6(4):229-34.
Status: Cross-over trial; Cross-over trial

Gormley N, Lyons D, Howard R. Behavioural management of aggression in dementia: A randomized controlled trial. Age Ageing 2001 Mar; 30(2):141-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Gortelmeyer R, Erbler H. Memantine in the treatment of mild to moderate dementia syndrome. A double-blind placebo-controlled study. Arzneimittelforschung 1992 Jul; 42(7):904-13.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gotestam KG, Ljunghall S, Olsson B. A double-blind comparison of the effects of haloperidol and cis-cloperthixol in senile dementia. *Acta Psychiatr Scand* 1981; 294(Suppl):46-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gotestam KG, Melin L. The effect of prompting and reinforcement of activity in elderly demented inpatients. *Scand J Psychol* 1990; 31(1):2-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gottfries CG, Karlsson I, Nyth AL. Treatment of depression in elderly patients with and without dementia disorders. *Int Clin Psychopharmacol* 1992; 6(Suppl 5):55-64.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gracon SI, Hoover TM, Lewis KW, et al. Tacrine in Alzheimer's disease efficacy and safety in a parallel group study. *Alzheimer's Disease Advances in Clinical and Basic Research* 1993; 549-57.
Status : Not included because not a full article

Gracon SI. Evaluation of tacrine hydrochloride (Cognexregistered trade mark)in two parallel-group studies. *Acta Neurol Scand Suppl* 1996; 93(165):114-22.
Status: Companion of an included article

Gracon SI, Knapp MJ, Berghoff WG, et al. Safety of tacrine: Clinical trials, treatment IND, and postmarketing experience. *Alzheimer Dis Assoc Disord* 1998 Jun; 12(2):93-101.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Graf A, Wallner C, Schubert V, et al. The effects of light therapy on mini-mental state examination scores in demented patients. *Biol Psychiatry* 2001 Nov 1; 50(9):725-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Graham JE, Rockwood K, Beattie BL, et al. Standardization of the diagnosis of dementia in the Canadian Study of Health and Aging. *Neuroepidemiology* 1996; 15(5):246-56.
Status: Background article

Graham JE, Rockwood K, Beattie BL, et al. Prevalence and severity of cognitive impairment with and without dementia in an elderly

population. *Lancet* 1997 Jun 21; 349(9068):1793-6.
Status: Background article

Granier S. Does estrogen prevent Alzheimer's progression. *Ochsner Journal* 2000; 2(4):238-9.
Status: Article not retrievable

Grasser A, Gotthardt U, Heuser-Link M, et al. Behavioral effects of an ACTH4-9-fragment in a mixed sample of depressed patients and patients with Alzheimer's disease. *Pharmacopsychiatry* 1992; Vol 25:102
Status: Not included because not a full article

Graves AB, van Duijn CM, Chandra V, et al. Occupational exposures to solvents and lead as risk factors for Alzheimer's disease: a collaborative re-analysis of case-control studies. EURODEM Risk Factors Research Group. *Int J Epidemiol* 1991; 20 Suppl 2:S58-S61
Status: Background article

Green J, McDonald WM, Vitek JL, et al. Neuropsychological and psychiatric sequelae of pallidotomy for PD: Clinical trial findings. *Neurology* 2002; 58(6):858-65.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Green MF, Marder SR, Glynn SM, et al. The neurocognitive effects of low-dose haloperidol: A two-year comparison with risperidone. *Biol Psychiatry* 2002; 51(12):972-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Green RC, Goldstein FC, Auchus AP, et al. Treatment trial of oxiracetam in Alzheimer's disease. *Arch Neurol* 1992 Nov; 49(11):1135-6.
Status: Not included because dementia population not randomized to treatment

Greenberg SM, Tennis MK, Brown LB, et al. Donepezil therapy in clinical practice: A randomized crossover study. *Arch Neurol* 2000 Jan; 57(1):94-9.
Status: Cross-over trial;
Cross-over trial

Greendyke RM, Kanter DR, Schuster DB, et al. Propranolol treatment of assaultive patients with organic brain disease. A double-blind crossover, placebo-controlled study. *J Nerv Ment Dis* 1986 May; 174(5):290-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Greendyke RM, Kanter DR. Therapeutic effects of pindolol on behavioral disturbances associated with organic brain disease: A double-blind study. *J Clin Psychiatry* 1986 Aug; 47(8):423-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Greendyke RM, Berkner JP, Webster JC, et al. Treatment of behavioral problems with pindolol. *Psychosomatics* 1989; 30(2):161-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Greer KL, Pustay KA, Zaun TC, et al. Comparison of the effects of toys versus live animals on the communication of patients with dementia of the Alzheimer's type. *Clin Gerontol* 2001; (3-4):157-82.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grieve AP. Issues for statisticians in pharmacoeconomic evaluations. *Stat Med* 1998 Aug 15; 17(15-16):1715-23.

Status: Background article

Grioli S, Lomeo C, Quattropiani MC, et al. Pyroglutamic acid improves the age associated memory impairment. *Fundam Clin Pharmacol* 1990; 4(2):169-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gross RJ, Eisdorfer CE, Schiller HS, et al. Effect of ergot alkaloids on serum prolactin in non-psychotic organic brain syndrome of the elderly. *Exp Aging Res* 1979 Aug; 5(4):293-302.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grossberg GT, Stahelin HB, Messina JC, et al. Lack of adverse pharmacodynamic drug interactions with rivastigmine and twenty-two classes of medications. *Int J Geriatr Psychiatry* 2000 Mar; 15(3):242-7.

Status: Not included because dementia population not randomized to treatment

Grossman M, Mickanin J, Onishi K, et al. An aspect of sentence processing in Alzheimer's disease: Quantifier-noun agreement. *Neurology* 1995 Jan; 45(1):85-91.

Status: Not included because does not meet criteria for treatment for dementia patients

Grossmann WM, Standl A, May U, et al. Naftidrofuryl in the treatment of mild senile dementia. A double-blind study.

Pharmacopsychiatry 1990 Nov; 23(6):265-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grothe DR, Piscitelli SC, Dukoff R, et al. Penetration of tacrine into cerebrospinal fluid in patients with Alzheimer's disease. *J Clin Psychopharmacol* 1998; 18(1):78-81.

Status: Not included because dementia population not randomized to treatment

Group for the Advancement of Psychiatry and Committee on Aging. Impact of tacrine in the care of patients with Alzheimer's disease: What we know one year after FDA approval. *Am J Geriatr Psychiatry* 1994; 2(4):285-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Growdon JH, Corkin S, Huff FJ, et al. Piracetam combined with lecithin in the treatment of Alzheimer's disease. *Neurobiol Aging* 1986 Jul; 7(4):269-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gu P, Jing QW, Ding XS. Observation of effect on Alzheimer's disease treated by Galanthamine hydrobromide capules. *Practical Geriatrics* 2000; (6):307-8.

Status: Article not retrievable

Guerzoni A, Santambrogio S. Efficacy of dihydroergocristine 20 mg once daily in patients with organic brain psychosyndrome. A 3-month randomised, double-blind, placebo-controlled study. *Clin Drug Investig* 1995; 10(1):1-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Guimon J, Blanco J, Caso C. L-Dopa carbidopa treatment of senile dementia: A control study. *Eur J Psychiatry* 1995; (1):29-36.

Status: Not included because Jadad Quality Scale score less than three

Gustafson L, Risberg J, Johanson M, et al. Effects of piracetam on regional cerebral blood flow and mental functions in patients with organic dementia. *Psychopharmacologia* 1978 Mar 1;

56(2):115-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gustafson L, Edvinsson L, Dahlgren N, et al. Intravenous physostigmine treatment of Alzheimer's disease evaluated by psychometric testing, regional cerebral blood flow (rCBF) measurement, and EEG. *Psychopharmacologia* 1987; 93(1):31-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gustafson L. Physostigmine and tetrahydroaminoacridine treatment of Alzheimer's disease. *Acta Neurol Scand Suppl* 1993; 149(Rand):39-41.

*Status: Cross-over trial;
Cross-over trial*

Gutzmann H, Kuhl KP, Kanowski S, et al. Measuring the efficacy of psychopharmacological treatment of psychomotoric restlessness in dementia: clinical evaluation of tiapride. *Pharmacopsychiatry* 1997 Jan; 30(1):6-11.

Status: Included

Gutzmann H, Hadler D. Sustained efficacy and safety of idebenone in the treatment of Alzheimer's disease: Update on a 2-year double-blind multicentre study. *J Neural Transm Suppl* 1998; 54:301-10.

Status: Included

Gutzmann H, Kuhl KP, Hadler D, et al. Safety and efficacy of idebenone versus tacrine in patients with Alzheimer's disease: Results of a randomized, double-blind, parallel-group multicenter study. *Pharmacopsychiatry* 2002 Jan; 35(1):12-8.

Status: Included

Haaland KY, Harrington DL, O'Brien S, et al. Cognitive-motor learning in Parkinson's disease. *Neuropsychology* 1997; 11(2):180-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hackman BW, Galbraith. Replacement therapy with piperazine oestrone sulphate (harmogen) and its effect on memory. *Curr Med Res Opin* 1976; 4(4):303-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hackman BW, Galbraith. Six-month pilot study of oestrogen replacement therapy and piperazine oestrone sulphate ('Harmogen') and its effect on memory. *Curr Med Res Opin* 1977; 4(3):21-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hadas-Lidor N, Katz N, Tyano S, et al. Effectiveness of dynamic cognitive intervention in rehabilitation of clients with schizophrenia. *Clin Rehabil* 2001; 15(4):349-59.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Haffmans PM, Sival RC, Lucius SA, et al. Bright light therapy and melatonin in motor restless behaviour in dementia: A placebo-controlled study. *Int J Geriatr Psychiatry* 2001 Jan; 16(1):106-10.

*Status: Cross-over trial;
Cross-over trial*

Hagstadius S, Gustafson L, Risberg J. The effects of bromvincamine and vincamine on regional cerebral blood flow and mental functions in patients with multi-infarct dementia. *Psychopharmacologia* 1984; 83(4):321-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Haley WE, Levine EG, Brown SL, et al. Stress, appraisal, coping, and social support as predictors of adaptational outcome among dementia caregivers. *Psychol Aging* 1987 Dec; 2(4):323-30.

Status: Background article

Haley WE, Brown SL, Levine EG. Experimental evaluation of the effectiveness of group intervention for dementia caregivers. *Gerontologist* 1987; 27(3):376-82.

Status: Not included because does not meet criteria for treatment for dementia patients

Hall CB, Lipton RB, Sliwinski M, et al. A change point model for estimating the onset of cognitive decline in preclinical Alzheimer's disease. *Stat Med* 2000 Jun 15; 19(11-12):1555-66.

Status: Background article

Hall P, Harcup M. A trial of lipotropic enzymes in atheromatous ("arteriosclerotic") dementia. *Angiology* 1969 May; 20(5):287-300.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Halliday GM, Shepherd CE, McCann H, et al. Effect of anti-inflammatory medications on neuropathological findings in Alzheimer's disease. Arch Neurol 2000 Jun; 57(6):831-6.
Status: Not included because dementia population not randomized to treatment

Hammeke TA, Haughton VM, Grogan JP, et al. A preliminary study of cognitive and affective alterations following intrathecal administration of iopamidol or metrizamide. Invest Radiol 1984; 19(Suppl 5):S268-S271.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hammen CL, Jacobs M, Mayol A, et al. Dysfunctional cognitions and the effectiveness of skills and cognitive-behavioral assertion training. J Consult Clin Psychol 1980 Dec; 48(6):685-95.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hanley IG, McGuire RJ, Boyd WD. Reality orientation and dementia: A controlled trial of two approaches. Br J Psychiatry 1981; 138:10-4.
Status: Not included because dementia population not randomized to treatment

Hansson L. Antihypertensive treatment and the prevention of dementia: Further insights from the Syst-Eur trial. J Hypertens 1999; 17(3):307-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hanyu H, Tanaka Y, Sakurai H, et al. Atrophy of the substantia innominata on magnetic resonance imaging and response to donepezil treatment in Alzheimer's disease. Neurosci Lett 2002 Feb 8; 319(1):33-6.
Status: Not included because dementia population not randomized to treatment

Harbaugh RE, Roberts DW, Coombs DW, et al. Preliminary report: Intracranial cholinergic drug infusion in patients with Alzheimer's disease. Neurosurgery 1984 Oct; 15(4):514-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harbaugh RE. Intracerebroventricular cholinergic drug administration in Alzheimer's disease: Preliminary results of a double-blind study. J Neural Transm Suppl 1987; 24:271-7.
*Status: Cross-over trial;
Cross-over trial*

Harbaugh RE, Reeder TM, Senter HJ, et al. Intracerebroventricular bethanechol chloride infusion in Alzheimer's disease. Results of a collaborative double-blind study. J Neurosurg 1989 Oct; 71(4):481-6.
*Status: Cross-over trial;
Cross-over trial*

Harding GF, Hall P, Young J, et al. Multifocal infarct dementia treated by cyclandelate and monitored by quantitative EEG. Angiology 1978 Feb; 29(2):139-40.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harding GFA, Hall P, Young J, et al. Multi focal infarct dementia (cerebral arteriosclerosis) treated by cyclandelate and monitored by quantitative electroencephalography. IRCS Med Sci Biomed Technol 1977; 5(3-4):118-4.
Status: Article not retrievable

Harenko A. A comparison between chlormethiazole and nitrazepam as hypnotics in psycho-geriatric patients. Curr Med Res Opin 1974 Jul; 2(10):657-63.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harkins SW, Taylor JR, Mattay VS. Response to tacrine in patients with dementia of the Alzheimer's type: cerebral perfusion change is related to change in mental status. Int J Neurosci 1996 Feb; 84(1-4):149-4.
Status: Not included because dementia population not randomized to treatment

Harrell LE, Callaway R, Morere D, et al. The effect of long-term physostigmine administration in Alzheimer's disease. Neurology 1990; 40(9):1350-4.
*Status: Cross-over trial;
Cross-over trial*

Harrell LE, Jope RS, Falgout J, et al. Biological and neuropsychological characterization of physostigmine responders and nonresponders in Alzheimer's disease. J Am Geriatr Soc 1990 Feb; 38(2):113-22.
*Status: Cross-over trial;
Cross-over trial*

Harrell TH, Ryon NB. Cognitive-behavioral assessment of depression: Clinical validation of the automatic thoughts questionnaire. J Consult Clin Psychol 1983 Oct; 51(5):721-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harris RP, Helfand M, Woolf SH, et al. Current methods of the US Preventive Services Task Force: a review of the process. *Am J Prev Med* 2001 Apr; 20(3 Suppl):21-35.

Status: Background article

Harris S, Dowson JH. The effects of Meclofenoxate on cognitive performance in elderly individuals with memory impairment: A placebo-controlled study. *Int J Geriatr Psychiatry* 1986; 1:93-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hart BD, Wells DL. The effects of language used by caregivers on agitation in residents with dementia. *Clin Nurse Spec* 1997 Jan; 11(1):20-3. *Status: Not included because dementia population not randomized to treatment*

Hart S, Smith CM, Swash M, et al. Word fluency in patients with early dementia of Alzheimer's type. *Br J Clin Psychol* 1988; 46(Pt 2):115-24, 1592-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hartmann A. Cerebral blood flow in patients with cerebrovascular disorders: study with pentoxifylline. *Ric Clin Lab* 1981; 11(Suppl 1):243-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hartmann A. Comparative randomized study of cerebral blood flow after long-term administration of pentoxifylline and co-dergocrine mesylate in patients with chronic cerebrovascular disease. *Curr Med Res Opin* 1985; 9(7):475-9. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Hartmann A, Tsuda Y. A controlled study on the effect of pentoxifylline and an ergot alkaloid derivative on regional cerebral blood flow in patients with chronic cerebrovascular disease. *Angiology* 1988 May; 39(5):449-57. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Harvey PD, Moriarty PJ, Serper MR, et al. Practice-related improvement in information processing with novel antipsychotic treatment.

Schizophr Res 2000; 46(2-3):139-48.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harwart D. The treatment of chronic cerebrovascular insufficiency. A double-blind study with pentoxifylline ('Trental' 400). *Curr Med Res Opin* 1979; 6(2):73-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hauber AB, Gnanasakthy A, Mauskopf JA. Savings in the cost of caring for patients with Alzheimer's disease in Canada: An analysis of treatment with rivastigmine. *Clin Ther* 2000 Apr; 22(4):439-51.

Status: Not included because does not meet criteria for treatment for dementia patients

Hauer K, Marburger C, Oster P. Motor performance deteriorates with simultaneously performed cognitive tasks in geriatric patients. *Arch Phys Med Rehabil* 2002 Feb; 83(2):217-23. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Haupt M, Karger A, Janner M. Improvement of agitation and anxiety in demented patients after psychoeducative group intervention with their caregivers. *Int J Geriatr Psychiatry* 2000 Dec; 15(12):1125-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Hebert R, Leclerc G, Bravo G, et al. Efficacy of a support group programme for care-givers of demented patients in the community: A randomized controlled trial. *Arch Gerontol Geriatr* 1994; 18(1):1-14.

Status: Not included because does not meet criteria for treatment for dementia patients

Hebert R, Girouard D, Leclerc G, et al. The impact of a support group programme for care-givers on the institutionalisation of demented patients. *Arch Gerontol Geriatr* 1995; 20(2):129-34.

Status: Not included because does not meet criteria for treatment for dementia patients

Hebert R, Lindsay J, Verreault R, et al. Vascular dementia : incidence and risk factors in the Canadian study of health and aging. *Stroke* 2000 Jul; 31(7):1487-93.

Status: Background article

Heinonen EH, Savijarvi M, Kotila M, et al. Effects of monoamine oxidase inhibition by selegiline on concentrations of noradrenaline and monoamine metabolites in CSF of patients with Alzheimer's disease. *J Neural Transm Park Dis Dement Sect* 1993; 5(3):193-202.

*Status: Cross-over trial;
Cross-over trial*

Heinze B, Karrass W, Peters T. Pharmacopsychological effects of flunarizine in geriatric patients with light brainorganic psychosyndrome. Preliminary communication. *Eur Neurol* 1986; 25(Suppl 1):115-21.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heiss WD, Kessler J, Slansky I, et al. Activation PET as an instrument to determine therapeutic efficacy in Alzheimer's disease. *Ann N Y Acad Sci* 1993; 695(Sep 24):327-13.
Status: Not included because Jadad Quality Scale score less than three

Heiss WD, Kessler J, Mielke R, et al. Long-term effects of phosphatidylserine, pyritinol, and cognitive training in Alzheimer's disease. A neuropsychological, EEG, and PET investigation. *Dementia* 1994 Mar; 5(2):88-98.
Status: Not included because Jadad Quality Scale score less than three

Helkala EL, Koivisto K, Hanninen T, et al. Stability of age-associated memory impairment during a longitudinal population-based study. *J Am Geriatr Soc* 1997 Jan; 45(1):120-2.
Status: Not included because not a full article

Hemmeter U, Annen B, Bischof R, et al. Polysomnographic effects of adjuvant ginkgo biloba therapy in patients with major depression medicated with trimipramine. *Pharmacopsychiatry* 2001; 34(2):50-9.
Status: Background article

Henderson VW, Roberts E, Wimer C, et al. Multicenter trial of naloxone in Alzheimer's disease. *Ann Neurol* 1989 Apr; 25(4):404-6.
*Status: Cross-over trial;
Cross-over trial*

Henderson VW, Paganini-Hill A, Miller BL, et al. Estrogen for Alzheimer's disease in women: Randomized, double-blind, placebo-controlled trial. *Neurology* 2000 Jan 25; 54(2):295-301.
Status: Included

Henderson VW, Paganini-Hill A, Miller BL, et al. A randomized controlled trial of estrogen for the treatment of Alzheimer's disease in women. *Neurology* 2000; 54(Suppl 3):A470.
Status: Not included because not a full article

Henke CJ, Burchmore MJ. The economic impact of tacrine in the treatment of Alzheimer's disease. *Clin Ther* 1997; 19(2):330-45.
Status: Companion of an included article

Herrmann N. Cognitive pharmacotherapy of Alzheimer's disease and other dementias. *Can J Psychiatry* 2002 Oct; 47(8):715-22.
Status: Background article

Herrmann WM, Kern U, Rohmel J. On the effects of pyritinol on functional deficits of patients with organic mental disorders. *Pharmacopsychiatry* 1986 Sep; 19(5):378-85.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Kern U, Rohmel J. Contribution to the search for vigilance-indicative EEG variables. Results of a controlled, double-blind study with pyritinol in elderly patients with symptoms of mental dysfunction. *Pharmacopsychiatry* 1986 Mar; 19(2):75-83.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Dietrich B, Hiersemenzel R. Pharmacoelectroencephalographic and clinical effects of the cholinergic substance--acetyl-L-carnitine--in patients with organic brain syndrome. *Int J Clin Pharmacol Res* 1990; 10(1-2):81-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Stephan K. Moving from the question of efficacy to the question of therapeutic relevance: an exploratory reanalysis of a controlled clinical study of 130 inpatients with dementia syndrome taking piracetam. *Int Psychogeriatr* 1992; 4(1):25-44.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Stephan K, Gaede K, et al. A multicenter randomized double-blind study on the efficacy and safety of nicergoline in patients with multi-infarct dementia. *Dement Geriatr Cogn Disord* 1997 Jan; 8(1):9-17.
Status: Included

Herz LR, Volicer L, Ross V, et al. A single-case-study method for treating resistiveness in patients with Alzheimer's disease. *Hospital & Community Psychiatry* 1992 Jul; 43(7):720-4.

*Status: Cross-over trial;
Cross-over trial*

Heseker H, Kubler W, Pudel V, et al. Interaction of vitamins with mental performance. *Bibl Nutr Dieta* 1995; (52):43-55.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heseltine PN, Goodkin K, Atkinson JH, et al. Randomized double-blind placebo-controlled trial of peptide T for HIV-associated cognitive impairment. *Arch Neurol* 1998 Jan; 55(1):41-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heuser I, Heuser-Link M, Gotthardt U, et al. Behavioral effects of a synthetic corticotropin 4-9 analog in patients with depression and patients with Alzheimer's disease. *J Clin Psychopharmacol* 1993 Jun; 13(3):171-4.
Status: Not included because dementia population not randomized to treatment

Hewawasam L. Floor patterns limit wandering of people with Alzheimer's. *Nurs Times* 1996 May 29; 92(22):41-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heyman A, Schmechel D, Wilkinson W, et al. Failure of long term high-dose lecithin to retard progression of early-onset Alzheimer's disease. *J Neural Transm Suppl* 1987; Vol 24:279-86.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hickey P, Stewart N, Price J. Pentoxifylline for dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software
Status: Background article

Higgins JP, Flicker L. Lecithin for dementia and cognitive impairment. In: *The Cochrane Library*, 2000. Issue 4. Oxford: Update Software
Status: Background article

Hinchliffe AC, Katona C, Livingston G. The assessment and management of behavioural manifestations of dementia: A review and results of a controlled trial. *Int J Psychiatry Clin Pract* 1997; 1(3):157-68.

Status: Not included because does not meet criteria for treatment for dementia patients

Hinchliffe AC, Hyman IL, Blizard B, et al. Behavioural complications of dementia - Can they be treated? *Int J Geriatr Psychiatry* 1995; 10(10):839-47.

Status: Not included because does not meet criteria for treatment for dementia patients

Hindle JV, Meara RJ, Sharma JC, et al. Prescribing pergolide in the elderly - An open label study of pergolide in elderly patients with Parkinson's disease. *Int J Geriatr Psychopharmacol* 1998; (2):78-81.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hindmarch I, Fuchs HH, Erzigkeit H. Efficacy and tolerance of vinpocetine in ambulant patients suffering from mild to moderate organic psychosyndromes. *Int Clin Psychopharmacol* 1991; 6(1):31-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hinkin CH, Castellon SA, Hardy DJ, et al. Methylphenidate improves HIV-1-associated cognitive slowing. *J Neuropsychiatry Clin Neurosci* 2001; 13(2):248-54.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hirohata S, Suda H, Hashimoto T. Low-dose weekly methotrexate for progressive neuropsychiatric manifestations in Behcet's disease. *JNS* 1998 Aug 14; 159(2):181-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hjorth A, Browne E, Jakobsen K, et al. Organic brain syndrome treated with oxiracetam. A double-blind randomized controlled trial. *Acta Neurol Scand* 1987 Apr; 75(4):271-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hock C, Maddalena A, Heuser I, et al. Treatment with the selective muscarinic agonist talsaclidine decreases cerebrospinal fluid levels of total amyloid beta-peptide in patients with Alzheimer's disease. *Ann N Y Acad Sci* 2000; 920:285-91.
Status: Not included because no extractable data relevant to review

Hodges, Graham JR, K.S. A reversal of the temporal gradient for famous person knowledge in semantic dementia: implications for the neural organisation of long-term memory. *Neuropsychologia* 1998; 36(8):803-25.

Status: Not included because does not meet criteria for treatment for dementia patients

Hoeffler B, Rader J, McKenzie D, et al. Reducing aggressive behavior during bathing cognitively impaired nursing home residents. *J Gerontol Nurs* 1997 May; 23(5):16-23.

Status: Not included because dementia population not randomized to treatment

Hofferberth B. The efficacy of EGb 761 in patients with senile dementia of the Alzheimer type, a double-blind, placebo-controlled study on different levels of investigation. *Hum Psychopharmacol* 1994; (3):215-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hogervorst E, Yaffe K, Richards M, et al. Hormone replacement therapy to maintain cognitive function in women with dementia. In: *The Cochrane Library*, 2002. Issue 3. Oxford: Update Software

Status : Background article

Hogervorst E, Yaffe K, Richards M, et al. Hormone replacement therapy for cognitive function in postmenopausal women (Cochrane Review). In: *The Cochrane Library*, 2002. Issue 4. Oxford: Update Software

Status: Background article

Holden M, Kelly C. Use of cholinesterase inhibitors in dementia. *Adv Psychiatr Treat* 2002; 8(2):89-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hollander E, Davidson M, Mohs RC, et al. RS 86 in the treatment of Alzheimer's disease: cognitive and biological effects. *Biol Psychiatry* 1987 Sep; 22(9):1067-78.

*Status: Cross-over trial;
Cross-over trial*

Holliman DC, Orgassa UC, Forney JP. Developing an interactive physical activity group in a geriatric psychiatry facility. *Activities Adaptation Aging* 2001; 26(1):57-69.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Holm A, Michel M, Stern GA, et al. The outcomes of an inpatient treatment program for geriatric patients with dementia and dysfunctional behaviors. *Gerontologist* 1999 Dec; 39(6):668-76.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Holmes C, Hopkins V, Hensford C, et al. Lavender oil as a treatment for agitated behaviour in severe dementia: A placebo controlled study. *Int J Geriatr Psychiatry* 2002 Apr; 17(4):305-8.
Status: Not included because dementia population not randomized to treatment

Holzman D. Ginkgo biloba for Alzheimer's disease. *Altern Complement Ther* 1998; (5):361-3.

Status: Not included because not a full article

Homma A, Takeda M, Imai Y, et al. Clinical efficacy and safety of donepezil on cognitive and global function in patients with Alzheimer's disease. A 24-week, multicenter, double-blind, placebo-controlled study in Japan. *E2020 Study Group. Dement Geriatr Cogn Disord* 2000 Nov; 11(6):299-313.

Status: Not included because Jadad Quality Scale score less than three

Honjo H, Ogino Y, Naitoh K, et al. In vivo effects by estrone sulfate on the central nervous system-senile dementia (Alzheimer's type). *J Steroid Biochem* 1989; 34(1-6):521-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Honjo H, Ogino Y, Tanaka K. An effect of conjugated estrogen to cognitive impairment in women with senile dementia-Alzheimer's type: a placebo-controlled, double-blind study. *Journal of the Japanese Menopause Society* 1993; 1:167-71.

Status: Article not retrievable

Honjo H, Tanaka T, Urabe M, et al. Senile-dementia Alzheimer's type and estrogen. *Hormone & Metabolic Research* 1995; 27(4):204-7.

Status: Not included because dementia population not randomized to treatment

Hopman-Rock M, Staats PG, Tak EC, et al. The effects of a psychomotor activation programme for use in groups of cognitively impaired people in homes for the elderly. *Int J Geriatr Psychiatry* 1999 Aug; 14(8):633-42.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hornig P. Commentary on A 30-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. *Nursing Scan in Research* 1994; 7(4):6

Status: Not included because not a full article

Hossain M, Jhee SS, Shiovitz T, et al. Estimation of the absolute bioavailability of rivastigmine in patients with mild to moderate dementia of the Alzheimer's type. *Clin Pharmacokinet* 2002; 41(3):225-34.

Status: Not included because no extractable data relevant to review

Hoyer S, Oesterreich K, Stoll KD. Effects of Pyritinol HCl on blood flow and oxidative metabolism of the brain in patients with dementia. *Arzneimittelforschung* 1977; 27(3):671-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hozumi S, Hori H, Okawa M, et al. Favorable effect of transcranial electrostimulation on behavior disorders in elderly patients with dementia: a double-blind study. *Int J Neurosci* 1996 Nov; 88(1-2):1-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Huang PX, Yang ZM, Huang M, et al. Effect of Chifukang oral liquor on senile dementia. *Journal of Guangzhou University of Traditional Chinese Medicine* 1997; (3):145-9.

Status: Article not retrievable

Hudson S, Tabet N. Acetyl-L-carnitine for dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software.

Status: Background article

Huff FJ, Antuono P, Murphy M, et al. Potential clinical use of an adrenergic/cholinergic agent (HP 128) in the treatment of Alzheimer's disease. *Ann N Y Acad Sci* 1991; 640:263-7.

Status: Included

Huff FJ, Shipley J, Somerville NJ, et al. Besipirdine (HP 749) treatment trial in Alzheimer's disease. *Neurology* 1995; 45(Suppl 4):A288.

Status: Not included because not a full article

Huff FJ. Preliminary evaluation of besipirdine for the treatment of Alzheimer's disease. *Ann N Y Acad Sci* 1996; 777:410-4.

Status: Not included because Jadad Quality Scale score less than three

Huff FJ, Antuono PG, Delagandara JE, et al. A treatment and withdrawal trial of besipirdine in Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1996; 10(2):93-102.

Status: Not included because Jadad Quality Scale score less than three

Hui SL, Gao S. Spacing of follow-up waves in incidence studies. *Stat Med* 2000 Jun 15; 19(11-12):1567-75.

Status: Background article

Huppert FA, van Niekerk JK, Herbert J. Dehydroepiandrosterone (DHEA) supplementation for cognition and well-being. In: *The Cochrane Library*, 2000. Issue 2. Oxford: Update Software.

Status: Background article

Husereau D, Wolfson C, Shukla VK. Drug treatment for Alzheimer's disease: Efficacy, outcome measurements and cost-effectiveness. *Technology Overview no. 4*. Ottawa: Canadian Coordinating Office for Health Technology Assessment (CCOHTA). 2001.

Status: Background article

Huster WJ, Enas GG. A framework establishing clear decision criteria for the assessment of drug efficacy. *Stat Med* 1998 Aug 15; 17(15-16):1829-38.

Status: Background article

Huusko TM, Karppi P, Avikainen V, et al. Randomised, clinically controlled trial of intensive geriatric rehabilitation in patients with hip fracture: Subgroup analysis of patients with dementia. *BMJ* 2000 Nov 4; 321(7269):1107-11.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hyman BT, Eslinger PJ, Damasio AR. Effect of naltrexone on senile dementia of the Alzheimer type. *J Neurol Neurosurg Psychiatry* 1985; 48(11):1169-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hyman BT, Gomez-Isla T, Rebeck GW, et al. Epidemiological, clinical, and neuropathological

study of apolipoprotein E genotype in Alzheimer's disease. *Ann N Y Acad Sci* 1996 Dec 16; 802:1-5.

Status: Background article

Ikeda S, Yamada Y, Ikegami N. Economic evaluation of donepezil treatment for Alzheimer's disease in Japan. *Dement Geriatr Cogn Disord* 2002; 13(1):33-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Imbimbo BP, Perini M, Verdelli G, et al. Two year treatment of Alzheimer's disease with eptastigmine. *JNS* 1997; 150(Suppl 1):S154.

Status: Companion of an included article

Imbimbo BP, Nicoli M, Martini C, et al. Acetylcholinesterase assay may predict cognitive response of Alzheimer patients to eptastigmine treatment. *Eur J Clin Pharmacol* 1998 Nov; 54(9-10):809-10.

Status: Not included because not a full article

Imbimbo BP, Lucca U, Lucchelli F, et al. A 25-week placebo-controlled study of eptastigmine in patients with Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1998 Dec; 12(4):313-22.

Status: Not included because Jadad Quality Scale score less than three

Imbimbo BP, Martelli P, Troetel WM, et al. Efficacy and safety of eptastigmine for the treatment of patients with Alzheimer's disease. *Neurology* 1999 Mar 10; 52(4):700-8.

Status: Included

Imbimbo BP, Verdelli G, Martelli P, et al. Two-year treatment of Alzheimer's disease with eptastigmine. *Dement Geriatr Cogn Disord* 1999; 10(2):139-47.

Status: Not included because dementia population not randomized to treatment

Imbimbo BP, Troetel WM, Martelli P, et al. A 6-month, double-blind, placebo-controlled trial of eptastigmine in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2000 Jan; 11(1):17-24.

Status: Not included because Jadad Quality Scale score less than three

In't Veld BA, Ruitenberg A, Hofman A, et al. Antihypertensive drugs and incidence of dementia: The Rotterdam Study. *Neurobiol Aging* 2001; 22(3):407-12.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ingersoll-Dayton B, Schroepfer T, Pryce J. The effectiveness of a solution-focused approach for problem behaviors among nursing home residents. *J Gerontol Soc Work* 1999; 32(3):49-64.

Status: Not included because does not meet criteria for treatment for dementia patients

Innes EH. Efficacy and tolerance of flurbiprofen in the elderly using liquid and tablet formulations. *Curr Med Res Opin* 1977; 5(1):122-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Inzitari D, Pantoni L. Subcortical vascular dementia: therapeutical aspects. *Arch Gerontol Geriatr* 1998; 263-8.

Status : Not included because not a full article

Ioannidis JP, Lau J. Completeness of safety reporting in randomized trials: An evaluation of 7 medical areas. *JAMA* 2001 Jan 24; 285(4):437-43.

Status: Background article

Ioannidis JP, Lau J. Improving safety reporting from randomised trials. *Drug Saf* 2002; 25(2):77-84.

Status: Background article

Iribar MC, Montes J, Gonzalez MR, et al. Alanyl-aminopeptidase activity decrease in cerebrospinal fluid of Alzheimer patients. *Dement Geriatr Cogn Disord* 1998 Jan; 9(1):44-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Ishizaki J, Meguro K, Ohe K, et al. Therapeutic psychosocial intervention for elderly subjects with very mild Alzheimer disease in a community: The tajiri project. *Alzheimer Dis Assoc Disord* 2002; 16(4):261-9.

Status: Not included because dementia population not randomized to treatment

Israel L, Melac M, Milinkevitch D, et al. Drug therapy and memory training programs: a double-blind randomized trial of general practice patients with age-associated memory impairment. *Int Psychogeriatr* 1994; 6(2):155-70.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Israel L, Myslinski M, Kozarevic D. Nootropic treatment and combined therapy in age-associated memory impairment. Arch Gerontol Geriatr 1998; 27(Suppl 6):269-74.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Itil T, Martorano D. Natural substances in psychiatry (Ginkgo biloba in dementia). Psychopharmacol Bull 1995; 31(1):147-58.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Itil TM, Mukherjee S, Michael ST. Clinical and electrophysiological effects of suloctidil in elderly patients with multi-infarct dementia (A double-blind placebo-controlled study). Psychopharmacol Bull 1983; 19(4):730-3.
Status: Not included because Jadad Quality Scale score less than three

Itil TM, Eralp E, Ahmed I, et al. The pharmacological effects of Ginkgo biloba, a plant extract, on the brain of dementia patients in comparison with tacrine. Psychopharmacol Bull 1998; 34(3):391-7.
Status: Not included because dementia population not randomized to treatment

Izmirlian G, Brock D, White L. Estimating incidence of dementia subtypes: assessing the impact of missed cases. Stat Med 2000 Jun 15; 19(11-12):1577-91.
Status: Background article

Jacobs EA, Winter PM, Alvis HJ, et al. Hyperoxygenation effect on cognitive functioning in the aged. N Engl J Med 1969 Oct 2; 281(14):753-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jani J, Prettyman R. Use of a prescribing protocol in routine clinical practice: Experience following the introduction of donepezil. Psychiatr Bull 2001; 25(5):174-7.
Status: Not included because dementia population not randomized to treatment

Jann MW, Cyrus PA, Eisner LS, et al. Efficacy and safety of a loading-dose regimen versus a no-loading-dose regimen of metrifonate in the symptomatic treatment of Alzheimer's disease: A randomized, double-masked, placebo-controlled trial. Clin Ther 1999; 21(1):88-102.
Status: Included

Jann MW. Rivastigmine, a new-generation cholinesterase inhibitor for the treatment of Alzheimer's disease. Pharmacotherapy 2000; 20(1 I):1-12.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jansen W, Bruckner GW, Jansen P. The treatment of senile dementia associated with cerebrovascular insufficiency: a comparative study of buflomedil and dihydrogenated ergot alkaloids. J Int Med Res 1985; 13(1):48-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jansen W, O'Connolly M, Lehmann E, et al. Experimental clinical studies on the effect of eburnamonine in cerebrovascular disorders. Pharmacopsychiatry 1986 Sep; 19(5):389-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Janssen Pharmaceutica INC. Low dose risperidone helps overcome burden of Alzheimer's disease. Pharmaco economics and outcomes news 1997; 143:1.
Status: Article not retrievable

Jean-Louis G, von Gizycki H, Zizi F. Melatonin effects on sleep, mood, and cognition in elderly with mild cognitive impairment. J Pineal Res 1998 Oct; 25(3):177-83.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jellinger K, Flament H, Riederer P. Levodopa in the treatment of (pre) senile dementia. Mech Ageing Dev 1980; 14(1-2):253-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jenike MA, Albert MS, Heller H, et al. Combination therapy with lecithin and ergoloid mesylates for Alzheimer's disease. J Clin Psychiatry 1986 May; 47(5):249-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jenike MA, Albert M, Baer L, et al. Ergot mesylates for Alzheimer's disease: A year-long double-blind trial of 3 mg vs 12 mg daily. Int J Geriatr Psychiatry 1990; 5(6):375-80.
Status: Not included because Jadad Quality Scale score less than three

Jenike MA, Albert MS, Heller H, et al. Oral physostigmine treatment for patients with presenile and senile dementia of the Alzheimer's type: a double-blind placebo-controlled trial. *J Clin Psychiatry* 1990 Jan; 51(1):3-7.

Status: Not included because dementia population not randomized to treatment

Jenike MA, Albert M, Baer L, et al. Oral physostigmine as treatment for primary degenerative dementia: A double-blind placebo-controlled inpatient trial. *J Geriatr Psychiatry Neurol* 1990 Jan; 3(1):13-6.

Status: Article not retrievable

Jennekens-Schinkel A, Wintzen AR, Lanser JB. A clinical trial with desglycinamide arginine vasopressin for the treatment of memory disorders in man. *Prog Neuropsychopharmacol Biol Psychiatry* 1985; 9(3):273-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jeste DV, Lacro JP, Bailey A, et al. Lower incidence of tardive dyskinesia with risperidone compared with haloperidol in older patients. *J Am Geriatr Soc* 1999 Jun; 47(6):716-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jeste DV, Okamoto A, Napolitano J, et al. Low incidence of persistent tardive dyskinesia in elderly patients with dementia treated with risperidone. *Am J Psychiatry* 2000 Jul; 157(7):1150-5.

Status: Companion of an included article

Jhee SS, Shiovitz T, Hartman RD, et al. Centrally acting antiemetics mitigate nausea and vomiting in patients with Alzheimer's disease who receive rivastigmine. *Clin Neuropharmacol* 2002 Mar; 25(2):122-3.

Status: Not included because not a full article

Jobe JB, Smith DM, Ball K, et al. ACTIVE: a cognitive intervention trial to promote independence in older adults. *Control Clin Trials* 2001 Aug; 22(4):453-79.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Joffres C, Bucks RS, Haworth J, et al. Patterns of clinically detectable treatment effects with galantamine: A qualitative analysis. *Dement Geriatr Cogn Disord* 2003; 15(1):26-33.

Status: Background article

Johnson SA, Simmon VF. Randomized, double-blind, placebo-controlled international clinical trial of the AMPAKINE CX516 in elderly participants with mild cognitive impairment. A progress report. *J Mol Neurosci* 2002; 19(1-2):197-200.

Status: Not included because no extractable data relevant to review

Johnstone EC, Owens DG, Crow TJ, et al. Does a four-week delay in the introduction of medication alter the course of functional psychosis? *J Psychopharmacol (Oxf)* 1999; 13(3):238-44.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jolkkonen JT, Soininen HS, Riekkinen PJ. The effect of ACTH4-9 analog (Org2766) on some cerebrospinal fluid parameters in patients with Alzheimer's disease. *Life Sci* 1985 Aug 19; 37(7):585-90.

Status: Not included because does not meet criteria for treatment for dementia patients

Jones GM, Sahakian BJ, Levy R, et al. Effects of acute subcutaneous nicotine on attention, information processing and short-term memory in Alzheimer's disease. *Psychopharmacologia* 1992; 108(4):485-94.

Status: Not included because dementia population not randomized to treatment

Jonsson A, Korfitzen EM, Heltberg A, et al. Effects of neuropsychological treatment in patients with multiple sclerosis. *Acta Neurol Scand* 1993 Dec; 88(6):394-400.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jonsson L, Lindgren P, Wimo A, et al. The cost-effectiveness of donepezil therapy in Swedish patients with Alzheimer's disease: A Markov model. *Clin Ther* 1999; 21(7):1230-40.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jonsson L, Gerth W, Fastbom J. The potential economic consequences of cognitive improvement with losartan. *Blood Press* 2002; 11(1):46-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jorgenson S, Bille A, Andersen J, et al. Fluvoxamine treatment of dementia: Tryptophan levels. *Biol Psychiatry* 1993; 34(8):587-8.

Status: Not included because not a full article

Jorissen BL, Brouns F, Van Boxtel MP, et al. The influence of soy-derived phosphatidylserine on cognition in age-associated memory impairment. *Nutr Neurosci* 2001; 4(2):121-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jorissen BL, Brouns F, Van Boxtel MPJ, et al. Safety of soy-derived phosphatidylserine in elderly people. *Nutr Neurosci* 2002; 5(5):337-43.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jorm AF, Korten AE, Henderson AS. The prevalence of dementia: a quantitative integration of the literature. *Acta Psychiatr Scand* 1987 Nov; 76(5):465-79.

Status: Background article

Jotkowitz S. Lack of clinical efficacy of chronic oral physostigmine in Alzheimer's disease. *Ann Neurol* 1983 Dec; 14(6):690-1.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jönsson B, Jönsson L, Wimo A. Costs of dementia: A review. In: Maj M, Sartorius N, editors. *Dementia*, Volume 3. Chichester: John Wiley & Sons; 2000. Chapter 6 p. 335-63.

Status: Background article

Judge KS, Camp CJ, Orsulic-Jeras S. Use of Montessori-based activities for clients with dementia in adult day care: Effects on engagement. *Am J Alzheimers Dis* 2000; 15(1):42-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Judge TG, Urquhart A. Naftidrofuryl: a double blind cross-over study in the elderly. *Curr Med Res Opin* 1972; 1(3):166-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Juvars KA, McDonald C. A controlled trial of butyrylperazine ("Randolectil") in senile dementia. *Med J Aust* 1967 Feb 18; 1(7):334-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kahana E, Kahana B. Therapeutic potential of age integration. Effects of age-integrated hospital environments on elderly psychiatric patients. *Arch Gen Psychiatry* 1970 Jul; 23(1):20-9.

Status: Not included because no extractable data relevant to review

Kanamori M, Suzuki M, Yamamoto K, et al. A day care program and evaluation of animal-assisted therapy (AAT) for the elderly with senile dementia. *Am J Alzheimers Dis* 2001; 16(4):234-9.

Status: Not included because dementia population not randomized to treatment

Kanowski S, Fischhof PK, Grobe-Einsler R, et al. Efficacy of xantinolnicotinate in patients with dementia. *Pharmacopsychiatry* 1990 May; 23(3):118-24.

Status: Included

Kanowski S, Kinzler E, Lehmann E, et al. Confirmed clinical efficacy of Actovegin in elderly patients with organic brain syndrome. *Pharmacopsychiatry* 1995 Jul; 28(4):125-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kanowski S, Herrmann WM, Stephan K, et al. Proof of efficacy of the Ginkgo biloba special extract EGb 761 in outpatients suffering from mild to moderate primary degenerative dementia of the Alzheimer type or multi-infarct dementia. *Pharmacopsychiatry* 1996 Mar; 29(2):47-56.

Status: Included

Kapur N, Ironside J, Abbott P, et al. A neuropsychological-neuropathological case study of variant Creutzfeldt-Jakob disease. *Neurocase* 2001; 7(3):261-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kario K, Matsuo T, Hoshida S, et al. Effect of thrombin inhibition in vascular dementia and silent cerebrovascular disease. An MR spectroscopy study. *Stroke* 1999 May; 30(5):1033-7.

Status: Not included because dementia population not randomized to treatment

Karlsson I, Brane G, Melin E, et al. Effects of environmental stimulation on biochemical and psychological variables in dementia. *Acta Psychiatr Scand* 1988; 77(2):207-13.

Status: Not included because dementia population not randomized to treatment

Karlsson I, Godderis J, Augusto De Mendonca LC, et al. A randomised, double-blind comparison of the efficacy and safety of citalopram compared to mianserin in elderly, depressed patients with or

without mild to moderate dementia. *Int J Geriatr Psychiatry* 2000 Apr; 15(4):295-305.
Status: Included

Karlsson J, Hallgren P, Kral J, et al. Predictors and effects of long-term dieting on mental well-being and weight loss in obese women. *Appetite* 1994 Aug; 23(1):15-26.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Karoutas G, Milonas I, Artemis N, et al. The pharmacological effects of buflomedil in patients with multi-cerebral infarcts dementia: An open, preliminary assessment. *Curr Med Res Opin* 1986; Vol 10:380-9.
Status: Not included because dementia population not randomized to treatment

Kasckow JW, McElroy SL, Cameron RI. A pilot study on the use of divalproex sodium in the treatment of behavioral agitation in elderly patients with dementia: assessment with the BEHAVE-AD and CGI rating scales. *Current therapeutic research, clinical and experimental* 1997; 58(981):989.
Status: Not included because dementia population not randomized to treatment

Katona CL, Hunter BN, Bray J. A double-blind comparison of the efficacy and safety of paroxetine and imipramine in the treatment of depression with dementia. *Int J Geriatr Psychiatry* 1998 Feb; 13(2):100-8.
Status: Included

Katz IR, Jeste DV, Mintzer JE, et al. Comparison of risperidone and placebo for psychosis and behavioral disturbances associated with dementia: A randomized, double-blind trial. *J Clin Psychiatry* 1999; 60(2):107-15.
Status: Included

Katzman R, Zhang M, YChen PJ, et al. Effects of apolipoprotein E on dementia and aging in the Shanghai Survey of Dementia. *Neurology* 1997; 49(3):779-85.
Status: Not included because does not meet criteria for treatment for dementia patients

Kaufer DI. Pharmacologic therapy of dementia with Lewy bodies. *J Geriatr Psychiatry Neurol* 2002; 15(4):224-32.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kawas C, Resnick S, Morrison A, et al. A prospective study of estrogen replacement therapy and the risk of developing Alzheimer's disease: the Baltimore Longitudinal Study of Aging. *Neurology* 1997 Jun; 48(6):1517-21.
Status: Background article

Kawas CH, Brookmeyer R. Aging and the public health effects of dementia. *N Engl J Med* 2001 Apr 12; 344(15):1160-1.
Status: Background article

Kawas CH. Clinical practice. Early Alzheimer's disease. *N Engl J Med* 2003 Sep 11; 349(11):1056-63.
Status: Background article

Kaye WH, Sitaram N, Weingartner H, et al. Modest facilitation of memory in dementia with combined lecithin and anticholinesterase treatment. *Biol Psychiatry* 1982; 17(2):275-80.
Status: Cross-over trial; Cross-over trial

Kazui H, Mori E, Hashimoto M, et al. Impact of emotion on memory. Controlled study of the influence of emotionally charged material on declarative memory in Alzheimer's disease. *Br J Psychiatry* 2000; 177:343-7.
Status: Not included because does not meet criteria for treatment for dementia patients

Kempenaar L, McNamara C, Creaney B. Sensory stimulation work with carers in the community. *Journal of Dementia Care* 2001; 9(1):16.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kennedy JS, Zagar A, Bymaster F, et al. The central cholinergic system profile of olanzapine compared with placebo in Alzheimer's disease. *Int J Geriatr Psychiatry* 2001; 16(Suppl 1):S24-S32
Status: Companion of an included article

Kertesz A. Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: A randomized trial. *Curr Neurol Neurosci Rep* 2002 Nov; 2(6):503-4.
Status: Not included because not a full article

Khan A, Mirolo MH, Claypoole K, et al. Low-dose thyrotropin-releasing hormone effects in cognitively impaired alcoholics. *Alcohol Clin Exp Res* 1993 Aug; 17(4):791-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Khan A. Are placebo controls necessary to test new antidepressants and anxiolytics? *Int J Neuropsychopharmacol* 2002; 5(3):193-7.
Status: Background article

Khan KS, Daya S, Collins JA, et al. Empirical evidence of bias in infertility research: Overestimation of treatment effect in crossover trials using pregnancy as the outcome measure. *Fertil Steril* 1996 May; 65(5):939-45.
Status: Background article

Kiebertz K, Schifitto G, McDermott M, et al. Safety and tolerability of the antioxidant OPC-14117 in HIV-associated cognitive impairment. *Neurology* 1997; 49(1):142-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kiebertz K, Schifitto G, McDermott M, et al. A randomized, double-blind, placebo-controlled trial of deprenyl and thioctic acid in human immunodeficiency virus-associated cognitive impairment. *Neurology* 1998; 50(3):645-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kim JM, Shin IS, Yoon JS. Correlates of dropout, efficacy, and adverse events in treatment with acetylcholinesterase inhibitors in Korean patients with Alzheimer's disease. *Int Psychogeriatr* 2002; 14(2):187-95.
Status: Not included because Jadad Quality Scale score less than three

Kimura M, Robinson RG, Kosier JT. Treatment of cognitive impairment after poststroke depression: A double-blind treatment trial. *Stroke* 2000 Jul; 31(7):1482-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

King AC, Brassington G. Enhancing physical and psychological functioning in older family caregivers: The role of regular physical activity. *Ann Behav Med* 1997; (No. 2):91-100.
Status: Not included because does not meet criteria for treatment for dementia patients

Kinosian BP, Stallard E, Lee JH, et al. Predicting 10-year care requirements for older people with suspected Alzheimer's disease. *J Am Geriatr Soc*

2000 Jun; 48(6):631-8.
Status: Background article

Kirby M, Denihan A, Bruce I, et al. Benzodiazepine use among the elderly in the community. *Int J Geriatr Psychiatry* 1999; 14(4):280-4.
Status: Not included because dementia population not randomized to treatment

Kirchner V, Kelly CA, Harvey RJ. Thioridazine for dementia. In: *The Cochrane Library*, 2000. Issue 2. Oxford: Update Software.
Status: Background article

Kirrane RM, Mitropoulou V, Nunn M, et al. Physostigmine and cognition in schizotypal personality disorder. *Schizophr Res* 2001 Mar 1; 48(1):1-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kishimoto T, Hiraoka Y, Oribe H, et al. Auditory P300 event-related potentials and mini mental state examination performance in dementia; effects of l-debenone and vinpocetine. *J Nara Med Assoc* 1995; 46(3):259-66.
Status: Not included because dementia population not randomized to treatment

Kittner B. Using a combined randomized start/withdrawal design to assess propentofylline's effect on disease progression in Alzheimer's disease and vascular dementia. *J Neural Transm Gen Sect* 2000; 107:XV.
Status: Not included because does not meet criteria for treatment for dementia patients

Kittner B. Propentofylline for the treatment of vascular dementia: A 48-week, placebo-controlled study examining safety, efficacy, and impact on disease progression. *J Cereb Blood Flow Metab* 1999; 19(Suppl 1):16.
Status: Not included because not a full article

Kittner B. Using a combined randomized start/withdrawal design to assess propentofylline's effects on disease progression in Alzheimer's disease and vascular dementia: Results of clinical studies. *J Eur Coll Neuropsychopharmacol* 1999; 9(Suppl 5):S320
Status: Not included because not a full article

Kittner B. Clinical trials of propentofylline in vascular dementia. *European/Canadian Propentofylline Study Group. Alzheimer Dis*

Assoc Disord 1999; 13(Suppl 3):S166-S171.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kittner B, De Deyn PP, Erkinjuntti T. Investigating the natural course and treatment of vascular dementia and Alzheimer's disease. Parallel study populations in two randomized, placebo-controlled trials. Ann N Y Acad Sci 2000 Apr; 903:535-41.
Status: Not included because Jadad Quality Scale score less than three

Knapp MJ, Gracon SI, Davis CS, et al. Efficacy and safety of high-dose tacrine: A 30-week evaluation. Alzheimer Dis Assoc Disord 1994a; 8(Suppl 2):S22-S31
Status: Companion of an included article

Knapp MJ, Knopman DS, Solomon PR, et al. A 30-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. The Tacrine Study Group. JAMA 1994b; 271(13):985-91.
Status : Included

Knezevic S, Mubrin Z, Risberg J, et al. Pyritinol treatment of SDAT patients: evaluation by psychiatric and neurological examination, psychometric testing and rCBF measurements. Int Clin Psychopharmacol 1989 Jan; 4(1):25-38.
Status: Cross-over trial;
Cross-over trial

Knezevic S. European Pentoxifylline Multi-Infarct Dementia Study. Eur Neurol 1996; 36(5):315-21.
Status: Included

Knopman D. Long-term retention of implicitly acquired learning in patients with Alzheimer's disease. J Clin Exp Neuropsychol 1991; 13(6):880-94.
Status: Not included because does not meet criteria for treatment for dementia patients

Knopman D, Schneider L, Davis K, et al. Long-term tacrine (Cognex) treatment: Effects on nursing home placement and mortality. Neurology 1996 Jul; 47(1):166-77.
Status: Companion of an included article

Knopman D, Schneider L, Davis K, et al. Long-term tacrine treatment effects. Neurology 1998 Feb; 50(2):567-8.
Status: Not included because dementia population not randomized to treatment

Knopman D, Boland LL, Mosley T, et al. Cardiovascular risk factors and cognitive decline in middle-aged adults. Neurology 2001 Jan 9; 56(1):42-8.
Status: Background article

Knopman DS. Metrifonate for Alzheimer's disease: Is the next cholinesterase inhibitor better? Neurology 1998; 50(5):1203-5.
Status: Not included because dementia population not randomized to treatment

Knopman DS, DeKosky ST, Cummings JL, et al. Practice parameter: diagnosis of dementia (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. Neurology 2001 May 8; 56(9):1143-53.
Status: Background article

Knopman D, Gracon S. Observations on the short-term "natural history" of probable Alzheimer's disease in a controlled clinical trial. Neurology 1994; 44(2):260-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Knott V, Engeland C, Mohr E, et al. Acute nicotine administration in Alzheimer's disease: An exploratory EEG study. Neuropsychobiology 2000; 41(4):210-20.
Status: Not included because dementia population not randomized to treatment

Knott V, Mohr E, Mahoney C, et al. Pharmacologic EEG test dose response predicts cholinesterase inhibitor treatment outcome in Alzheimer's disease. Methods & Findings in Experimental & Clinical Pharmacology 2000 Mar; 22(2):115-22.
Status: Not included because dementia population not randomized to treatment

Knox J, Hindmarch I, Wallace M. Effects of twice standard dosage of neuroactive drugs in dementia. A preliminary report. Br J Clin Pract 1984 Sep; 38(9):313-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Koch GG, Davis SM, Anderson RL. Methodological advances and plans for improving regulatory success for confirmatory studies. Stat Med 1998 Aug 15; 17(15-16):1675-90.
Status: Background article

Kodjian A, Barriaga C, Turcot G. Double-blind study of pimozide in senile dementia patients. *Curr Ther Res Clin Exp* 1986; 40(4):694-701.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Koepp R, Miles SH. Meta-analysis of tacrine for Alzheimer disease: The influence of industry sponsors. *JAMA* 1999 Jun 23; 281(24):2287-8.
Status: Background article

Kofler B, Erhart C, Erhart P, et al. A multidimensional approach in testing nootropic drug effects (Cerebrolysin (R)). *Arch Gerontol Geriatr* 1990; 10(2):129-40.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kogan EA, Korczyn AD, Virchovsky RG, et al. EEG changes during long-term treatment with donepezil in Alzheimer's disease patients. *J Neural Transm Gen Sect* 2001; 108(10):1167-73.
Status: Not included because dementia population not randomized to treatment

Koh K, Ray R, Lee J, et al. Dementia in elderly patients: Can the 3R mental stimulation programme improve mental status? *Age Ageing* 1994; 23(3):195-9.
Status: Not included because dementia population not randomized to treatment

Koivisto, Portin M, Seinela R, et al. Automatic influences of memory in Alzheimer's disease. *Cortex* 1998; 34(2):209-19.
Status: Not included because does not meet criteria for treatment for dementia patients

Koivisto K, Helkala EL, Hanninen T, et al. Long-term selegiline treatment reduces the progression of Alzheimer's disease: Results after three years' follow-up. *Alzheimers Res* 1995; 1(Suppl 1):28.
Status: Not included because not a full article

Koltai DC, Welsh-Bohmer KA, Schmechel DE. Influence of anosognosia on treatment outcome among dementia patients. *Neuropsychol Rehab* 2001; 11(3-4):455-75.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Koltringer P, Langsteger W, Eber O. Dose-dependent hemorheological effects and microcirculatory modifications following intravenous administration of Ginkgo biloba special extract EGb 761. *Clin Hemorheol* 1995;

15(4):649-56.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kompoliti K, Goetz CG, Boeve BF, et al. Clinical presentation and pharmacological therapy in corticobasal degeneration. *Arch Neurol* 1998 Jul; 55(7):957-61.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kontush A, Mann U, Arlt S, et al. Influence of vitamin E and C supplementation on lipoprotein oxidation in patients with Alzheimer's disease. *Free Radic Biol Med* 2001 Aug 1; 31(3):345-54.
Status: Not included because dementia population not randomized to treatment

Kosten TR, Rosen MI, McMahon TL, et al. Treatment of early AIDS dementia in intravenous drug users: High versus low dose peptide T. *Am J Drug Alcohol Abuse* 1997 Nov; 23(4):543-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kountouris D. Therapeutic effects of piracetam combined with intravenous immunoglobulin premature of Alzheimer type. *J Neural Transm Gen Sect* 2000; 107:XVIII.
Status: Not included because not a full article

Kragh-Sorensen P, Olsen RB, Lund S, et al. Neuropeptides: ACTH-peptides in dementia. *Prog Neuropsychopharmacol Biol Psychiatry* 1986; 10(3-5):479-92.
Status: Included

Krebs Roubicek EM. Group therapy with demented elderly. *Alzheimers Dis Relat Disord* 1989; 1:1261-72.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kristensen V, Olsen M, Theilgaard A. Levodopa treatment of presenile dementia. *Acta Psychiatr Scand* 1977 Jan; 55(1):41-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kruck TPA, Fisher EA, McLachlan DRC. A predictor for side effects in patients with Alzheimer's disease treated with deferoxamine mesylate. *Clin Pharmacol Ther* 1993; 53(1):30-7.
Status: Not included because dementia population not randomized to treatment

Kruck TPA, Fisher EA, McLachlin DRC. Prediction of side effects in Alzheimer's disease patients on longterm deferoxamine-mesylate treatment. *Alzheimers Dis Relat Disord Adv Biosci* 1993; 87:257-8.

Status: Not included because no extractable data relevant to review

Krystio RJ, Schmitt FA. Foreword. *Stat Med* 2000; 19(11, 12):1389-91.

Status: Background article

Kuhl DE, Minoshima S, Frey KA, et al. Limited donepezil inhibition of acetylcholinesterase measured with positron emission tomography in living Alzheimer cerebral cortex. *Ann Neurol* 2000 Sep; 48(3):391-5.

Status: Not included because dementia population not randomized to treatment

Kuhn DR, de Leon CFM. Evaluating an educational intervention with relatives of persons in the early stages of Alzheimer's disease. *Res Soc Work Pract* 2001 Sep; 11(5):531-48.

Status: Not included because does not meet criteria for treatment for dementia patients

Kulisevsky J, Garcia-Sanchez C, Berthier ML, et al. Chronic effects of dopaminergic replacement on cognitive function in Parkinson's disease: A two-year follow-up study of previously untreated patients. *Mov Disord* 2000 Jul; 15(4):613-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kumar V, Smith RC, Sherman KA, et al. Cortisol responses to cholinergic drugs in Alzheimer's disease. *Int J Clin Pharmacol Ther Toxicol* ; 26(10):471-6.

Status: Not included because dementia population not randomized to treatment

Kumar V, Brecher M. Psychopharmacology of atypical antipsychotics and clinical outcomes in elderly patients. *J Clin Psychiatry* 1999; 60(Suppl 13):5-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kumar V, Anand R, Messina J, et al. An efficacy and safety analysis of Exelon in Alzheimer's disease patients with concurrent vascular risk factors. *Eur J Neurol* 2000 Mar; 7(2):159-69.

Status: Companion of an included article

Kuskowski MA, Morley G, Malone SM, et al. Hydergine treatment and psychophysiological measures in primary degenerative dementia. *J Geriatr Psychiatry Neurol* 1990 Jan; 3(1):41-7.

Status: Not included because dementia population not randomized to treatment

Kwiecinski H, Lusakowska A, Mieszkowski J. Improvement in concentration following treatment with Ginseng/Ginkgo biloba combination in patients with chronic cerebrovascular disorders: A double-blind, placebo-controlled study. *Eur J Clin Res* 1997; 9:59-67.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kwok T, Tang C, Woo J, et al. Randomized trial of the effect of supplementation on the cognitive function of older people with subnormal cobalamin levels. *Int J Geriatr Psychiatry* 1998 Sep; 13(9):611-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kyomen HH, Satlin A, Hennen J, et al. Estrogen therapy and aggressive behavior in elderly patients with moderate-to-severe dementia: Results from a short-term, randomized, double-blind trial. *Am J Geriatr Psychiatry* 1999; 7(4):339-48.

Status: Included

Kyomen HH, Hennen J, Gottlieb GL, et al. Estrogen therapy and noncognitive psychiatric signs and symptoms in elderly patients with dementia. *Am J Psychiatry* 2002; 159(7):1225-7.

Status: Companion of an included article

LaBarge E, Rosenman LS, Leavitt K, et al. Counseling Clients with Mild Senile Dementia of the Alzheimer's Type: A Pilot Study. *J Neurol Rehabil* 1988; 2(4):167-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LaBrecque DC, Goldberg RI. A double-blind study of pentylentetrazol combined with niacin in senile patients. *Curr Ther Res Clin Exp* ; 9(12):611-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lacomblez L, Chatellier L. Tetrahydroaminoacridine (THA) in Alzheimer's disease (AD): A double-blind pilot study. *Fundam Clin Pharmacol* 1989; 3(2):172.

Status: Not included because not a full article

Lacomblez L, Chatellier L. A multicenter trial of tetrahydroaminoacridine in senile dementia of the Alzheimer type. *Fundam Clin Pharmacol* 1990; 4(4):456.

Status: Not included because not a full article

Lamb HM, Faulds D. Metrifonate. *Drugs Aging* 1997; 11(6):490-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lanctot KL, Herrmann N, Van Reekum R, et al. Gender, aggression and serotonergic function are associated with response to sertraline for behavioral disturbances in Alzheimer's disease. *Int J Geriatr Psychiatry* 2002; 17(6):531-41.

*Status: Cross-over trial;
Cross-over trial*

Lanctot KL, Herrmann N, Yau KK, et al. Efficacy and safety of cholinesterase inhibitors in Alzheimer's disease: a meta-analysis. *CMAJ* 2003 Sep 16; 169(6):557-64.

Status: Background article

Landi F, Bernabei R, Russo A, et al. Predictors of rehabilitation outcomes in frail patients treated in a geriatric hospital. *J Am Geriatr Soc* 2002; 50(4):679-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lange KW, Robbins TW, Marsden CD, et al. L-dopa withdrawal in Parkinson's disease selectively impairs cognitive performance in tests sensitive to frontal lobe dysfunction. *Psychopharmacologia* 1992; 107(2-3):394-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lange KW, Paul GM, Naumann M, et al. Dopaminergic effects on cognitive performance in patients with Parkinson's disease. *J Neural Transm Suppl* 1995; 46:423-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Langlais PJ, Mair RG, Whalen PJ, et al. Memory effect of DL-threo-3,4-dihydroxyphenylserine (DOPS) in human Korsakoff's disease. *Psychopharmacologia* 1988; 95(2):250-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LaPorte DJ, Lahti AC, Koffel B, et al. Absence of ketamine effects on memory and other cognitive

functions in schizophrenia patients. *J Psychiatr Res* 1996 Sep; 30(5):321-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Larrieu S, Letenneur L, Orgogozo JM, et al. Incidence and outcome of mild cognitive impairment in a population-based prospective cohort. *Neurology* 2002 Nov 26; 59(10):1594-9.

Status: Background article

Lasagna L. Balancing risks versus benefits in drug therapy decisions. *Clin Ther* 1998; 20 Suppl C:C72-C79

Status: Background article

Lavie P, Aharon-Peretz J, Klein F, et al. Sleep quality in geriatric depressed patients: Comparison with elderly demented patients and normal controls and the effects of moclobemide. *Dementia* 1992; 3(5-6):360-6.

Status: Article not retrievable

Lawlor BA, Mellow AM, Sunderland T, et al. A pilot study of serotonergic system responsivity in Alzheimer's disease. *Psychopharmacol Bull* 1988; 24(1):127-9.

Status: Not included because Jadad Quality Scale score less than three

Lawlor BA, Sunderland T, Mellow AM, et al. Hyperresponsivity to the serotonin agonist m-chlorophenylpiperazine in Alzheimer's disease. A controlled study. *Arch Gen Psychiatry* 1989 Jun; 46(6):542-9.

*Status: Cross-over trial;
Cross-over trial*

Lawlor BA, Sunderland T, Mellow AM, et al. A pilot placebo-controlled study of chronic m-CPP administration in Alzheimer's disease. *Biol Psychiatry* 1991 Jul 15; 30(2):140-4.

*Status: Cross-over trial;
Cross-over trial*

Lawlor BA, Radcliffe J, Molchan SE, et al. A pilot placebo-controlled study of trazodone and buspirone in Alzheimer's disease. *Int J Geriatr Psychiatry* 1994; 9(1):55-9.

*Status: Cross-over trial;
Cross-over trial*

Lawlor BA, Aisen PS, Green C, et al. Selegiline in the treatment of behavioural disturbance in Alzheimer's disease. *Int J Geriatr Psychiatry* 1997 Mar; 12(3):319-22.

*Status: Cross-over trial;
Cross-over trial*

Laws SM, Clarnette RM, Taddei K, et al. APOE-epsilon4 and APOE-491A polymorphisms in individuals with subjective memory loss. *Mol Psychiatry* 2002; 7(7):768-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lawton MP, Van Haitsma K, Klapper J, et al. A stimulation-retreat special care unit for elders with dementing illness. *Int Psychogeriatr* 1998 Dec; 10(4):379-95.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lawton MP, Brody EM, Saperstein AB, et al. Respite services for caregivers: Research findings for service planning. *Home Health Care Serv Q* 1989; 10(1-2):5-32.

Status: Not included because does not meet criteria for treatment for dementia patients

Lawton MP, Brody EM, Saperstein AR. Controlled study of respite service for caregivers of Alzheimer's patients. *Gerontologist* 1989 Feb; (1):8-16.

Status: Not included because does not meet criteria for treatment for dementia patients

Lazar PA. Ginkgo is not a smart pill. *J Fam Pract* 2002 Nov; 51(11):912.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Le Bars PL, Katz MM, Berman N, et al. A placebo-controlled, double-blind, randomized trial of an extract of Ginkgo biloba for dementia. *JAMA* 1997; (16):1327-32.

Status: Included

Le Bars PL, Kieser M, Itil KZ. A 26-week analysis of a double-blind, placebo-controlled trial of the Ginkgo biloba extract EGb 761 in dementia. *Dement Geriatr Cogn Disord* 2000 Jul; 11(4):230-7.

Status: Companion of an included article

Le Bars PL, Velasco FM, Ferguson JM, et al. Influence of the severity of cognitive impairment on the effect of the Ginkgo biloba extract EGb 761 in Alzheimer's disease. *Neuropsychobiology* 2002; 45(1):19-26.

Status: Companion of an included article

Leber P. Guidelines for the clinical evaluation of anti-dementia drugs. 1st draft. Rockville,MD: US Food and Drug Administration; 1990.

Status: Background article

Leblhuber F, Neubauer C, Peichl M, et al. Age and sex differences of dehydroepiandrosterone sulfate (DHEAS) and cortisol (CRT) plasma levels in normal controls and Alzheimer's disease (AD). *Psychopharmacologia* 1993; 111(1):23-6.

Status: Not included because dementia population not randomized to treatment

Leblhuber F, Walli J, Widner B, et al. Homocysteine and B vitamins in dementia. *Am J Clin Nutr* 2001; 73(1):127-8.

Status: Not included because not a full article

Lebowitz BD, Pollock BG, Schneider LS. Estrogen in geriatric psychopharmacology. *Psychopharmacol Bull* 1997; 33(2):287-8.

Status: Not included because not a full article

Lechner H, Walzl M, Walzl B, et al. HELP application in multi-infarct dementia. *Journal of Stroke and Cerebrovascular Diseases* 1992; 2:228-31.

Status: Not included because Jadad Quality Scale score less than three

Leckman J, Ananth JV, Ban TA, et al. Pentylentetrazol in the treatment of geriatric patients with disturbed memory function. *J Clin Pharmacol New Drugs* 1971 Jul; 11(4):301-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lefebvre C, Clarke MJ. Identifying randomised trials. In: Egger M, Smith DG, Altman DG, editors. *Systematic Reviews in Health Care: Meta-analysis in context*, 2nd Edition London: BMJ Books; 2001. Chapter 4 p. 69-86.

Status: Background article

Lehmann HE, Ban TA, Saxena BM. Nicotinic acid, thioridazine, fluoxymesterone and their combinations in hospitalized geriatric patients: A systematic clinical study. *Can Psychiatr Assoc J* 1972 Aug; 17(4):315-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Leighman L, Stotsky BA, Cole JO. A controlled study of drugs in long-term geriatric psychiatric patients: A double-blind comparison of pentylentetrazol, papaverine, and niacin. *Arch*

Gen Psychiatry 1971 Sep; 25:284-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lerner V, Miodownik C, Kaptan A, et al. Vitamin B6 as add-on treatment in chronic schizophrenic and schizoaffective patients: a double-blind, placebo-controlled study. J Clin Psychiatry 2002 Jan; 63(1):54-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Leszek J, Inglot AD, Janusz M, et al. Colostrin proline-rich polypeptide complex from ovine colostrum - A long-term study of its efficacy in Alzheimer's disease. Med Sci Monit 2002; 8(10):I93-I96

Status: Not included because dementia population not randomized to treatment

Levin HS. Cognitive rehabilitation. Unproved but promising. Arch Neurol 1990 Feb; 47(2):223-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Levine FM, Spitalnik R, Dobos C. Caudate nucleus effects on geriatric senility: Effect of belladonna on learning and memory of geriatric patients. Percept Mot Skills 1973 Dec; 37(3):1003-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LeVine H, III, Scholten JD. Screening for pharmacologic inhibitors of amyloid fibril formation. Methods Enzymol 1999; 309:467-76.

Status: Not included because dementia population not randomized to treatment

Levinson B, Wright P, Barklem S. Effect of buflomedil on behaviour, memory, and intellectual capacity in patients with dementia. A placebo-controlled study. S Afr Med J 1985 Aug 31; 68(5):302-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Levkovitz Y, Bloch Y, Kaplan D, et al. Fluvoxamine for psychosis in Alzheimer's disease. J Nerv Ment Dis 2001 Feb; 189(2):126-9.

*Status: Cross-over trial;
Cross-over trial*

Levy A, Brandeis R, Treves TA, et al. Transdermal physostigmine in the treatment of Alzheimer's disease. Alzheimer Dis Assoc Disord

1994; 8(1):15-21.

Status: Not included because dementia population not randomized to treatment

Levy MA, Burgio LD, Sweet R, et al. A trial of buspirone for the control of disruptive behaviors in community-dwelling patients with dementia. Int J Geriatr Psychiatry 1994; 9(10):841-8.

Status: Not included because dementia population not randomized to treatment

Levy R, Little A, Chuaqui P, et al. Early results from double-blind, placebo controlled trial of high dose phosphatidylcholine in Alzheimer's disease. Lancet 1983 Apr 30; 1(8331):987-8.

Status: Not included because not a full article

Lewis C, Ballinger BR, Presly AS. Trial of levodopa in senile dementia. BMJ 1978 Mar 4; 1(6112):550.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Li C, Jiang Z, Wu D, et al. Clinical study on treating vascular dementia by muskiness injection in points. Int J Clin Acupunct 2002; 13(1):1-7.

Status: Not included because Jadad Quality Scale score less than three

Lim A, Tsuang D, Kukull W, et al. Clinico-neuropathological correlation of Alzheimer's disease in a community-based case series. J Am Geriatr Soc 1999 May; 47(5):564-9.

Status: Background article

Lincoln NB, Dent A, Harding J, et al. Evaluation of cognitive assessment and cognitive intervention for people with multiple sclerosis. J Neurol Neurosurg Psychiatry 2002; 72(1):93-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lindner K, Bedard M, William MD, et al. Changes in medication use and functional status of community-dwelling Alzheimer's patients after consultation at a memory clinic. Clin Gerontol 2001; 22(3-4):13-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Linn MW, Gurel L, Williford WO, et al. Nursing home care as an alternative to psychiatric hospitalization. A Veterans Administration cooperative study. Arch Gen Psychiatry 1985 Jun; 42(6):544-51.

Status: Not included because does not meet criteria for treatment for dementia patients

Lipinska B, Backman L. Encoding-retrieval interactions in mild Alzheimer's disease: The role of access to categorical information. *Brain Cogn* 1997 Jul; 34(2):274-86.

Status: Background article

Lipper S, Tuchman MM. Treatment of chronic post-traumatic organic brain syndrome with dextroamphetamine: First reported case. *J Nerv Ment Dis* 1976 May; 162(5):366-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lipton SA, Dafni U, Simpson D, et al. Double blind randomised placebo-controlled trial of the calcium channel antagonist Nimodipine for the neurological manifestations of acquired immunodeficiency syndrome, including dementia and painful neuropathy. *Ann Neurol* 1995; 38:347

Status: Not included because not a full article

Little A, Levy R, Chuaqui-Kidd P, et al. A double-blind, placebo controlled trial of high-dose lecithin in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 1985; 48(8):736-42.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Litvan I, Gomez C, Atack JR, et al. Physostigmine treatment of progressive supranuclear palsy. *Ann Neurol* 1989 Sep; 26(3):404-7.

*Status: Cross-over trial;
Cross-over trial*

Litvan I, Blesa R, Clark K, et al. Pharmacological evaluation of the cholinergic system in progressive supranuclear palsy. *Ann Neurol* 1994 Jul; 36(1):55-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Litvan I, Sirigu A, Toothman J, et al. What can preservation of autobiographic memory after muscarinic blockade tell us about the scopolamine model of dementia? *Neurology* 1995 Feb; 45(2):387-9.

Status: Not included because dementia population not randomized to treatment

Litvan I, FitzGibbon EJ. Can tropicamide eye drop response differentiate patients with progressive supranuclear palsy and Alzheimer's disease from

healthy control subjects? *Neurology* 1996; 47(5):1324-6.

Status: Not included because does not meet criteria for treatment for dementia patients

Litvan I, Phipps M, Pharr VL, et al. Randomized placebo-controlled trial of donepezil in patients with progressive supranuclear palsy. *Neurology* 2001 Aug 14; 57(3):467-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Liu X, Teresi JA, Waternaux C. Modelling the decline pattern in functional measures from a prevalent cohort study. *Stat Med* 2000 Jun 15; 19(11-12):1593-606.

Status: Background article

Livingston GA, Sax KB, McClenahan Z, et al. Acetyl-L-carnitine in dementia. *Int J Geriatr Psychiatry* 1991; 6(12):853-60.

Status: Included

Llorente AM, van Gorp WG, Stern MJ, et al. Long-term effects of high-dose zidovudine treatment on neuropsychological performance in mildly symptomatic HIV-positive patients: Results of a randomized, double-blind, placebo-controlled investigation. *J Int Neuropsychol Soc* 2001 Jan; 7(1):27-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lloyd-Evans S, Brocklehurst JC, Palmer MK. Piracetam in chronic brain failure. *Curr Med Res Opin* 1979; 6(5):351-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lofgren B, Nyberg L, Mattsson M, et al. Three years after in-patient stroke rehabilitation: A follow-up study. *Cardiovasc Dis* 1999 May; 9(3):163-70.

Status: Background article

Loher TJ, Krauss JK, Wielepp JP, et al. Pallidal deep brain stimulation in a parkinsonian patient with late-life dementia: Sustained benefit in motor symptoms but not in functional disability. *Eur Neurol* 2002; 47(2):122-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Londos E, Passant U, Gustafson L. Blood pressure and drug treatment in clinically diagnosed Lewy body dementia and Alzheimer's

disease. Arch Gerontol Geriatr 2000; 30(1):35-46.
Status: Not included because does not meet criteria for treatment for dementia patients

Lopez Arrieta JM, Birks J. Nimodipine for primary degenerative, mixed and vascular dementia. In: The Cochrane Library, 2000. Issue 2. Oxford: Update Software.
Status: Background article

Lopez Arrieta JM. Role of meta-analysis of clinical trials for Alzheimer's disease. Drug Dev Res 2002; 56(3):401-11.
Status: Background article

Lopez Arrieta JM, Schneider L. Metrifonate for Alzheimer's disease (Cochrane Protocol). In: The Cochrane Library, 2002. Issue 3. Oxford: Update Software.
Status: Background article

Lopez Arrieta JM, Rodriguez JL, Sanz F. Efficacy and safety of Nicotine on Alzheimer's disease patients. In: The Cochrane Library, 2003. Issue 1. Oxford: Update Software.
Status: Background article

Lopez OL, Becker JT, Wisniewski S, et al. Cholinesterase inhibitor treatment alters the natural history of Alzheimer's disease. J Neurol Neurosurg Psychiatry 2002 Mar; 72(3):310-4.
Status: Not included because dementia population not randomized to treatment

Lord TR, Garner JE. Effects of music on Alzheimer patients. Percept Mot Skills 1993 Apr; 76(2):451-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lott IT, Osann K, Doran E, et al. Down syndrome and Alzheimer disease: Response to donepezil. Arch Neurol 2002; 59(7):1133-6.
Status: Not included because dementia population not randomized to treatment

Lovett WC, Stokes DK, Taylor LB, et al. Management of behavioral symptoms in disturbed elderly patients: Comparison of trifluoperazine and haloperidol. J Clin Psychiatry 1987 Jun; 48(6):234-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lucca U, Tettamanti M, Forloni G, et al. Nonsteroidal antiinflammatory drug use in Alzheimer's disease. Biol Psychiatry 1994 Dec; 36(12):854-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lucca U, Lucchelli F, Alberoni M, et al. Reliability and correlation measures of cognitive function and behavioural scales in controlled clinical trial of eptastigmine in Alzheimer's disease patients. J Neurol 1995; 242:S106.
Status: Not included because not a full article

Lupien SJ, Wilkinson CW, Briere S, et al. Acute modulation of aged human memory by pharmacological manipulation of glucocorticoids. J Clin Endocrinol Metab 2002; 87(8):3798-807.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Luqman WA, Riesenber D, Oakley N, et al. Tacrine for Alzheimer's disease. JAMA 1994; (24):1896-8.
Status: Not included because not a full article

Lyford J. Statins reduce dementia risk by 70%. Curr Control Trials Cardiovasc Med 2000; 1(3):172.
Status: Not included because not a full article

Lyketsos CG, Lindell VL, Baker A, et al. A randomized, controlled trial of bright light therapy for agitated behaviors in dementia patients residing in long-term care. Int J Geriatr Psychiatry 1999 Jul; 14(7):520-5.
Status: Cross-over trial; Cross-over trial

Lyketsos CG, Sheppard JM, Steele CD, et al. Randomized, placebo-controlled, double-blind clinical trial of sertraline in the treatment of depression complicating Alzheimer's disease: Initial results from the Depression in Alzheimer's Disease study. Am J Psychiatry 2000 Oct; 157(10):1686-9.
Status: Included

MacGowan SH, Wilcock GK, Scott M. Effect of gender and apolipoprotein E genotype on response to anticholinesterase therapy in Alzheimer's disease. Int J Geriatr Psychiatry 1998 Sep; 13(9):625-30.
Status: Not included because dementia population not randomized to treatment

MacMahon S, Kermode S. A clinical trial of the effect of aromatherapy on motivational behaviour in a dementia care setting using a single subject design. *Aust J Holist Nurs* 1998 Oct; 5(2):47-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Madhusoodanan S, Brenner R, Cohen CI. Role of atypical antipsychotics in the treatment of psychosis and agitation associated with dementia. *CNS Drugs* 1999; 12(2):135-50.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Maestri NE, Brusilow SW, Clissold DB, et al. Long-term treatment of girls with ornithine transcarbamylase deficiency. *N Engl J Med* 1996 Sep 19; 335(12):855-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Magai C, Kennedy G, Cohen CI, et al. A controlled clinical trial of sertraline in the treatment of depression in nursing home patients with late-stage Alzheimer's disease. *Am J Geriatr Psychiatry* 2000; 8(1):66-74.
Status: Included

Magai C, Cohen CI, Gomberg D. Impact of training dementia caregivers in sensitivity to nonverbal emotion signals. *Int Psychogeriatr* 2002 Mar; 14(1):25-38.
Status: Not included because does not meet criteria for treatment for dementia patients

Magnus RV. A controlled trial of chlormethiazole in the management of symptoms of the organic dementias in the elderly. *Clin Ther* 1978; 1(6):387-96.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Maidment R, Livingston G, Katona C. 'Just keep taking the tablets': Adherence to antidepressant treatment in older people in primary care. *Int J Geriatr Psychiatry* 2002; 17(8):752-7.
Status: Not included because does not meet criteria for treatment for dementia patients

Maina G, Fiori L, Torta R, et al. Oxiracetam in the treatment of primary degenerative and multi-infarct dementia: A double-blind, placebo-controlled study. *Neuropsychobiology* 1989; 21(3):141-5.
Status: Included

Mair RG, McEntee WJ. Cognitive enhancement in Korsakoff's psychosis by clonidine: A comparison with L-dopa and ephedrine. *Psychopharmacologia* 1986; 88(3):374-80.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Malaguarnera M, Pistone G, Vinci M, et al. Tacrine treatment of Alzheimer's disease: Many expectations, few certainties. *Neuropsychobiology* 1998 Nov; 38(4):226-31.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Malaret B. Use of music to decrease aggressive behaviours in people with dementia. *J Am Geriatr Soc* 1998; 46(12):1586.
Status: Not included because not a full article

Malsch U, Dennler. Treatment of Alzheimer's disease: Tolerability of rivastigmine during the initial period. *Psycho* 2001; (6):337-42.
Status: Article not retrievable

Maltby N, Broe GA, Creasey H, et al. Efficacy of tacrine and lecithin in mild to moderate Alzheimer's disease: Double blind trial. *BMJ* 1994 Apr 2; 308(6933):879-83.
Status: Included

Mancini GB, Schulzer M. Reporting risks and benefits of therapy by use of the concepts of unqualified success and unmitigated failure: applications to highly cited trials in cardiovascular medicine. *Circulation* 1999 Jan 26; 99(3):377-83.
Status: Background article

Manes F, Jorge R, Morcuende M, et al. A controlled study of repetitive transcranial magnetic stimulation as a treatment of depression in the elderly. *Int Psychogeriatr* 2001; 13(2):225-31.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mangoni A, Perin C, Smirne S, et al. A double-blind, placebo-controlled study with oxiracetam in demented patients administered the Luria-Nebraska Neuropsychological Battery. *Drug Dev Res* 1988; 14(3-4):217-4.
Status: Included

Mangoni A, Grassi MP, Frattola L, et al. Effects of a MAO-B inhibitor in the treatment of Alzheimer disease. *Eur Neurol* 1991; 31(2):100-7.
Status: Included

Mann AH, Schneider J, Mozley CG, et al. Depression and the response of residential homes to physical health needs. *Int J Geriatr Psychiatry* 2000; 15(12):1105-12.

Status: Not included because does not meet criteria for treatment for dementia patients

Mann K, Gunther A, Stetter F, et al. Rapid recovery from cognitive deficits in abstinent alcoholics: A controlled test-retest study. *Alcohol* 1999 Jul; 34(4):567-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mantero MA, Barbero M, Giannini R, et al. Acetyl-L-carnitine as a therapeutic agent for mental deterioration in geriatric patients. (Double-blind placebo controlled study). *New Trends in Clinical Neuropharmacology* 1989; (1):17-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marcusson J, Rother M, Kittner B, et al. A 12-month, randomized placebo-controlled trial of propentofylline (HWA 285) in patients with dementia according to DSM III-R. *Dement Geriatr Cogn Disord* 1997; 8(5):320-8.

Status: Included

Marder K, Tang MX, Alfaro B, et al. Postmenopausal estrogen use and Parkinson's disease with and without dementia. *Neurology* 1998 Apr; 50(4):1141-3.

Status: Not included because does not meet criteria for treatment for dementia patients

Marigliano V, Abate G, Barbagallo-Sangiorgi G, et al. Randomized, double-blind, placebo controlled, multicentre study of idebenone in patients suffering from multi-infarct dementia. *Arch Gerontol Geriatr* 1992; 15(3):239-48.

Status: Included

Marin DB, Bierer LM, Lawlor BA, et al. L-deprenyl and physostigmine for the treatment of Alzheimer's disease. *Psychiatry Res* 1995 Oct 16; 58(3):181-9.

*Status: Cross-over trial;
Cross-over trial*

Marini G, Caratti C, Peluffo F, et al. Placebo-controlled double-blind study of pramiracetam (CI-879) in the treatment of elderly subjects with memory impairment. *Adv Ther* 1992; 9(3):136-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marra CM, Boutin P, McArthur JC, et al. A pilot study evaluating ceftriaxone and penicillin G as treatment agents for neurosyphilis in human immunodeficiency virus-infected individuals. *Clin Infect Dis* 2000 Mar; 30(3):540-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marriott A, Donaldson C, Tarrier N, et al. Effectiveness of cognitive-behavioural family intervention in reducing the burden of care in carers of patients with Alzheimer's disease. *Br J Psychiatry* 2000 Jun; 176:557-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Marsh L, Lyketsos C, Reich SG. Olanzapine for the treatment of psychosis in patients with Parkinson's disease and dementia. *Psychosomatics* 2001 Nov; 42(6):477-81.

Status: Not included because dementia population not randomized to treatment

Martignoni E, Bono G, Blandini F, et al. Monoamines and related metabolite levels in the cerebrospinal fluid of patients with dementia of Alzheimer type. Influence of treatment with L-deprenyl. *J Neural Transm Park Dis Dement Sect* 1991; 3(1):15-25.

Status: Not included because dementia population not randomized to treatment

Martin JC. Effect of a synthetic peptide, ORG 2766, on inpatients with severe senile dementia. *Acta Psychiatr Scand* 1983; 67(3):205-7.

*Status: Cross-over trial;
Cross-over trial*

Martin PR, Adinoff B, Eckardt MJ, et al. Effective pharmacotherapy of alcoholic amnesic disorder with fluvoxamine. Preliminary findings. *Arch Gen Psychiatry* 1989 Jul; 46(7):617-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Martin PR, Adinoff B, Lane E, et al. Fluvoxamine treatment of alcoholic amnesic disorder. *Eur Neuropsychopharmacol* 1995 Mar; 5(1):27-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Martin RM, Rink E, Wilkinson DG, et al. Did knowledge, opinions, background, and health authority advice influence early prescribing of the novel Alzheimer's disease drug donepezil in general practice? - National postal survey.

Pharmacoepidemiol Drug Saf 1999; 8(6):413-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Martinez Perez JA, Chavida GF, Sanchez-Seco HP, et al. Epidemiology of cognitive impairment in Spain. Eur J Gen Pract 2000; 6(2):52-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marx J. Alzheimer's congress. Drug shows promise for advanced disease. Science 2000 Jul 21; 289(5478):375-7.
Status: Not included because not a full article

Marx JL. Alzheimer's drug trial put on hold. Science 1987; 238(4830):1041-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Masaki KH, Losonczy KG, Izmirlian G, et al. Association of vitamin E and C supplement use with cognitive function and dementia in elderly men. Neurology 2000; 54(6):1265-72.
Status: Not included because dementia population not randomized to treatment

Masuda Y, Akagawa Y, Hishikawa Y. Effect of serotonin 1A agonist tandospirone on depression symptoms in senile patients with dementia. Hum Psychopharmacol 2002 Jun; 17(4):191-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Matthews HP, Korbey J, Wilkinson DG, et al. Donepezil in Alzheimer's disease: Eighteen month results from Southampton Memory Clinic. Int J Geriatr Psychiatry 2000 Aug; 15(8):713-20.
Status: Not included because dementia population not randomized to treatment

Maurer I, Moller HJ, Saletu B. Treatment with propentofylline in dementia. Pharmacopsychiatry 1993; 26:179.
Status: Not included because not a full article

Maurer K, Ihl R, Dierks T, et al. Clinical efficacy of Ginkgo biloba special extract EGb 761 in dementia of the Alzheimer type. J Psychiatr Res 1997 Nov; 31(6):645-55.
Status: Included

Maxwell CJ, Hogan DB, Ebly EM. Calcium-channel blockers and cognitive function in elderly people: results from the Canadian Study of Health

and Aging.[comment][erratum appears in CMAJ 1999 Nov 30;161(11):1396]. CMAJ 1999 Sep 7; 161(5):501-6.
Status: Background article

Mayeux R, Saunders AM, Shea S, et al. Utility of the apolipoprotein E genotype in the diagnosis of Alzheimer's disease. Alzheimer's Disease Centers Consortium on Apolipoprotein E and Alzheimer's Disease. N Engl J Med 1998 Feb 19; 338(8):506-11.
Status: Background article

Mayuex R, Sano M. Drug therapy: Treatment of Alzheimer's disease. N Engl J Med 1999; 341(22):1670-9.
Status: Background article

McCaffrey RJ, Steckler RA, Gansler DA, et al. An experimental evaluation of the efficacy of suloctidil in the treatment of primary degenerative dementia. Arch Clin Neuropsychol 1987; (2):155-61.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McCallion P, Toseland RW, Freeman K. An evaluation of a family visit education program. J Am Geriatr Soc 1999 Feb; 47(2):203-14.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McCarten JR, Kovera C, Maddox MK, et al. Triazolam in Alzheimer's disease: Pilot study on sleep and memory effects. Pharmacol Biochem Behav ; 52(2):447-52.
Status: Not included because dementia population not randomized to treatment

McClendon MJ, Bass DM, Brennan PF, et al. A computer network for Alzheimer's caregivers and use of support group services. J Ment Health Aging 1998; (4):403-20.
Status: Not included because dementia population not randomized to treatment

McConnachie RW. A clinical trial comparing 'Hydergine' with placebo in the treatment of cerebrovascular insufficiency in elderly patients. Curr Med Res Opin 1973; 1(8):463-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McConnachie RW. The clinical assessment of brain failure in the elderly. Pharmacology 1978; Vol 16(Suppl 1):27-35.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McCusker J, Verdon J, Tousignant P, et al. Rapid emergency department intervention for older people reduces risk of functional decline: Results of a multicenter randomized trial. *J Am Geriatr Soc* 2001; 49(10):1272-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McDermott MP, Hall WJ, Oakes D, et al. Design and analysis of two-period studies of potentially disease-modifying treatments. *Control Clin Trials* 2002 Dec; 23(6):635-49.

Status: Background article

McDonald C, Mowbray RM, Wilson JM. A sequential trial of amylodbarbitone sodium used as sedation for confused female psychogeriatric patients. *Gerontol Clin (Basel)* 1970; 12(6):335-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McEntee WJ, Crook TH, Jenkyn LR, et al. Treatment of age-associated memory impairment with guanfacine. *Psychopharmacol Bull* 1991; 27(1):41-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McGeer PL, McGeer EG. Anti-inflammatory drugs in the fight against Alzheimer's disease. *Ann N Y Acad Sci* 1996; 777:213-20.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McKeith I, Del Ser T, Anand Rao. Rivastigmine provides symptomatic benefit in dementia with lewy bodies: Findings from a placebo-controlled international multicenter study. *Neurology* 2000; 54(Suppl 3):A450.
Status: Not included because not a full article

McKeith I, Del Ser T, Spano P, et al. Efficacy of rivastigmine in dementia with Lewy bodies: A randomised, double-blind, placebo-controlled international study. *Lancet* 2000 Dec 16; 356(9247):2031-6.
Status: Included

McKeith I, Del Ser T, Anand R. Erratum: Rivastigmine provides symptomatic benefit in dementia with lewy bodies: Findings from a placebo-controlled international multicenter study.

Expert Opin Pharmacother 2001; 2(5):907.
Status: Not included because not a full article

McKeith IG. The clinical trial protocol of the Metrifonate in Alzheimer's Trial (MALT). *Dement Geriatr Cogn Disord* 1998; Vol 9(Suppl 2):2-7.
Status: Companion of an included article

McKeith IG, Grace JB, Walker Z, et al. Rivastigmine in the treatment of dementia with Lewy bodies: Preliminary findings from an open trial. *Int J Geriatr Psychiatry* 2000 May; 15(5):387-92.
Status: Not included because dementia population not randomized to treatment

McKhann G, Drachman D, Folstein M, et al. Clinical diagnosis of Alzheimer's disease: Report of the NINCDS-ADRDA Work Group under the auspices of Department of Health and Human Services Task Force on Alzheimer's Disease. *Neurology* 1984 Jul; 34(7):939-44.
Status: Background article

McLachlan DR, Smith WL, Kruck TP. Desferrioxamine and Alzheimer's disease: Video home behavior assessment of clinical course and measures of brain aluminum. *Ther Drug Monit* 1993 Dec; 15(6):602-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McLean A, Jr., Stanton KM, Cardenas DD, et al. Memory training combined with the use of oral physostigmine. *Brain Inj* 1987 Oct; 1(2):145-59.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McLean A, Jr., Cardenas DD, Burgess D, et al. Placebo-controlled study of pramiracetam in young males with memory and cognitive problems resulting from head injury and anoxia. *Brain Inj* 1991 Oct; 5(4):375-80.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McLennan J, Findlay DJ, Sharma J, et al. Prolactin response to withdrawal of thioridazine in dementia. *Int J Geriatr Psychiatry* 1992; 7(10):739-42.
Status: Not included because Jadad Quality Scale score less than three

McNamara MJ, Gomez-Isla T, Hyman BT. Apolipoprotein E genotype and deposits of Abeta40 and Abeta42 in Alzheimer disease. *Arch*

Neurol 1998 Jul; 55(7):1001-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McNeil JK. Neuropsychological characteristics of the dementia syndrome of depression: Onset, resolution, and three-year follow-up. Clin Neuropsychol 1999; 13(2):136-46.

Status: Not included because does not meet criteria for treatment for dementia patients

Mcpherson A, Furniss FG, Sdogati C, et al. Effects of individualized memory aids on the conversation of persons with severe dementia a pilot study. Aging Ment Health 2001; 5(3):289-94.
Status: Not included because dementia population not randomized to treatment

McRae T, Griesing T, Whalen E. Donepezil and sertraline for the management of behavioral symptoms in patients with Alzheimer's disease. Neurology 2000; 54(Suppl 3):A416.
Status: Not included because not a full article

Mead MG, Castleden CM. Confusion and hypnotics in demented patients. J R Coll Gen Pract 1982 Dec; 32(245):763-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meador K, Loring D, Nichols M, et al. Preliminary findings of high-dose thiamine in dementia of Alzheimer's type. J Geriatr Psychiatry Neurol 1993 Oct; 6(4):222-9.
Status: Cross-over trial;
Cross-over trial

Mecocci P, Grossi E, Buscema M, et al. Use of artificial networks in clinical trials: A pilot study to predict responsiveness to donepezil in Alzheimer's disease. J Am Geriatr Soc 2002 Nov; 50(11):1857-60.
Status: Not included because dementia population not randomized to treatment

Medina A, Bodick N, Goldberger AL, et al. Effects of central muscarinic-1 receptor stimulation on blood pressure regulation. Hypertension 1997 Mar; 29(3):828-34.
Status: Not included because dementia population not randomized to treatment

Meehan KM, Wang H, David SR, et al. Comparison of rapidly acting intramuscular olanzapine, lorazepam, and placebo: A double-blind, randomized study in acutely agitated

patients with dementia.

Neuropsychopharmacology 2002 Apr; 26(4):494-504.

Status: Included

Mega MS, Dinov ID, Lee L, et al. Orbital and dorsolateral frontal perfusion defect associated with behavioral response to cholinesterase inhibitor therapy in Alzheimer's disease. J Neuropsychiatry Clin Neurosci 2000; 12(2):209-18.

Status: Not included because dementia population not randomized to treatment

Mega MS, Cummings JL, O'Connor SM, et al. Cognitive and metabolic responses to metrifonate therapy in Alzheimer disease. Neuropsychiatry Neuropsychol Behav Neurol 2001 Jan; 14(1):63-8.
Status: Not included because dementia population not randomized to treatment

Meier DE, Ahronheim JC, Morris J, et al. High short-term mortality in hospitalized patients with advanced dementia: Lack of benefit of tube feeding. Arch Intern Med 2001; 161(4):594-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meier U, Kintzel D. Clinical experiences with different valve systems in patients with normal-pressure hydrocephalus: Evaluation of the Miethke dual-switch valve. Childs Nerv Syst 2002 Jul; 18(6-7):288-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Melin L, Gotestam KG. Effects of rearranging Ward routines on communication and eating behaviors of psychogeriatric patients. J Appl Behav Anal 1981; (No. 1):47-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mellow AM, Sunderland T, Cohen RM, et al. Acute effects of high-dose thyrotropin releasing hormone infusions in Alzheimer's disease. Psychopharmacologia 1989; 98(3):403-7.
Status: Cross-over trial;
Cross-over trial

Mendez MF, Perryman KM. Neuropsychiatric features of frontotemporal dementia: Evaluation of consensus criteria and review. J Neuropsychiatry Clin Neurosci 2002; 14(4):424-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mendiondo MS, Ashford JW, Kryscio RJ, et al. Modelling mini mental state examination changes in Alzheimer's disease. *Stat Med* 2000 Jun 15; 19(11-12):1607-16.

Status: Background article

Merrick BJ. A geriatric care management approach to a treatment plan for dementia. *J Geriatr Psychiatry* 2001; 34(2):233-45.

Status: Not included because dementia population not randomized to treatment

Mervis RJ, Ganzell S, Fitten LJ, et al. Comparison of Carbamazepine and Trazodone in the control of aggression / agitation in demented institutionalized patients. *J Am Geriatr Soc* 1991; 39(8):A75

Status: Not included because not a full article

Meszaros Z, Borcsiczky D, Mate M, et al. Platelet MAO-B activity and serotonin content in patients with dementia: Effect of age, medication, and disease. *Neurochem Res* 1998 Jun; 23(6):863-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Meyer JS, Rogers RL, McClintic K, et al. Controlled clinical trial of daily aspirin therapy in multi-infarct dementia. *Stroke* 1988; 19(1):148

Status: Not included because not a full article

Meyer JS, Rogers RL, McClintic K, et al. Randomized clinical trial of daily aspirin therapy in multi-infarct dementia. A pilot study. *J Am Geriatr Soc* 1989 Jun; 37(6):549-55.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meyer JS, Lotfi J, Martinez G, et al. Effects of medical and surgical treatment on cerebral perfusion and cognition in patients with chronic cerebral ischemia. *Surg Neurol* 1990 Nov; 34(5):301-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meyer JS, Li Y, Xu G, et al. Feasibility of treating mild cognitive impairment with cholinesterase inhibitors. *Int J Geriatr Psychiatry* 2002 Jun; 17(6):586-8.

Status: Not included because dementia population not randomized to treatment

Meythaler JM, Depalma L, Devivo MJ, et al. Sertraline to improve arousal and alertness in severe traumatic brain injury secondary to motor

vehicle crashes. *Brain Inj* 2001 Apr; 15(4):321-31.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Micca JL, Sky AJ, Uhrig-Hitchcock LG. Quality care: A practical guide to managing behavioral symptoms of dementia. *Journal of the American Medical Directors Association* 2002; 3(Suppl 4):H21-H25

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miceli G, Caltagirone C, Gainotti G. Gangliosides in the treatment of mental deterioration. A double-blind comparison with placebo. *Acta Psychiatr Scand* 1977 Feb; 55(2):102-10.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mielke R, Kittner B, Ghaemi M, et al. Propentofylline improves regional cerebral glucose metabolism and neuropsychologic performance in vascular dementia. *JNS* 1996 Sep 15; 141(1-2):59-2.

Status: Included

Mielke R, Moller HJ, Erkinjuntti T, et al. Propentofylline in the Treatment of Vascular Dementia and Alzheimer-Type Dementia: Overview of Phase I and Phase II Clinical Trials. *Alzheimer Dis Assoc Disord* 1998; 12:S29-S35.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mielke R, Ghaemi M, Kessler J, et al. Propentofylline enhances cerebral metabolic response to auditory memory stimulation in Alzheimer's disease. *JNS* 1998 Jan 21; 154(1):76-82.

Status: Included

Miguel-Hidalgo JJ. Rivastigmine Novartis AG. *Curr Opin Cent Peripher Nerv Syst Invest Drugs* 2000; 2(4):438-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Milders M, Deelman B, Berg I. Rehabilitation of memory for people's names. *Memory* 1998 Jan; 6(1):21-36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miller E. On the nature of the memory disorder in presenile dementia. *Neuropsychologia* 1971; 9(1):75-81.

Status: Not included because does not meet criteria for treatment for dementia patients

Miller E. Efficiency of coding and the short-term memory defect in presenile dementia. *Neuropsychologia* 1972; 10(1):133-6.

Status: Not included because does not meet criteria for treatment for dementia patients

Miller IW, Norman WH, Keitner GI. Treatment response of high cognitive dysfunction depressed inpatients. *Compr Psychiatry* 1990 Jan; 31(1):62-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miller PA, Butin D. The role of occupational therapy in dementia - C.O.P.E. (Caregiver Options for Practical Experiences). *Int J Geriatr Psychiatry* 2000; 15(1):86-9.

Status: Not included because dementia population not randomized to treatment

Miller R, Newcomer R, Fox P. Effects of the Medicare Alzheimer's disease demonstration on nursing home entry. *Health Serv Res* 1999 Aug; 34(3):691-714.

Status: Not included because does not meet criteria for treatment for dementia patients

Miller RG. Leptazol and meso-inositol hexanicotinate in the treatment of chronic cerebrovascular degenerative disorders. A double blind study. *Gerontol Clin (Basel)* 1995 Oct; 5:95-102.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miller TP, Fong K, Tinklenberg JR. An ACTH 4-9 analog (Org 2766) and cognitive performance: High-dose efficacy and safety in dementia of the Alzheimer's type. *Biol Psychiatry* 1993 Feb 15; 33(4):307-9.

Status: Not included because no extractable data relevant to review

Milliken JK, Edland SD. Mixed effect models of longitudinal Alzheimer's disease data: a cautionary note. *Stat Med* 2000 Jun 15; 19(11-12):1617-29.

Status: Background article

Minger SL, Esiri MM, McDonald B, et al. Cholinergic deficits contribute to behavioral disturbance in patients with dementia. *Neurology* 2000; 55(10):1460-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Minthon L, Gustafson L, Dalfelt G, et al. Oral tetrahydroaminoacridine treatment of Alzheimer's disease evaluated clinically and by regional cerebral blood flow and EEG. *Dementia* 1993 Jan; 4(1):32-42.

*Status: Cross-over trial;
Cross-over trial*

Minthon L, Edvinsson L, Gustafson L. Tacrine treatment modifies cerebrospinal fluid neuropeptide levels in Alzheimer's disease. *Dementia* 1994 Nov; 5(6):295-301.

Status: Not included because no extractable data relevant to review

Minthon L, Nilsson K, Edvinsson L, et al. Long-term effects of tacrine on regional cerebral blood flow changes in Alzheimer's disease. *Dementia* 1995 Sep; 6(5):245-51.

Status: Not included because dementia population not randomized to treatment

Mintzer J, Faison W, Street JS, et al. Olanzapine in the treatment of anxiety symptoms due to Alzheimer's disease: A post hoc analysis. *Int J Geriatr Psychiatry* 2001; 16 Suppl 1:S71-S77

Status: Companion of an included article

Mintzer JE, Brawman MO, Mirski DF, et al. Anxiety in the behavioral and psychological symptoms of dementia. *Int Psychogeriatr* 2000; 12(Suppl 1):139-42.

Status: Not included because dementia population not randomized to treatment

Mintzer JE, Madhusoodanan S, Brenner Ronald. Risperidone in dementia. *Psychiatr Ann* 2000; 30(3):181-7.

Status: Not included because dementia population not randomized to treatment

Mishima K, Okawa M, Hishikawa Y, et al. Morning bright light therapy for sleep and behavior disorders in elderly patients with dementia. *Acta Psychiatr Scand* 1994; 89(1):1-7.

Status: Not included because dementia population not randomized to treatment

Mishima K, Hishikawa Y, Okawa M. Randomized, dim light controlled, crossover test of morning bright light therapy for rest-activity rhythm disorders in patients with vascular dementia and dementia of Alzheimer's type. *Chronobiol Int*

1998 Nov; 15(6):647-54.

*Status: Cross-over trial;
Cross-over trial*

Mitchell A, Drachman DA, O'Connell B, et al. Oral physostigmine in Alzheimer's disease. *Neurology* 1986; 36(Suppl 1):295.

Status: Not included because not a full article

Mitchell A, Maercklein LA. The effect of individualized special instruction on the behaviors of nursing home residents diagnosed with dementia. *Am J Alzheimers Dis* 1996; (1):23
Status: Not included because does not meet criteria for treatment for dementia patients

Mitchell S. Aromatherapy's Effectiveness in Disorders Associated With Dementia. *Int J Aromatherapy* 1993; (2):20-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mitnitski AB, Graham JE, Mogilner AJ, et al. Vector diagnostics in dementia derived from Bayes' theorem. *Am J Epidemiol* 1997 Oct 15; 146(8):665-71.
Status: Background article

Mittelman MS, Ferris SH, Steinberg G, et al. An intervention that delays institutionalization of Alzheimer's disease patients: Treatment of spouse-caregivers. *Gerontologist* 1993 Dec; 33(6):730-40.
Status: Not included because does not meet criteria for treatment for dementia patients

Mittelman MS, Ferris SH, Shulman E, et al. A family intervention to delay nursing home placement of patients with Alzheimer disease. A randomized controlled trial. *JAMA* 1996 Dec 4; 276(21):1725-31.
Status: Not included because does not meet criteria for treatment for dementia patients

Miyamoto Y, Ito H, Otsuka T, et al. Caregiver burden in mobile and non-mobile demented patients: A comparative study. *Int J Geriatr Psychiatry* 2002 Aug; 17(8):765-73.
Status: Not included because does not meet criteria for treatment for dementia patients

Mobius HJ, Stoffler A. New approaches to clinical trials in vascular dementia: Memantine in small vessel disease. *Cardiovasc Dis* 2002; 13 Suppl 2:61-6.
Status: Background article

Moffoot A, O'Carroll RE, Murray C, et al. Clonidine infusion increases uptake of 99mTc-Exametazime in anterior cingulate cortex in Korsakoff's psychosis. *Psychol Med* 1994 Feb; 24(1):53-61.

Status: Not included because dementia population not randomized to treatment

Moglia A, Arrigo A, Bono G. Citicoline in patients with chronic cerebrovascular diseases (CCVD): Quantitative EEG study. *Curr Ther Res Clin Exp* 1984; 36(2):309-13.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohide EA. A cognitive behavioural family intervention reduced psychiatric morbidity in caregivers of patients with Alzheimer's disease. *Evid Based Ment Health* 2001 May; 4(2):50
Status: Background article

Mohr E, Bruno G, Foster N, et al. GABA-agonist therapy for Alzheimer's disease. *Clin Neuropharmacol* 1986; 9(3):257-63.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohr E, Schlegel J, Fabbrini G, et al. Clonidine treatment of Alzheimer's disease. *Arch Neurol* 1989 Apr; 46(4):376-8.
*Status: Cross-over trial;
Cross-over trial*

Mohr E, Knott V, Sampson M, et al. Cognitive and quantified electroencephalographic correlates of cycloserine treatment in Alzheimer's disease. *Clin Neuropharmacol* 1995 Feb; 18(1):28-38.
Status: Not included because Jadad Quality Scale score less than three

Mohr E, Walker D, Randolph C, et al. Utility of clinical trial batteries in the measurement of Alzheimer's and Huntington's dementia. *Int Psychogeriatr* 1996; 8(3):397-411.
Status: Not included because does not meet criteria for treatment for dementia patients

Mohr E, Nair NP, Sampson M, et al. Treatment of Alzheimer's disease with sabeluzole: Functional and structural correlates. *Clin Neuropharmacol* 1997 Aug; 20(4):338-45.
Status: Included

Mohs R, Doody R, Morris J, et al. Donepezil preserves functional status in Alzheimer's disease patients: Results from a 1-year prospective

placebo-controlled attrition study. J Eur Coll Neuropsychopharmacol 1999; 9(Suppl 5):S328
Status: Not included because not a full article

Mohs R, Doody R, Morris J, et al. Donepezil preserves activities of daily living in Alzheimer's disease patients results from a one-year placebo-controlled functional survival study. Neurology 2000 Apr; 54(Suppl 3):1
Status: Not included because not a full article

Mohs RC, Davis KL. A signal detectability analysis of the effect of physostigmine on memory in patients with Alzheimer's disease. Neurobiol Aging 1982; 3(2):105-10.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Davis BM, Johns CA, et al. Oral physostigmine treatment of patients with Alzheimer's disease. Am J Psychiatry 1985 Jan; 142(1):28-33.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Davis BM, Greenwald BS. Clinical studies of the cholinergic deficit in Alzheimer's disease. II. Psychopharmacologic studies. J Am Geriatr Soc 1985; 33(11):749-57.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Ferris SH. Measuring response to treatment in Alzheimer's disease: What constitutes meaningful change? Int J Geriatr Psychopharmacol 1998; 1(Suppl 1):S7-S14.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Schmeidler J, Aryan M. Longitudinal studies of cognitive, functional and behavioural change in patients with Alzheimer's disease. Stat Med 2000 Jun 15; 19(11-12):1401-9.
Status: Background article

Mohs RC, Doody RS, Morris JC, et al. A 1-year, placebo-controlled preservation of function survival study of donepezil in AD patients. Neurology 2001 Aug 14; 57(3):481-8.
Status: Included

Molchan SE, Mellow AM, Hill JL, et al. The effects of thyrotropin-releasing hormone and scopolamine in Alzheimer's disease and normal volunteers. J Psychopharmacol (Oxf) 1992; 6(4):489-500.

*Status: Cross-over trial;
Cross-over trial*

Molchan SE, Manji H, Chen G, et al. Effects of chronic lithium treatment on platelet PKC isozymes in Alzheimer's and elderly control subjects. Neurosci Lett 1993; 162(1-2):187-2.
Status: Not included because dementia population not randomized to treatment

Molchan SE, Hill JL, Minichiello M, et al. Scopolamine effects on the pressor response to thyrotropin-releasing hormone in humans. Life Sci 1994; 54(13):933-8.
Status: Not included because no extractable data relevant to review

Moller HJ, Maurer I, Saletu B. Placebo-controlled trial of the xanthine derivative propentofylline in dementia. Pharmacopsychiatry 1994 Jul; 27(4):159-65.
Status: Companion of an included article

Moller HJ, Hampel H, Hegerl U, et al. Double-blind, randomized, placebo-controlled clinical trial on the efficacy and tolerability of a physostigmine patch in patients with senile dementia of the Alzheimer type. Pharmacopsychiatry 1999 May; 32(3):99-106.
Status: Included

Moller HJ, Hartmann A, Kessler C, et al. Naftidrofuryl in the treatment of vascular dementia. Eur Arch Psychiatry Clin Neurosci 2001; 251(6):247-54.
Status: Included

Molloy DW, Cape RD. Acute effects of oral pyridostigmine on memory and cognitive function in SDAT. Neurobiol Aging 1989 Mar; 10(2):199-204.
*Status: Cross-over trial;
Cross-over trial*

Molloy DW, Guyatt GH, Wilson DB, et al. Effect of tetrahydroaminoacridine on cognition, function and behaviour in Alzheimer's disease. CMAJ 1991 Jan 1; 144(1):29-34.
*Status: Cross-over trial;
Cross-over trial*

Molloy DW, Guyatt G, Brown G. Effects of oxiracetam on cognition, behaviour and activities of daily living in dementia. J Clin Exp Gerontol 1992; (3-4):217-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Molloy DW, Guyatt GH, Standish T, et al. Effect of a new nootropic agent, CGS 5649B, on cognition, function, and behavior in dementia. *J Gen Intern Med* 1993 Aug; 8(8):444-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Molloy DW, Standish TI. Clinical experience with Cerebrolysin. *J Neural Transm Suppl* 2000; 59:293-300.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Molsa PK, Marttila RJ, Rinne UK. Survival and cause of death in Alzheimer's disease and multi-infarct dementia. *Acta Neurol Scand* 1986 Aug; 74(2):103-7.

Status: Background article

Moniz-Cook E, Agar S, Gibson G, et al. A preliminary study of the effects of early intervention with people with dementia and their families in a memory clinic. *Aging Ment Health* 1998 Aug; 2(3):199-211.

Status: Not included because dementia population not randomized to treatment

Monreal M, Lafoz E, Olive A, et al. Comparison of subcutaneous unfractionated heparin with a low molecular weight heparin (Fragmin (R)) in patients with venous thromboembolism and contraindications to coumarin. *Thromb Haemost* 1994; 71(1):7-11.

Status: Not included because does not meet criteria for treatment for dementia patients

Monteverde A, Gnemmi P, Rossi F, et al. Selegiline in the treatment of mild to moderate Alzheimer-type dementia. *Clin Ther* 1990 Jul; 12(4):315-22.

Status: Not included because Jadad Quality Scale score less than three

Montgomery P, Erickson GK. Neuropsychological perspectives in amyotrophic lateral sclerosis. *Neurol Clin* 1987; 5(1):61-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Montori VM, Guyatt GH. Intention-to-treat principle. *CMAJ* 2001 Nov 13; 165(10):1339-41.

Status: Background article

Moore MJ, Clipp EC. Alzheimer's disease and caregiver time. *Lancet* 1994 Jan 22; 343(8891):239-40.

Status: Not included because not a full article

Moore S, Sandman CA, McGrady K, et al. Memory training improves cognitive ability in patients with dementia. *Neuropsychol Rehab* 2001; 11(3-4):245-61.

Status: Not included because does not meet criteria for treatment for dementia patients

Moretti R, Torre P, Antonello RM, et al. Rivastigmine in subcortical vascular dementia: A comparison trial on efficacy and tolerability for 12 months follow-up. *Eur J Neurol* 2001 Jul; 8(4):361-2.

Status: Not included because not a full article

Moretti R, Torre P, Antonello RM, et al. An open-label pilot study comparing rivastigmine and low-dose aspirin for the treatment of symptoms specific to patients with subcortical vascular dementia. *Curr Ther Res Clin Exp* 2002; 63(7):443-58.

Status: Not included because dementia population not randomized to treatment

Moretti R, Torre P, Antonello RM, et al. Depression and Alzheimer's disease: Symptom or comorbidity? *Am J Alzheimers Dis Other Demen* 2002; 17(6):338-44.

Status: Not included because Jadad Quality Scale score less than three

Moretti R, Torre P, Antonello RM, et al. Frontotemporal dementia: Paroxetine as a possible treatment of behavior symptoms: A randomized, controlled, open 14-month study. *Eur Neurol* 2003; 49(1):13-9.

Status: Not included because Jadad Quality Scale score less than three

Morey LC, Ban TA, Cassano G, et al. Glycosaminoglycan polysulfate in old-age dementias: A factor-analytic study of change in psychopathologic symptoms. *Neuropsychobiology* 1988; 19(3):135-8.

Status: Not included because dementia population not randomized to treatment

Morganroth J, Graham S, Hartman R, et al. Electrocardiographic effects of rivastigmine. *J Clin Pharmacol* 2002 May; 42(5):558-68.

Status: Not included because does not meet criteria for treatment for dementia patients

Mori T, Inoue D, Kosugi S, et al. Effects of low dose L-triiodothyronine administration on mental, behavioural and thyroid states in elderly subjects. *Endocrinol Jpn* 1988 Aug; 35(4):585-92.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Morich FJ, Bieber F, Lewis JM, et al. Nimodipine in the treatment of probable Alzheimer's disease: Results of two multicentre trials. *Clin Drug Investig* 1996; (4):185-95.
Status: Not included because dementia population not randomized to treatment

Moroney JT, Tang MX, Berglund L, et al. Low-density lipoprotein cholesterol and the risk of dementia with stroke. *JAMA* 1999; 282(3):254-60.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Moroney JT, Tseng C-L, Paik MC, et al. Treatment for the secondary prevention of stroke in older patients: The influence of dementia status. *J Am Geriatr Soc* 1999; 47(7):824-9.
Status: Not included because dementia population not randomized to treatment

Morris JC, McKeel DW, Jr., Fulling K, et al. Validation of clinical diagnostic criteria for Alzheimer's disease. *Ann Neurol* 1988 Jul; 24(1):17-22.
Status: Background article

Morris JC, Cyrus PA, Orazem J, et al. Metrifonate benefits cognitive, behavioral, and global function in patients with Alzheimer's disease. *Neurology* 1998 May; 50(5):1222-30.
Status: Included

Morrison RS, Siu AL. A comparison of pain and its treatment in advanced dementia and cognitively intact patients with hip fracture. *J Pain Symptom Manage* 2000 Apr; 19(4):240-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mortimer JA, van Duijn CM, Chandra V, et al. Head trauma as a risk factor for Alzheimer's disease: a collaborative re-analysis of case-control studies. EURODEM Risk Factors Research Group. *Int J Epidemiol* 1991; 20 Suppl 2:S28-S35
Status: Background article

Moss DE, Berlanga P, Hagan MM, et al. Methanesulfonyl fluoride (MSF): A double-blind,

placebo-controlled study of safety and efficacy in the treatment of senile dementia of the Alzheimer type. *Alzheimer Dis Assoc Disord* 1999 Jan; 13(1):20-5.
Status: Not included because Jadad Quality Scale score less than three

Mouradian MM, Mohr E, Williams JA, et al. No response to high-dose muscarinic agonist therapy in Alzheimer's disease. *Neurology* 1988 Apr; 38(4):606-8.
Status: Not included because dementia population not randomized to treatment

Mouradian MM, Blin J, Giuffra M, et al. Somatostatin replacement therapy for Alzheimer dementia. *Ann Neurol* 1991 Oct; 30(4):610-3.
Status: Not included because dementia population not randomized to treatment

Mubrin Z, Knezevic S, Spilich G, et al. Normalization of rCBF pattern in senile dementia of the Alzheimer's type. *Psychiatry Res* 1989 Sep; 29(3):303-6.
Status: Not included because no extractable data relevant to review

Mucke HAM. Metrifonate. Treatment of Alzheimer's disease, acetylcholinesterase inhibitor. *Drugs of the Future* 1998; 23(5):491-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mulnard RA, Cotman CW, Kawas C, et al. Estrogen replacement therapy for treatment of mild to moderate Alzheimer disease: A randomized controlled trial. *JAMA* 2000; (8):1007-15.
Status: Included

Mulsant BH, Mazumdar S, Pollock BG, et al. Methodological issues in characterizing treatment response in demented patients with behavioral disturbances. *Int J Geriatr Psychiatry* 1997 May; (No. 5):537-47.
Status: Not included because dementia population not randomized to treatment

Munch-Petersen S, Pakkenberg H, Kornerup H, et al. RNA treatment of dementia. A double-blind study. *Acta Neurol Scand* 1974; 50(5):553-72.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mungas D, Reed BR. Application of item response theory for development of a global functioning

measure of dementia with linear measurement properties. *Stat Med* 2000 Jun 15; 19(11-12):1631-44.

Status : Background article

Murali DP, Kaiser L. Variability of the mini-mental state examination in dementia. *Neurology* 2000 Apr 11; 54(7):1538-9.

Status: Not included because not a full article

Muramoto O, Sugishita M, Sugita H, et al. Effect of physostigmine on constructional and memory tasks in Alzheimer's disease. *Arch Neurol* 1979 Aug; 36(8):501-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Muramoto O, Sugishita M, Ando K. Cholinergic system and constructional praxis: A further study of physostigmine in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 1984; 47(5):485-91.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Muratorio A, Bonuccelli U, Nuti A, et al. A neurotropic approach to the treatment of multi-infarct dementia using L-alpha-glycerylphosphorylcholine. *Curr Ther Res Clin Exp* 1992; 52(5):741-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Muresanu DF, Rainer M, Moessler H. Improved global function and activities of daily living in patients with AD: a placebo-controlled clinical study with the neurotrophic agent Cerebrolysin. *J Neural Transm Suppl* 2002; (62):277-85.

Status: Not included because Jadad Quality Scale score less than three

Murialdo G, Barreca A, Nobili F, et al. Dexamethasone effects on cortisol secretion in Alzheimer's disease: Some clinical and hormonal features in suppressor and nonsuppressor patients. *J Endocrinol Invest* 2000 Mar; 23(3):178-86.

Status: Not included because does not meet criteria for treatment for dementia patients

Murri L, Bardi C, Arena R. A comparison between lormetazepam and flunitrazepam in insomniac patients affected by chronic cerebrovascular disorders. *Curr Ther Res Clin Exp* 1984; 35(1):113-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nadeau SE, Malloy PF, Andrew ME. A crossover trial of bromocriptine in the treatment of vascular dementia. *Ann Neurol* 1988 Aug; 24(2):270-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nagaraja D, Jayashree S. Randomized study of the dopamine receptor agonist piribedil in the treatment of mild cognitive impairment. *Am J Psychiatry* 2001 Sep; 158(9):1517-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nair NP, Ban TA, Hontela S, et al. Trazodone in the treatment of organic brain syndromes, with special reference to psychogeriatrics. *Curr Ther Res Clin Exp* ; 15(10):769-75.

Status: Not included because dementia population not randomized to treatment

Nair NPV, Gauthier S, Etienne Pao. Lack of effect of a 3-month treatment with lecithin in Alzheimer's disease. *Prog Neuropsychopharmacol* 1981; 26:426-8.

Status: Article not retrievable

Nakamura H, Nakanishi M, Hamanaka T, et al. Semantic priming in patients with Alzheimer and semantic dementia. *Cortex* 2000 Apr; 36(2):151-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Nakamura K. Aniracetam: Its novel therapeutic potential in cerebral dysfunctional disorders based on recent pharmacological discoveries. *CNS Drug Rev* 2002; 8(1):70-89.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nakano S, Asada T, Matsuda H, et al. Donepezil hydrochloride preserves regional cerebral blood flow in patients with Alzheimer's disease. *J Nucl Med* 2001 Oct; 42(10):1441-5.

Status: Not included because Jadad Quality Scale score less than three

Namazi KH, Haynes SR. Sensory stimuli reminiscence for patients with Alzheimer's disease: Relevance and implications. *Clin Gerontol* 1994; 29-46.

Status: Not included because dementia population not randomized to treatment

Namazi KH, Johnson BD. Environmental effects on incontinence problems in Alzheimer's disease

patients. Am J Alzheimers Care Relat Disord 1991 Nov; (6):16-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nappi G, Bono G. Comparison between glycerophosphorylserine and oxiracetam in the treatment of patients with mental deterioration. Funct Neurol 1993; (Suppl 5):73-82.

Status: Article not retrievable

Nappi G, Bono G, Merlo P, et al. Long-term nicergoline treatment of mild to moderate senile dementia. Results of a multicentre, double-blind, placebo-controlled study. Clin Drug Investig 1997; (6):308-16.

Status: Included

Naritomi H, Murata S, Shimizu T, et al. Long-term effects of bifemelane hydrochloride on post-stroke deterioration of cognitive function and cerebral blood flow. Curr Ther Res Clin Exp 1995; Vol 56(231):238

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nasrallah HA, Varney N, Coffman JA, et al. Effects of naloxone on cognitive deficits following electroconvulsive therapy. Psychopharmacol Bull 1985; 21(1):89-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Navia BA, Dafni U, Simpson D, et al. A phase I/II trial of nimodipine for HIV-related neurologic complications. Neurology 1998 Jul; 51(1):221-8.

Status: Not included because dementia population not randomized to treatment

Nebes RD, Pollock BG, Mulsant BH, et al. Cognitive effects of paroxetine in older depressed patients. J Clin Psychiatry 1999; 60(Suppl):9

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Netz Y, Yaretzki A, Salganik I, et al. The effect of supervised physical activity on cognitive and affective state of geriatric and psychogeriatric in-patients. Clin Gerontol 1994; 15(1):47-56.

Status: Not included because does not meet criteria for treatment for dementia patients

Neumann PJ, Hermann RC, Kuntz KM, et al. Cost-effectiveness of donepezil in the treatment of mild or moderate Alzheimer's disease. Neurology

1999; 52(6):1138-45.

Status: Companion of an included article

Neumeister A, Graf A, Willeit M, et al. Effects of Light Therapy in dementia. Biol Psychiatry 2000; Vol 47:S536

Status: Not included because not a full article

Newcomer R, Spitalny M, Fox P, et al. Effects of the Medicare Alzheimer's disease demonstration on the use of community-based services. Health Serv Res 1999 Aug; 34(3):645-67.

Status: Not included because does not meet criteria for treatment for dementia patients

Newhouse PA, Sunderland T, Tariot PN, et al. Intravenous nicotine in Alzheimer's disease: A pilot study. Psychopharmacologia 1988; 95(2):171-5.

Status: Not included because dementia population not randomized to treatment

Newhouse PA, Sunderland T, Narang PK, et al. Neuroendocrine, physiologic, and behavioral responses following intravenous nicotine in nonsmoking healthy volunteers and in patients with Alzheimer's disease.

Psychoneuroendocrinology 1990; 15(5-6):471-6.

Status: Not included because dementia population not randomized to treatment

Niemann H, Ruff RM, Baser CA. Computer-assisted attention retraining in head-injured individuals: A controlled efficacy study of an outpatient program. J Consult Clin Psychol 1990 Dec; 58(6):811-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nijhuis E, Hinloopen B, van Duijn C, et al. Decreased sensitivity to dexamethasone in lymphocytes from patients with Alzheimer's disease. Clin Immunol Immunopathol 1994 Oct; 73(1):45-52.

Status: Not included because does not meet criteria for treatment for dementia patients

Nikolova G, Traykov L. Efficacy of donepezil in patients with Alzheimer's disease - Results of 12-week open clinical trial. Acta Med Bulg 2001; 28:70-5.

Status: Not included because dementia population not randomized to treatment

Nimodipine Clinical Study Group. Effects of Nimodipine on vascular dementia. Cardiovasc

Dis 1996; 6(Suppl 2):70.

Status: Not included because not a full article

Nirenberg TD. Relocation of institutionalized elderly. J Consult Clin Psychol 1983 Oct; 51(5):693-701.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nobili F, Vitali P, Canfora M, et al. Effects of long-term Donepezil therapy on rCBF of Alzheimer's patients. Clin Neurophysiol 2002; 113(8):1241-8.
Status: Not included because dementia population not randomized to treatment

Nobili F, Koulibaly M, Vitali P, et al. Brain perfusion follow-up in Alzheimer's patients during treatment with acetylcholinesterase inhibitors. J Nucl Med 2002 Aug; 43(8):983-90.
Status: Not included because dementia population not randomized to treatment

Noel G, Jeanmart M, Reinhardt B. Treatment of the organic brain syndrome in the elderly. A double-blind comparison on the effects of a neurotropic drug and placebo. Neuropsychobiology 1983; 10(2-3):90-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nolan KA, Black RS, Sheu KF, et al. A trial of thiamine in Alzheimer's disease. Arch Neurol 1991 Jan; 48(1):81-3.
Status: Included

Nolan KA, Lino MM, Seligmann AW, et al. Absence of vascular dementia in an autopsy series from a dementia clinic. J Am Geriatr Soc 1998 May; 46(5):597-604.
Status: Background article

Norbergh KG, Hellzen O, Sandman PO, et al. The relationship between organizational climate and the content of daily life for people with dementia living in a group-dwelling. J Clin Nurs 2002 Mar; 11(2):237-46.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Novo FP, Ryan RP, Frazier EL. Dihydroergotoxine mesylate in treatment of symptoms of idiopathic cerebral dysfunction in geriatric patients. Clin Ther 1978; 1(5):359-69.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nyback H, Hassan M, Junthe T, et al. Clinical experiences and biochemical findings with tacrine (THA). Acta Neurol Scand Suppl 1993; 88(149):36-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nygaard H, Bakke K, Brudvik E, et al. Zuclopenthixol and melperon in the treatment of elderly patients: A double-blind, controlled, multi-centre study. Pharmatherapeutica 1987; 5(3):152-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nygaard HA, Bakke K, Brudvik E, et al. Dosing of neuroleptics in elderly demented patients with aggressive and agitated behaviour: A double-blind study with zuclopenthixol. Curr Med Res Opin 1994; 13(4):222-32.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nyth AL, Gottfries CG. The clinical efficacy of citalopram in treatment of emotional disturbances in dementia disorders. A Nordic multicentre study. Br J Psychiatry 1990 Dec; 157:894-901.
Status: Included

Nyth AL, Gottfries CG, Lyby K, et al. A controlled multicenter clinical study of citalopram and placebo in elderly depressed patients with and without concomitant dementia. Acta Psychiatr Scand 1992 Aug; 86(2):138-45.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Brien BJ, Goeree R, Hux M, et al. Economic evaluation of donepezil for the treatment of Alzheimer's disease in Canada. J Am Geriatr Soc 1999; 47(5):570-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Brien JT, Schweitzer I, Ames D, et al. The function of the hypothalamic-pituitary-adrenal axis in Alzheimer's disease. Response to insulin hypoglycaemia. Br J Psychiatry 1994 Nov; 165(5):650-7.
Status: Not included because dementia population not randomized to treatment

O'Carroll RE, Moffoot A, Ebmeier KP, et al. Korsakoff's syndrome, cognition and clonidine. Psychol Med 1993 May; 23(2):341-7.

Status: Not included because dementia population not randomized to treatment

O'Carroll RE, Moffoot AP, Ebmeier KP, et al. Effects of fluvoxamine treatment on cognitive functioning in the alcoholic Korsakoff syndrome. *Psychopharmacologia* 1994 Sep; 116(1):85-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Connolly M, Dierdorf D, Greb WH, et al. Efficacy of denbutylline in patients with multi-infarct dementia. *Drug Dev Res* 1988; 14(3-4):195-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Connor DW, Pollitt PA, Brook CP, et al. Does early intervention reduce the number of elderly people with dementia admitted to institutions for long term care? *BMJ* 1991; 302(6781):871-5.
Status: Not included because dementia population not randomized to treatment

O'Donnell VM, Pitts WM, Fann WE. Noradrenergic and cholinergic agents in Korsakoff's syndrome. *Clin Neuropharmacol* 1986; 9(1):65-70.
Status: Background article

O'Keeffe ST, Lavan JN. Subcutaneous fluids in elderly hospital patients with cognitive impairment. *Gerontology* 1996; 42(1):36-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Neill RT. Biostatistical considerations in pharmacovigilance and pharmacoepidemiology: linking quantitative risk assessment in pre-market licensure application safety data, post-market alert reports and formal epidemiological studies. *Stat Med* 1998 Aug 15; 17(15-16):1851-8.
Status: Background article

Oakley F, Sunderland T. Assessment of motor and process skills as a measure of IADL functioning in pharmacologic studies of people with Alzheimer's disease: A pilot study. *Int Psychogeriatr* 1997 Jun; 9(2):197-206.
Status: Cross-over trial;
Cross-over trial

Obonsawin MC, Robertson A, Crawford JR, et al. Non-mnemonic cognitive function in the scopolamine model of Alzheimer's disease. *Hum Psychopharmacol* 1998; 13:439-49.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ohno Y. The effects of magnetized mineral water on memory loss delay in Alzheimer's disease. *The centre for frontier sciences* 1997; (6):38-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Oishi M, Mochizuki Y, Takasu T, et al. Effectiveness of traditional Chinese medicine in Alzheimer disease. *Alzheimer Dis Assoc Disord* 1998 Sep; 12(3):247-50.
Status: Not included because dementia population not randomized to treatment

Okajima Y. Quantitative pharmaco-EEG study of indeloxazine hydrochloride in psychogeriatric patients. *Integr Psychiatry* 1993; 9(1):25-33.
Status: Not included because dementia population not randomized to treatment

Olafsson K, Jorgensen S, Jensen HV, et al. Fluvoxamine in the treatment of demented elderly patients: A double-blind, placebo-controlled study. *Acta Psychiatr Scand* 1992 Jun; 85(6):453-6.
Status: Included

Olin J, Schneider L, Novit A, et al. Hydergine for dementia. In: *The Cochrane Library*, 2000. Issue 2. Oxford: Update Software
Status: Background article

Olin JT, Schneider LS. Assessing response to tacrine using the factor analytic structure of the Alzheimer's Disease Assessment Scale (ADAS) - Cognitive subscale. *Int J Geriatr Psychiatry* 1995; 10(9):753-6.
Status: Not included because Jadad Quality Scale score less than three

Olin JT, Fox LS, Pawluczyk S, et al. A pilot randomized trial of carbamazepine for behavioral symptoms in treatment-resistant outpatients with Alzheimer's disease. *Am J Geriatr Psychiatry* 2001; 9(4):400-5.
Status: Included

Onofrij M, Thomas A, Luciano AL, et al. Donepezil versus vitamin E in Alzheimer's disease: Part 2: Mild versus moderate-severe Alzheimer's disease. *Clin Neuropharmacol* 2002 Jul; 25(4):207-15.
Status: Not included because Jadad Quality Scale score less than three

Opie J, Rosewarne R, O'Connor DW. The efficacy of psychosocial approaches to behaviour disorders in dementia: a systematic literature review. Aust N Z J Psychiatry 1999 Dec; 33(6):789-99.

Status: Background article

Opie J, Doyle C, O'Connor DW. Challenging behaviours in nursing home residents with dementia: A randomized controlled trial of multidisciplinary interventions. Int J Geriatr Psychiatry 2002 Jan; 17(1):6-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Oremus M, Perrault A, Demers L, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Psychometric properties of global scales. J Geriatr Psychiatry Neurol 2000; 13(4):197-205.

Status: Background article

Oremus M, Wolfson C, Perrault A, et al. Interrater reliability of the modified Jadad quality scale for systematic reviews of Alzheimer's disease drug trials. Dement Geriatr Cogn Disord 2001 May; 12(3):232-6.

Status: Background article

Orengo CA, Kidwell K, Kunik ME, et al. The effect of risperidone on cognitive performance in elderly psychotic and aggressive patients with dementia; a pilot study. Int J Geriatr Psychopharmacol 1998; 1(4):193-6.

Status: Not included because dementia population not randomized to treatment

Orgogozo J, Rigaud AS, Stoffler A, et al. Efficacy and safety of memantine in patients with mild to moderate vascular dementia. Stroke 2002; 33:1834-9.

Status: Included

Orten JD, Allen M, Cook J. Reminiscence Groups with Confused Nursing Center Residents: An Experimental Study. Soc Work Health Care 1989; 14(1):73-86.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ostwald SK, Hepburn KW, Caron W, et al. Reducing caregiver burden: A randomized psychoeducational intervention for caregivers of persons with dementia. Gerontologist 1999 Jun; 39(3):299-309.

Status: Not included because does not meet criteria for treatment for dementia patients

Ott BR, Lapane KL. Tacrine therapy is associated with reduced mortality in nursing home residents with dementia. J Am Geriatr Soc 2002 Jan; 50(1):35-40.

Status: Background article

Ousset PJ, Viallard G, Puel M, et al. Lexical Therapy and episodic word learning in dementia of the Alzheimer type. Brain Lang 2002; 80(1):14-20.

Status: Not included because does not meet criteria for treatment for dementia patients

Owensworth TL, McFarland K. Memory remediation in long-term acquired brain injury: Two approaches in diary training. Brain Inj 1999 Aug; 13(8):605-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Paire JA, Karney RJ. The effectiveness of sensory stimulation for geropsychiatric inpatients. Am J Occup Ther 1984 Aug; 38(8):505-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Palmer GC. Neuroprotection by NMDA receptor antagonists in a variety of neuropathologies. Curr Drug Targets 2001; 2(3):241-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Palmieri G, Palmieri R, Inzoli MR. Double-blind controlled trial of phosphatidylserine in patients with senile mental deterioration. Clin Trials J 1987; 24(1):73-83.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Panisset M, Gauthier S, Moessler H, et al. Cerebrolysin in Alzheimer's disease: A randomized, double-blind, placebo-controlled trial with a neurotrophic agent. J Neural Transm Gen Sect 2002; 109(7-8):1089-104.

Status: Included

Pantev M, Ritter R, Gortelmeyer R. Therapy responder analysis of dementia study with memantine. Pharmacopsychiatry 1993; 26:185.

Status: Not included because not a full article

Pantoni L, Rossi R, Inzitari D, et al. Efficacy and safety of nimodipine in subcortical vascular dementia: A subgroup analysis of the Scandinavian Multi-Infarct Dementia Trial. *JNS* 2000b; 175(2):124-34.

Status: Companion of an included article

Pantoni L, Bianchi C, Beneke M, et al. The Scandinavian Multi-Infarct Dementia Trial: A double-blind, placebo-controlled trial on nimodipine in multi-infarct dementia. *JNS* 2000a; 175(2):116-23.

Status: Included

Parkes JD, Marsden CD, Rees JE. Parkinson's disease, cerebral arteriosclerosis, and senile dementia. Clinical features and response to levodopa. *Q J Med* 1974; 43(169):49-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Ciuffetti G, Mercuri M, et al. Haemorheological pattern in initial mental deterioration: Results of a long-term study using piracetam and pentoxifylline. *Arch Gerontol Geriatr* 1985 Jul; 4(2):141-55.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Ciuffetti G, Mercuri M, et al. Relationship between haemorheological factors and initial mental deterioration in the elderly. A preliminary study. *Clin Hemorheol* 1985; 5(4):361-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Ciuffetti G, Mercuri M, et al. The role of haemorheological factors in the ageing brain: Long-term therapy with pentoxifylline ('Trental' 400) in elderly patients with initial mental deterioration. *Pharmatherapeutica* 1986; 4(10):617-27.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Bartorelli L, Bonaiuto S, et al. Aniracetam (Ro 13-5057) for the treatment of senile dementia of Alzheimer type: Results of a multicentre clinical study. *Dementia* 1991; 2(5):262-7.

Status: Article not retrievable

Parnetti L, Abate G, Bartorelli L, et al. Multicentre study of l-alpha-glyceryl-phosphorylcholine vs ST200 among patients with probable senile

dementia of Alzheimer's type. *Drugs Aging* 1993 Mar; 3(2):159-64.

Status: Not included because Jadad Quality Scale score less than three

Parnetti L, Senin U, Carosi M, et al. Mental deterioration in old age: results of two multicenter, clinical trials with nimodipine. The Nimodipine Study Group. *Clin Ther* 1993 Mar; 15(2):394-406.

Status: Not included because dementia population not randomized to treatment

Parnetti L, Ambrosoli L, Abate G, et al. Posatirelin for the treatment of late-onset Alzheimer's disease: A double-blind multicentre study vs citicoline and ascorbic acid. *Acta Neurol Scand* 1995 Aug; 92(2):135-40.

Status: Included

Parnetti L, Ambrosoli L, Agliati G, et al. Posatirelin in the treatment of vascular dementia: A double-blind multicentre study vs placebo. *Acta Neurol Scand* 1996 Jun; 93(6):456-63.

Status: Included

Parnetti L, Mari D, Abate G, et al. Vascular dementia Italian sulodexide study (VA.D.I.S.S.). Clinical and biological results. *Thromb Res* 1997 Jul 15; 87(2):225-33.

Status: Included

Parnetti L, Amici S, Lanari A, et al. Cerebrospinal fluid levels of biomarkers and activity of acetylcholinesterase (AChE) and butyrylcholinesterase in AD patients before and after treatment with different AChE inhibitors. *Neurol Sci* 2002; 23(Suppl 2):S95-S96.

Status: Not included because not a full article

Partanen J, Wolters EC, van Duijn H. Clinical neurophysiological correlates of vasopressin derivative therapy in Alzheimer's disease. *Clin Neurol Neurosurg* 1987; 2:34.

Status: Not included because not a full article

Partanen JV, Soininen H, Riekkinen PJ. Does an ACTH derivative (Org 2766) prevent deterioration of EEG in Alzheimer's disease? *Electroencephalogr Clin Neurophysiol* 1986 Jun; 63(6):547-51.

Status: Companion of an included article

Passeri M, Cucinotta D, de Mello M, et al. Minaprine for senile dementia. *Lancet* 1985 Apr 6; 1(8432):824.

Status: Not included because not a full article

Passeri M, Cucinotta D, de Mello M, et al. Comparison of minaprine and placebo in the treatment of Alzheimer's disease and multi-infarct dementia. *Int J Geriatr Psychiatry* 1987; 2(2):97-103.

Status : Included

Passeri M, Cucinotta D. Ateroid in the clinical treatment of multi-infarct dementia. *Mod Probl Pharmacopsychiatry* 1989; 23:85-94.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Passeri M, Cucinotta D, Bonati PA, et al. Acetyl-L-carnitine in the treatment of mildly demented elderly patients. *Int J Clin Pharmacol Res* 1990; 10(1-2):75-82.

Status: Not included because Jadad Quality Scale score less than three

Passeri M, Cucinotta D, Abate G, et al. Oral 5'-methyltetrahydrofolic acid in senile organic mental disorders with depression: Results of a double-blind multicenter study. *Aging (Milano)* 1993 Feb; 5(Milano):63-71.

Status: Included

Paterson J, Hamilton MM, Grant H. The effectiveness of the Hierarchic Dementia Scale in tailoring interventions to reduce problem behaviours in people with Alzheimer's disease. *Aust Occup Ther J* 2000; 47(3):134-40.

Status: Not included because does not meet criteria for treatment for dementia patients

Pathy J, Menon G, Reynolds A, et al. Betahistine hydrochloride (Serc) in cerebrovascular disease: A placebo-controlled study. *Age Ageing* 1977 Aug; 6(3):179-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Patterson C, Gauthier S. Canadian consensus conference on dementia: Two years later. *Can J Neurol Sci* 2001 Feb; 28 Suppl 1:S1-S2

Status: Background article

Patterson CJ, Gauthier S, Bergman H, et al. The recognition, assessment and management of dementing disorders: Conclusions from the Canadian Consensus Conference on Dementia. *CMAJ* 1999 Jun; 160(Suppl 12):S1-15.

Status: Background article

Patterson CJ, Gass DA. Screening for cognitive impairment and dementia in the elderly. *Can J*

Neurol Sci 2001 Feb; 28(Suppl):S42-S51.

Status: Background article

Peabody CA, Thiemann S, Pigache R, et al. Desglycinamide-9-arginine-8-vasopressin (DGA VP, Organon 5667) in patients with dementia. *Neurobiol Aging* 1985; 6(2):95-100.

Status: Not included because dementia population not randomized to treatment

Peabody CA, Davies H, Berger PA, et al. Desamino-D-arginine-vasopressin (DDAVP) in Alzheimer's disease. *Neurobiol Aging* 1986 Jul; 7(4):301-3.

Status: Included

Peabody CA, Deblois TE, Tinklenberg JR. Thyrotropin-releasing hormone (TRH) and Alzheimer's disease. *Am J Psychiatry* 1986; 143(2):262-3.

Status: Not included because not a full article

Penn RD, Martin EM, Wilson RS, et al. Intraventricular bethanechol infusion for Alzheimer's disease: Results of double-blind and escalating-dose trials. *Neurology* 1988 Feb; 38(2):219-22.

Status: Cross-over trial; Cross-over trial

Pepping J. Phosphatidylserine. *Am J Health Syst Pharm* 1999; 56(20):2038-44.

Status: Not included because not a full article

Pepping J. Alternative therapies. Huperzine A: A potent and selective acetylcholinesterase inhibitor. *Am J Health Syst Pharm* 2000 Mar 15; 57(6):530-4.

Status: Not included because dementia population not randomized to treatment

Perini M, Montanini R, Casucci R. Effects of eptastigmine: A new cholinesterase inhibitor on regional cerebral blood flow in Alzheimer patients. *J Neurol* 1995; 242:S57.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Perrault A, Oremus M, Demers L, et al. Review of outcome measurement instruments in Alzheimer's disease drug trials: Psychometric properties of behavior and mood scales. *J Geriatr Psychiatry Neurol* 2000; 13(4):181-96.

Status: Background article

Perryman KM, Fitten LJ. Quantitative EEG during a double-blind trial of THA and lecithin in patients with Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1991 Jul; 4(3):127-33.

Status: Not included because no extractable data relevant to review

Peskind ER, Wingerson D, Murray S, et al. Effects of Alzheimer's disease and normal aging on cerebrospinal fluid norepinephrine responses to yohimbine and clonidine. *Arch Gen Psychiatry* 1995 Sep; 52(9):774-82.

Status: Not included because does not meet criteria for treatment for dementia patients

Peters BH, Levin HS. Memory enhancement after physostigmine treatment in the amnesic syndrome. *Arch Neurol* 1977 Apr; 34(4):215-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Peters BH, Levin HS. Effects of physostigmine and lecithin on memory in Alzheimer's disease. *Ann Neurol* 1979 Sep; 6(3):219-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Peters BH. Memory augmentation following physostigmine or physostigmine plus lecithin administration in amnesic syndrome and Alzheimer's disease. *Int J Neurosci* 1981; Vol 12:178.

Status: Article not retrievable

Petersen RC, Smith GE, Waring SC, et al. Aging, memory, and mild cognitive impairment. *Int Psychogeriatr* 1997; 9 Suppl 1:65-9.

Status: Background article

Petersen RC, Stevens JC, Ganguli M, et al. Practice parameter: Early detection of dementia: Mild cognitive impairment (an evidence-based review). Report of the Quality Standards Subcommittee of the American Academy of Neurology. *Neurology* 2001 May; 56(9):1133-42.

Status: Background article

Peterson LG, Bongar B. Navane versus Haldol. Treatment of acute organic mental syndromes in the general hospital. *Gen Hosp Psychiatry* 1989 Nov; 11(6):412-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Petit D, Montplaisir J, Lorrain D, et al. THA does not affect sleep or EEG spectral power in

Alzheimer's disease. *Biol Psychiatry* 1993; 33(10):753-4.

Status: Not included because dementia population not randomized to treatment

Petracca G, Teson A, Chemerinski E, et al. A double-blind placebo-controlled study of clomipramine in depressed patients with Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1996; 8(3):270-5.

Status: Cross-over trial; Cross-over trial

Petracca GM, Chemerinski E, Starkstein SE. A double-blind, placebo-controlled study of fluoxetine in depressed patients with Alzheimer's disease. *Int Psychogeriatr* 2001 Jun; 13(2):233-40.

Status: Included

Petrie WM, Ban TA, Berney S, et al. Loxapine in psychogeriatrics: A placebo- and standard-controlled clinical investigation. *J Clin Psychopharmacol* 1982 Apr; 2(2):122-6.

Status: Included

Pettegrew JW, Klunk WE, Panchalingam K, et al. Clinical and neurochemical effects of acetyl-L-carnitine in Alzheimer's disease. *Neurobiol Aging* 1995 Jan; 16(1):1-4.

Status: Not included because dementia population not randomized to treatment

Pettegrew JW, Levine J, McClure RJ. Acetyl-L-carnitine physical-chemical, metabolic, and therapeutic properties: Relevance for its mode of action in Alzheimer's disease and geriatric depression. *Mol Psychiatry* 2000; 5(6):616-32.

Status: Not included because does not meet criteria for treatment for dementia patients

Pettegrew JW, McClure RJ. Acetyl-L-carnitine as a possible therapy for Alzheimer's disease. *Expert Rev Neurother* 2002; 2(5):647-54.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pettigrew LC, Bieber F, Lettieri J, et al. Pharmacokinetics, pharmacodynamics, and safety of metrifonate in patients with Alzheimer's disease. *J Clin Pharmacol* 1998 Mar; 38(3):236-45.

Status: Included

Petursson H. A controlled study of Moclobemide in elderly depression with cognitive decline.

Psychopharmacologia 1993; Vol 111:B5
Status: Not included because not a full article

Pérodeau G, Lauzon S, Lévesque L, et al. Mental health stress correlates and psychotropic drug use or non user among aged caregivers to elders with dementia. Aging Ment Health 2001; 5(3):225-34.
Status: Not included because does not meet criteria for treatment for dementia patients

Pfefferbaum A, Davis KL, Coulter CL, et al. EEG effects of physostigmine and choline chloride in humans. Psychopharmacologia 1979 Apr 25; 62(3):225-33.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Phanjoo AL, Link C. Remoxipride versus thioridazine in elderly psychotic patients. Acta Psychiatr Scand Suppl 1990; Vol 358:181-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Phillips CD, Spry KM, Sloane PD, et al. Use of physical restraints and psychotropic medications in Alzheimer special care units in nursing homes. Am J Public Health 2000; 90(1):92-6.
Status: Not included because dementia population not randomized to treatment

Piccinin GL, Finali G, Piccirilli M. Neuropsychological effects of L-deprenyl in Alzheimer's type dementia. Clin Neuropharmacol 1990 Apr; 13(2):147-63.
*Status: Cross-over trial;
Cross-over trial*

Piccoli F, Battistini N, Carbonin P, et al. CDP-choline in the treatment of chronic cerebrovasculopathies. Arch Gerontol Geriatr 1994; 18(3):161-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pillemer K, Jill SJ. Peer support for Alzheimer's caregivers: Is it enough to make a difference? Res Aging 2002; 24(2):171-92.
Status: Not included because does not meet criteria for treatment for dementia patients

Pillo G. Double-blind computerized EEG study on the effect of mesoglycan in patients with chronic cerebrovascular insufficiency. Riv Neurobiol 1988; 17(3-4):75-4.
Status: Article not retrievable

Pincus MM, Kilander L, Ohrvall M. Alpha-tocopherol and Alzheimer's disease. N Engl J Med 1997; 337(8):572-3.
Status: Not included because not a full article

Pisvejc J, Hyrman V, Sikora J, et al. A comparison of brief and ultrabrief pulse stimuli in unilateral ECT. J ECT 1998 Jun; 14(2):68-75.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pittera A, Ciancitto S. Effect of oral nimodipine on cerebral blood flow in patients with chronic cerebrovascular disorders. A supra-aortic Doppler ultra-sound open study. Curr Ther Res Clin Exp 1990; 48(4):716-29.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pocock SJ, Abdalla M. The hope and the hazards of using compliance data in randomized controlled trials. Stat Med 1998 Feb 15; 17(3):303-17.
Status: Background article

Pohjasvaara T, Mantyla R, Ylikoski R, et al. Comparison of different clinical criteria (DSM-III, ADDTC, ICD-10, NINDS-AIREN, DSM-IV) for the diagnosis of vascular dementia. National Institute of Neurological Disorders and Stroke-Association Internationale pour la Recherche et l'Enseignement en Neurosciences. Stroke 2000 Dec; 31(12):2952-7.
Status: Background article

Poitrenaud J, Piette F, Malbezin M, et al. Almitrine-raubasine and cognitive impairment in the elderly: Results of a 6-month controlled multicenter study. Clin Neuropharmacol 1990; Vol 13(Suppl 3):S100-S108.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Polinsky RJ. Clinical pharmacology of rivastigmine: A new-generation acetylcholinesterase inhibitor for the treatment of Alzheimer's disease. Clin Ther 1998; 20(4):634-47.
Status: Not included because dementia population not randomized to treatment

Pollock BG, Mulsant BH, Rosen J, et al. Comparison of citalopram, perphenazine, and placebo for the acute treatment of psychosis and behavioral disturbances in hospitalized, demented patients. Am J Psychiatry 2002; 159(3):460-5.
Status: Included

Pomara N, Block R, Abraham J. Combined cholinergic precursor treatment and dihydroergotoxine mesylate in Alzheimer's disease. *IRCS Med Sci* 1983; (12):1048-9.
Status: Not included because dementia population not randomized to treatment

Pomara N, Block R, Moore N, et al. Combined Piracetam and cholinergic precursor treatment for primary degenerative dementia. *IRCS Med Sci Psychol Psychiatry* 1984; (5-6):388-6.
Status: Cross-over trial;
Cross-over trial

Pomara N, Roberts R, Rhiew HB, et al. Multiple, single-dose naltrexone administrations fail to effect overall cognitive functioning and plasma cortisol in individuals with probable Alzheimer's disease. *Neurobiol Aging* 1985; 6(3):233-6.
Status: Cross-over trial;
Cross-over trial

Pomara N, Deptula D, Singh R. Pretreatment postural blood pressure drop as a possible predictor of response to the cholinesterase inhibitor velnacrine (HP 029) in Alzheimer's disease. *Psychopharmacol Bull* 1991; 27(3):301-7.
Status: Not included because dementia population not randomized to treatment

Pomara N, Doraiswamy PM, Tun H, et al. Mifepristone (RU 486) for Alzheimer's disease. *Neurology* 2002; 58(9):1436.
Status: Not included because Jadad Quality Scale score less than three

Pomeroy VM. The effect of physiotherapy input on mobility skills of elderly people with severe dementing illness. *Clin Rehabil* 1993; 7(2):163-70.
Status: Not included because does not meet criteria for treatment for dementia patients

Pomeroy VM. Immobility and severe dementia: When is physiotherapy treatment appropriate? *Clin Rehabil* 1994; 8:226-32.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pomeroy VM, Warren CM, Honeycombe C, et al. Mobility and dementia: Is physiotherapy treatment during respite care effective? *Int J Geriatr Psychiatry* 1999 May; 14(5):389-97.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Popa R, Schneider F, Mihalas G, et al. Antagonic-Stress superiority versus meclofenoxate in gerontopsychiatry. *Arch Gerontol Geriatr* 1994; 18(Suppl 4):197-206.
Status: Included

Por CP, Evans MF. Does Ginkgo help delay dementia? *Can Fam Physician* 1998 May; 44:997-9.
Status: Companion of an included article

Porsteinsson AP, Tariot PN, Erb R, et al. Placebo-controlled study of divalproex sodium for agitation in dementia. *Am J Geriatr Psychiatry* 2001; 9(1):58-66.
Status: Included

Portegies P, Enting RH, de Jong MD, et al. AIDS dementia complex and didanosine. *Lancet* 1994 Sep 10; 344(8924):759
Status: Not included because not a full article

Porter RJ, Lunn BS, Walker LL, et al. Cognitive deficit induced by acute tryptophan depletion in patients with Alzheimer's disease. *Am J Psychiatry* 2000 Apr; 157(4):638-40.
Status: Not included because dementia population not randomized to treatment

Porter RJ, Marshall EF, O'Brien JT. Effects of rapid tryptophan depletion on salivary and plasma cortisol in Alzheimer's disease and the healthy elderly. *J Psychopharmacol (Oxf)* 2002 Mar; 16(1):73-8.
Status: Not included because does not meet criteria for treatment for dementia patients

Postiglione A, Soricelli A, Cicerano U, et al. Effect of acute administration of L-acetyl carnitine on cerebral blood flow in patients with chronic cerebral infarct. *Pharmacol Res* 1991 Apr; 23(3):241-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Potkin SG, Fleming K, Jin Y, et al. Clozapine enhances neurocognition and clinical symptomatology more than standard neuroleptics. *J Clin Psychopharmacol* 2001; 21(5):479-83.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Potkin SG, Anand R, Fleming K, et al. Brain metabolic and clinical effects of rivastigmine in Alzheimer's disease. *Int J*

Neuropsychopharmacol 2001 Sep; 4(3):223-30.
Status: Included

Potkin SG, Alva G, Keator D, et al. Brain metabolic effects of Neotrofin in patients with Alzheimer's disease. Brain Res 2002; 951(1):87-95.
Status: Not included because dementia population not randomized to treatment

Potkin SG, Anand R, Hartman R, et al. Impact of Alzheimer's disease and rivastigmine treatment on activities of daily living over the course of mild to moderately severe disease. Prog Neuropsychopharmacol Biol Psychiatry 2002; 26(4):713-20.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Potter A, Corwin J, Lang J, et al. Acute effects of the selective cholinergic channel activator (nicotinic agonist) ABT-418 in Alzheimer's disease. Psychopharmacologia 1999 Mar; 142(4):334-42.
*Status: Cross-over trial;
Cross-over trial*

Prasher VP, Huxley A, Haque MS, et al. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Down syndrome and Alzheimer's disease: Pilot study. Int J Geriatr Psychiatry 2002 Mar; 17(3):270-8.
Status: Included

Pratt RD, Perdomo CA, Surick IW, et al. Donepezil: Tolerability and safety in Alzheimer's disease. Int J Clin Pract 2002 Nov; 56(9):710-7.
Status: Not included because dementia population not randomized to treatment

Pratt RD. Patient populations in clinical trials of the efficacy and tolerability of donepezil in patients with vascular dementia. J Neurol Sci 2002 Nov 15; 203-204:57-65.:57-65.
Status: Not included because no extractable data relevant to review

Pratt RD, Perdomo CA. Donepezil-treated patients with probable vascular dementia demonstrate cognitive benefits. Ann N Y Acad Sci 2002 Nov; 977:513-22.:513-22.
Status: Included;

Predescu V, Riga D, Riga S, et al. Antagonistic-stress. A new treatment in gerontopsychiatry and for a healthy productive life. Ann N Y Acad Sci

1994; 717(Jun 30):315-31.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Prentice N, Van BM, Dougall NJ, et al. A double-blind, placebo-controlled study of tacrine in patients with Alzheimer's disease using SPET. J Psychopharmacol (Oxf) 1996; 10(3):175-81.
Status: Included

Preston GC, Brazell C, Ward C, et al. The scopolamine model of dementia: Determination of central cholinomimetic effects of physostigmine on cognition and biochemical markers in man. J Psychopharmacol (Oxf) 1988; 2(2):67-79.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Preston GC, Ward C, Lines CR, et al. Scopolamine and benzodiazepine models of dementia: Cross-reversals by Ro 15-1788 and physostigmine. Psychopharmacologia 1989; 98(4):487-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Proctor R, Burns A, Powell HS, et al. Behavioural management in nursing and residential homes: a randomised controlled trial. Lancet 1999 Jul 3; 354(9172):26-9.
Status: Not included because does not meet criteria for treatment for dementia patients

Pryse-Phillips W. A drug to control behavioural disturbances of dementia. Mature Medicine Canada 2000; 3(2):58.
Status : Companion of an included article

Puchler K, Schaffler K, Plenker A. The comparative effects of single and multiple doses of RS-8359, moclobemide and placebo on psychomotor function in healthy subjects. Int Clin Psychopharmacol 1997; Vol 12(Suppl 5):S17-S23.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Purandare N, Bloom C, Page S, et al. The effect of anticholinesterases on personality changes in Alzheimer's disease. Aging Ment Health 2002; 6(4):350-4.
Status: Not included because dementia population not randomized to treatment

Puri BK, Bydder GM, Counsell SJ, et al. MRI and neuropsychological improvement in Huntington

disease following ethyl-EPA treatment. NeuroReport 2002; 13(1):123-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Putt ME, Ravina B. Randomized, placebo-controlled, parallel group versus crossover study designs for the study of dementia in Parkinson's disease. Control Clin Trials 2002 Apr; 23(2):111-26.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Qizilbash N, Whitehead A, Higgins J, et al. Cholinesterase inhibition for Alzheimer disease: a meta-analysis of the tacrine trials. Dementia Trialists' Collaboration.[comment]. JAMA 1998 Nov 25; 280(20):1777-82.
Status: Background article

Quayhagen MP, Quayhagen M, Corbeil RR, et al. A dyadic remediation program for care recipients with dementia. Nurs Res 1995 May; 44(3):153-9.
Status: Not included because does not meet criteria for treatment for dementia patients

Quayhagen MP, Quayhagen M, Corbeil RR, et al. Coping with dementia: Evaluation of four nonpharmacologic interventions. Int Psychogeriatr 2000 Jun; 12(2):249-65.
Status: Not included because does not meet criteria for treatment for dementia patients

Quayhagen MP, Quayhagen M. Testing of a cognitive stimulation intervention for dementia caregiving dyads. Neuropsychol Rehab 2001; 11(Special issue (3-4)):319-22.
Status: Not included because does not meet criteria for treatment for dementia patients

Quinlivan R, Hough R, Crowell A, et al. Service utilization and costs of care for severely mentally ill clients in an intensive case management program. Psychiatr Serv 1995 Apr; 46(4):365-71.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rabey JM, Neufeld MY, Treves TA, et al. Cognitive effects of scopolamine in dementia. J Neural Transm Gen Sect 1996; 103(7):873-81.
Status: Cross-over trial;
Cross-over trial

Rada RT, Kellner R. Thiothixene in the treatment of geriatric patients with chronic organic brain syndrome. J Am Geriatr Soc 1976 Mar;

24(3):105-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rader MA, Alston JB, Ellis DW. Sensory stimulation of severely brain-injured patients. Brain Inj 1989 Apr; 3(2):141-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Raffaele KC, Berardi A, Asthana S, et al. Effects of long-term continuous infusion of the muscarinic cholinergic agonist arecoline on verbal memory in dementia of the Alzheimer type. Psychopharmacol Bull 1991; 27(3):315-9.
Status: Not included because dementia population not randomized to treatment

Raffaele KC, Asthana S, Berardi A, et al. Differential response to the cholinergic agonist arecoline among different cognitive modalities in Alzheimer's disease. Neuropsychopharmacology 1996; 15(2):163-70.
Status: Not included because dementia population not randomized to treatment

Ragneskog H, Kihlgren M, Karlsson I, et al. Dinner music for demented patients: Analysis of video-recorded observations. Clin Nurs Res 1996 Aug; 5(3):262-77, 278-82.
Status: Not included because dementia population not randomized to treatment

Ragneskog H, Brane G, Karlsson I, et al. Influence of dinner music on food intake and symptoms common in dementia. Scand J Caring Sci 1996; 10(1):11-7.
Status: Not included because dementia population not randomized to treatment

Ragneskog H, Asplund K, Kihlgren M, et al. Individualized music played for agitated patients with dementia: Analysis of video-recorded sessions. Int J Nurs Pract 2001 Jun; 7(3):146-55.
Status: Not included because dementia population not randomized to treatment

Rai G, Wright G, Scott L, et al. Double-blind, placebo-controlled study of acetyl-L-carnitine in patients with Alzheimer's disease. Curr Med Res Opin 1990; 11(10):638-47.
Status: Included

Rai GS, Shovlin C, Wesnes KA. A double-blind, placebo controlled study of Ginkgo biloba extract ('tanakan') in elderly outpatients with mild to

moderate memory impairment. *Curr Med Res Opin* 1991; 12(6):350-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rained M, Mucke HAM. Long-term cognitive benefit from galanthamine in Alzheimer's disease. *Int J Geriatr Psychopharmacol* 1998; 1(4):197-201.

Status: Not included because dementia population not randomized to treatment

Rainer M, Mucke HA, Chwatal K, et al. Alcohol-induced organic cerebral psychosyndromes: Partial reversal of cognitive impairments assisted by dihydroergocristine. *Psychopharmacologia* 1996 Oct; 127(4):365-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Randolph C, Roberts JW, Tierney MC, et al. D-cycloserine treatment of Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1994; 8(3):198-205.

Status: Cross-over trial;
Cross-over trial

Ransmayr G, Plorer S, Gerstenbrand F, et al. Double-blind placebo-controlled trial of phosphatidylserine in elderly patients with arteriosclerotic encephalopathy. *Clin Trials J* 1987; 24(1):62-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rao DB, Georgiev EL, Paul PD, et al. Cycandelate in the treatment of senile mental changes: A double-blind evaluation. *J Am Geriatr Soc* 1977 Dec; 25(12):548-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rapp S, Brenes G, Marsh AP. Memory enhancement training for older adults with mild cognitive impairment: A preliminary study. *Aging Ment Health* 2002 Feb; 6(1):5-11.

Status: Not included because does not meet criteria for treatment for dementia patients

Rascol O, Sieradzan K, Peyro-Saint-Paul H, et al. Efaroxan, an alpha-2 antagonist, in the treatment of progressive supranuclear palsy. *Mov Disord* 1998 Jul; 13(4):673-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Raskind MA, Sadowsky CH, Sigmund WR, et al. Effect of tacrine on language, praxis, and noncognitive behavioral problems in Alzheimer's disease. *Arch Neurol* 1997 Jul; 54(7):836-40.

Status: Companion of an included article

Raskind MA, Cyrus PA, Ruzicka BB, et al. The effects of metrifonate on the cognitive, behavioral, and functional performance of Alzheimer's disease patients. *J Clin Psychiatry* 1999; 60(5):318-25.

Status: Included

Raskind MA, Peskind ER, Wessel T, et al. Galantamine in AD: A 6-month randomized, placebo-controlled trial with a 6-month extension. The Galantamine USA-1 Study Group. *Neurology* 2000 Jun 27; 54(12):2261-8.

Status: Included

Rasmusen L, Yan B, Robillard A, et al. Effects of washout and dose-escalation periods on the efficacy, safety, and tolerability of galantamine in patients previously treated with donepezil: Ongoing clinical trials. *Clin Ther* 2001; 23(Suppl):A25-A30.

Status: Not included because dementia population not randomized to treatment

Ray PG, Meador KJ, Loring DW, et al. Effects of scopolamine on visual evoked potentials in aging and dementia. *Electroencephalogr Clin Neurophysiol* 1991 Sep; 80(5):347-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ray WA, Taylor JA, Meador KG, et al. Reducing antipsychotic drug use in nursing homes. A controlled trial of provider education. *Arch Intern Med* 1993; 153(6):713-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Raz N, Torres IJ, Briggs SD, et al. Selective neuroanatomic abnormalities in Down's syndrome and their cognitive correlates: Evidence from MRI morphometry. *Neurology* 1995 Feb; 45(2):356-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Reddy H, De Stefano N, Mortilla M, et al. Functional reorganization of motor cortex increases with greater axonal injury from CADASIL. *Stroke* 2002 Feb; 33(2):502-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Reeve W. Use of environmental manipulation and classroom and modified informal reality orientation with institutionalized, confused elderly patients. *Age Ageing* 1985; 14(2):119-21.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Regland B, Lehmann W, Abedini I, et al. Treatment of Alzheimer's disease with clioquinol. *Dement Geriatr Cogn Disord* 2001 Nov; 12(6):408-14.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rehman SA. Two trials comparing 'Hydergine' with placebo in the treatment of patients suffering from cerebrovascular insufficiency. *Curr Med Res Opin* 1973; 1(8):456-62.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Reichman WE. New therapeutic options for patients with Alzheimer's Disease. *Ann Gen Hosp Psychiatry* 2003; 2(1):1-42.
Status: Background article

Reid WG. The evolution of dementia in idiopathic Parkinson's disease: Neuropsychological and clinical evidence in support of subtypes. *Int Psychogeriatr* 1992; 4(Suppl 2):147-60.
Status: Background article

Reifler BV, Teri L, Raskind M, et al. Double-blind trial of imipramine in Alzheimer's disease patients with and without depression. *Am J Psychiatry* 1989 Jan; 146(1):45-9.
Status: Included

Reisberg B, Ferris SH, Schneck MK. Piracetam in the treatment of cognitive impairment in the elderly. *Drug Dev Res* 1982; 2(5):475-80.
*Status: Cross-over trial;
Cross-over trial*

Reisberg B, Ferris SH, Anand R. Effects of naloxone in senile dementia: A double-blind trial. *N Engl J Med* 1983; 308(12):721-2.
Status: Not included because not a full article

Reisberg B, Ferris SH, de Leon MJ, et al. Global Deterioration Scale (GDS). *Psychopharmacol*

Bull 1988; 24(4):661-3.
Status: Background article

Remington R. Calming music and hand massage with agitated elderly. *Nurs Res* 2002 Sep; 51(5):317-23.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Richard F, Helbecque N, Neuman E, et al. APOE genotyping and response to drug treatment in Alzheimer's disease. *Lancet* 1997 Feb 22; 349(9051):539
Status: Not included because Jadad Quality Scale score less than three

Riedel-Heller SG, Busse A, Aurich C, et al. Prevalence of dementia according to DSM-III-R and ICD-10: Results of the Leipzig Longitudinal Study of the Aged (LEILA75+) Part 1. *Br J Psychiatry* 2001 Sep; 179:250-4.
Status: Background article

Riekkinen M, Laakso MP, Jakala P, et al. Clonidine impairs sustained attention and memory in Alzheimer's disease. *Neuroscience* 1999; 92(3):975-82.
Status: Not included because dementia population not randomized to treatment

Riekkinen P, Jr., Kuikka J, Soininen H, et al. Tetrahydroaminoacridine modulates technetium-99m labelled ethylene dicysteinate retention in Alzheimer's disease measured with single photon emission computed tomography imaging. *Neurosci Lett* 1995 Jul 28; 195(1):53-6.
Status: Not included because no extractable data relevant to review

Riekkinen P, Jr., Riekkinen M, Soininen H, et al. Frontal dysfunction blocks the therapeutic effect of THA on attention in Alzheimer's disease. *NeuroReport* 1997 May 27; 8(8):1845-9.
Status: Not included because dementia population not randomized to treatment

Riekkinen PJ, Koivisto K, Helkala EL, et al. Long-term, double-blind trial of selegiline in Alzheimer's disease. *Neurobiol Aging* 1994; 15(Suppl 1):67.
Status: Not included because not a full article

Riekkinen PJ, Soininen H, Helkala E-L, et al. Hippocampal atrophy, acute THA treatment and memory in Alzheimer's disease. *NeuroReport* 1995; 6(9):1297-300.

Status: Not included because dementia population not randomized to treatment

Riekkinen PJr, Paakkonen A, Karhu J, et al. THA disrupts mismatch negativity in Alzheimer's disease. *Psychopharmacologia* 1997 Sep; 133(2):203-6.

Status: Not included because no extractable data relevant to review

Riekkinen PJr, Soininen H, Partanen J, et al. The ability of THA treatment to increase cortical alpha waves is related to apolipoprotein E genotype of Alzheimer disease patients. *Psychopharmacologia* 1997 Feb; 129(3):285-8.
Status: Not included because dementia population not randomized to treatment

Riekkinen PJr, Riekkinen M. THA improves word priming and clonidine enhances fluency and working memory in Alzheimer's disease. *Neuropsychopharmacology* 1999 Apr; 20(4):357-64.
Status: Not included because dementia population not randomized to treatment

Rigaud A-S, Traykov L, Latour F, et al. Presence or absence of at least one epsilon4 allele and gender are not predictive for the response to donepezil treatment in Alzheimer's disease. *Pharmacogenetics* 2002; 12(5):415-20.
Status: Not included because dementia population not randomized to treatment

Rigaud AS, Andre G, Vellas B, et al. No additional benefit of HRT on response to rivastigmine in menopausal women with AD. *Neurology* 2003; 60(1):148-50.
Status: Not included because Jadad Quality Scale score less than three

Rimon R, Rakkolainen V. Lithium iodide in the treatment of confusional states. *Br J Psychiatry* 1968 Jan; 114(506):109-10.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rinne JO, Portin R, Ruottinen H, et al. Cognitive impairment and the brain dopaminergic system in Parkinson's disease: Fluorodopa positron emission tomographic study. *Arch Neurol* 2000 Apr; 57(4):470-5.
Status: Not included because dementia population not randomized to treatment

Rinsky JR, Wikler A, Way JG, et al. 'Mental set' in controls, postalcoholics, chronic schizophrenics, and 'organics'. *Biol Psychiatry* 1979; 14(6):881-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Riordan JM, Bennet AV. An evaluation of an augmented domiciliary service to older people with dementia and their carers. *Aging Ment Health* 1998; 2(2):137-43.
Status: Not included because dementia population not randomized to treatment

Ritchie K, Artero S, Touchon J. Classification criteria for mild cognitive impairment: a population-based validation study. *Neurology* 2001 Jan 9; 56(1):37-42.
Status: Background article

Rivera VM, Meyer JS, Baer PE, et al. Vertebrobasilar arterial insufficiency with dementia. Controlled trials of treatment with betahistine hydrochloride. *J Am Geriatr Soc* 1974 Sep; 22(9):397-406.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Riviere S, Gillette-Guyonnet S, Voisin T, et al. A nutritional education program could prevent weight loss and slow cognitive decline in Alzheimer's disease. *J Nutr Health Aging* ; 5(4):295-9.
Status: Not included because dementia population not randomized to treatment

Rizzo JA, Bogardus ST, Jr., Leo-Summers L, et al. Multicomponent targeted intervention to prevent delirium in hospitalized older patients: What is the economic value? *Med Care* 2001 Jul; 39(7):740-52.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rizzo M, Ficola U, Marozzi P, et al. Effects of aniracetam on clinical pattern and cerebral blood flow of old patients with degenerative dementia (SDAT). *J Neurol* 1994; 241:S126.
Status: Not included because not a full article

Robbins TW, Semple J, Kumar R, et al. Effects of scopolamine on delayed-matching-to-sample and paired associates tests of visual memory and learning in human subjects: Comparison with diazepam and implications for dementia. *Psychopharmacologia* 1997 Nov; 134(1):95-106.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Robert PH, Allain H. Clinical management of agitation in the elderly with tiapride. *Eur Psychiatry* 2001 Jan; 16(Suppl):47S.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Roberts CJ, Ford JM, Truman CA, et al. Assessment of the value of therapeutic monitoring of tacrine in Alzheimer's disease. *Eur J Clin Pharmacol* 1998 Nov; 54(9-10):721-10.

Status: Not included because dementia population not randomized to treatment

Roberts RJ, Musick BS, Olley B, et al. Data management in a longitudinal cross-cultural study. *Stat Med* 2000 Jun 15; 19(11-12):1645-9.

Status: Background article

Robichaud L, Hebert R, Desrosiers J. Efficacy of a sensory integration program on behaviors of inpatients with dementia. *Am J Occup Ther* 1994 Apr; 48(4):355-60.

Status: Not included because does not meet criteria for treatment for dementia patients

Rocca WA, van Duijn CM, Clayton D, et al. Maternal age and Alzheimer's disease: a collaborative re-analysis of case-control studies. EURODEM Risk Factors Research Group. *Int J Epidemiol* 1991; 20 Suppl 2:S21-S27

Status: Background article

Rochtchina I, Gavrilova S, Kolykhalov I, et al. Neuropsychological assessment of modification of efficacy of cholinergic therapy by preceding cerebrolysin treatment in Alzheimer's disease. *J Eur Coll Neuropsychopharmacol* 1999; (Suppl 5):S330.

Status: Not included because not a full article

Rockwood K, Stolee P, Howard K, et al. Use of Goal Attainment Scaling to measure treatment effects in an anti-dementia drug trial. *Neuroepidemiology* 1996; 15(6):330-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rockwood K, Beattie BL, Eastwood MR, et al. A randomized, controlled trial of linopirdine in the treatment of Alzheimer's disease. *Can J Neurol Sci* 1997 May; 24(2):140-5.

Status: Included

Rockwood K, Stolee P. Responsiveness of outcome measures used in an antidementia drug trial. *Alzheimer Dis Assoc Disord* 2000 Jul; 14(3):182-5.

Status: Companion of an included article; Background article

Rockwood K, Hogan DB, MacKnight C. Conceptualisation and measurement of frailty in elderly people. *Drugs Aging* 2000 Oct; 17(4):295-302.

Status: Background article

Rockwood K, Mintzer J, Truyen L, et al. Effects of a flexible galantamine dose in Alzheimer's disease: a randomised, controlled trial. *J Neurol Neurosurg Psychiatry* 2001; 71(5):589-95.

Status: Included

Rockwood K, MacKnight C. Assessing the clinical importance of statistically significant improvement in anti-dementia drug trials. *Neuroepidemiology* 2001 May; 20(2):51-6.

Status: Background article

Rockwood K, Graham JE, Fay S, et al. Goal setting and attainment in Alzheimer's disease patients treated with donepezil. *J Neurol Neurosurg Psychiatry* 2002 Nov; 73(5):500-7.

Status: Not included because dementia population not randomized to treatment

Rockwood K, Joffres C. Improving clinical descriptions to understand the effects of dementia treatment: Consensus recommendations. *Int J Geriatr Psychiatry* 2002; 17(11):1006-11.

Status: Background article

Rodriguez-Martin JL, Lopez Arrieta JM, Qizilbash N. Thiamine for Alzheimer's disease. In: *The Cochrane Library*, 2000. Issue 2. Oxford: Update Software.

Status: Background article

Rogers J, Kirby LC, Hempelman SR, et al. Clinical trial of indomethacin in Alzheimer's disease. *Neurology* 1993 Aug; 43(8):1609-11.

Status: Included

Rogers JD, Sanchez-Saffon A, Frol AB, et al. Elevated plasma homocysteine levels in patients treated with levodopa: Association with vascular disease. *Arch Neurol* 2003; 60(1):59-64.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rogers JF, Morrison AL, Nafziger AN, et al. Flumazenil reduces midazolam-induced cognitive impairment without altering pharmacokinetics. *Clin Pharmacol Ther* 2002; 72(6):711-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rogers SL, Doody R, Mohs R, et al. E2020 produces both clinical, global and cognitive test improvement in patients with mild to moderately severe Alzheimer's disease: Results of a 30-week phase III trial. *Neurology* 1996; 46(Suppl):217
Status: Not included because not a full article

Rogers SL, Friedhoff LT. The efficacy and safety of donepezil in patients with Alzheimer's disease: Results of a US Multicentre, Randomized, double-blind, placebo-controlled trial. The Donepezil Study Group. *Dementia* 1996 Nov; 7(6):293-303.
Status: Included

Rogers SL, Cooper NM, Sukovaty R, et al. Pharmacokinetic and pharmacodynamic profile of donepezil HCl following multiple oral doses. *Br J Clin Pharmacol* 1998 Nov; 46(Suppl 1):7-12.
Status: Background article

Rogers SL, Doody RS, Mohs RC, et al. Donepezil improves cognition and global function in Alzheimer disease: A 15-week, double-blind, placebo-controlled study. *Arch Intern Med* 1998a; 158(9):1021-31.
Status: Included

Rogers SL, Friedhoff LT. Long-term efficacy and safety of donepezil in the treatment of Alzheimer's disease: An interim analysis of the results of a US multicentre open label extension study. *Eur Neuropsychopharmacol* 1998c; 8(1):67-75.
Status: Companion of an included article

Rogers SL, Farlow MR, Doody RS, et al. A 24-week, double-blind, placebo-controlled trial of donepezil in patients with Alzheimer's disease. *Neurology* 1998b; 50(1):136-45.
Status: Included

Rogers SL, Doody RS, Pratt RD, et al. Long-term efficacy and safety of donepezil in the treatment of Alzheimer's disease: Final analysis of a US multicentre open-label study. *Eur Neuropsychopharmacol* 2000 May; 10(3):195-203.
Status: Companion of an included article

Roman GC, Tatemichi TK, Erkinjuntti T, et al. Vascular dementia: diagnostic criteria for research studies. Report of the NINDS-AIREN International Workshop. *Neurology* 1993 Feb; 43(2):250-60.
Status: Background article

Rombouts SA, Barkhof F, Van Meel CS, et al. Alterations in brain activation during cholinergic enhancement with rivastigmine in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2002 Dec; 73(6):665-71.
Status: Not included because dementia population not randomized to treatment

Ronnberg L. Quality of life in nursing-home residents: An intervention study of the effect of mental stimulation through an audiovisual programme. *Age Ageing* 1998; 27(3):393-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosen HJ. Mental decline in the elderly: Pharmacotherapy (ergot alkaloids versus papaverine). *J Am Geriatr Soc* 1975 Apr; 23(4):169-74.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosen WG, Mohs RC, Davis KL. A new rating scale for Alzheimer's disease. *Am J Psychiatry* 1984 Nov; 141(11):1356-64.
Status: Background article

Rosenberg SJ, Ryan JJ, Prifitera A. Rey Auditory-Verbal Learning Test performance of patients with and without memory impairment. *J Clin Psychol* 1984; (3):785-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosenzweig P, Patat A, Zieleniuk I, et al. Cognitive performance in elderly subjects after a single dose of befloxatone, a new reversible selective monoamine oxidase A inhibitor. *Clin Pharmacol Ther* 1998 Aug; 64(2):211-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosewarne R, Bruce A, McKenna M. Dementia programme effectiveness in long-term care. *Int J Geriatr Psychiatry* 1997 Feb; (2):173-82.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosler, M., Dennler, et al. A double-blind placebo controlled study of ENA 713 in Alzheimer's

disease (DAT). *Pharmacopsychiatry* 1997; 30:212.

Status: Article not retrievable

Rosler M, Retz W, Retz-Junginger P, et al. Effects of two-year treatment with the cholinesterase inhibitor rivastigmine on behavioural symptoms in Alzheimer's disease. *Behav Neurol* 1998; (4):211-6.

Status: Companion of an included article

Rosler M, Anand R, Cicin-Sain A, et al. Efficacy and safety of rivastigmine in patients with Alzheimer's disease: International randomised controlled trial. *BMJ* 1999 Mar 6; 318(7184):633-8.

Status: Included

Rosler M. Erratum: Efficacy and safety of rivastigmine in patients with Alzheimer's disease: International randomised controlled trial. *BMJ* 2001; 322(7300):1456.

Status: Companion of an included article

Rossi R, Inzitari D, Pantoni L, et al. Nimodipine in subcortical vascular dementia trial. *Alzheimer Dis Assoc Disord* 1999; 13(Suppl 3):S159-S165.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rostow CD. The effect of self- vs. external-monitoring and locus of control upon the pacing and general adjustment of psychiatric inpatients. *Behav Res Ther* 1980; 18(6):541-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Roth M, Mountjoy CQ, Amrein R. Moclobemide in elderly patients with cognitive decline and depression: An international double-blind, placebo-controlled trial. *Br J Psychiatry* 1996 Feb; 168(2):149-57.

Status: Included

Rother M, Kittner B, Rudolphi K, et al. HWA 285 (propentofylline) - A new compound for the treatment of both vascular dementia and dementia of the Alzheimer type. *Ann N Y Acad Sci* 1996; 777:404-9.

Status: Not included because dementia population not randomized to treatment

Rother M. Long-term effects of propentofylline in patients with Alzheimer's disease: A 72-week, placebo-controlled study assessing safety, efficacy, and impact on disease progression. *J*

Cereb Blood Flow Metab 1999; 19(Suppl 1):18.

Status: Not included because not a full article

Rouy JM, Douillon AM, Compan B, et al. Ergoloid mesylates ('Hydergine') in the treatment of mental deterioration in the elderly: A 6-month double-blind, placebo-controlled trial. *Curr Med Res Opin* 1989; 11(6):380-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rovner BW, Steele CD, Shmueli Y, et al. A randomized trial of dementia care in nursing homes. *J Am Geriatr Soc* 1996 Jan; 44(1):7-13.

Status: Not included because does not meet criteria for treatment for dementia patients

Royall DR, Lauterbach EC, Cummings JL, et al. Executive control function: a review of its promise and challenges for clinical research. A report from the Committee on Research of the American Neuropsychiatric Association. *J Neuropsychiatry Clin Neurosci* 2002; 14(4):377-405.

Status: Background article

Rozzini R, Zanetti O, Bianchetti A. Effectiveness of oxiracetam therapy in the treatment of cognitive deficiencies secondary to primary degenerative dementia. *Acta Neurol (Napoli)* 1992 Apr; 14(2):117-26.

Status: Included

Rozzini R, Zanetti O, Bianchetti A. Treatment of cognitive impairment secondary to degenerative dementia. Effectiveness of oxiracetam therapy. *Acta Neurol (Napoli)* 1993 Feb; 15(1):44-52.

Status: Companion of an included article

Rozzini R, Ferrucci L, Losonczy K, et al. Protective effect of chronic NSAID use on cognitive decline in older persons. *J Am Geriatr Soc* 1996; 44(9):1025-9.

Status: Not included because dementia population not randomized to treatment

Ruberg S, Cairns V. Providing evidence of efficacy for a new drug. *Stat Med* 1998 Aug 15; 17(15-16):1813-23.

Status: Background article

Rudman D, Racette D, Rudman IW, et al. Hyponatremia in tube-fed elderly men. *J Chronic Dis* 1986; 39(2):73-80.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ruether E, Ritter R, Apecechea M, et al. Efficacy of the peptidergic nootropic drug cerebrolysin in patients with senile dementia of the Alzheimer type (SDAT). *Pharmacopsychiatry* 1994 Jan; 27(1):32-40.

Status: Included

Ruether E, Husmann R, Kinzler E, et al. A 28-week, double-blind, placebo-controlled study with cerebrolysin in patients with mild to moderate Alzheimer's disease. *Int Clin Psychopharmacol* 2001; 16(6):372

Status: Not included because not a full article

Ruether E, Husmann R, Kinzler E, et al. A 28-week, double-blind, placebo-controlled study with Cerebrolysin in patients with mild to moderate Alzheimer's disease. *Int Clin Psychopharmacol* 2001 Sep; 16(5):253-63.

Status: Included

Ruether E, Alvarez XA, Rainer M, et al. Sustained improvement of cognition and global function in patients with moderately severe Alzheimer's disease: A double-blind, placebo-controlled study with the neurotrophic agent Cerebrolysin. *J Neural Transm Suppl* 2002; (62):265-75.

Status: Companion of an included article

Ruggiero, Ovallesco U. Clozapine and validation therapy in the treatment of dementia. *Funct Neurol* 1997; 3-4(12):240-412.

Status: Not included because dementia population not randomized to treatment

Ruther E, Glaser A, Bleich S, et al. A prospective PMS study to validate the sensitivity for change of the D-scale in advanced stages of dementia using the NMDA-antagonist memantine.

Pharmacopsychiatry 2000 May; 33(3):103-8.

Status: Article not retrievable

Ruther E, Ritter R, Apecechea M, et al. Sustained improvements in patients with dementia of Alzheimer's type (DAT) 6 months after termination of Cerebrolysin therapy. *J Neural Transm Gen Sect* 2000; 107(7):815-29.

Status: Companion of an included article

Ryden MB, Snyder M, Gross CR, et al. Value-added outcomes: The use of advanced practice nurses in long-term care facilities. *Gerontologist* 2000 Dec; 40(6):654-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Ryynanen OP, Myllykangas M, Kinnunen J, et al. Doctors' willingness to refer elderly patients for elective surgery. *Fam Pract* 1997 Jun; 14(3):216-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saarela T, Kiviharju U. Evaluating the usefulness of training in psychogeriatrics. *Int J Geriatr Psychiatry* 1995 Dec; (12):1019-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sabe L, Kuzis G, Garcia CA, et al. A randomized, double-blind, placebo-controlled study of idebenone in Alzheimer's disease (AD). *JNS* 1997; 150(Suppl):S296.

Status: Not included because not a full article

Sacktor N, Schifitto G, McDermott MP, et al. Transdermal selegiline in HIV-associated cognitive impairment: Pilot, placebo-controlled study. *Neurology* 2000 Jan 11; 54(1):233-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sacktor N, McDermott MP, Marder K, et al. HIV-associated cognitive impairment before and after the advent of combination therapy. *J Neurovirol* 2002; 8(2):136-42.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sahakian BJ, Coull JT. Tetrahydroaminoacridine (THA) in Alzheimer's disease: An assessment of attentional and mnemonic function using CANTAB. *Acta Neurol Scand Suppl* 1993; Vol 149:29-35.

Status: Cross-over trial;

Cross-over trial

Sahakian BJ, Owen AM, Morant NJ, et al. Further analysis of the cognitive effects of tetrahydroaminoacridine (THA) in Alzheimer's disease: Assessment of attentional and mnemonic function using CANTAB. *Psychopharmacologia* 1993; 110(4):395-401.

Status: Cross-over trial;

Cross-over trial

Sahakian BJ, Coull JT. Nicotine and tetrahydroaminoacridine: Evidence for improved attention in patients with dementia of the Alzheimer type. *Drug Dev Res* 1994; 31(1):80-8.

Status: Not included because dementia population not randomized to treatment

Sahin HA, Gurvit IH, Bilgic B, et al. Therapeutic effects of an acetylcholinesterase inhibitor (donepezil) on memory in Wernicke-Korsakoff's disease. Clin Neuropharmacol 2002 Jan; 25(1):16-20.

Status: Not included because dementia population not randomized to treatment

Saine K, Cullum CM, Martin-Cook K, et al. Comparison of functional and cognitive donepezil effects in Alzheimer's disease. Int Psychogeriatr 2002; 14(2):181-5.

Status: Not included because dementia population not randomized to treatment

Sajatovic M, Mullen JA, Sweitzer DE. Efficacy of quetiapine and risperidone against depressive symptoms in outpatients with psychosis. J Clin Psychiatry 2002; 63(12):1156-63.

Status: Not included because no extractable data relevant to review

Saldmann F, Funel A, Jacquet P. Efficacy of naftidrofuryl in patients with moderate senile dementia. Curr Med Res Opin 1991; 12(6):379-89.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saletu B, Linzmayer L, Grunberger J, et al. Double-blind, placebo-controlled, clinical, psychometric and neurophysiological investigations with oxiracetam in the organic brain syndrome of late life. Neuropsychobiology 1985; 13(1-2):44-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saletu B, Gruenberger J, Linzmayer L, et al. EEG brain mapping and psychometry in age-associated memory impairment after acute and 2-week infusions with the hemoderivative Actovegin®: Double-blind, placebo-controlled trials. Neuropsychobiology 1990; 24(3):135-48.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saletu B, Semlitsch HV, Anderer P, et al. On acute central effects of amantadine infusion in mild dementia: Double-blind placebo-controlled EEG/ERP-brain mapping and psychometric studies. Pharmacopsychiatry 1990; 25:90.

Status: Article not retrievable

Saletu B, Moller HJ, Grunberger J, et al. Propentofylline in adult-onset cognitive disorders:

double-blind, placebo-controlled, clinical, psychometric and brain mapping studies. Neuropsychobiology 1990 Sep; 24(4):173-84.
Status: Included

Saletu B, Anderer P, Semlitsch HV, et al. Amantadine infusions in mild dementia: Acute double-blind placebo-controlled EEG mapping and psychometric studies. Arch Gerontol Geriatr 1992; 15(1):43-58.

*Status: Cross-over trial;
Cross-over trial*

Saletu B, Anderer P, Fischhof PK, et al. EEG mapping and psychopharmacological studies with denbufylline in SDAT and MID. Biol Psychiatry 1992 Oct 15; 32(8):668-81.

Status: Not included because Jadad Quality Scale score less than three

Saletu B, Paulus E, Linzmayer L, et al. Nicergoline in senile dementia of Alzheimer type and multi-infarct dementia: A double-blind, placebo-controlled, clinical and EEG/ERP mapping study. Psychopharmacologia 1995 Feb; 117(4):385-95.

Status: Included

Saletu B, Anderer P, Semlitsch HV. Relations between symptomatology and brain function in dementias: Double-blind, placebo-controlled, clinical and EEG/ERP mapping studies with nicergoline. Dement Geriatr Cogn Disord 1997; Vol 8(Suppl 1):12-21.

Status: Included

Saletu M, Grunberger J, Saletu B, et al. Accelerated remission of the alcoholic organic brain syndrome with EMD 21657. Double-blind clinical and psychometric trials. Arzneimittelforschung 1978; 28(9):1525-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Salib E, Sheridan T, Allington M. Ventricular measurements in computed tomography of responders and non-responders to donepezil in the treatment of Alzheimer's disease. Int J Psychiatry Clin Pract 2001; 5(3):189-94.
Status: Not included because dementia population not randomized to treatment

Salvioli G, Neri G. L-acetylcarnitine treatment of mental decline in the elderly. Drugs Exp Clin Res 1994; 20(4):169-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Samorajski T, Voulis GA, Smith RC. Piracetam plus lecithin trials in senile dementia of the Alzheimer type. *Ann N Y Acad Sci* 1985; 444:478-81.

Status: Not included because dementia population not randomized to treatment

Samuel W, Caligiuri M, Galasko D, et al. Better cognitive and psychopathologic response to donepezil in patients prospectively diagnosed as dementia with Lewy bodies: A preliminary study. *Int J Geriatr Psychiatry* 2000 Sep; 15(9):794-802.
Status: Not included because dementia population not randomized to treatment

Sanders DS, Carter MJ, D'Silva J, et al. Survival analysis in percutaneous endoscopic gastrostomy feeding: A worse outcome in patients with dementia. *Am J Gastroenterol* 2000; 95(6):1472-5.
Status: Not included because dementia population not randomized to treatment

Sano M, Stern Y, Marder K, et al. A controlled trial of piracetam in intellectually impaired patients with Parkinson's disease. *Mov Disord* 1990; 5(3):230-4.
Status: Not included because no extractable data relevant to review

Sano M, Bell K, Cote L, et al. Double-blind parallel design pilot study of acetyl levocarnitine in patients with Alzheimer's disease. *Arch Neurol* 1992 Nov; 49(11):1137-41.
Status: Included

Sano M, Bell K, Marder K, et al. Safety and efficacy of oral physostigmine in the treatment of Alzheimer's disease. *Clin Neuropharmacol* 1993 Feb; 16(1):61-9.
*Status: Cross-over trial;
Cross-over trial*

Sano M, Growdon J, Klauber M, et al. Expanding the severity range of patients in clinical trials for Alzheimer's disease: A multicentre clinical trial of Selegiline and alpha-tocopherol. *Neurology* 1995; 45(Suppl 4):A289.
Status: Not included because not a full article

Sano M, Ernesto C, Klauber MR, et al. Rationale and design of a multicenter study of selegiline and alpha-tocopherol in the treatment of Alzheimer's

disease using novel clinical outcomes. *Alzheimer Dis Assoc Disord* 1996; 10(3):132-40.
Status: Companion of an included article

Sano M, Ernesto C, Thomas RG, et al. A controlled trial of selegiline, alpha-tocopherol, or both as treatment for Alzheimer's disease. The Alzheimer's Disease Cooperative Study. *N Engl J Med* 1997; 336(17):1216-22.
Status: Included

Sano M, Ernesto C, Thomas RG, et al. Effects of Selegiline and alpha-Tocopherol on cognitive and functional outcome measures in moderately impaired patients with Alzheimer's disease. *Neurology* 1997; 48(Suppl):A377-A378
Status: Not included because not a full article

Sano M, Bell K, Jacobs D. Cognitive effects of estrogens in women with cardiac disease: What we do not know. *Am J Med* 2002; 113(7):612-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sansom G. Comparing multisensory stimulation with (1) tactile stimulation and (2) themed reminiscence. *Journal of Dementia Care* 2002; 10(4):38
Status: Not included because not a full article

Satlin A, Volicer L, Ross V, et al. Bright light treatment of behavioral and sleep disturbances in patients with Alzheimer's disease. *Am J Psychiatry* 1992; 149(8):1028-32.
Status: Not included because dementia population not randomized to treatment

Satlin A, Bodick N, Offen WW, et al. Brain proton magnetic resonance spectroscopy (1H-MRS) in Alzheimer's disease: Changes after treatment with xanomeline, an M1 selective cholinergic agonist. *Am J Psychiatry* 1997 Oct; 154(10):1459-61.
Status: Companion of an included article

Sato K, Kamiya S, Okawa M, et al. Effect of transcranial electrostimulation on EEG component waves of elderly patients with dementia. *J Brain Sci* 1998; 24(1-2):65-72.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saver JL, Kalafut M. Combination therapies and the theoretical limits of evidence-based medicine. *Neuroepidemiology* 2001 May; 20(2):57-64.
Status: Background article

Savoldi F, Nappi G, Martignoni E, et al. Brain phospholipids in the treatment of chronic cerebrovascular insufficiency. *Curr Ther Res Clin Exp* 1978; 24(2):209.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Scarzella L, Bono G, Bergamasco B. Dihydroergocryptine in the management of senile psycho-organic syndrome. *Int J Clin Pharmacol Res* 1992; 12(1):37-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Scharf S, Mander A, Ugoni A, et al. A double-blind, placebo-controlled trial of diclofenac/misoprostol in Alzheimer's disease. *Neurology* 1999 Jul 13; 53(1):197-201.

Status: Included

Scharlach AE. A comparison of employed caregivers of cognitively impaired and physically impaired elderly persons. *Res Aging* 1989 Jun; 11(2):225-43.

Status: Background article

Schellenberg R, Todorova A, Wedekind W, et al. Pathophysiology and psychopharmacology of dementia: A new study design. *Neuropsychobiology* 1997; 35(3):132-42.

Status: Included

Scherder E, Bouma A, Steen L. Effects of simultaneously applied short-term transcutaneous electrical nerve stimulation and tactile stimulation on memory and affective behaviour of patients with probable Alzheimer's disease. *Behav Neurol* 1995; (1):3-13.

Status: Not included because dementia population not randomized to treatment

Scherder EJ, Bouma A, Steen L. Influence of transcutaneous electrical nerve stimulation on memory in patients with dementia of the Alzheimer type. *J Clin Exp Neuropsychol* 1992 Nov; 14(6):951-60.

Status: Not included because dementia population not randomized to treatment

Scherder EJ, Bouma A, Steen AM. Effects of short-term transcutaneous electrical nerve stimulation on memory and affective behaviour in patients with probable Alzheimer's disease. *Behav Brain Res* 1995; 67(2):211-9.

Status: Not included because dementia population not randomized to treatment

Scherder EJ, Bouma A, Steen LM. Effects of "isolated" transcutaneous electrical nerve stimulation on memory and affective behavior in patients with probable Alzheimer's disease. *Biol Psychiatry* 1998 Mar 15; 43(6):417-24.

Status: Not included because does not meet criteria for treatment for dementia patients

Scherder EJ, Bouma A. Effects of transcutaneous electrical nerve stimulation on memory and behavior in Alzheimer's disease may be stage-dependent. *Biol Psychiatry* 1999 Mar 15; 45(6):743-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Scherder EJ, Van Someren EJ, Swaab DF. Transcutaneous electrical nerve stimulation (TENS) improves the rest-activity rhythm in midstage Alzheimer's disease. *Behav Brain Res* 1999 May; 101(1):105-7.

Status: Not included because no extractable data relevant to review

Scherder EJ, Van Someren EJ, Bouma A, et al. Effects of transcutaneous electrical nerve stimulation (TENS) on cognition and behaviour in aging. *Behav Brain Res* 2000 Jun 15; 111(1-2):223-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Scherder EJ, Deijen JB, Vreeswijk SH, et al. Cranial electrostimulation (CES) in patients with probable Alzheimer's disease. *Behav Brain Res* 2002 Jan 22; 128(2):215-7.

Status: Not included because Jadad Quality Scale score less than three

Scherder E, Bouma A, Steen L. Effects of peripheral tactile stimulation on memory in patients with probable Alzheimer's disease. *Am J Alzheimers Dis* 1995 May; (3):15-21.

Status: Not included because dementia population not randomized to treatment

Scherder E, Bouma A, Steen L. Effects of peripheral tactile nerve stimulation on affective behavior of patients with probable Alzheimer's disease. *Am J Alzheimers Dis* 1998 Mar; (2):61-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Schiffmann R, Heyes MP, Aerts JM, et al. Prospective study of neurological responses to

treatment with macrophage-targeted glucocerebrosidase in patients with type 3 Gaucher's disease. *Ann Neurol* 1997 Oct; 42(4):613-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schifitto G, Sacktor N, Marder K, et al. Randomized trial of the platelet-activating factor antagonist lexipafant in HIV-associated cognitive impairment. *Neurological AIDS Research Consortium. Neurology* 1999 Jul 22; 53(2):391-6. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Schlegel J, Mohr E, Williams J, et al. Guanfacine treatment of Alzheimer's disease. *Clin Neuropharmacol* 1989; 12(2):124-8. *Status: Cross-over trial; Cross-over trial*

Schmechel DE, Schmitt F, Horner J, et al. Lack of effect of oral physostigmine and lecithin in patients with probable Alzheimer's disease. *Neurology* 1984; 34(Suppl 1):280. *Status: Not included because not a full article*

Schmidt IK, Fastbom J. Quality of drug use in Swedish nursing homes. A follow-up study. *Clin Drug Investig* 2000; 20(6):433-46. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Schmitt FA, Miller JP, Kryscio RJ. Promoting interactions with basic scientists and clinicians: summary of the panel session. *Stat Med* 2000 Jun 15; 19(11-12):1463-8. *Status: Background article*

Schmitt R, Capo T, Frazier H, et al. Cranial electrotherapy stimulation treatment of cognitive brain dysfunction in chemical dependence. *J Clin Psychiatry* 1962 Mar; 45(2):60-1. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Schneider F, Popa R, Mihalas G, et al. Superiority of antagonistic-stress composition versus nicergoline in gerontopsychiatry. *Ann N Y Acad Sci* 1994 Jun; 717:332-42. *Status: Included*

Schneider LS, Gleason RP. Carbamazepine in behaviorally disturbed primary dementia patients. *J Clin Psychiatry* 1990; 51(12):524 *Status: Not included because not a full article*

Schneider LS, Lyness SA, Pawluczyk S, et al. Do blood pressure and age predict response to tacrine (THA) in Alzheimer's disease? A preliminary report. *Psychopharmacol Bull* 1991; 27(3):309-14.

Status: Not included because no extractable data relevant to review

Schneider LS, Olin JT, Pawluczyk S. A double-blind crossover pilot study of l-deprenyl (selegiline) combined with cholinesterase inhibitor in Alzheimer's disease. *Am J Psychiatry* 1993 Feb; 150(2):321-3. *Status: Cross-over trial; Cross-over trial*

Schneider LS, Farlow MR, Henderson VW, et al. Estrogen replacement therapy may enhance response to tacrine in women with Alzheimer's disease. *Neurology* 1995; 45:288. *Status: Not included because not a full article*

Schneider LS, Farlow MK. Severity of Alzheimer's disease and response to cholinergic therapy. *Eur J Neurol* 1996; 3:238. *Status: Not included because not a full article*

Schneider LS, Farlow MR, Henderson VW, et al. Effects of estrogen replacement therapy on response to tacrine in patients with Alzheimer's disease. *Neurology* 1996 Jun; 46(6):1580-4. *Status: Companion of an included article*

Schneider LS, Farlow M. Combined tacrine and estrogen replacement therapy in patients with Alzheimer's disease. *Ann N Y Acad Sci* 1997 Sep; 826:317-22. *Status: Companion of an included article*

Schneider LS, Farlow MR, Pogoda JM. Potential role for estrogen replacement in the treatment of Alzheimer's dementia. *Am J Med* 1997 Sep 22; 103(3A):46S-50S. *Status: Not included because Jadad Quality Scale score less than three*

Schneider LS, Tariot PN, Lyketsos CG, et al. National Institute of Mental Health Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE): Alzheimer's disease trial methodology. *Am J Geriatr Psychiatry* 2001; 9(4):346-60. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Schneider LS. Cholinesterase inhibitors for Alzheimer's disease. *JAMA* 2003 May 14;

289(18):2359-60.

Status: Background article

Schneider LS, Pollock VE, Lyness SA. A metaanalysis of controlled trials of neuroleptic treatment in dementia. *J Am Geriatr Soc* 1990; 38(5):553-63.

Status: Background article

Schredl M, Weber B, Braus D, et al. The effect of rivastigmine on sleep in elderly healthy subjects. *Exp Gerontol* 2000; 35(2):243-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schreiber M, Schweizer A, Lutz K, et al. Potential of an interactive computer-based training in the rehabilitation of dementia: An initial study. *Neuropsychol Rehab* 1999; (2):155-67.

Status: Not included because dementia population not randomized to treatment

Schrijnemaekers V, van Rossum E, Candel M, et al. Effects of emotion-oriented care on elderly people with cognitive impairment and behavioral problems. *Int J Geriatr Psychiatry* 2002 Oct; 17(10):926-37.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schroder J, Kratz B, Pantel J, et al. Prevalence of mild cognitive impairment in an elderly community sample. *J Neural Transm Suppl* 1998; 54:51-9.

Status: Background article

Schuck S, Lebreton S. Aminotransferase levels and silymarin in de novo tacrine-treated Alzheimer's disease patients. *Fundam Clin Pharmacol* 1999; 13(3):422.

Status: Not included because not a full article

Schulz M. Intensive geriatric rehabilitation reduced hospital stay and time to independent living in hip fracture patients with mild to moderate dementia... commentary on Huusko TM, Karppi P, Avikainen V, et al. Randomised, clinically controlled trial of intensive geriatric rehabilitation in patients with hip fracture: subgroup analysis of patients with dementia. *BMJ* 2000 Nov 4;321:1107-11. *Evid Based Nurs* 2001 Apr; 4(2):54

Status: Background article

Schulz R, O'Brien A, Czaja S, et al. Dementia caregiver intervention research: in search of clinical significance. *Gerontologist* 2002 Oct;

42(5):589-602.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schwartz AS, Kohlstaedt EV. Physostigmine effects in Alzheimer's disease: Relationship to dementia severity. *Life Sci* 1986 Mar 17; 38(11):1021-8.

Status: Not included because no extractable data relevant to review

Schwartz BL, Hashtroudi S, Herting RL, et al. D-Cycloserine enhances implicit memory in Alzheimer patients. *Neurology* 1996 Feb; 46(2):420-4.

Status: Not included because no extractable data relevant to review

Schweiger C. A 48-week, placebo-controlled study examining propentofylline's safety, efficacy, and impact on disease progression in patients with vascular dementia. *J Eur Coll Neuropsychopharmacol* 1999; (Suppl 5):S319.

Status: Not included because not a full article

Scott HD, Laake K. Statins for the reduction of risk of Alzheimer's disease. In: *The Cochrane Library*, 2001. Issue 3. Oxford: Update Software

Status: Background article

Seipel JH, Fisher R, Blatchley RJ, et al. Rheoencephalographic and other studies of betahistine in humans. IV. Prolonged administration with improvement in arteriosclerotic dementia. *J Clin Pharmacol* 1977 Feb; 17(2-3):140-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Seipel JH, Fisher R, Floam JE, et al. Rheoencephalographic and other studies of betahistine in humans. III. Improved methods of diagnosis and selection in arteriosclerotic dementia. *J Clin Pharmacol* 1977 Jan; 17(1):63-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sekijima Y, Ikeda S, Tokuda T, et al. Prevalence of dementia of Alzheimer type and apolipoprotein E phenotypes in aged patients with Down's syndrome. *Eur Neurol* 1998; 39(4):234-7.

Status: Background article

Semlitsch HV, Anderer P, Saletu B, et al. Topographic mapping of cognitive event-related

potentials in a double-blind, placebo-controlled study with the hemoderivative Actovegin in age-associated memory impairment. *Neuropsychobiology* 1990 Sep; 24(1):49-56.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Semlitsch HV, Anderer P, Saletu B. Topographic mapping of long latency "cognitive" event-related potentials (P 300): A double-blind, placebo-controlled study with amantadine in mild dementia. *J Neural Transm Park Dis Dement Sect* 1992; 4:319-36.
Status: Not included because no extractable data relevant to review

Semlitsch HV, Anderer P, Saletu B, et al. Cognitive psychophysiology in nootropic drug research: Effects of Ginkgo biloba on event-related potentials (P300) in age-associated memory impairment. *Pharmacopsychiatry* 1995 Jul; 28(4):134-42.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Senin U, Abate G, Fieschi C, et al. Aniracetam (Ro 13-5057) in the treatment of senile dementia of Alzheimer type (SDAT): Results of a placebo controlled multicentre clinical study. *Eur Neuropsychopharmacol* 1991 Dec; 1(4):511-7.
Status: Included

Senin U, Parnetti L, Barbagallo-Sangiorgi G, et al. Idebenone in senile dementia of Alzheimer type: A multicentre study. *Arch Gerontol Geriatr* 1992; 15(3):249-60.
Status: Not included because Jadad Quality Scale score less than three

Serafetinides EA, Willis D, Clark ML. The EEG effects of zinc in geriatric psychiatric patients. *Int Pharmacopsychiatry* 1974; 9(2):95-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Serby M, Angrist B, Corwin J, et al. Cholecystokinin octapeptide in dementia. *Psychopharmacol Bull* 1984; 20(3):546-7.
Status: Not included because dementia population not randomized to treatment

Serby M, Resnick R, Jordan B, et al. Naltrexone and Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 1986; 10(3-5):587-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Serfaty M, Kennell-Webb S, Warner J, et al. Double blind randomised placebo controlled trial of low dose melatonin for sleep disorders in dementia. *Int J Geriatr Psychiatry* 2002; 17(12):1120-7.
*Status: Cross-over trial;
Cross-over trial*

Seux M-L, Forette F, Staessen JA, et al. Treatment of isolated systolic hypertension and dementia prevention in older patients. Results of the Systolic Hypertension in Europe trial (SYST-EUR) vascular dementia project. *Eur Heart J Suppl* 1999; 1(M):M6-M12.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sevush S, Guterman A, Villalon AV. Improved verbal learning after outpatient oral physostigmine therapy in patients with dementia of the Alzheimer type. *J Clin Psychiatry* 1991 Jul; 52(7):300-3.
Status: Not included because dementia population not randomized to treatment

Shaw FE, Bond J, Richardson DA, et al. Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: Randomised controlled trial. *BMJ* 2003; 326(7380):73-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Shaw TG, Meyer JS. Double-blind trial of oral papaverine in chronic cerebrovascular ischemia. *Angiology* 1978 Nov; 29(11):839-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sheikh JI, Hill RD, Yesavage JA. Long-term efficacy of cognitive training for age-associated memory impairment: A six-month follow-up study. *Dev Neuropsychol* 1986; 2(4):413-21.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Shelton P, Schraeder C, Dworak D, et al. Caregivers' utilization of health services: results from the Medicare Alzheimer's Disease Demonstration, Illinois site. *J Am Geriatr Soc* 2001 Dec; 49(12):1600-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Sherwin BB. Estrogen and cognitive functioning in men with Mild Cognitive Impairment. *J Mol Neurosci* 2002; 19(1-2):219-23.

Status: Not included because no extractable data relevant to review

Shimada Y, Terasawa K, Yamamoto T, et al. A well-controlled study of Choto-san and placebo in the treatment of vascular dementia. *J Tradit Med* 1994; 11:246-55.

Status: Included

Shrotriya RC, Cutler NR, Sramek JJ, et al. Efficacy and safety of BMY 21,502 in Alzheimer's disease. *Ann Pharmacother* 1996 Dec; 30(12):1376-80.

Status: Included

Shua-Haim JR, Shua-Haim V, Sabo M, et al. Case report: Donepezil in the treatment of advanced Alzheimer's disease. *Ann Long Term Care* 1999; 7(2):67-71.

Status: Not included because dementia population not randomized to treatment

Shua-Haim JR, Shua-Haim V, Comsti E, et al. Donepezil (Aricept(TM)) treatment of multi infarct dementia: The caregivers and clinical impression. *Am J Alzheimers Dis* 2000; 15(4):201-11.

Status: Not included because dementia population not randomized to treatment

Shukla VK, Otten N, Coyle D. Drug treatments for Alzheimer's disease III. A review of published pharmacoeconomic evaluations. Technology report no. 11. Ottawa: Canadian Coordinating Office for Health Technology Assessment. 2000.

Status: Background article

Shumaker SA, Reboussin BA, Espeland MA, et al. The Women's Health Initiative Memory Study (WHIMS): A trial of the effect of estrogen therapy in preventing and slowing the progression of dementia. *Control Clin Trials* 1998 Dec; 19(6):604-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sidtis JJ, Gatsonis C, Price RW, et al. Zidovudine treatment of the AIDS dementia complex: Results of a placebo-controlled trial. *AIDS Clinical Trials Group. Ann Neurol* 1993 Apr; 33(4):343-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Silverberg GD, Levinthal E, Sullivan EV, et al. Assessment of low-flow CSF drainage as a treatment for AD: Results of a randomized pilot study. *Neurology* 2002 Oct 22; 59(8):1139-45.

Status: Not included because does not meet criteria for treatment for dementia patients

Simons M, Schwarzler F, Lutjohann D, et al. Treatment with simvastatin in normocholesterolemic patients with Alzheimer's disease: A 26-week randomized, placebo-controlled, double-blind trial. *Ann Neurol* 2002; 52(3):346-50.

Status: Included

Sinforiani E, Iannuccelli M, Mauri M, et al. Neuropsychological changes in demented patients treated with acetyl-L-carnitine. *Int J Clin Pharmacol Res* 1990; 10(1-2):69-74.

Status: Not included because Jadad Quality Scale score less than three

Sival RC, Haffmans PM, Jansen PA, et al. Sodium valproate in the treatment of aggressive behavior in patients with dementia: A randomized placebo controlled clinical trial. *Int J Geriatr Psychiatry* 2002; 17(6):579-85.

Status: Cross-over trial; Cross-over trial

Skelly J, Flint A, Brunt S, et al. Treatment of urinary incontinence in dementia using low dose oxybutynin chloride. *Age Ageing* 1995; 24(S):19.

Status: Not included because not a full article

Slooter AJC, Houwing-Duistermaat JJ, van Harskamp F, et al. Apolipoprotein E genotype and progression of Alzheimer's disease: The Rotterdam Study. *J Neurol* 1999; 246(4):304-8.

Status: Not included because dementia population not randomized to treatment

Smallwood J, Brown R, Coulter F, et al. Aromatherapy and behaviour disturbances in dementia: A randomized controlled trial. *Int J Geriatr Psychiatry* 2001 Oct; 16(10):1010-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

SMID Group. Phosphatidylserine in the treatment of clinically diagnosed Alzheimer's disease. *J Neural Transm Gen Sect* 1987; 24(Suppl):287-92.

Status: Companion of an included article

Smid HG, Trumper BG, Pottag G, et al. Differentiation of hypoglycaemia induced cognitive

impairments. An electrophysiological approach. Brain 1997 Jun; 120(Pt 6):1041-56.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smirne S, Truci G, Peri E, et al. Efficacy and tolerability of oxiracetam in Alzheimer's disease: A double-blind, six month study. Clin Neurol Neurosurg 1987; 2(19):

Status: Article not retrievable

Smirne S, Palazzi S, Grassi MP, et al. L-deprenyl in Alzheimer's patients of different severity possible preventive use. Alzheimers Dis Relat Disord Adv Biosci 1993; 87:451-5.

Status: Companion of an included article

Smith CM, Swash M, Exton-Smith AN, et al. Choline therapy in Alzheimer's disease. Lancet 1978 Aug 5; 2(8084):318.

Status: Not included because not a full article

Smith CM, Swash M. Physostigmine in Alzheimer's disease. Lancet 1979 Jan 6; 1(8106):42.

Status: Not included because not a full article

Smith DF, Stromgren E, Petersen HN. Lack of effect of tryptophan treatment in demented gerontopsychiatric patients. A double-blind, crossover-controlled study. Acta Psychiatr Scand 1984; 70(5):470-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith DJ, Yuhnevich S. Adverse reactions to rivastigmine in three cases of dementia. Aust N Z J Psychiatry 2001; 35(5):694-5.

Status: Not included because not a full article

Smith F, Talwalker S, Gracon S, et al. The use of survival analysis techniques in evaluating the effect of long-term tacrine (Cognex) treatment on nursing home placement and mortality in patients with Alzheimer's disease. J Biopharm Stat 1996 Nov; 6(4):395-409.

Status: Companion of an included article

Smith F. Mixed-model analysis of incomplete longitudinal data from a high-dose trial of tacrine (Cognex) in Alzheimer's patients. J Biopharm Stat 1996 Mar; 6(1):59-67.

Status: Not included because Jadad Quality Scale score less than three

Smith F, Gracon S, Knopman D, et al. Survival analysis to evaluate the effect of long-term tacrine (Cognex) treatment on nursing home placement in Alzheimer's patients. J Neurol 1997; 244(Suppl 3):S88.

Status: Not included because not a full article

Smith F, Gracon S, Knopman D, et al. Tacrine treatment and nursing home placement: Application of the Cox proportional hazards model with time-dependent covariates. Drug Inf J 1998; 32(3):729-35.

Status: Not included because dementia population not randomized to treatment

Smith GR, Taylor CW, Linkous P. Haloperidol versus thioridazine for the treatment of psychogeriatric patients: A double-blind clinical trial. Psychosomatics: Journal of Consultation Liaison Psychiatry 1974; 15(3):134-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith MK. The effects of participatory music on the reality orientation and sociability of Alzheimer's residents in a long-term-care setting. Activities Adaptation Aging 1994; 18(2):41-55.

Status: Not included because dementia population not randomized to treatment

Smith P, Loy C, Wong M. Naftidrofuryl (nafronyl, Praxilene) for cognitive impairment (Cochrane Protocol). In: The Cochrane Library, 2002. Issue 2. Oxford: Update Software.

Status: Background article

Smith RC, Voulis G, Johnson R, et al. Comparison of therapeutic response to long-term treatment with lecithin versus piracetam plus lecithin in patients with Alzheimer's disease. Psychopharmacol Bull 1984; 20(3):542-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith WL, Lowrey JB, Davis JA. The effects of cyclandelate on psychological test performance in patients with cerebral vascular insufficiency. Curr Ther Res Clin Exp 1968; 10(12):613-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Snaedal J, Johannesson T, Jonsson JE, et al. The effects of nicotine in dermal plaster on cognitive functions in patients with Alzheimer's disease. Dementia 1996 Jan; 7(1):47-52.

Status: Not included because dementia population not randomized to treatment

Snowdon DA, Greiner LH, Mortimer JA, et al. Brain infarction and the clinical expression of Alzheimer disease. The Nun Study. JAMA 1997 Mar 12; 277(10):813-7.

Status: Background article

Snyder M, Egan E, Burns K. Interventions to decrease disruptive behaviors in persons with dementia. Minn Nurs Accent 1993; 65(9):4.

Status: Article not retrievable

Snyder M, Egan EC, Burns KR. Interventions for decreasing agitation behaviors in persons with dementia. J Gerontol Nurs 1995 Jul; 21(7):34-40.

Status: Not included because dementia population not randomized to treatment

Snyder M, Tseng Y, Brandt C, et al. A glider swing intervention for people with dementia. Geriatr Nurs (Minneap) 2001 Mar; 22(2):86-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sobel BP. Bingo vs. physical intervention in stimulating short-term cognition in Alzheimer's disease patients. Am J Alzheimers Dis Other Demen 2001 Mar; 16(2):115-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sobow TM, Maczkiewicz M, Kloszewska I. Tianeptine versus fluoxetine in the treatment of depression complicating Alzheimer's disease. Int J Geriatr Psychiatry 2001; 16(11):1108-9.

Status: Not included because not a full article

Soininen H, Koskinen T, Helkala E-L. Synthetic ACTH 4-9 (ORG2766) in treatment of Alzheimer's disease. Acta Neurol Scand 1984; 69(Suppl 98):236-7.

Status: Companion of an included article

Soininen H, Koskinen T, Helkala EL, et al. Treatment of Alzheimer's disease with a synthetic ACTH 4-9 analog. Neurology 1985 Sep; 35(9):1348-51.

Status: Included

Solomon PR, Knapp MJ, Gracon SI, et al. Long-term tacrine treatment in patients with Alzheimer's disease. Lancet 1996 Jul 27; 348(9022):275-6.

Status: Not included because not a full article

Soncrant TT, Raffaele KC, Asthana S, et al. Memory improvement without toxicity during chronic, low dose intravenous arecoline in Alzheimer's disease. Psychopharmacologia 1993; 112(4):421-7.

Status: Cross-over trial;

Cross-over trial

Sourander LB, Portin R, Molsa P, et al. Senile dementia of the Alzheimer type treated with aniracetam: A new nootropic agent. Psychopharmacologia 1987; 91(1):90-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Spagnoli A, Lucca U, Menasce G, et al. Long-term acetyl-L-carnitine treatment in Alzheimer's disease. Neurology 1991 Nov; 41(11):1726-32.

Status: Included

Spagnolo C, Dallasta D, Iannuccelli M. A controlled double-blind trial comparing etoperidone with thioridazine in the management of severe senile dementia. Drugs Exp Clin Res 1983; 9(12):873-80.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sparano N. Donepezil for Alzheimer's disease. J Fam Pract 1998 May; 46(5):356

Status: Companion of an included article

Spector A, Orrell M, Davies S, et al. Can reality orientation be rehabilitated? Development and piloting of an evidence-based programme of cognition-based therapies for people with dementia. Neuropsychol Rehab 2001; 11(3-4):377-97.

Status: Not included because does not meet criteria for treatment for dementia patients

Spilich GJ, Wannenmacher W, Duarte A, et al. Efficacy of pyritinol versus hydergine upon cognitive performance in patients with senile dementia of the Alzheimer's type: A double-blind multi-center trial. Alzheimers Res 1996; (3):79-84.

Status: Included

Sramek JJ, Cutler NR, Hurley DJ, et al. The utility of salivary amylase as an evaluation of M3 muscarinic agonist activity in Alzheimer's disease. Prog Neuropsychopharmacol Biol Psychiatry 1995 Jan; 19(1):85-91.

Status: Not included because does not meet criteria for treatment for dementia patients

Sramek JJ, Block GA, Reines SA, et al. A multiple-dose safety trial of eptastigmine in Alzheimer's disease, with pharmacodynamic observations of red blood cell cholinesterase. *Life Sci* 1995; 56(5):319-26.

Status: Not included because no extractable data relevant to review

Sramek JJ, Viereck C, Huff FJ, et al. A "bridging" (safety/tolerance) study of besipirdine hydrochloride in patients with Alzheimer's disease. *Life Sci* 1995; 57(12):1241-8.

Status: Not included because no extractable data relevant to review

Sramek JJ, Hurley DJ, Wardle TS, et al. The safety and tolerance of xanomeline tartrate in patients with Alzheimer's disease. *J Clin Pharmacol* 1995 Aug; 35(8):800-6.

Status: Not included because no extractable data relevant to review

Sramek JJ, Sedman AJ, Reece PA, et al. Safety and tolerability of CI-979 in patients with Alzheimer's disease. *Life Sci* 1995; 57(5):503-10.

Status: Not included because no extractable data relevant to review

Sramek JJ, Anand R, Wardle TS, et al. Safety/tolerability trial of SDZ ENA 713 in patients with probable Alzheimer's disease. *Life Sci* 1996; 58(15):1201-7.

Status: Not included because no extractable data relevant to review

Sramek JJ, Forrest M, Mengel H, et al. A bridging study of LU 25-109 in patients with probable Alzheimer's disease. *Life Sci* 1998; 62(3):195-202.

Status: Not included because dementia population not randomized to treatment

Sramek JJ, Hourani J, Jhee SS, et al. NXX-066 in patients with Alzheimer's disease: A bridging study. *Life Sci* 1999; 64(14):1215-21.

Status: Not included because no extractable data relevant to review

St Clair D, Norrman J, Perry R, et al. Apolipoprotein E epsilon 4 allele frequency in patients with Lewy body dementia, Alzheimer's disease and age-matched controls. *Neurosci Lett* 1994 Jul 18; 176(1):45-6.

Status: Not included because dementia population not randomized to treatment

Stapleton JM, Eckardt MJ, Martin P, et al. Treatment of alcoholic organic brain syndrome with the serotonin reuptake inhibitor fluvoxamine: A preliminary study. *Adv Alcohol Subst Abuse* 1988; 7(3-4):47-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Starr JM, Whalley LJ. Hypertensive Old People in Edinburgh (HOPE) Study: Electrocardiographic changes after captopril or bendroflumazide treatment. *Age Ageing* 1993 Sep; 22(5):343-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Starr JM, Whalley LJ, Inch S, et al. A double-blind trial of captopril or bendroflumazide in newly diagnosed senile hypertension. *Curr Med Res Opin* 1994; 13(4):214-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Starr JM, Whalley LJ, Deary IJ. The effects of antihypertensive treatment on cognitive function: Results from the HOPE study. *J Am Geriatr Soc* 1996 Apr; 44(4):411-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Steele C, Lucas MJ, Tune L. Haloperidol versus thioridazine in the treatment of behavioral symptoms in senile dementia of the Alzheimer's type: Preliminary findings. *J Clin Psychiatry* 1986 Jun; 47(6):310-2.

Status: Not included because dementia population not randomized to treatment

Steele LS, Glazier RH. Is donepezil effective for treating Alzheimer's disease? *Can Fam Physician* 1999 Apr; 45:917-9.

Status: Companion of an included article

Stegink AJ. The clinical use of piracetam, a new nootropic drug. The treatment of symptoms of senile involution. *Arzneimittelforschung* 1972 Jun; 22(6):975-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Steiger William A. Effects of naloxone in treatment of senile dementia. *J Am Geriatr Soc* 1985; 33(2):155.

Status: Not included because not a full article

Stern FH. Management of chronic brain syndrome secondary to cerebral arteriosclerosis, with

special reference to papaverine hydrochloride. J Am Geriatr Soc 1970 Jun; 18(6):507-12.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Stern RG, Mohs RC, Davidson M, et al. A longitudinal study of Alzheimer's disease: measurement, rate, and predictors of cognitive deterioration. Am J Psychiatry 1994 Mar; 151(3):390-6.
Status: Background article

Stern Y, Sano M, Mayeux R. Effects of oral physostigmine in Alzheimer's disease. Ann Neurol 1987 Sep; 22(3):306-10.
Status: Not included because dementia population not randomized to treatment

Stern Y, Sano M, Mayeux R. Long-term administration of oral physostigmine in Alzheimer's disease. Neurology 1988 Dec; 38(12):1837-41.
*Status: Cross-over trial;
Cross-over trial*

Stevermer JJ, Lindbloom EJ. Ginkgo biloba for dementia. J Fam Pract 1998; 46(1):20.
Status: Not included because not a full article

Stewart A, Phillips R, Dempsey G. Pharmacotherapy for people with Alzheimer's disease: A Markov-cycle evaluation of five years' therapy using donepezil. Int J Geriatr Psychiatry 1998 Jul; 13(7):445-53.
Status: Not included because dementia population not randomized to treatment

Stewart WF, Kawas C, Corrada M, et al. Risk of Alzheimer's disease and duration of NSAID use. Neurology 1997 Mar; 48(3):626-32.
Status: Not included because does not meet criteria for treatment for dementia patients

Stoppe G, Sandholzer H, Staedt J, et al. Sleep disturbances in the demented elderly: Treatment in ambulatory care. Sleep 1995; 18(10):844-8.
Status: Not included because dementia population not randomized to treatment

Stotsky BA, Cole JO, Tang YT, et al. Sodium butabarbital (butisol sodium) as an hypnotic agent for aged psychiatric patients with sleep disorders. J Am Geriatr Soc 1971 Oct; 19(10):860-70.
Status: Not included because no extractable data relevant to review

Stotsky BA, Cole JO, Lu LM, et al. A controlled study of the efficacy of pentylentetrazol (Metrazol) with hard-core hospitalized psychogeriatric patients. Am J Psychiatry 1972 Oct; 129(4):387-91.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Stracciari A, Ciucci G, Loreta R, et al. Multi-infarct dementia: Clinical trial with acetyl-L-carnitine vs placebo. J Neurol 1988; 235(Suppl):92.
Status: Not included because not a full article

Street J, Clark WS, Gannon KS, et al. Reduction of psychotic symptoms in patients with Lewy Body-like symptoms treated with olanzapine. J Eur Coll Neuropsychopharmacol 1999; 9(Suppl 5):S331.
Status: Not included because not a full article

Street JS, Clark WS, Gannon KS, et al. Olanzapine treatment of psychotic and behavioral symptoms in patients with Alzheimer disease in nursing care facilities: A double-blind, randomized, placebo-controlled trial. The HGEU Study Group. Arch Gen Psychiatry 2000 Oct; 57(10):968-76.
Status: Included

Street JS, Clark WS, Kadam DL, et al. Long-term efficacy of olanzapine in the control of psychotic and behavioral symptoms in nursing home patients with Alzheimer's dementia. Int J Geriatr Psychiatry 2001 Dec; 16(Suppl):S62-S70
Status: Companion of an included article

Streim JE, Oslin DW, Katz IR, et al. Drug treatment of depression in frail elderly nursing home residents. Am J Geriatr Psychiatry 2000; 8(2):150-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sudilovsky A, Cutler NR, Sramek JJ, et al. A pilot clinical trial of the angiotensin-converting enzyme inhibitor ceranapril in Alzheimer's disease. Alzheimer Dis Assoc Disord 1993; 7(2):105-11.
*Status: Cross-over trial;
Cross-over trial*

Sultzer DL, Gray KF, Gunay I, et al. A double-blind comparison of trazodone and haloperidol for treatment of agitation in patients with dementia. Am J Geriatr Psychiatry 1997; 5(1):60-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sultzer DL, Gray KF, Gunay I, et al. Does behavioral improvement with haloperidol or trazodone treatment depend on psychosis or mood symptoms in patients with dementia? J Am Geriatr Soc 2001 Oct; 49(10):1294-300.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sumiyoshi T, Matsui M, Nohara S, et al. Enhancement of cognitive performance in schizophrenia by addition of tandospirone to neuroleptic treatment. Am J Psychiatry 2001; 158(10):1722-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Summers WK, Majovski LV, Marsh GM, et al. Oral tetrahydroaminoacridine in long-term treatment of senile dementia, Alzheimer type. N Engl J Med 1986 Nov 13; 315(20):1241-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sunderland T, Tariot P, Murphy DL, et al. Scopolamine challenges in Alzheimer's disease. Psychopharmacologia 1985; 87(2):247-9.

*Status: Cross-over trial;
Cross-over trial*

Sunderland T, Tariot PN, Cohen RM, et al. Anticholinergic sensitivity in patients with dementia of the Alzheimer type and age-matched. Arch Gen Psychiatry 1987; 44(5):418-26.

Status: Not included because Jadad Quality Scale score less than three

Sunderland T, Weingartner H, Cohen RM, et al. Low-dose oral lorazepam administration in Alzheimer subjects and age-matched controls. Psychopharmacologia 1989; 99(1):129-33.

*Status: Cross-over trial;
Cross-over trial*

Sunderland T, Molchan S, Lawlor B, et al. A strategy of "combination chemotherapy" in Alzheimer's disease: Rationale and preliminary results with physostigmine plus deprenyl. Int Psychogeriatr 1992; 4(Suppl 2):291-309.

Status: Not included because dementia population not randomized to treatment

Swanson EA, Maas ML, Buckwalter KC. Alzheimer's residents' cognitive and functional measures: Special and traditional care unit comparison. Clin Nurs Res 1994 Feb; 3(1):27-41.

Status: Not included because does not meet criteria for treatment for dementia patients

Swanson EA, Maas ML, Buckwalter KC. Catastrophic reactions and other behaviors of Alzheimer's residents: Special unit compared with traditional units. Arch Psychiatr Nurs 1993; 7(5):292-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Swartz JR, Miller BL, Lesser IM, et al. Frontotemporal dementia: Treatment response to serotonin selective reuptake inhibitors. J Clin Psychiatry 1997; 58(5):212-6.

Status: Not included because dementia population not randomized to treatment

Szatmari Sz, Whitehouse PJ. Vinpocetine for cognitive impairment and dementia (Cochrane Protocol). In: The Cochrane Library, 2002. Issue 2. Oxford: Update Software.

Status: Background article

Tabet N, Birks J, Grimley EJ. Vitamin E for Alzheimer's disease. In: The Cochrane Library, 2000. Issue 4. Oxford: Update Software.

Status: Background article

Tabet N, Mantle D, Walker Z, et al. Dietary and endogenous antioxidants in dementia. Int J Geriatr Psychiatry 2001; 16(6):639-41.

Status: Not included because not a full article

Tabet N, Mantle D, Walker Z, et al. Endogenous antioxidant activities in relation to concurrent vitamins A, C, and E intake in dementia. Int Psychogeriatr 2002; 14(1):7-15.

Status: Not included because does not meet criteria for treatment for dementia patients

Tabet N, Feldman H. Indomethacin for the treatment of Alzheimer's disease patients. In: The Cochrane Library, 2002. Issue 2. Oxford: Update Software.

Status: Background article

Tabourne CE. The effects of a life review program on disorientation, social interaction and self-esteem of nursing home residents. Int J Aging Hum Dev 1995; 41(3):251-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Talwalker S. Analysis of repeated measurements with dropouts among Alzheimer's disease patients using summary measures and meta-analysis. J Biopharm Stat 1996; 6(1):49-58.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Talwalker S, Overall JE, Srirama MK, et al. Cardinal features of cognitive dysfunction in Alzheimer's disease: A factor-analytic study of the Alzheimer's Disease Assessment Scale. J Geriatr Psychiatry Neurol 1996 Jan; 9(1):39-46.

Status: Not included because does not meet criteria for treatment for dementia patients

Talwalker S. The cardinal features of cognitive and noncognitive dysfunction and the differential efficacy of tacrine in Alzheimer's disease patients. J Biopharm Stat 1996 Nov; 6(4):443-56.

Status: Not included because no extractable data relevant to review

Tang MX, Jacobs D, Stern Y, et al. Effect of oestrogen during menopause on risk and age at onset of Alzheimer's disease. Lancet 1996 Aug 17; 348(9025):429-32.

Status: Background article

Tappen RM. The effect of skill training on functional abilities of nursing home residents with dementia. Res Nurs Health 1994 Jun; 17(3):159-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tappen RM, Roach KE, Applegate EB, et al. Effect of a combined walking and conversation intervention on functional mobility of nursing home residents with Alzheimer disease. Alzheimer Dis Assoc Disord 2000 Oct; 14(4):196-201.

Status: Not included because does not meet criteria for treatment for dementia patients

Tappen RM, Williams CL, Barry C, et al. Conversation intervention with Alzheimer's patients: Increasing the relevance of communication. Clin Gerontol 2002; 24(3-4):63-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Taragano FE, Lyketsos CG, Mangone CA, et al. A double-blind, randomized, fixed-dose trial of fluoxetine vs. amitriptyline in the treatment of major depression complicating Alzheimer's disease. Psychosomatics 1997 May; 38(3):246-

52.

Status: Included

Targum SD, Wieland S, Glasky MS, et al. Evaluation of AIT-082 in patients with mild to moderate senile dementia of the Alzheimer's type. J Eur Coll Neuropsychopharmacol 1999; (Suppl 5):S320.

Status : Not included because not a full article

Tariot P, Parys W, Kershaw P. The efficacy and tolerability of galantamine in Alzheimer's disease a 5 month placebo controlled study with slow dose escalation. Neurology 2000; 54(Suppl 3):A415.

Status: Not included because not a full article

Tariot PN, Sunderland T, Weingartner H, et al. Low- and high-dose naloxone in dementia of the Alzheimer type. Psychopharmacol Bull 1985; 21(3):680-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Sunderland T, Weingartner H, et al. Naloxone and Alzheimer's disease. Cognitive and behavioral effects of a range of doses. Arch Gen Psychiatry 1986 Aug; 43(8):727-32.

*Status: Cross-over trial;
Cross-over trial*

Tariot PN, Sunderland T, Murphy DL, et al. Design and interpretation of opiate antagonist trials in dementia. Prog Neuropsychopharmacol Biol Psychiatry 1986; 10(3-5):611-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Sunderland T, Weingartner H, et al. Cognitive effects of L-deprenyl in Alzheimer's disease. Psychopharmacologia 1987; 91(4):489-95.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Cohen RM, Sunderland T, et al. L-deprenyl in Alzheimer's disease. Preliminary evidence for behavioral change with monoamine oxidase B inhibition. Arch Gen Psychiatry 1987 May; 44(5):427-33.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Sunderland T, Cohen RM, et al. Tranylcypromine compared with L-deprenyl in Alzheimer's disease. J Clin Psychopharmacol 1988 Feb; 8(1):23-7.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Cohen RM, Welkowitz JA, et al. Multiple-dose arecoline infusions in Alzheimer's disease. Arch Gen Psychiatry 1988 Oct; 45(10):901-5.

*Status: Cross-over trial;
Cross-over trial*

Tariot PN, Gross M, Sunderland T, et al. High-dose naloxone in older normal subjects: Implications for Alzheimer's disease. J Am Geriatr Soc 1988 Aug; 36(8):681-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Erb R, Leibovici A, et al. Carbamazepine treatment of agitation in nursing home patients with dementia: A preliminary study. J Am Geriatr Soc 1994 Nov; 42(11):1160-6.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Leibovici A, Erb R, et al. Carbamazepine therapy for agitation in dementia: Status report. Am J Geriatr Psychiatry 1994; 2(3):257-9.

Status: Not included because not a full article

Tariot PN, Frederiksen K, Erb R, et al. Lack of carbamazepine toxicity in frail nursing home patients: A controlled study. J Am Geriatr Soc 1995 Sep; 43(9):1026-9.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Goldstein B, Podgorski CA, et al. Short-term administration of selegiline for mild-to-moderate dementia of the Alzheimer's type. Am J Geriatr Psychiatry 1998; 6(2):145-54.

*Status: Cross-over trial;
Cross-over trial*

Tariot PN, Erb R, Podgorski CA, et al. Efficacy and tolerability of carbamazepine for agitation and aggression in dementia. Am J Psychiatry 1998 Jan; 155(1):54-61.

Status: Included

Tariot PN, Upadhyaya A, Sunderland T, et al. Physiologic and neuroendocrine responses to intravenous naloxone in subjects with Alzheimer's disease and age-matched controls. Biol Psychiatry 1999 Aug 1; 46(3):412-9.

Status: Not included because no extractable data relevant to review

Tariot PN, Solomon PR, Morris JC, et al. A 5-month, randomized, placebo-controlled trial of galantamine in AD. The Galantamine USA-10 Study Group. Neurology 2000 Jun 27; 54(12):2269-76.

Status: Included

Tariot PN, Salzman C, Yeung PP, et al. Long-Term use of quetiapine in elderly patients with psychotic disorders. Clin Ther 2000 Sep; 22(9):1068-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Cummings JL, Katz IR, et al. A randomised, double-blind, placebo-controlled study of the efficacy and safety of Donepezil in patients with Alzheimer's disease in the nursing home setting. J Am Geriatr Soc 2001a; 49(12):1590-9.

Status: Included

Tariot PN, Schneider LS, Mintzer JE, et al. Safety and tolerability of divalproex sodium in the treatment of signs and symptoms of mania in elderly patients with dementia: Results of a double-blind, placebo-controlled trial. Curr Ther Res Clin Exp 2001b; 62(1):51-67.

Status: Included

Tekin S, Aykut BC, Tanridag T, et al. Antiglutamatergic therapy in Alzheimer's disease: effects of Lamotrigine. J Neural Transm Gen Sect 1998; 105(2-3):295-303.

*Status: Cross-over trial;
Cross-over trial*

Templeton L, Barker A, Wesnes K, et al. A double-blind, placebo-controlled single dose trial of intravenous flumazenil in Alzheimer's disease. Hum Psychopharmacol 1999; 14(4):239-45.

*Status: Cross-over trial;
Cross-over trial*

Tennant FS. Preliminary observations on naltrexone for treatment of Alzheimer's type dementia. J Am Geriatr Soc 1987; 35(4):369-70.

Status: Not included because not a full article

ter Haar HW. A comparison of chlormethiazole and haloperidol in the treatment of elderly patients with confusion of organic and psychogenic origin: A double-blind crossover study.

Pharmatherapeutica 1977; 1(9):563-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Terano T, Fujishiro S, Ban T, et al.

Docosahexaenoic acid supplementation improves the moderately severe dementia from thrombotic cerebrovascular diseases. *Lipids* 1999; 34(Suppl):S345-S346.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Terasawa K, Shimada Y, Kita T, et al. Choto-san in the treatment of vascular dementia: A double-blind, placebo-controlled study. *Phytomedicine* 1997; 4(1):15-22.

Status: Included

Teresi JA, Kleinman M, Ocepek-Welikson K. Modern psychometric methods for detection of differential item functioning: application to cognitive assessment measures. *Stat Med* 2000 Jun 15; 19(11-12):1651-83.

Status: Background article

Teri L, Reifler BV, Veith RC, et al. Imipramine in the treatment of depressed Alzheimer's patients: Impact on cognition. *J Gerontol* 1991 Nov; 46(6):P372-P377.

Status: Not included because Jadad Quality Scale score less than three

Teri L. Behavioral treatment of depression patients with dementia. *Alzheimer Dis Assoc Disord* 1994; 8(Suppl 3):66-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Teri L, Logsdon RG, Uomoto J, et al. Behavioral treatment of depression in dementia patients: A controlled clinical trial. *J Gerontol B Psychol Sci Soc Sci* 1997 Jul; 52(4):159-66.

Status: Not included because does not meet criteria for treatment for dementia patients

Teri L, Logsdon RG, Whall AL, et al. Treatment for agitation in dementia patients: A behavior management approach. *Psychotherapy: Theory, Research, Practice, Training* 1998; 35(4):354-443.

Status: Not included because dementia population not randomized to treatment

Teri L, Logsdon RG, Peskind E, et al. Treatment of agitation in AD: A randomized, placebo-controlled clinical trial. *Neurology* 2000 Nov 14;

55(9):1271-8.

Status: Included

Tewfik GI, Jain VK, Harcup M, et al.

Effectiveness of various tranquilisers in the management of senile restlessness. *Gerontol Clin (Basel)* 1970; 12(6):351-9.

Status: Not included because dementia population not randomized to treatment

Thal LJ, Rosen W, Sharpless NS, et al. Choline chloride fails to improve cognition in Alzheimer's disease. *Neurobiol Aging* 1981; (3):205-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thal LJ, Fuld PA, Masur DM, et al. Oral physostigmine and lecithin improve memory in Alzheimer disease. *Ann Neurol* 1983; 13(5):491-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thal LJ, Masur DM, Sharpless NS, et al. Acute and chronic effects of oral physostigmine and lecithin in Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 1986; 10(3-5):627-36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thal LJ, Carta A, Clarke WR, et al. A 1-year multicenter placebo-controlled study of acetyl-L-carnitine in patients with Alzheimer's disease. *Neurology* 1996a; 47(3):705-11.

Status: Included

Thal LJ, Schwartz G, Sano M, et al. A multicenter double-blind study of controlled-release physostigmine for the treatment of symptoms secondary to Alzheimer's disease. *Neurology* 1996b; 47(6):1389-95.

Status: Included

Thal LJ, Ferguson JM, Mintzer J, et al. A 24-week randomized trial of controlled-release physostigmine in patients with Alzheimer's disease. *Neurology* 1999 Apr 12; 52(6):1146-52.

Status: Included;

Thal LJ, Forrest M, Loft H, et al. Lu 25-109, a muscarinic agonist, fails to improve cognition in Alzheimer's disease. *Lu25-109 Study Group. Neurology* 2000b; 54(2):421-6.

Status: Included

Thal LJ, Calvani M, Amato A, et al. A 1-year controlled trial of acetyl-L-carnitine in early-onset AD. *Neurology* 2000a; 55(6):805-10.
Status: Included

The Canadian Study of Health and Aging Working Group. Patterns of caring for people with dementia in Canada: The Canadian Study of Health and Aging. *Can J Aging* 1994; 13:470-87.
Status: Background article

The Cochrane Non Randomized Studies Method Group: Adverse effects subgroup. Proposed draft addition to Cochrane Handbook: Including adverse effects in Cochrane reviews. Internet. <http://www.dsru.org/wwwboard/latestdraft.pdf>
Status: Background article

Thibault A. A double-blind evaluation of 'Hydergine' and placebo in the treatment of patients with organic brain syndrome and cerebral arteriosclerosis in a nursing home. *Curr Med Res Opin* 1974; 2(8):482-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thienhaus OJ, Wheeler BG, Simon S. A controlled double-blind study of high-dose dihydroergotoxine mesylate (Hydergine(R)) in mild dementia. *J Am Geriatr Soc* 1987; 35(3):219-23.
Status: Not included because Jadad Quality Scale score less than three

Thomas A, Iacono D, Bonanni L, et al. Donepezil, rivastigmine, and vitamin E in Alzheimer disease: A combined P300 event-related potentials/neuropsychologic evaluation over 6 months. *Clin Neuropharmacol* 2001 Jan; 24(1):31-42.
Status: Included

Thomas RG, Berg JD, Sano M, et al. Analysis of longitudinal data in an Alzheimer's disease clinical trial. *Stat Med* 2000 Jun 15; 19(11-12):1433-40.
Status: Background article

Thompson SA, Hodges JR. Mild cognitive impairment: A clinically useful but currently ill-defined concept? *Neurocase* 2002; 8(6):405-10.
Status: Background article

Thompson TL, Filley CM, Mitchell WD, et al. Lack of efficacy of hydergine in patients with Alzheimer's disease. *N Engl J Med* 1990 Aug 16; 323(7):445-8.
Status: Included

Thorpe L, Middleton J, Russell G, et al. Bright light therapy for demented nursing home patients with behavioural disturbance. *Am J Alzheimers Dis* 2000; 15(1):18-26.
Status: Not included because dementia population not randomized to treatment

Tian J, Du H, Yang H, et al. A clinical study on compound Da Huang (Radix et Rhizoma Rhei) preparations for improvement of senile persons' memory ability. *J Tradit Chin Med* 1997 Sep; 17(3):168-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tian J, Yin J, Liu H, et al. Jian Nao Ning for treatment of memory impairment in patients with mild to moderate multi-infarct dementia. *J Tradit Chin Med* 2002 Dec; 22(4):247-51.
Status: Not included because Jadad Quality Scale score less than three

Tierney MC, Fisher RH, Lewis AJ, et al. The NINCDS-ADRDA Work Group criteria for the clinical diagnosis of probable Alzheimer's disease: a clinicopathologic study of 57 cases. *Neurology* 1988 Mar; 38(3):359-64.
Status: Background article

Tignol J, Pujol-Domenech J, Chartres JP, et al. Double-blind study of the efficacy and safety of milnacipran and imipramine in elderly patients with major depressive episode. *Acta Psychiatr Scand* 1998 Feb; 97(2):157-65.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tinetti ME, Baker D, Gallo WT, et al. Evaluation of restorative care vs usual care for older adults receiving an acute episode of home care. *JAMA* 2002 Apr 24; 287(16):2098-105.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tollefson GD. Short-term effects of the calcium channel blocker nimodipine (Bay-e-9736) in the management of primary degenerative dementia. *Biol Psychiatry* 1990 May 15; 27(10):1133-42.
Status: Not included because Jadad Quality Scale score less than three

Toseland RW, Diehl M, Freeman K, et al. The impact of validation group therapy on nursing home residents with dementia. *J Appl Gerontol* 1997; 16(1):31-50.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Trappler B, Cohen CI. Use of SSRIs in "very old" depressed nursing home residents. *Am J Geriatr Psychiatry* 1998; 6(1):83-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Trappler B, Cohen CI. Using fluoxetine in "very old" depressed nursing home residents. *Am J Geriatr Psychiatry* 1996; 4(3):258-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Treves TA, Korczyn AD. Denbufylline in dementia: a double-blind controlled study. *Dement Geriatr Cogn Disord* 1999 Nov; 10(6):505-10.

Status: Included

Tsai GE, Falk WE, Gunther J. A preliminary study of D-cycloserine treatment in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1998; 10(2):224-6.

Status: Cross-over trial;
Cross-over trial

Tsai GE, Falk WE, Gunther J, et al. Improved cognition in Alzheimer's disease with short-term D-cycloserine treatment. *Am J Psychiatry* 1999 Mar; 156(3):467-9.

Status: Cross-over trial;
Cross-over trial

Tsiskaridze A, Vashadze A. Comparison study of Amiridine and Piracebral effects on cognition in patients with primary dementias. *J Neural Transm Gen Sect* 1996; 103:LII.

Status: Not included because not a full article

Tsolaki M, Kapinas K, Kazis A. Nimodipine vs dihydroergocristine in the treatment of old age dementias: A blind, randomized, placebo-controlled study. *Review of Clinical Pharmacology and Pharmacokinetics, International Edition* ; (1):16-24.

Status: Article not retrievable

Tsuang MM, Lu LM, Stotsky BA, et al. Haloperidol versus thioridazine for hospitalized psychogeriatric patients: Double-blind study. *J Am Geriatr Soc* 1971 Jul; 19(7):593-600.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tully MP, Cantrill JA. Subjective outcome measurement--a primer. *Pharm World Sci* 1999 Jun; 21(3):101-9.

Status: Background article

Tune L, Brandt J, Frost JJ, et al. Physostigmine in Alzheimer's disease: Effects on cognitive functioning, cerebral glucose metabolism analyzed by positron emission tomography and cerebral blood flow analyzed by single photon emission tomography. *Acta Psychiatr Scand Suppl* 1991; 366:61-5.

Status: Cross-over trial;
Cross-over trial

Tune LE, Steele C, Cooper T. Neuroleptic drugs in the management of behavioral symptoms of Alzheimer's disease. *Psychiatr Clin North Am* 1991 Jun; 14(2):353-73.

Status: Not included because dementia population not randomized to treatment

Tuokko H, Frerichs R J, Kristjansson E. Cognitive loss without dementia: Comparing clinicians judgement to cognitive test performance. Poster session presented at the 52nd annual scientific meeting of the Gerontological Society of America. In San Francisco: 1999.

Reason for exclusion: *Background article*

Turek I, Kurland AA, Ota KY, et al. Effects of pipradrol hydrochloride on geriatric patients. *J Am Geriatr Soc* 1969 Apr; 17(4):408-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tweedy JR, Garcia CA. Lecithin treatment of cognitively impaired Parkinson's patients. *Eur J Clin Invest* 1982 Feb; 12(1):87-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tyas SL, Koval JJ, Pederson LL. Does an interaction between smoking and drinking influence the risk of Alzheimer's disease? Results from three Canadian data sets. *Stat Med* 2000 Jun 15; 19(11-12):1685-96.

Status: Background article

Udani JK. Extract of ginkgo biloba for treatment of dementia. *Integr Med* 1998; (1):43-4.

Status: Not included because not a full article

Ulmar G, Weickelt G. Anticholinergic treatment, cognition and the acetylcholine hypothesis of Alzheimer's disease (AD). *J Neural Transm Park*

Dis Dement Sect 1989; 1:144.

Status: Not included because not a full article

Underhill JA. Exercise for older people with dementia. Age Ageing 1993; 22(Suppl 3):35.

Status: Not included because not a full article

Uney JB, Jones GM, Rebeiro A, et al. The effect of long-term high dose lecithin on erythrocyte choline transport in Alzheimer patients. Biol Psychiatry 1992 Mar 15; 31(6):630-3.

Status: Not included because no extractable data relevant to review

van Asselt DZ, Pasman JW, van Lier HJ, et al. Cobalamin supplementation improves cognitive and cerebral function in older, cobalamin-deficient persons. J Gerontol A Biol Sci Med Sci 2001 Dec; 56(12):M775-M779.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

van Belle G, Arnold A. Reliability of cognitive tests used in Alzheimer's disease. Stat Med 2000 Jun 15; 19(11-12):1411-20.

Status: Background article

van Diepen E, Baillon SF, Redman J, et al. A pilot study of the physiological and behavioural effects of snoezelen in dementia. Br J Occup Ther 2002; 65(2):61-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

van Dongen MC, van Rossum E, Kessels AG, et al. The efficacy of ginkgo for elderly people with dementia and age-associated memory impairment: New results of a randomized clinical trial. J Am Geriatr Soc 2000 Oct; 48(10):1183-94.

Status: Not included because no extractable data relevant to review

van Dongen MCJM, van Rossum E, Kessels A. Effectiveness of ginkgo biloba in elderly people with beginning dementia and memory loss: Results of a randomised experiment. Tijdschr Soc Geneesk 1998; 76(Suppl):

Status: Article not retrievable

van Duijn CM, Clayton D, Chandra V, et al. Familial aggregation of Alzheimer's disease and related disorders: a collaborative re-analysis of case-control studies. EURODEM Risk Factors Research Group. Int J Epidemiol 1991; 20 Suppl 2:S13-S20

Status: Background article

van Duijn CM, Stijnen T, Hofman A. Risk factors for Alzheimer's disease: overview of the EURODEM collaborative re-analysis of case-control studies. EURODEM Risk Factors Research Group. Int J Epidemiol 1991; 20 Suppl 2:S4-12.

Status: Background article

van Duijn H, Wolters EC, Beckmann MKF. Desglycinamide arginine vasopressin (DGAVP) in Alzheimer's disease. Clin Neurol Neurosurg 1987; 2:21

Status: Not included because not a full article

Van Dyck CH, Lin CH, Robinson R, et al. The acetylcholine releaser linopirdine increases parietal regional cerebral blood flow in Alzheimer's disease. Psychopharmacologia 1997 Aug; 132(3):217-26.

Status: Included

Van Dyck CH, McMahon TJ, Rosen MI, et al. Sustained-release methylphenidate for cognitive impairment in HIV-1-infected drug abusers: A pilot study. J Neuropsychiatry Clin Neurosci 1997; 9(1):29-36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Van Dyck CH, Newhouse P. Extended-release physostigmine in Alzheimer's disease: A multicenter, double-blind, 12-week study with dose enrichment. Arch Gen Psychiatry 2000; 57(2):157-64.

Status: Included

Van Gool WA. Erratum: effect of hydroxychloroquine on progression of dementia in early alzheimer's disease: An 18-month randomised, double-blind, placebo-controlled study. Lancet 2001; 358(9288):1188.

Status: Not included because not a full article

Van Gool WA, Weinstein HC, Scheltens PK, et al. Effect of hydroxychloroquine on progression of dementia in early Alzheimer's disease: An 18-month randomised, double-blind, placebo-controlled study. Lancet 2001 Aug 11; 358(9280):455-60.

Status: Included

Van Gool WA, Lemstra AW. Cholinesterase inhibitors for Alzheimer's disease. JAMA 2003 May 14; 289(18):2359.

Status: Background article

van Heugten CM, Dekker J, Deelman BG, et al. Rehabilitation of stroke patients with apraxia: The role of additional cognitive and motor impairments. *Disabil Rehabil* 2000 Aug 15; 22(12):547-54.

Status: Background article

Van Loveren-Huyben CM, Engelaar HFWJ, Hermans MBM. Double-blind clinical and psychologic study of ergoloid mesylates (Hydergine®) in subjects with senile mental deterioration. *J Am Geriatr Soc* 1984; 32(8):584-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Van Reekum R, Clarke D, Conn D, et al. A randomized, placebo-controlled trial of the discontinuation of long-term antipsychotics in dementia. *Int Psychogeriatr* 2002; 14(2):197-210.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Van Someren EJ, Scherder EJ, Swaab DF. Transcutaneous electrical nerve stimulation (TENS) improves circadian rhythm disturbances in Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1998 Jun; 12(2):114-8.

Status: Not included because no extractable data relevant to review

Vangtorp A, Simmelsgaard H, Mellegaard M. Experience with a new butyrophenone derivative (Buronil). *Acta Psychiatr Scand Suppl* 1968 Jan 1; 203:235-8.

Status: Not included because dementia population not randomized to treatment

Vecchi GP, Chiari G, Cipolli C, et al. Acetyl-L-carnitine treatment of mental impairment in the elderly: Evidence from a multicentre study. *Arch Gerontol Geriatr* 1991; (Suppl 2):159-68.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vellas B, Inglis F, Potkin S, et al. Interim results from an international clinical trial with rivastigmine evaluating a 2-week titration rate in mild to severe Alzheimer's disease patients. *Int J Geriatr Psychopharmacol* 1998; (3):140-4.

Status: Not included because dementia population not randomized to treatment

Velligan DI, Newcomer J, Pultz J, et al. Does cognitive function improve with quetiapine in comparison to haloperidol? *Schizophr Res* 2002

Jan 15; 53(3):239-48.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vennerica A, Shanks MF, Staff RT, et al. Cerebral blood flow and cognitive responses to rivastigmine treatment in Alzheimer's disease. *NeuroReport* 2002; 13(1):83-7.

Status: Not included because dementia population not randomized to treatment

Verhey FR, Lodder J, Rozendaal N, et al. Comparison of seven sets of criteria used for the diagnosis of vascular dementia. *Neuroepidemiology* 1996; 15(3):166-72.

Status: Background article

Veroff AE, Bodick NC, Offen WW, et al. Efficacy of xanomeline in Alzheimer's disease: Cognitive improvement measured using the Computerized Neuropsychological Test Battery (CNTB). *Alzheimer Dis Assoc Disord* 1998 Dec; 12(4):304-12.

Status: Companion of an included article

Versiani M, da Silva JA, Mundim FD. Loxapine versus thioridazine in the treatment of organic psychosis. *J Int Med Res* 1980; 8(1):22-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vespa A, Gori G, Spazzafumo L. Evaluation of non-pharmacological intervention on antisocial behavior in patients suffering from Alzheimer's disease in a day care center. *Arch Gerontol Geriatr* 2002; 34(1):1-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Vida S, Gauthier L, Gauthier S. Canadian collaborative study of tetrahydroaminoacridine (THA) and lecithin treatment of Alzheimer's disease: Effect on mood. *Can J Psychiatry* 1989; 34(3):165-70.

Status: Not included because dementia population not randomized to treatment

Villardita C, Parini J, Grioli S, et al. Clinical and neuropsychological study with oxiracetam versus placebo in patients with mild to moderate dementia. *J Neural Transm Suppl* 1987; 24:293-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Villardita C, Grioli S, Lomeo C, et al. Clinical studies with oxiracetam in patients with dementia of Alzheimer type and multi-infarct dementia of mild to moderate degree. *Neuropsychobiology* 1992; 25(1):24-8.

Status: Included

Viukari M, Linnoila M. Effect of methyldopa on tardive dyskinesia in psychogeriatric patients. *Curr Ther Res Clin Exp* ; 18(3):417-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Viukari M, Salo H, Lamminsivu U, et al. Tolerance and serum levels of haloperidol during parenteral and oral haloperidol treatment in geriatric patients. *Acta Psychiatr Scand* 1982 Apr; 65(4):301-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Viukari M, Jaatinen P, Kylmamaa T. Flunitrazepam, nitrazepam and psychomotor skills in psychogeriatric patients. *Curr Ther Res Clin Exp* 1983; 33(5):828-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vojtechovsky M, Sobotkova J, Wodniakova J. Secatoxin Spofa-super(R) in the treatment of chronic organic psychosyndrome. *Act Nerv Super (Praha)* 1981; 23(3):232-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Volicer L, Stelly M, Morris J, et al. Effects of dronabinol on anorexia and disturbed behavior in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 1997 Sep; 12(9):913-9.

*Status: Cross-over trial;
Cross-over trial*

Vroulis GA, Smith RC, Brinkman S, et al. The effects of lecithin on memory in patients with senile dementia of the Alzheimer's type. *Psychopharmacol Bull* 1981; 17(1):127-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Waegemans T, Wilsher CR, Danniau A, et al. Clinical efficacy of piracetam in cognitive impairment: A meta-analysis. *Dement Geriatr Cogn Disord* 2002; 13(4):217-24.

Status: Background article

Wakelin JS. Fluvoxamine in the treatment of the older depressed patient; Double-blind, placebo-controlled data. *Int Clin Psychopharmacol* 1986 Jul; 1(3):221-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Walikonis JE, Lennon VA. Radioimmunoassay for glutamic acid decarboxylase (GAD65) autoantibodies as a diagnostic aid for stiff-man syndrome and a correlate of susceptibility to type 1 diabetes mellitus. *Mayo Clin Proc* 1998 Dec; 73(12):1161-6.

Status: Background article

Walker C. Ergoloid mesylates vs. Alzheimer's: The latest round. *Geriatrics* 1990; 45(12):22, 24.

Status: Not included because not a full article

Walsh JS, Welch HG, Larson EB. Survival of outpatients with Alzheimer-type dementia. *Ann Intern Med* 1990 Sep 15; 113(6):429-34.

Status: Background article

Walshe TM. The use of nimodipine in vascular dementia. *J Neurol* 1985; Vol 232:74

Status: Not included because not a full article

Walzl M, Walzl B, Lechner H. Results of a two-month follow-up after single heparin-induced extracorporeal LDL precipitation in vascular dementia. *National Stroke Association* 1994; (3):179-83.

Status : Not included because no extractable data relevant to review

Walzl M. A promising approach to the treatment of multi-infarct dementia. *Neurobiol Aging* 2000 Mar; 21(2):283-7.

Status: Not included because Jadad Quality Scale score less than three

Wang D, Huang X, Du S. A clinical trial on yu cong tang in treatment of senile dementia. *J Tradit Chin Med* 1999 Mar; 19(1):32-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wang PN, Liao SQ, Liu RS, et al. Effects of estrogen on cognition, mood, and cerebral blood flow in AD: A controlled study. *Neurology* 2000 Jun 13; 54(11):2061-6.

Status: Included

Wang X, Xie Z. A clinical study on the effect of reinforcement of kidney on senile brain functions. *J Tradit Chin Med* 1997 Jun; 17(2):92-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ward CR, Los Kamp. The effects of participation in an intergenerational program on the behavior of residents with dementia. *Activities Adaptation Aging* 1996; 20(4):61-76.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wardle J, Rogers P, Judd P, et al. Randomized trial of the effects of cholesterol-lowering dietary treatment on psychological function. *Am J Med* 2000 May; 108(7):547-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Waring SC, Rocca WA, Petersen RC, et al. Postmenopausal estrogen replacement therapy and risk of AD: A population-based study. *Neurology* 1999; 52(5):965-70.
Status: Not included because dementia population not randomized to treatment

Warren PA, Dunn L, Jackson-Clark A. The Medicare Alzheimer's Project in Portland, Oregon. *Pride Inst J Long Term Home Health Care* 1991; 10(2):20-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Watkins PB, Zimmerman HJ, Knapp MJ, et al. Hepatotoxic effects of tacrine administration in patients with Alzheimer's disease. *JAMA* 1994 Apr 6; 271(13):992-8.
Status: Not included because dementia population not randomized to treatment

Watson NM, Wells TJ, Cox C. Rocking chair therapy for dementia patients: Its effect on psychosocial well-being and balance. *Am J Alzheimers Dis* 1998; (6):296-308.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Weiner MF, Tractenberg RE, Sano M, et al. No long-term effect of behavioral treatment on psychotropic drug use for agitation in Alzheimer's disease patients. *J Geriatr Psychiatry Neurol* 2002; 15(2):95-8.
Status: Not included because does not meet criteria for treatment for dementia patients

Weingartner H, Kaye W, Gold P, et al. Vasopressin treatment of cognitive dysfunction in progressive dementia. *Life Sci* 1981; 29(26):2721-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Weingartner H, Buchsbaum MS, Linnoila M. Zimelidine effects on memory impairments produced by ethanol. *Life Sci* 1983 Nov 28; 33(22):2159-63.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Weinstein HC, Teunisse S, Van Gool WA. Tetrahydroaminoacridine and lecithin in the treatment of Alzheimer's disease. Effect on cognition, functioning in daily life, behavioural disturbances and burden experienced by the carers. *J Neurol* 1991 Feb; 238(1):34-8.
Status: Included

Weiser M, Rotmensch HH, Korczyn AD, et al. A pilot, randomized, open-label trial assessing safety and pharmacokinetic parameters of co-administration of rivastigmine with risperidone in dementia patients with behavioral disturbances. *Int J Geriatr Psychiatry* 2002 Apr; 17(4):343-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wells D, Dawson P, Sidani S, et al. The benefits of abilities-focused morning care for residents with dementia and their caregivers. *Perspectives (Montclair)* 2000; 24(1):17.
Status: Not included because does not meet criteria for treatment for dementia patients

Wentzel C, Rose H, Rockwood K. Measurement of the influence of the physical environment on adverse health outcomes: Technical report from the Canadian Study of Health and Aging. *Int Psychogeriatr* 2001; 13(Suppl 1):215-21.
Status: Background article

Wesensten NJ, Balkin TJ, Belenky GL. Effects of daytime administration of zolpidem versus triazolam on memory. *Eur J Clin Pharmacol* 1995; 48(2):115-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wesensten NJ, Balkin TJ, Davis HQ, et al. Reversal of triazolam- and zolpidem-induced memory impairment by flumazenil. *Psychopharmacologia* 1995 Sep; 121(2):242-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wesnes K, Simmons D, Rook M, et al. A double-blind placebo-controlled trial of Tanakan in the treatment of idiopathic cognitive impairment in the elderly. *Hum Psychopharmacol* 1987; 2:159-69.
Status : Not included because dementia population not defined by DSM, NINCDS or ICD

Wesnes KA, McKeith IG, Ferrara R, et al. Effects of rivastigmine on cognitive function in dementia with lewy bodies: A randomised placebo-controlled international study using the cognitive drug research computerised assessment system. *Dement Geriatr Cogn Disord* 2002; 13(3):183-92.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

West S, Kling V, Carey TS et al. Systems to rate the strength of scientific evidence. AHRQ publication No. 02-E016. Rockville, MD: Agency for Healthcare Research and Quality US Department of Health and Human Services. 2002.
Status: Background article

Westreich G, Alter M, Lundgren S. Effect of cyclandelate on dementia. *Stroke* 1975 Sep; 6(5):535-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wettstein A. No effect from double-blind trial of physostigmine and lecithin in Alzheimer disease. *Ann Neurol* 1983; 13(2):210-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wettstein A, Spiegel R. Clinical trial with the cholinergic drug RS 86 in Alzheimer's disease (AD) and senile dementia of the Alzheimer type (SDAT). *Psychopharmacologia* 1984; 84(4):572-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Weyer G, Erzigkeit H, Hadler D, et al. Efficacy and safety of idebenone in the long-term treatment of Alzheimer's disease: A double-blind, placebo controlled multicentre study. *Hum Psychopharmacol* 1996; (1):53-65.
Status: Companion of an included article

Weyer G, Babej-Dolle RM, Hadler D, et al. A controlled study of 2 doses of idebenone in the treatment of Alzheimer's disease.

Neuropsychobiology 1997; 36(2):73-82.
Status: Included

Weyer G, Eul A, Milde K, et al. Cyclandelate in the treatment of patients with mild to moderate primary degenerative dementia of the Alzheimer type or vascular dementia: Experience from a placebo controlled multi-center study. *Pharmacopsychiatry* 2000 May; 33(3):89-97.
Status: Included

White CM, Dicks RS. Rivastigmine: An acetylcholinesterase inhibitor for patients with Alzheimer's disease. *Hosp Formul* 1999; 34(6):493-9.
Status: Not included because dementia population not randomized to treatment

White HK, Levin ED. Four-week nicotine skin patch treatment effects on cognitive performance in Alzheimer's disease. *Psychopharmacologia* 1999 Apr; 143(2):158-65.
Status: Cross-over trial; Cross-over trial

White JC, Christensen JF, Singer CM. Methylphenidate as a treatment for depression in acquired immunodeficiency syndrome: an n-of-1 trial. *J Clin Psychiatry* 1992 May; 53(5):153-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

White L, Crovello JN, Rosenberg SN, et al. Evaluation of isobaric oxygenation for the aged with cognitive impairment: Pilot study. *J Am Geriatr Soc* 1975 Feb; 23(2):80-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Whitehead A, Whitehead J. A general parametric approach to the meta-analysis of randomized clinical trials. *Stat Med* 1991 Nov; 10(11):1665-77.
Status: Background article

Whitehead J, Matsushita T. Stopping clinical trials because of treatment ineffectiveness: a comparison of a futility design with a method of stochastic curtailment. *Stat Med* 2003 Mar 15; 22(5):677-87.
Status: Background article

Whitehouse PJ, Winblad B, Shostak D, et al. First International Pharmacoeconomic Conference on Alzheimer's Disease: report and summary. *Alzheimer Dis Assoc Disord* 1998 Dec; 12(4):266-

80.

Status: Background article

Whitehouse PJ, Kittner B, Roessner M, et al. Clinical trial designs for demonstrating disease-course-altering effects in dementia. *Alzheimer Dis Assoc Disord* 1998 Dec; 12(4):281-94.

Status: Background article

Whitlatch CJ, Zarit SH, von Eye A. Efficacy of interventions with caregivers: A reanalysis. *Gerontologist* 1991 Feb; (1):9-14.

Status: Not included because does not meet criteria for treatment for dementia patients

Whitlatch CJ, Zarit SH, Goodwin PE, et al. Influence of the success of psychoeducational interventions on the course of family care. *Clin Gerontol* 1995; (1):17-30.

Status: Not included because does not meet criteria for treatment for dementia patients

Wiener PK, Kiessens DN, Klimstra S, et al. A short-term inpatient program for agitated demented nursing home residents. *Int J Geriatr Psychiatry* 2001; 16(9):866-72.

Status: Not included because dementia population not randomized to treatment

Wilcock G, Mobius HJ, Stoffler A. A double-blind, placebo-controlled multicentre study of memantine in mild to moderate vascular dementia (MMM500). *Int Clin Psychopharmacol* 2002; 17(6):297-305.

Status: Included

Wilcock GK, Scott M, Pearsall T, et al. Galanthamine and the treatment of Alzheimer's disease. *Int J Geriatr Psychiatry* 1993; 8(9):781-2.

Status: Not included because not a full article

Wilcock GK, Surmon DJ, Scott M, et al. An evaluation of the efficacy and safety of tetrahydroaminoacridine (THA) without lecithin in the treatment of Alzheimer's disease. *Age Ageing* 1993; 22(5):316-24.

*Status: Cross-over trial;
Cross-over trial*

Wilcock GK, Lilienfeld S, Gaens E. Efficacy and safety of galantamine in patients with mild to moderate Alzheimer's disease: Multicentre randomised controlled trial. Galantamine International-1 Study Group. *BMJ* 2000 Dec 9;

321(7274):1445-9.

Status: Included

Wilcock GK. Erratum: Efficacy and safety of galantamine in patients with mild to moderate Alzheimer's disease: Multicentre randomised controlled trial. *BMJ* 2001; 322(7278):90.

Status: Companion of an included article

Wild R, Pettit T, Burns A. Cholinesterase inhibitors for Lewy Body Dementia (Cochrane Protocol). In: *The Cochrane Library*, 2002. Issue 2. Oxford: Update Software

Status: Background article

Wilkinson D. Clinical experience with Donepezil (Aricept) in the UK. *J Neural Transm Suppl* 1998; 54:311-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wilkinson D, Srikumar S, Shaw K, et al. Drama and movement therapy in dementia: A pilot study. *Arts in Psychotherapy* 1998; (3):195-201.

Status: Not included because dementia population not randomized to treatment

Wilkinson D, Murray J. Galantamine: A randomized, double-blind, dose comparison in patients with Alzheimer's disease. *Int J Geriatr Psychiatry* 2001 Sep; 16(9):852-7.

Status: Included

Wilkinson DG, Hock C, Farlow M, et al. Galantamine provides broad benefits in patients with 'advanced moderate' Alzheimer's disease (MMSE < or = 12) for up to six months. *Int J Clin Pract* 2002 Sep; 56(7):509-14.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wilkinson DG, Passmore AP, Bullock R, et al. A multinational, randomised, 12-week, comparative study of donepezil and rivastigmine in patients with mild to moderate Alzheimer's disease. *Int J Clin Pract* 2002 Jul; 56(6):441-6.

Status: Not included because Jadad Quality Scale score less than three

Williams PS, Rands G, Orrel M, et al. Aspirin for vascular dementia. In: *The Cochrane Library*, 2000. Issue 4. Oxford: Update Software.

Status: Background article

Williams R, Reeve W, Ivison D, et al. Use of environmental manipulation and modified informal reality orientation with institutionalized, confused elderly subjects: A replication. *Age Ageing* 1987; 16(5):315-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Williams R. Optimal dosing with risperidone: Updated recommendations. *J Clin Psychiatry* 2001 Apr; 62(4):282-9.

Status: Background article

Willis SL, Schaie KW. Training the elderly on the ability factors of spatial orientation and inductive reasoning. *Psychol Aging* 1986 Sep; 1(3):239-47.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wilson AL, Langley LK, Monley J, et al. Nicotine patches in Alzheimer's disease: Pilot study on learning, memory, and safety. *Pharmacol Biochem Behav* ; 51(2-3):509-3.

Status: Not included because dementia population not randomized to treatment

Wilson RS, Martin EM. New intrathecal drugs in Alzheimer's disease and psychometric testing. *Ann N Y Acad Sci* 1988; 531:180-6.

Status: Not included because dementia population not randomized to treatment

Winblad B, Poritis N. Memantine in severe dementia: Results of the 9M-Best Study (Benefit and efficacy in severely demented patients during treatment with memantine). *Int J Geriatr Psychiatry* 1999 Feb; 14(2):135-46.

Status: Included

Winblad B, Bonura ML, Rossini BM, et al. Nicergoline in the treatment of mild-to-moderate alzheimer's disease: A European multicentre trial. *Clin Drug Investig* 2001a; (9):621-32.

Status: Included

Winblad B, Brodaty H, Gauthier S, et al. Pharmacotherapy of Alzheimer's disease: Is there a need to redefine treatment success? *Int J Geriatr Psychiatry* 2001 Jul; 16(7):653-66.

Status: Background article

Winblad B, Engedal K, Soininen H, et al. A 1-year, randomized, placebo-controlled study of donepezil in patients with mild to moderate AD. *Neurology* 2001b; 57(3):489-95.

Status: Included

Wincor MZ. Ginkgo biloba for dementia: A reasonable alternative? *J Am Pharm Assoc (Wash)* 1999 May; 39(3):415-6.

Status: Not included because dementia population not randomized to treatment

Winther K, Randlov C, Rein E, et al. Effects of Ginkgo biloba extract on cognitive function and blood pressure in elderly subjects. *Curr Ther Res Clin Exp* 1998; 59(12):881-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wishart L, Macerollo J, Loney P, et al. "Special steps": An effective visiting/walking program for persons with cognitive impairment. *Can J Nurs Res* 2000 Mar; 31(4):57-71.

Status: Not included because does not meet criteria for treatment for dementia patients

Wolfgang SA. Olanzapine in whole, not half, tablets for psychosis from Alzheimer's dementia. *Am J Health Syst Pharm* 1999 Nov 1; 56(21):2245-6.

Status: Not included because not a full article

Wolfson C, Moride Y, Perrault A et al. Technology Report: Drug treatments for Alzheimer's disease. I. A comparative analysis of clinical trials. Ottawa: 2000.

Status: Background article

Wolfson C, Moride Y, Perrault A et al. Drug treatments for Alzheimer's disease. II. A review of outcome measures in clinical trials. Canadian Coordinating Office for Health Technology Assessment. 2000.

Status: Background article

Wolfson C, Wolfson DB, Asgharian M, et al. A reevaluation of the duration of survival after the onset of dementia. *N Engl J Med* 2001 Apr 12; 344(15):1111-6.

Status: Background article

Wolters EC, Riekkinen P, Lowenthal A, et al. DGAVP (Org 5667) in early Alzheimer's disease patients: An international double-blind, placebo-controlled, multicenter trial. *Neurology* 1990 Jul; 40(7):1099-101.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wong WJ, Liu HC, Fuh JL, et al. A double-blind, placebo-controlled study of tacrine in Chinese patients with Alzheimer's disease. *Dement*

Geriatr Cogn Disord 1999 Jul; 10(4):289-94.
Status: Included

Wong YT, Packer H. Penicillin versus penicillin-malaria in the treatment of dementia paralytica. Br J Vener Dis 2000; 25(1):39.
Status: Not included because not a full article

Wood PC, Castleden CM. A double-blind, placebo controlled, multicentre study of tacrine for Alzheimer's disease. Int J Geriatr Psychiatry 1994; 9(8):649-54.
Status: Included

Woods B. Promoting well-being and independence for people with dementia. Int J Geriatr Psychiatry 1999; 14(2):97-109.
Status: Not included because dementia population not randomized to treatment

Wooltorton E. Risperidone (Risperdal): Increased rate of cerebrovascular events in dementia trials. CMAJ 2002 Nov 26; 167(11):1269-70.
Status: Not included because not a full article

World Health Organization. The tenth revision of the International Classification of Diseases and relative health problems (ICD-10). Geneva: WHO; 1992.
Status: Background article

Wouters-Wesseling W, Wouters AE, Kleijer CN, et al. Study of the effect of a liquid nutrition supplement on the nutritional status of psycho-geriatric nursing home patients. Eur J Clin Nutr 2002 Mar; 56(3):245-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wright LK, Litaker M, Laraia MT, et al. Continuum of care for Alzheimer's disease: A nurse education and counseling program. Issues Ment Health Nurs 2001 Apr; 22(3):231-52.
Status: Not included because does not meet criteria for treatment for dementia patients

Wroblewski T, Silvestre J, Castaner J. JTP-4819. Cognition enhancer, prolyl endopeptidase inhibitor. Drugs of the Future 1998; 23(4):384-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wu SF, Xie SZ, Lu WJ, et al. Comparison of the clinical effects of capsules and tablets in a randomized double-blind treatment of mild and

moderate Alzheimer's disease. Shanghai Medical & Pharmaceutical Journal 1999; (8):36-8.
Status: Article not retrievable

Xiao S, Yan H, Yao P, et al. The efficacy of cerebrolysin in patients with vascular dementia: Results of a Chinese multicentre, randomised, double-blind, placebo-controlled trial. Hong Kong Journal of Psychiatry 1999; (2):13-9.
Status: Included

Xiao S, Yan H, Yao P, et al. Efficacy of FPF 1070 (cerebrolysin) in patients with Alzheimer's disease: A multicentre, randomised, double-blind, placebo-controlled trial. Clin Drug Investig 2000; (1):43-53.
Status: Included

Xiao SF, Yan HE, Yao BF, et al. A multi-center, double blind, placebo-controlled trial of the efficacy of cerebrolysin in treatment of vascular dementia. J Clin Psychol Med 1999; (1):1-3.
Status: Article not retrievable

XiaoSF, Yan HQ, Yao PF, et al. The efficacy of cerebrolysin in patients with Alzheimer disease. Mod Rehab 2000; (11):1624-5.
Status: Article not retrievable

Xu H, Shao N, Cui D, et al. A clinical study of yi zhi capsules in prevention of vascular dementia. J Tradit Chin Med 2000 Mar; 20(1):10-3.
Status: Not included because Jadad Quality Scale score less than three

Xu SS, Gao ZX, Weng Z, et al. Efficacy of tablet huperzine-A on memory, cognition, and behavior in Alzheimer's disease. Zhongguo Yao Li Xue Bao/Acta Pharmacologica Sinica 1995 Sep; 16(5):391-5.
Status: Included

Xu SS, Cai ZY, Qu ZW, et al. Huperzine-A in capsules and tablets for treating patients with Alzheimer disease. Zhongguo Yao Li Xue Bao/Acta Pharmacologica Sinica 1999; 20(6):486-90.
Status: Included

Xu SS, Zhi X, Weng Z, et al. Efficacy of tablet huperzine-A on memory, cognition and behavior in Alzheimer's disease. Int Med J 1997; (2):127-31.
Status: Duplicate publication

Yaffe K, Krueger K, Sarkar S, et al. Cognitive function in postmenopausal women treated with raloxifene. *N Engl J Med* 2001 Apr 19; 344(16):1207-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yamamoto Y, Akiguchi I, Oiwa K, et al. Twenty-four-hour blood pressure changes in the course of lacunar disease. *Cardiovasc Dis* 2001; 11(2):100-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yehuda S, Rabinovitz S, Carasso RL, et al. Essential fatty acids preparation (SR-3) improves Alzheimer's patients quality of life. *Int J Neurosci* 1996 Nov; 87(3-4):141-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yesavage JA, Hollister LE, Burian E. Dihydroergotoxine: 6-mg versus 3-mg dosage in the treatment of senile dementia. Preliminary report. *J Am Geriatr Soc* 1979 Feb; 27(2):80-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yesavage JA, Westphal J, Rush L. Senile dementia: Combined pharmacologic and psychologic treatment. *J Am Geriatr Soc* 1981; 29(4):164-71.

Status: Not included because dementia population not randomized to treatment

Yesavage JA, Tinklenberg JR, Hollister LE, et al. Effect of nafronyl on lactate and pyruvate in the cerebrospinal fluid of patients with senile dementia. *J Am Geriatr Soc* 1982; 30(2):105-8.

Status: Not included because dementia population not randomized to treatment

Zanetti O, Frisoni GB, De Leo D, et al. Reality orientation therapy in Alzheimer disease: Useful or not? A controlled study. *Alzheimer Dis Assoc Disord* 1995; 9(3):132-8.

Status: Not included because dementia population not randomized to treatment

Zanetti O, Bianchetti A, Trabucchi M. Cost effectiveness of non pharmacological interventions in Alzheimer's disease. *J Am Geriatr Soc* 1998 Nov; 46(11):1481

Status: Not included because not a full article

Zanetti O, Zanieri G, Di Giovanni G, et al. Effectiveness of procedural memory stimulation in mild Alzheimer's disease patients: A controlled study. *Neuropsychol Rehab* 2001; (3-4):263-4.

Status: Not included because dementia population not randomized to treatment

Zank S, Schacke C. Evaluation of geriatric day care units: effects on patients and caregivers. *J Gerontol B Psychol Sci Soc Sci* 2002 Jul; 57(4):348-57.

Status: Not included because does not meet criteria for treatment for dementia patients

Zappoli R, Arnetoli G, Paganini M, et al. Contingent negative variation and reaction time in patients with presenile idiopathic cognitive decline and presenile Alzheimer-type dementia.

Preliminary report on long-term nicergoline treatment. *Neuropsychobiology* 1987; 18(3):149-54.

Status: Not included because Jadad Quality Scale score less than three

Zappoli R, Arnetoli G, Paganini M, et al. Topographic bit-mapped event-related neurocognitive potentials and clinical status in patients with primary presenile mental decline chronically treated with nicergoline. *Curr Ther Res Clin Exp* 1991; 49(6):1078-97.

Status: Not included because Jadad Quality Scale score less than three

Zarit SH, Zarit JM, Reeve KE. Memory training for severe memory loss: Effects on senile dementia patients and their families. *Gerontologist* 1982; 22(4):373-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Zec RF, Trivedi MA. The effects of estrogen replacement therapy on neuropsychological functioning in postmenopausal women with and without dementia: A critical and theoretical review. *Neuropsychol Rev* 2002; 12(2):65-109.

Status: Background article

Zemlan FP, Folks DG, Goldstein BJ, et al. Velnacrine for the treatment of Alzheimer's disease: A double-blind, placebo-controlled trial. *J Neural Transm Gen Sect* 1996; 103(8-9):1105-16.

Status: Included

Zemlan FP, Keys M, Richter RW, et al. Double-blind placebo-controlled study of velnacrine in

Alzheimer's disease. Life Sci 1996; 58(21):1823-32.

Status: Not included because Jadad Quality Scale score less than three

Zhang LX, Wang TH, Zhong XS. Comparison of effects between malloryl and chlorpromazine on type I dementia praecox. J Guandong Med Coll 1999; (2):122-3.

Status: Article not retrievable

Zhou XH, Higgs RE. Assessing the relative accuracies of two screening tests in the presence of verification bias. Stat Med 2000 Jun 15; 19(11-12):1697-705.

Status: Background article

Ziemba C, Foster G, Neufeld R, et al. Haloperidol holiday: Is it a beneficial vacation for some nursing home residents? Clin Gerontol 1997; 17(3):15-24.

Status: Not included because no extractable data relevant to review

Zissis NP, Alevizos V, Dontas AS. Flunarizine, an inhibitor of Casup +sup 2-induced vascular constriction in geriatric patients. Curr Ther Res Clin Exp 1991; 29(31):395-400.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Zivadinov R, Rudick RA, De Masi R, et al. Effects of IV methylprednisolone on brain atrophy in relapsing-remitting MS. Neurology 2001 Oct 9; 57(7):1239-47.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Zwerling I, Plutchik R, Hotz M, et al. Effects of a procaine preparation (Gerovital H3) in hospitalized geriatric patients: A double-blind study. J Am Geriatr Soc 1975 Aug; 23(8):355-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Zill P, Burger K, Behrens S, et al. Polymorphisms in the alpha-2 macroglobulin gene in psychogeriatric patients. Neurosci Lett 2000 Nov 17; 294(2):69-72.

Status: Not included because does not meet criteria for treatment for dementia patients

Zimmer JG, Eggert GM, Chiverton P. Individual versus team case management in optimizing community care for chronically ill patients with dementia. J Aging Health 1990; 2(3):357-72.

Status: Not included because does not meet criteria for treatment for dementia patients

Zisselman MH, Rovner BW, Shmueli Y, et al. A pet therapy intervention with geriatric psychiatry inpatients. Am J Occup Ther 1996 Jan; 50(1):47-51.

Pharmacological Treatment of Dementia

Appendixes

- Appendix A: Search Strings
- Appendix B: Sample Data Extraction Forms
- Appendix C: Evidence Tables
- Appendix D: Peer Reviewers
- Appendix E: Outcome measures
- Appendix F: List of excluded studies

Prepared for:

Agency for Healthcare Research and Quality
U.S. Department of Health and Human Services
2101 East Jefferson St.
Rockville, MD 20852
<http://www.ahrq.gov>
Contract No. 290-02-0020 Task Order #1

Prepared by:

McMaster University
Evidence-based Practice Center based at
McMaster University, Faculty of Health Sciences
Hamilton, Ontario, Canada

Appendix A

Search strategies

Appendix A.

Database: Cochrane Central Register of Controlled Trials <4th Quarter 2002> on OVID

Search strategy executed on February 3, 2003

Same search executed on February 4th, 2003 on the online version of the Cochrane Library for the new references <1st Quarter 2003>:

- 1 (mild cognitive impairment or MCI).tw.
- 2 ((cognitive impairment not dementia) or CIND).tw.
- 3 ((cognitive loss not dementia) or CLOND).tw.
- 4 Delirium, Dementia, Amnestic, Cognitive Disorders/
5 exp Amnesia/
6 Cognition Disorders/
7 exp Dementia/
8 exp tauopathies/
9 dement:.tw.
- 10 Alzheimer:.tw.
- 11 Huntington disease/
12 Lewy: ajd8 bod:.tw.
- 13 ((cognit: or memory or mental:) adj8 (decli: or impair: or los: or deteriorat:)).tw.
- 14 (chronic adj8 cerebrovascular).tw.
- 15 supra-nuclear palsy.tw.
- 16 (normal pressure hydrocephalus adj8 shunt:).tw.
- 17 benign senescent forgetfulness.tw.
- 18 (cerebr: adj8 deteriorat:).tw.
- 19 cerebr: ajd8 insufficien:.tw.
- 20 (confusion: or confused).tw.
- 21 (pick: adj8 disease).tw.
- 22 (creutzfeldt: or JCD: or CJD:).tw.
- 23 (Huntington: or Huntingdon).tw.
- 24 Binswanger:.tw.
- 25 brain atrophy.tw.
- 26 exp Cerebral Amyloid Angiopathy/
27 neurofibrillary tangles/
28 senile plaques/
29 neuropil threads/
30 spongiform encephalopathy.tw.
- 31 exp Hypothyroidism/
32 neurosyphilis/
33 exp amyloid beta-protein/ not (Down syndrome/ or trisomy 21.tw.)

- 34 (CADASIL or cerebral autosomal dominant ischemia with subcortical leukoencephalopathy).tw.
- 35 (corticobasal ganglionic degeneration or cortical basal degeneration or corticobasal ganglionic degeneration).tw.
- 36 multisystem atrophy.tw.
- 37 exp alcohol amnestic disorder/
- 38 (alcohol adj3 amnestic).tw.
- 39 or/1-38

- 1 randomized controlled trial.pt.
- 2 controlled clinical trial.pt.
- 3 controlled clinical trials/
- 4 (clinical trials, phase II or clinical trials, phase III or clinical trials, phase IV or multicenter studies).sh.
- 5 random allocation.sh.
- 6 double blind method.sh.
- 7 cross-over studies.sh.
- 8 single-blind method.sh.
- 9 clinical trial.pt.
- 10 (clin: adj25 trial:).ti,ab.
- 11 ((singl: or doubl: or trebl: or tripl:) adj25 (blind: or mask:)).ti,ab.
- 12 placebos.sh.
- 13 placebo:.ti,ab.
- 14 random:.ti,ab.
- 15 or/1-14
- 16 comparative study.sh.
- 17 exp evaluation studies/
- 18 follow up studies.sh.
- 19 prospective studies.sh.
- 20 or/16-19
- 21 (tu or th).xs.
- 22 treatment outcome/
- 23 exp therapeutics/
- 24 or/21-23
- 25 20 and 24
- 26 15 or 25
- 27 (mild cognitive impairment or MCI).tw.
- 28 ((cognitive impairment not dementia) or CIND).tw.
- 29 ((cognitive loss not dementia) or CLOND).tw.
- 30 exp dementia/
- 31 exp tauopathies/
- 32 (dement: or alzheimer:).tw.
- 33 amentia.tw.
- 34 frontotemporal lobar degeneration.tw.
- 35 hiv-associated cognitive motor complex.tw.
- 36 encephalopathy, aids.tw.
- 37 encephalopathy, hiv.tw.
- 38 mesulam syndrome.tw.
- 39 progressive nonfluent aphasia.tw.
- 40 binswanger disease.tw.
- 41 binswanger encephalopathy.tw.
- 42 leukoencephalopathy, subcortical.tw.

43 subcortical arteriosclerotic encephalopathy.tw.
 44 chronic progressive subcortical encephalopathy.tw. or alcohol amnestic disorder/ or
 alcohol induced disorders, nervous system/ or (alcohol adj3 amnestic).tw. or (alcohol
 adj2 dysmestic).tw. or (ethanol adj3 nervous system disorders).tw. or ethyl alcohol abuse
 neurologic syndromes.tw.
 45 (lewy: bod: adj8 disease).tw.
 46 brain atrophy, circumscribed lobar.tw.
 47 (pick: adj8 disease).tw.
 48 exp amyloid beta-protein/ not (down syndrome/ or trisomy 21.tw.)
 49 exp cerebral amyloid angiopathy/
 50 neurofilament proteins/
 51 tau proteins/
 52 neurofibrillary tangles/
 53 neuropil threads/
 54 senile plaques/
 55 (Corticobasil ganglionic degeneration or cortical basal degeneration or cortica).tw.
 56 (CADASIL or Cerebral autosomal dominant ischemia with subcortical
 leukoencephalopathy).tw.
 57 Multisystems atrophy.tw.
 58 huntington disease/
 59 hydrocephalus, normal pressure/
 60 Creutzfeldt-Jakob syndrome/
 61 spongiform encephalopathy.tw.
 62 (cjd or jcd).tw.
 63 Creutzfeldt-Jakob disease.tw.
 64 spongiform encephalopathy.tw.
 65 exp Hypothyroidism/
 66 exp Vitamin B 12 Deficiency/
 67 exp Neurosyphilis/
 68 or/27-67
 69 26 and 68
 70 animal.sh.
 71 69 not 70
 72 71 not (comment or editorial or news or letter).pt.
 73 72 and eng.la.
 74 limit 73 to yr=1998-2003

Database: EMBASE <1996 to 2003 Week 5> on OVID
Search strategy executed on February 6, 2003

-
- 1 (mild cognitive impairment or MCI).tw.
 - 2 ((cognitive impairment not dementia) or CIND).tw.
 - 3 ((cognitive loss not dementia) or CLOND).tw.
 - 4 exp dementia/
 - 5 (dement: or alzheimer:).tw.
 - 6 amentia.tw.
 - 7 frontotemporal lobar degeneration.tw.
 - 8 hiv-associated cognitive motor complex.tw.
 - 9 encephalopathy, aids.tw.
 - 10 encephalopathy, hiv.tw.
 - 11 mesulam syndrome.tw.
 - 12 progressive nonfluent aphasia.tw.
 - 13 binswanger disease.tw.
 - 14 binswanger encephalopathy.tw.
 - 15 leukoencephalopathy, subcortical.tw.
 - 16 subcortical arteriosclerotic encephalopathy.tw.
 - 17 chronic progressive subcortical encephalopathy.tw.
 - 18 exp Korsakoff psychosis/ or exp Wernicke Korsakoff syndrome/
 - 19 (alcohol adj3 amnestic).tw.
 - 20 (alcohol adj2 dysmnesic).tw.
 - 21 (ethanol adj3 nervous system disorders).tw.
 - 22 ethyl alcohol abuse neurologic syndromes.tw.
 - 23 (Lewy: bod: adj8 disease).tw.
 - 24 brain atrophy, circumscribed lobar.tw.
 - 25 (Pick: adj8 disease).tw.
 - 26 exp brain atrophy/ or exp brain cortex atrophy/ or exp brain degeneration/ or exp corticobasal degeneration/ or exp lewy body/ or exp neurofibrillary tangle/ or exp neuropil thread/ or exp organic brain syndrome/
 - 27 exp amyloid beta-protein/ not (exp Down syndrome/ or trisomy 21.tw.)
 - 28 exp Vascular Amyloidosis/
 - 29 exp Neurofilament Protein/
 - 30 Tau Protein/
 - 31 Neurofibrillary Tangle/
 - 32 Neuropil Thread/
 - 33 Senile Plaque/
 - 34 (corticobasal ganglionic degeneration or cortical basal degeneration or corticobasal degeneration).tw.
 - 35 (CADASIL or Cerebral autosomal dominant ischemia with subcortical leukoencephalopathy).tw.
 - 36 multisystems atrophy.tw.
 - 37 Normotensive Hydrocephalus/
 - 38 Creutzfeldt Jakob Disease/
 - 39 exp Brain Spongiosis/

40 spongiform encephalopathy.tw.
 41 (CJD or JCD).tw.
 42 Creutzfeldt-Jakob disease.tw.
 43 exp Hypothyroidism/
 44 Cyanocobalamin Deficiency/
 45 Neurosyphilis/
 46 or/1-45
 47 multicenter study/ or phase 2 clinical trial/ or phase 3 clinical trial/ or phase 4
 clinical trial/ or randomized controlled trial/ or exp postmarketing surveillance/
 48 randomization/
 49 crossover procedure/ or double blind procedure/ or experimental design/ or latin
 square design/ or parallel design/ or single blind procedure/
 50 (clin: adj25 trial:).ti,ab.
 51 ((singl: or doubl: or trebl: or tripl:) adj25 (blin: or mask:)).ti,ab.
 52 Placebo/
 53 placebo:.ti,ab.
 54 random:.ti,ab.
 55 exp comparative study/ or exp drug comparison/
 56 exp "evaluation and follow up"/
 57 longitudinal study/ or major clinical study/ or prospective study/
 58 or/47-54
 59 or/55-57
 60 (tu or th).fs.
 61 exp treatment outcome/
 62 exp therapy/
 63 or/60-62
 64 59 and 63
 65 58 or 64
 66 65 and 46
 67 exp animal/
 68 66 not 67
 69 68 not (comment or editorial or news or letter or conference paper).pt.
 70 limit 69 to English language
 71 limit 70 to yr=1998-2003

Database: AMED (Allied and Complementary Medicine) <1985 to February 2003> on OVID

Search strategy executed on March 4, 2003

-
- 1 exp clinical trials/ or double blind method/ or random allocation/
 - 2 clinical trial.pt.
 - 3 (clin: adj25 trial:).ti,ab.
 - 4 ((singl: or doubl: or trebl: or tripl:) adj25 (blind or mask:)).ti,ab.
 - 5 placebos.sh.
 - 6 placebo:.ti,ab.
 - 7 random:.ti,ab.
 - 8 or/1-7
 - 9 (mild cognitive impairment or MCI).tw.
 - 10 ((cognitive impairment not dementia) or CIND).tw.
 - 11 ((cognitive loss not dementia) or CLOND).tw.
 - 12 exp dementia/
 - 13 (dement: or Alzheimer:).tw.
 - 14 amentia.tw.
 - 15 frontotemporal lobar degeneration.tw.
 - 16 hiv-associated cognitive motor complex.tw.
 - 17 (encephalopathy, aids or encephalopathy, hiv).tw.
 - 18 mesulam syndrome.tw.
 - 19 progressive nonfluent aphasia.tw.
 - 20 binswanger disease.tw.
 - 21 binswanger encephalopathy.tw.
 - 22 leukoencephalopathy, subcortical.tw.
 - 23 subcortical arteriosclerotic encephalopathy.tw.
 - 24 chronic progressive subcortical encephalopathy.tw.
 - 25 (alcohol adj3 amnestic).tw.
 - 26 (alcohol adj2 dysmnestic).tw.
 - 27 (ethanol adj3 nervous system disorders).tw.
 - 28 ethyl alcohol abuse neurologic syndromes.tw.
 - 29 (Lewy: bod: adj8 disease).tw.
 - 30 brain atrophy, circumscribed lobar.tw.
 - 31 (Pick: adj8 disease).tw.
 - 32 (corticobasil ganglionic degeneration or cortical basal degeneration or cortica).tw.
 - 33 (CADISIL or cerebral autosomal dominant ischemia with subcortical leukoencephalopathy).tw.
 - 34 Multisystems atrophy.tw.
 - 35 spongiform encephalopathy.tw.
 - 36 (cjd or Jcd).tw.
 - 37 Creutzfeldt-Jakob disease.tw.
 - 38 hypothyroidism/
 - 39 or/9-38
 - 40 8 and 39
 - 41 40 not (comment or editorial or news or letter).pt.

42 41 and English.lg.

Database: CINAHL <1982 to February Week 3 2003> on OVID
Search strategy executed on March 5, 2003

-
- 1 crossover design/ or empirical research/ or experimental studies/ or exp clinical trials/ or community trials/ or factorial design/ or quantitative studies/
 - 2 clinical trial.pt.
 - 3 (clin: adj25 trial:).ti,ab.
 - 4 ((singl: or doubl: or trebl: or tripl:) adj25 (blind: or mask:)).ti,ab.
 - 5 Placebos/
 - 6 placebo:.ti,ab.
 - 7 random:.ti,ab.
 - 8 Study Design/
 - 9 or/1-8
 - 10 (mild cognitive impairment or MCI).tw.
 - 11 ((cognitive impairment not dementia) or CIND).tw.
 - 12 ((cognitive loss not dementia) or CLOND).tw.
 - 13 exp Dementia/
 - 14 (dement: or Alzheimer:).tw.
 - 15 amentia.tw.
 - 16 frontotemporal lobar degeneration.tw.
 - 17 hiv-associated cognitive motor complex.tw.
 - 18 (encephalopathy, aids or encephalopathy, hiv).tw.
 - 19 mesulam syndrome.tw.
 - 20 progressive nonfluent aphasia.tw.
 - 21 Binswanger disease.tw.
 - 22 Binswanger encephalopathy.tw.
 - 23 leukoencephalopathy, subcortical.tw.
 - 24 (chronic progressive subcortical encephalopathy or (alcohol adj3 amnestic) or (alcohol adj2 dysmestic) or (ethanol adj3 nervous system disorders) or ethyl alcohol abuse neurologic syndromes).tw.
 - 25 Lewy body disease.tw.
 - 26 brain atrophy, circumscribed lobar.tw.
 - 27 Pick: disease.tw.
 - 28 (corticobasil ganglionic degeneration or cortical basal degeneration or cortica).tw.
 - 29 (CADASIL or cerebral autosomal dominant ischemia with subcortical leukoencephalopathy).tw.
 - 30 multisystems atrophy.tw.
 - 31 Huntington's Disease/
 - 32 Creutzfeldt-Jakob Syndrome/
 - 33 spongiform encephalopathy.tw.
 - 34 (cjd or jcd).tw.
 - 35 Creutzfeldt-Jakob disease.tw.
 - 36 spongiform encephalopathy.tw.
 - 37 exp Hypothyroidism/
 - 38 Neurosyphilis/
 - 39 or/10-38

- 40 9 and 39
- 41 40 not (editorial or letter or proceedings).pt.
- 42 limit 41 to English

Database: Ageline <1978 to December 2002> on SILVERPLATTER
Search strategy executed on March 6, 2003

- #1 randomized controlled trials in DE
- #2 controlled clinical trials in de
- #3 random allocation
- #4 controlled clinical trial*
- #5 randomized controlled trial*
- #6 random allocation
- #7 double blind method
- #8 single blind method
- #9 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8
- #10 clinical trial*
- #11 (clin* near trial*) in TI
- #12 (clin* near trial*) in AB
- #13 (singl* or doubl* or trebl* or tripl*) near (blind* or mask*)
- #14 (#13 in TI) or (#13 in AB)
- #15 placebo*
- #16 Placebo* in TI
- #17 placebo* in AB
- #18 random* in TI
- #19 random* in AB
- #20 research design
- #21 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or
#14 or #15 or #16 or #17 or #18 or #19 or #20
- #22 mild cognitive impairment
- #23 (cognitive impairment and dementia) or CIND
- #24 (cognitive loss not dementia) or CLOND
- #25 (explode 'Dementia-' in DE) or (explode 'Early-Onset-Dementia' in DE) or
(explode 'Vascular-Dementia' in DE)
- #26 dement* or alzheimer*
- #27 amentia*
- #28 frontotemporal lobar degeneration
- #29 hiv-associated cognitive motor complex
- #30 aids associated encephalopathy
- #31 hiv associated encephalopathy
- #32 mesulam syndrome
- #33 progressive nonfluent aphasia
- #34 binswanger disease
- #35 binswanger encephalopathy
- #36 leukoencephalopathy subcortical
- #37 subcortical arteriosclerotic encephalopathy
- #38 chronic progressive subcortical encephalopathy
- #39 alcohol near amnestic
- #40 alcohol amnestic disorder
- #41 alcohol induced disorders

#42 alcohol near dysmnestic
 #43 ethanol near nervous system disorders
 #44 lewy* bod* near disease
 #45 ethyl alcohol abuse neurologic syndromes
 #46 brain atrophy lobar
 #47 Pick* near disease*
 #48 cerebral amyloid angiopathy
 #49 neurofilament protein*
 #50 tau protein*
 #51 neurofibrillary tangles
 #52 neuropil threads
 #53 senile plaque*
 #54 corticobasil ganglionic degeneration or cortical basal degeneration or cortica
 #55 cadisil
 #56 cerebral autosomal dominant ischemia with subcortical leukoencephalopathy
 #57 multisystems atrophy
 #58 explode 'Huntingtons-Disease' in DE
 #59 normal pressure hydrocephalus
 #60 Creutzfeldt-Jakob syndrome
 #61 spongiform encephalopathy
 #62 cjd or jcd
 #63 Creutzfeldt-Jakob disease
 #64 hypothyroidism
 #65 vitamin b12 deficiency
 #66 neurosyphilis
 #67 #22 or #23 or #24 or #25 or #26 or #27 or #28 or #29 or #30 or #31 or #32 or #33
 or #34 or #35 or #36 or #37 or #38 or #39 or #40 or #41 or #42 or #43 or #44 or #45 or
 #46 or #47 or #48 or #49 or #50 or #51 or #52 or #53 or #54 or #55 or #56 or #57 or #58
 or #59 or #60 or #61 or #62 or #63 or #64 or #65 or #66
 #68 #67 and #21
 * #69 #68 and (DT=JOURNAL-ARTICLE)

Database: PsychINFO <1967 TO 2002/12> on SILVERPLATTER
Search strategy executed on March 7, 2003

- #1 randomized controlled trial in PT
- #2 controlled clinical trial in PT
- #3 controlled clinical trials
- #4 random allocation
- #5 (clinical trial*) in DE,SU
- #6 (random allocation) in DE,SU
- #7 (double blind method) in DE,SU
- #8 (cross-over studies) in DE,SU
- #9 (single-blind method) in DE,SU
- #10 (clinical trial) in PT
- #11 ((clin* near trial*)) in TI
- #12 (placebo*) in DE,SU
- #13 (placebo*) in TI
- #14 (random*) in TI
- #15 #1 or #2 or #3 or #4 or #5 or #6 or #7 or #8 or #9 or #10 or #11 or #12 or #13 or #14
- #16 mild cognitive impairment
- #17 MCI
- #18 (cognitive impairment not dementia) or CIND
- #19 (cognitive loss not dementia) or CLOND
- #20 (explode 'AIDS-Dementia-Complex' in DE) or (explode 'Alzheimers-Disease' in DE) or (explode 'Dementia-with-Lewy-Bodies' in DE) or (explode 'Dementia-' in DE) or (explode 'General-Paresis' in DE) or (explode 'Multi-Infarct-Dementia' in DE) or (explode 'Presenile-Dementia' in DE) or (explode 'Senile-Dementia' in DE) or (explode 'Vascular-Dementia' in DE)
- #21 dement* or Alzheimer*
- #22 amentia
- #23 HIV-associated cognitive motor complex
- #24 encephalopathy aids
- #25 encephalopathy hiv
- #26 mesulam syndrome
- #27 progressive nonfluent aphasia
- #28 binswanger disease
- #29 binswanger encephalopathy
- #30 leukoencephalopathy subcortical
- #31 subcortical arteriosclerotic encephalopathy
- #32 chronic progressive subcortical encephalopathy
- #33 alcohol near amnestic
- #34 alcohol near dysmnesic
- #35 ethanol near (nervous system disorders)
- #36 ethyl alcohol abuse neurologic syndromes
- #37 lewy* bod* near disease
- #38 brain atrophy circumscribed lobar

#39 Pick* near disease
 #40 cerebral amyloid angiopathy
 #41 (neurofilament proteins) in DE,SU
 #42 (tau proteins) in DE,SU
 #43 (neurofibrillary tangles) in DE,SU
 #44 (neuropil threads) in DE,SU
 #45 (senile plaque*) in DE,SU
 #46 (corticobasal ganglionic degeneration) or (cortical basal degeneration)
 #47 cadisil or (cerebral autosomal dominant ischemia with subcortical
 leukoencephalopathy) (0 records)
 #48 multisystems atrophy (0 records)
 #49 'Huntingtons-Disease' in DE (883 records)
 #50 hydrocephalus normal pressure (140 records)
 #51 'Creutzfeldt-Jakob-Syndrome' in DE (109 records)
 #52 spongiform encephalopathy (39 records)
 #53 cjd or jcd (80 records)
 #54 Creutzfeldt-Jakob disease (183 records)
 #55 explode 'Hypothyroidism-' in DE (260 records)
 #56 explode 'Neurosyphilis-' in DE (39 records)
 #57 vitamin B12 deficiency* (50 records)
 #58 frontotemporal lobar degeneration (15 records)
 #59 #16 or #17 or #18 or #19 or #20 or #21 or #22 or #23 or #24 or #25 or #26 or #27
 or #28 or #29 or #30 or #31 or #32 or #33 or #34 or #35 or #36 or #37 or #38 or #39 or
 #40 or #41 or #42 or #43 or #44 or #45 or #46 or #47 or #48 or #49 or #50 or #51 or #52
 or #53 or #54 or #55 or #56 or #57 or #58 (27527 records)
 #60 #15 and #59 (338 records)
 * #61 #60 and (LA=ENGLISH) and (PO=HUMAN) (322 records)

Appendix B

Forms

REF ID #

FIRST AUTHOR

SCREENER

EXCLUDE ☐ because: BUT..... KEEP ANYWAY as SPECIAL ☐

1. ☐ Language other than English (specify) _____
2. ☐ Not a full article ☐ Should check for full article
3. ☐ No Dementia subjects in population
4. ☐ Not a treatment for Dementia
5. ☐ Dementia population not randomized to treatment
6. ☐ No outcomes provided for Dementia subjects
7. ☐ Other reason (specify) _____

**** ☐ **** **CONSULTATION REQUIRED**INCLUDE ☐ for the following interests:

- ☒ Diagnosis of interest: ☐ Alzheimers ☐ AIDS dementia ☐ MCI ☐ CIND ☐ CLOND ☐ Aphasia
☐ Huntington's ☐ Parkinson's ☐ Alcohol ☐ Supranuclear Palsy
☐ Pick's ☐ Hypothyroidism ☐ Vitamin deficiency ☐ Vascular
☐ Corticobasal deterioration ☐ Lewy bodies ☐ Organic Brain

(other) ☐ _____

- ☒ Treatments randomized
☐ Placebo ☐ Tacrine ☐ Donepezil/Aricept ☐ Rivastigmine ☐ Acetylcholine/inhibitors
☐ Galantamine ☐ Metrifonate ☐ Memantine ☐ Ginko Bilboa ☐ Estrogens

(specify others) _____

- ☒ Population analyzed ☐ All ☐ Subgroup (specify) _____

- ☒ Outcomes reported (of randomized treatment on included population) _____

- ☒ Other _____

☐ Include by consensus
☐ Exclude by consensus

TREATMENT OF DEMENTIA – GUIDE TO FULLTEXT SCREENING FORM

USING THE FORM

- a) Be sure to fill in the DM ID#, the Name of the First Author and Your Initials in the three boxes at the top right of the form.
- b) If a paper should be excluded, fill in the “EXCLUDE” box and fill in the box for the reason for exclusion that occurs first in the list of 7 reasons for exclusion.
- c) The boxes for “KEEP ANYWAY as SPECIAL’ or “should check for full article” can also be checked if it is an excluded article but may be useful to our review as background or clarification of it appears to be a companion paper for another report that is likely in our review.
- d) If you choose to exclude the paper, the details for included papers do not need to be filled in.
- e) If a paper should be included, fill in the “INCLUDE” box and fill in the information for ONLY TWO of the subsequent categories listed: Diagnosis of interest and Treatments randomized. Ignore Population analyzed, Outcomes reported and Other.....we may use them later for grouping.
- f) If you are not sure if a paper qualifies for inclusion and want it to be looked at by our clinicians or methodologists, mark “CONSULTATION REQUIRED”
- g) If a paper is excluded because the Dementia population is not defined by DSM, NINCDS OR ICD-10 criteria, save it for consultation and mark “population’ beside the Consultation Required box.

EXCLUDING ARTICLES

1. Complete report must be in English to be included. If there is only an English abstract, exclude the article.
2. Only full reports will be included. If the article is a letter, comment, editorial, news, abstract, proceedings of a meeting or any other brief description, exclude the article. If it seems that the study would otherwise be included, check the box “Should check for full article”.
3. All dementia populations will be accepted at this stage if they are documented by DSM III, DSM III-R, DSM IV, NINCDS-ADRDA, ICD-9, ICD-10. The population studied may include those with mild cognitive impaired (MCI), cognitive impairment, not Dementia (CIND), cognitive loss, not Dementia (CLOND). If the author *cites the article by McKhann as the criteria for diagnosis, it can be included because that is the criteria for NINCDS.*
4. Articles included should look at treatment of disease, cognition, behaviour, or quality of life, time to deterioration, depression, falls etc. Exclude if outcomes reported are ONLY neurophysiologic or neuroimaging (eg EEG)
5. Exclude if not a report of a randomized controlled trial.
6. Outcomes reported should be for subjects with Dementia. If the entire population does not have Dementia, only data sub-grouped for Dementia will be examined. Exclude if there are no outcomes of interest reported for specifically Dementia subjects.
7. Any previously unmentioned, compelling reason to exclude the study should be specified.

INCLUDING ARTICLES

1. If you are unclear about whether the diagnosis is an included one, mark the referral box and pass along for a consult. If entire population is demented, mark boxes for all specific diagnoses included in study outcomes. If not all of population is demented, mark boxes for all specific diagnoses included in subgroup analysis of outcomes. If diagnosis is not listed as a choice, but is an included diagnosis, specify on line provided.
2. Specify treatments if they were randomly provided to the dementia population.

DETAIL ABOUT DISEASE TERMS (terms from the literature search)

NOT

- Not normal or healthy volunteers
- Not general population of elderly persons
- Not selected for depression (some may have dementia but not all)...BUT... If subgroup analysis may have been done, it should be marked "Retrieve".

INCLUDE

- Alzheimer's disease by DSM, NINCDS OR ICD
- Dementia defined by DSM, NINCDS OR ICD
- MCI – mild cognitive impairment
- CIND – cognitive impairment, not Dementia
- CLOND – cognitive loss, not Dementia

*Keep articles aside in a group if the intervention is directed toward the caregiver or caregiver/patient dyad.

TREATMENT OF DEMENTIA - FULLTEXT SCREENING FORM
SECONDARY EXCLUSION

REF ID #

FIRST AUTHOR

SCREENER

Article was included on first fulltext screening form**Crossover trials**Include ☐ there is data of interest to extract on first phase aloneExclude ☐ there is no data of interest to extract on first phase alone**Quality for all trials included on primary screen**Any blinding was done ☐ NO.....EXCLUDE
☐ YES...CONTINUEWithdrawals were enumerated for each arm ☐ YES....INCLUDE
☐ NOEXCLUDE

REFID _____ 1st AUTHOR _____ EXTRACTOR _____

QUALITY SCORE FOR JADAD SCALE AND FOR MODIFIED JADAD SCALE

CRITERIA	RESULT	SCORING	SCORE
Reported as randomized	<input type="checkbox"/> YES <input type="checkbox"/> NO	1 point for YES	
Randomization is appropriate	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NOT DESCRIBED	1 point for YES -1 point for NO	
Double blinding is reported	<input type="checkbox"/> YES <input type="checkbox"/> NO	1 point for YES	
Double blinding is appropriate	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NOT DESCRIBED	1 point for YES -1 point for NO	
Withdrawals are reported by number and reason per arm	<input type="checkbox"/> YES <input type="checkbox"/> NO	1 point for YES	
JADAD SCORE			/5 _____
Method used to assess adverse events is described	<input type="checkbox"/> YES <input type="checkbox"/> NO	1 point for YES	
Methods of statistical analysis are described	<input type="checkbox"/> YES <input type="checkbox"/> NO	1 point for YES	
Inclusion criteria reported	<input type="checkbox"/> YES <input type="checkbox"/> NO	1 point for YES in at least one of two criteria	
Exclusion criteria reported	<input type="checkbox"/> YES <input type="checkbox"/> NO		
JADAD IN AD SCORE			/8 _____
Intended allocation to tx group concealed from investigator	<input type="checkbox"/> YES <input type="checkbox"/> NO <input type="checkbox"/> NOT REPORTED		

Table A. Key Characteristics

REF ID #	Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified by patient characteristic

GUIDE TO DATA EXTRACTION TABLE

REF ID #	Enter number written on top of first page of article.
Author / Year	Enter last name of first author on one line and year of publication on the second line.
Funding Source	Enter one of the following codes: IF (Industry Funded) PI (Partially funded by Industry) IS (Industry provided Supplies) NI (Non-Industry funding source) NR (not reported) Use more than one code if necessary.
Quality score	Enter the Modified Jadad score for Alzheimer's Disease (out of 8 points)
Interventions	Enter name of drugs used in trial. Use the most commonly recognized name (eg use Tacrine instead of generic name). If more than one drug is used, put one on each line. If a dose response trial is reported, treat as one drug at the highest dose. If placebo is used, enter as first drug.
Criteria for Diagnosis	Indicate what criteria were used for diagnosis. It should be one of NINCDS, DSMIII, DSMIII-R, DSMIV, ICD-10
Diagnosis	Enter all dementia diagnoses included in the trial. PDD = PRIMARY DEGENERATIVE DEMENTIA MID = MULTI- INFARCT DEMENTIA AD = DAT = SDAT = ALZHEIMER'S DEMENTIA MIXED VaD = VASCULAR DEMENTIA DEMENTIA
Disease Severity	Use the descriptive terms as used in the paper.
Total Number randomized	Give the number in all groups that were initially randomized.
Number completing trial	Give the number in all groups that completed the treatment. If ITT population is given, report also.
Mean age (range) % Male (M) Population	Enter whatever information is given in paper for whole population Compute this figure if possible for those randomized at baseline Give any special inclusion criteria (or existing factor in population) which may affect the external validity (eg comorbid disorders, race, setting)
Highest Dose	Give the dose per day. If the dose was titrated up to individual doses, give the highest dose used and enter details of titration on the second line. Record as reported in paper - make sure to note if dose is by weight or a set amount and report how often given (preferably by day).
Treatment Period open extension	This should be the length of time for which the subjects received drug treatment. Use the longest period if there is more than one. Note here if there is an
Outcomes Measured	List all of the tests reported. If a battery of psychological tests were done, list the name of the battery. If physical tests are reported (e.g. blood tests, scans) list in very general terms (e.g. blood levels, EEG).
Outcome reports stratified by patient characteristic	Enter a Y if any of the data is reported in any way and is stratified by any patient characteristic (e.g. gender, age, race, genotype, education) and describe what the characteristic is. Otherwise, enter N.

Table B. Evidence Results

DRUG									
REF ID #	Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P value
				Baseline		Mid-Point: (specify)		Final: (specify) 24w	
				Baseline		Mid-Point: (specify)		Final: (specify)	

GUIDE TO DATA EXTRACTION TABLE B

REF ID #	Enter number written on top of first page of article.
Author / Year	Enter last name of first author on one line and year on the second line
Analysis groups and Interventions	<p>Use one line for each intervention that will be reported on. Number them as follows (1] 2] 3])</p> <p>Make the Control condition # 1 and Drug treatment conditions # 2 and so forth.</p> <p>If results are reported as change or relationship between baseline and later time-points or between placebo and drug conditions, these will be added as additional Analysis Groups (e.g. [3] Baseline vs 28w Placebo) would be entered as the third Analysis Group and would be for the score that reports the change in value between the one at baseline and the one at 28 weeks for the placebo group. The Analysis Group [5] Placebo vs Tactine 28w) would be the fifth Analysis Group and be used for the score that reports the change in value between the placebo condition and the drug condition at 28 weeks.</p> <p>If subgroup analysis has been reported, an Analysis Group can be created to report the results.</p> <p>For Analysis Groups that refer to the drug used, enter the name of drug used in trial. Use the most commonly recognized name (eg use Tactine instead of generic name). If more than one drug is used, put one on each line. If variable dosing is used within an arm, report the higher dose. Put the Placebo condition first in the list</p>
Test Used	List all psychological and functional tests used that have extractable data (cognition, behavior, functional, global). Do not report on physiological measurements such as blood levels. Enter the primary outcomes first in the list and bold the font on the test name.
Baseline, Mid-Point, Final	<p>Enter number of hours, days, weeks, months, years from baseline measurement to current measurement. Use abbreviations (h = hours, d = days, w = weeks, m = months, y = years)</p> <p>If more than 3 time points given, use the most central one for Time 2.</p>
Result Values	<p>Enter the value for Mean " Standard Deviation for each arm for each time-point for each test. If Standard Error is used, use an * .</p> <p>If % is used, enter the % sign after the value.</p> <p>If the test consists of subsections (eg a battery that also reports the total), just use the total score unless a sub-score is a primary outcome. Give P value if provided. If no P value available, but CI reported, put CI in P value column.</p>

Overall Summary Table Interpretation:

- SC = statistically significant CHANGE at the alpha = 0.05
- = based on the PRIMARY outcomes (no need to specify as 1^o because this is the default). If the paper does not specify primary or secondary ASSUME primary for all reported outcomes
 - = report secondary outcomes ONLY when there is NO primary variable for that domain AND indicate with (2^o) in front of the result code
 - = based on the ITT analyses results ONLY; if ITT results were not reported in the paper, then indicate with an asterick (*) located behind the result code (i.e. NS*)
 - = this change is ONLY relative to placebo (within group findings are reported in Table B..so not necessary to recapitulate this in summary table)
 - = in those instances where there is A PRIORI hypotheses for subgroup analyses and there are statistical results reported, then indicate with a symbol (i.e. # or ^) that the subgroup analyses were SC or NS and specify with respect to what factor (i.e. Vascular dementia versus not, or gender, etc)
- NS = not statistically significant effect for primary or secondary outcomes (some additional domains are tested with the secondary outcomes)
- NT = outcomes were not tested reflecting in this domain
- MX = mixed results for two primary outcomes (i.e one was significant and the other variable was not significant) and do not represent a SUBGROUP analysis
- = indicates that two measures within the same domain show conflicting results (one outcome is significant and the other is not significant)

Reporting Safety information in Randomized controlled trials (Ioannidis and Lau, 2002)

DOMAIN recommendation Ioannidis and Lau	OUR QUESTION	STATUS
FREQUENCY of WITHDRAWALS due to adverse events (AE)	Do the authors specify the number of patients withdrawn from the study due to AE per study arm and per type of AE that caused withdrawal	Y N Unclear Y N Unclear
FREQUENCY of AE (can be stated as a count or as a proportion for either CLINICAL AE or LABORATORY-DEFINED TOXICITY)	Do the authors provide the number of AE with respect to severity (reference to a known scale of toxicity such as mild, moderate, severe, life threatening or grades 1 to 3, etc) per study arm and per type of specific AE (i.e. diarrhea, headache, etc)	Y N Unclear Y N Unclear
Was the recording of the AE (i.e. surveillance) ACTIVE or PASSIVE	Is the surveillance ACTIVE (actively monitor the presence of absence of AE during the study...do not rely on methods that are PASSIVE (sometimes called spontaneous reporting).	Y N Unclear
Describe a SCHEDULE for collection of safety info	Optional 1) Do the authors specify the schedule for collection of safety information?	
FREQUENCY of SERIOUS AE (i.e. results in death, requires inpatient hospitalization, persistent or significant disability or is life threatening, WHO 2001)	Are exact numbers for high-grade (serious and life threatening) clinical AE laboratory toxicity reported.	Y N Unclear
SEVERITY of each AE (i.e. mild, moderate, or severe headache)	Have each of the AE been reported with respect to a severity continuum (i.e. mild diarrhea, severe headache, etc) ? Have some of the AE been reported with respect to a severity grade ?	Y N Unclear Optional: Y N Unclear
Description of UNUSUAL or NOT PREVIOUSLY RECORDED AE	Has a detailed description of cases of unusual or not previously recorded AE effects been presented?	Y N Unclear
STANDARDIZED SCALES used to capture AE.	Do the authors report the use of widely known, standardized scales for AE? Specify scale: If the scale is new , do the authors provide definitions for the grades of severity	Y N Unclear Optional: Y N Unclear
		Y N Unclear

Identify specific SAFETY TESTS or QUESTIONNAIRES used for data collection	Do the authors identify specific safety tests or questionnaires used for data collection	
---	--	--

THRESHOLD SCORING for ADVERSE EVENTS:

1) Scoring:

YES = 1

NO = 0

Unclear = consult with other rater to reach consensus

2) For the first 3 questions a **MINIMUM** score of **3** is required to proceed to the subsequent **5** questions

3) If all patients were accounted for (i.e. no withdrawals), then we assume a score of 2 for the **WITHDRAWAL** due to AE question

4) If no mention of serious (see definition) is mentioned in the paper, then we will **ASSUME** that they were **NOT** monitored (rather than not reported)

Appendix C

Evidence tables

Guide to the Results Tables

The results from all of the studies have been recorded in the following tables, which have been organized according to the intervention used in the trial. There are three sections:

- 1) Cholinergic neurotransmitter modifying agents (CNMA)
- 2) Non-cholinergic neurotransmitter/neuropeptide modifying agents (NCNNM)
- 3) Other pharmacological agents (OTHER)

Within each category, the results for drugs with three or more trials included are grouped together and presented in alphabetical order by drug name. There is a table showing the characteristics of the all of the studies using the drug. This table is followed by a separate table for each of the studies using that drug which shows the detailed results reported. These detailed tables are followed by a table containing adverse event information about each of the studies using that particular drug. Where there are only one or two trials included which use a drug, the tables for these studies are grouped together as various in tables as described above.

Following is a list of all of the drugs found in this review and the section in which they can be found. The drugs are ordered alphabetically within their section.

INTERVENTION	DRUG GROUP
5'-MTHF (FOLATE)	OTHER
ALAPROCLATE	NCNNM
ALPRAZOLAM	NCNNM
AMITRIPTYLINE	OTHER
ANAPSOS	NCNNM
ANIRACETAM	CNMA
ANTAGONIC STRESS	CNMA
ATEROID	OTHER
BMT	OTHER
BMV (NOOTROPIC)	NCNNM
BUFLOMEDIL	OTHER
CARBAMAZEPINE	CNMA
CARNITINE	CNMA
CEREBROLYSIN	OTHER
CHOTO-SAN (HERB)	OTHER
CITALOPRAM	NCNNM
CITICOLINE	OTHER
CYCLANDELATE	OTHER
DENBUFYLLINE	OTHER
DEFERRIOXAMINE (DFO)	OTHER
DICLOFENAC/MISOPROSTOL	OTHER
DIPHENHYDRAMINE	NCNNM
DIVALPROEX	NCNNM
DONEPEZIL	CNMA
EPTASTIGMINE	CNMA
ERGOKRYPTINE DEK (DIHYDROERGOKRYPTINE)	OTHER

INTERVENTION	DRUG GROUP
ESTROGENS	OTHER
FLUOXETINE	NCNNM
FLUVOXAMINE	NCNNM
GALANTAMINE	CNMA
GINKO BILOBA	OTHER
GLYCOSAMINOGLYCAN POLYSULFATE	OTHER
GUANFACINE	OTHER
HALOPERIDOL	NCNNM
HUPERZINE-A	CNMA
HYDERGINE	OTHER
HYDROXYCHLOROQUINE	OTHER
IDEBENONE	OTHER
IMIPRAMINE	NCNNM
INDOMETHACIN	OTHER
LINOPIRIDINE	CNMA
LISURIDE	NCNNM
LORAZEPAM	NCNNM
LOXAPINE	NCNNM
LU25-109	NCNNM
MAPROTILINE	NCNNM
MECLOFENOXATE	CNMA
MELPERONE	NCNNM
MEMANTINE	NCNNM
METRIFONATE	CNMA
MIANSERIN	NCNNM
MINAPRINE	NCNNM
MOCLOBEMIDE	NCNNM
MONOSIALOTETRAHEXOSYLGANGLIOSIDE (GM1)	OTHER
N-ACETYLCYSTEINE NAC	OTHER
NAFTIDROFURYL	NCNNM
NICERGOLINE	CNMA
NIMESULIDE (NSAID)	OTHER
NIMODIPINE	OTHER
NIZATIDINE	OTHER
NOOTROPIC	OTHER
OLANZAPINE	NCNNM
ORG 2766	OTHER
OXAZEPAM (Benzodiazapine)	NCNNM
OXIRACETAM	OTHER
PAROXETINE	NCNNM
PENTOXIFYLLINE	OTHER
PERPHENAZINE	NCNNM
PHOSPHATIDYLSERINE	NCNNM
PHYSOSTIGMINE	CNMA
PIRACETAM	OTHER

INTERVENTION	DRUG GROUP
POSATIRELIN	CNMA
PREDNISONE	OTHER
PROPENTOFYLLINE	OTHER
PYRITINOL	OTHER
RISPERIDONE	NCNNM
RIVASTIGMINE	CNMA
SABELUZOLE	CNMA
SELEGILINE (DEPRENYL)	NCNNM
SERTRALINE	NCNNM
SIMVASTATIN	OTHER
SULFOMUCOPOLYSACCHARIDES	OTHER
SULODEXIDE	OTHER
TACRINE	CNMA
THIAMINE	OTHER
THIORIDAZINE	NCNNM
TIAPRIDE	NCNNM
TRAZODONE	NCNNM
VASOPRESSIN (DDAVP)	OTHER
VELNACRINE	CNMA
VINCAMINE	OTHER
VITAMIN E	OTHER
XANOMELINE	NCNNM
XANTINOLNICOTINATE	OTHER

Appendix D

Peer review

Peer Reviewers for the Treatment of Dementia Report

1. Dr. Sandra Black, Sunnybrook Health Sciences Centre, Toronto, Ontario, Canada
2. Dr. Malaz Boustani, Indiana University School of Medicine, Indianapolis, Indiana, USA
3. Dr. Charles Cefalu (American Geriatrics Assoc)
4. Dr. Larry Chambers, University of Ottawa Institute for Health of the Elderly, Ottawa, Ontario, Canada
5. Dr. Tommy Cross (ACP/ASIM representative), Colorado Springs, Colorado, USA
6. Dr. Rachelle Doody, Baylor College of Medicine, Houston, Texas, USA
7. Dr. John Feightner, University of Western Ontario, Canadian Task Force for Periodic Health, London, Ontario, Canada
8. Dr. Serge Gauthier, Alzheimer's Disease Research Unit, McGill Centre for Studies in Aging, Verdun Quebec, Canada
9. Lisa Gwyther (American Geriatrics Assoc), MSW Duke University, Durham, North Carolina
10. Dr. David Hogan, Division of Geriatric Medicine, Calgary, Alberta, Canada
11. Dr. Rodney Hornbake, (ACP/ASIM representative), Hadlyme, Connecticut, USA
12. Dr. Roy Jones, St. Martin's Hospital, Bath, UK
13. Dr. David Royall, UTHSCSA, San Antonio Texas, USA
14. Dr. Lon Schneider (Cochrane), Keck School of Medicine, University of Southern California, Los Angeles, California
15. Dr. Jennie Ward-Robinson, US National Alzheimer's Association, Chicago, Illinois, USA
16. Dr. Peter Whitehouse (Cochrane), Case Western Reserve University, Cleveland, Ohio

Others reviewing the report:

Dr. Vincenza Snow – partner, ACSIM
Dr. Martin Kabongo - American Association of Family Practitioners
Dr. Marcel Morrison-Bogorad - Representative from the Institute for Aging (NIH)
David Atkins, Kate Rickard and AHRQ staff
Mary Grady – technical review (AHRQ)

Criticism Editor:

Dr. Patricia Huston, Ottawa, ON

Structured format for collecting referee comments

Thank you for agreeing to review this report. This is still in the draft stages and a thorough copy edit will take place before the publication of the final report. Please do not feel you need to spend your time correcting spelling and punctuation – we are relying on your expertise to address the questions below and provide insight that will assist us in improving the content and format of the report.

Problem Formulation

- Are review questions well formulated with specified key components?
- Are comparison groups clearly stated?
- Were major changes in review questions avoided during the review process?

Study Identification

- Is there a thorough search for relevant data using appropriate resources?
- Are there unbiased explicit searching strategies that are appropriately matched to the question?

Study Selection

- Are appropriate inclusion and exclusion criteria used to select articles?
- Are selection criteria applied in a manner that limits bias?
- Are efforts made to identified unpublished data, if this is appropriate?
- Are major changes in selection criteria avoided during the review process?
- Are reasons for excluding studies from the report stated?

Appraisal of Studies

- Is the validity of individual studies addressed in a reliable manner?
- Are important parameters (e.g., setting, study population, study design) that could affect study results systematically addressed?

Data Collection

- Is there a minimal amount of missing information regarding outcomes and other variables considered key to interpretation of results?
- Are efforts made to reduce bias in the data collection process?

Data Synthesis

- Are important parameters, such as study designs, considered in the synthesis?
- Are reasonable decisions made concerning whether and how to combine the data?
- Are results sensitive to changes in the way the analysis was done?
- Is precision of results reported?

Discussion

- Are limitations and inconsistencies of studies stated?
- Are limitations of the review process stated?
- Are review finding integrated within the context of relevant indirect evidence?
- Are implications for research discussed
- Are implications for practice discussed?

Conclusions

- Are conclusions supported by the data reviewed?
- Are plausible competing explanations of observed effects addressed?
- Is evidence appropriately interpreted as inconclusive (no evidence of effect) or as showing a particular strategy did not work (evidence of no effect)?
- Are important considerations for decision makers identified, including values and contextual factors that might influence decisions?
- Is a summary of pertinent findings provided?

Appendix E

Outcome measures

Appendix E. Outcome Measures

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
ABID	Agitated Behavior Inventory for Dementia	High	Behavior
ABS	Adaptive Behavior Scale	Low	Function
ABSR	Aggressive Behavior Scale Rating	High	Behavior
ACES	Agitation Calmness Evaluation Scale	Low	Behavior
ACPT	Auditory Continuous Performance Test		Specific cognitive test
ADAS-	Alzheimer's Disease Assessment Scale	High	Global assessment
ADAS-Cog	Cognitive Sections		General cognitive function
ADAS-Noncog	Non-Cognitive, behavioral section		Behavior
ADCS-CGIC	Alzheimer's Disease Cooperative Study – Clinical, Global Impression of Change	High	Global assessment
ADCS-ADL	Alzheimer's Disease Cooperative Study – Activities of Daily Living	Low	Function
ADFACS	Alzheimer's Disease Functional Assessment and Change Scale	High	Function
ADL	Activities of Daily Living	High	Function
ADL-C	Activities of Daily Living Checklist		Function
ADL-PDS	Activities of Daily Living Progressive Deterioration Scale		Function
ADS	Alzheimer's Deficit Scale		Global assessment
AFBS	Aversive Feeding Behavior Scale	High	Behavior
AGS- E	Assessment of Global Symptomatology - Elderly		Global assessment
AIMS	Abnormal Involuntary Movement Scale	High	Adverse Events, Dyskinesia
AMTS	Abbreviated Mental Test Score		General cognitive function

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
---	Barbizet Visuospatial	Low	Specific cognitive test
BARS, BAS	Barnes Akathisia Rating Scale	High	Adverse Events, Akathisia
BDI	Beck Depression Inventory	High	Behavior
BEHAVE – AD	Behavioral Pathology in Alzheimer's Disease Rating Scale	High	Behavior
BCRS	Brief Cognitive Rating Scale	High	General cognitive function
Bf-S	Self assessment according to Zerksen and Möller		Global assessment
BGP	Behavioral Rating Scale for Geriatric Patients		Global assessment
BI	Barthel Index		Function
Blessed-D or BDRS	Blessed Dementia Rating Scale	High	Global assessment
BNT	Boston Naming Test	Low	Specific cognitive test
BPRS	Brief Psychiatric Rating Scale	High	Behavior
BRMS	Bech-Rafaelsen Mania Scale		Behavior
BRSD	Behavioral Rating Scale for Dementia		Behavior
BSR	Babcock Story Recall Test	Low	Specific cognitive test
BSRT	Buschke Selective Reminding Test	Low	Specific cognitive test
BSV	Buschke Sentence Verification		Specific cognitive test
BLM	Buschke Letter Matching		Specific cognitive test
BVR	Benton Visual Retention – Number Correct Benton Visual Retention – Errors		Specific cognitive test
CamCOG	Cambridge Cognitive Schedule		General cognitive function
CAPE	Clifton Assessment Procedures for the Elderly	Low	Global assessment
CASI	Cognitive Abilities Screening Instrument	Low	General cognitive function

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
CAUST	Canadian Utilization of Service Tracking questionnaire	---	Health-care utilization and work productivity
---	Category Fluency	Low	Specific cognitive test
CDR-NH CDR-SB	Clinical Dementia Rating – Nursing Home Version Clinical Dementia Rating-Sum of Boxes	High	Global assessment
CDT	Clock Drawing Test		Specific cognitive test
CERAD- BRSD	Consortium to Establish a Registry for Alzheimer's Disease – Behavioral Rating Scale for Dementia	High	Behavior
CETM	---	---	General cognitive function
CCASSS	Computerized Cognitive Assessment System Speed Score / unweighted sum of reaction time	Low	Specific cognitive test
CATS	Caregivers' Activity Time Survey		Caregiver burden
CGAE	Clinical Global Assessment and Efficacy		Global assessment
CGI	Clinical Global Impression	High	Global assessment
CGIC	Clinical Global Impression of Change	High	Global assessment
CGRS	Clinician's Global Rating Scale		Global assessment
CIBIC	Clinician's Interview Based Impression of Change	High	Global assessment
CIBIC+	Clinician's Interview Based Impression of Change plus Caregiver	High	Global assessment
CDT	Clock Drawing Test	Low	Specific cognitive test
CMAI	Cohen Mansfield Agitation Inventory	High	Behavior
CNTB	Computerized Neuropsychological test battery		Specific cognitive test
COWAT	Controlled Oral Word Association Test	Low	Specific cognitive test
CS or CSDD	Cornell Scale for Depression in Dementia	NR	Behavior
CSS	Caregiver Stress Scale	High	Caregiver burden

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
CSI	University of Iowa Caregiver Stress Inventory		Caregiver burden
CVLT	California Verbal Learning Test		Specific cognitive test
---	Dependency Scale	High	Function
DAD	Disability Assessment for Dementia	Low	Function
DST	Digit Span Test		Specific cognitive test
DBDS	Dementia Behavioral Disturbance Scale		Global assessment
DMR	Dementia Questionnaire for Mentally Retarded Persons	High	Global assessment
DRS	Dementia Rating Scale	High	Global assessment
DSCS	Depressive Symptoms Collateral Source Cornell Scale		Behavior
DSS	Depressive Signs Scale	High	Behavior
DSST	Digit Symbol Substitution Test	Low	Specific cognitive test
EIS	Efficacy Index Score		Global assessment
EFRT	Emotional Face Recognition Test	Low	Specific cognitive test
ERP	Event-Related Potential (Amplitude)	Low	Response to stimuli
ESRS	Extrapyramidal Symptom Rating Scale	High	Adverse events/ Extrapyramidal symptoms
---	Facial Behavior	NR	Behavior
---	Finger Tapping Test	Low	Motor coordination
FAST	Functional Assessment Staging	High	Function
FCCA	Final Comprehensive Consensus Assessment		Global assessment
FCMT	Figure Copy/ Memory Test	Low	Specific cognitive test
FIGT	Figure detection test		Specific cognitive test

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
FIM	Functional Independence Measure	Low	Function
FOM	Fuld-Object Memory Evaluation		Specific cognitive test
FRS	Functional Rating Scale test	Low	Global assessment
GAGS	Guide to Adult Assessment Battery for Physical pharmacology		Specific cognitive test
GERRI	Geriatric Evaluation by Relative's Rating Instrument	High	Global assessment
GBS	Gottfries-Bråne-Steen Total Score Motor function subscale Intellectual subscale Emotional function subscale Symptoms subscale	High	Global assessment Function General cognitive function Behavior Behavior
GDS	Global Deterioration Scale	High	Global assessment
GIS	Global Impairment Scale (Adaptation of the CGIS)		Global assessment
GPI-E	General Psychiatric Impression – Elderly		Global assessment
GS	Gestalt Scale	High	Behavior
---	Grooved Pegboard Test	Low	Specific cognitive test
HAM-A	Hamilton Anxiety Scale	High	Behavior
HAM-D HDRS	Hamilton Rating Scale for Depression	High	Behavior
HDS	Hachinski Dementia Scale		Global assessment
HDS-R	Hasegawa's Dementia Scale – Revised		Behavior
HIS	Hachinski Ischemic Score	High	Global assessment
IADL	Instrumental Activities of Daily Living	High	Function
IDDD	Interview for Deterioration in Daily Living Activities in Dementia – complex task	High	Function
IPSC-E	Raskin's and Crook's Inventory of Psychic and Somatic Complaints for the Elderly	High	Behavior
IQCODE	Informant Questionnaire on Cognitive Decline in the Elderly	High	General cognitive function

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
---	Letter Cancellation	Low	Specific cognitive test
---	Letter Fluency	Low	Specific cognitive test
LMT	Logical Memory Test	Low	Specific cognitive test
LNNB	Luria-Nebraska Neuropsychological Battery		Specific cognitive test
LPRS	London Psychogeriatric Rating Scale		Behavior
MAACL-R	Multiple Affect Adjective Checklist-Revised	High	Behavior
MADRS	Montgomery-Asberg Depression Rating Scale	High	Behavior
MAE	Benton Multi-Lingual Aphasia Examination		Specific cognitive test
MCPT	Modified Continuous Performance Test		General cognitive function
MEMT	Memory test		Specific cognitive test
MMSE MMMSE SMMSE CMMSE	Mini-Mental Status Exam Modified MMSE Standardized MMSE Cantonese MMSE	Low	General cognitive function
MNLT	Modified Names Learning test		Specific Cognitive Test
MOSES			Behavior
MQ	Memory Quotient		General cognitive function
NAA	Scale from the Nuremberg Gerontopsychological inventory for assessing activities of daily living		Function
NAB	Nürnberg Alters-Beobachtungskala		Behavior
NAI	Nuremberg Age Inventory	Low	Function
NCT	Number Correction Test		Specific cognitive test

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
NDT	New Dot Test	Low	Specific cognitive test
NLT	Names Learning Test		Specific cognitive test
NMIC	Newcastle Memory, Information, and Concentration Test	Low	Specific cognitive test
NMS	Nowlis Mood Scale	High	Behavior
NOSGER	Nurses Observation Scale for Geriatric Patients	High	Global assessment
NOSGER-IADL	Nurses Observation Scale for Geriatric Patients – Instrumental Activities of Daily Living subscale	High	Function
NOSIE	Nurses Observation Scale for Inpatients	High	Global assessment
NPI (NPI-4, NPI-10)	Neuropsychiatric Inventory Subscores 4,10	Low	Behavior
NPI-NH	Neuropsychiatric Inventory – Nursing Home Version	High	Behavior
NSL			Behavior
NST	Number Symbol Test		Specific cognitive test
OARS – ADL	Older Americans Resource Scale	High	Function
OAS	Overt Aggression Scale		Behavior
OLT	Object Learning Test		Specific cognitive test
PANSS-EC	Positive and Negative Syndrome Scale		Behavior
PDRS	Psychogeriatric Dependency Rating Scale		Global assessment
PDS	Progressive Deterioration Scale	High	Function
PGIR	Patient's Global Improvement Rating		Global assessment
POMS	Profile of Mood States		Behavior
PSMS	Physical Self-Maintenance Scale	High	Function
PSQI	Pittsburgh Sleep Quality Index	High	Function

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
QoL	Quality of Life	Low	Function
QoL-P QoL-C	Patient-rated Quality of Life Caregiver-rated Quality of Life	Low	Function
RAGS	Relative's Assessment of Global Symptomatology		Global assessment
R-AVL	Rey Auditory-Verbal-Learning test	Low	Specific cognitive test
RGRS	Relatives' Global Rating Scale		Global assessment
RM RPM	Raven Matrices Raven's Progressive Matrices		Specific cognitive test
RMT	Randt Memory Test	Low	General cognitive function
RMBPC	Revised Memory and Behavior Problems Checklist	High	Behavior
RPT	Rivermead Profile Test		Behavior
RVM	Rey's Verbal Memory	Low	General cognitive function
SAS	Simpson-Angus Scale	High	Adverse effects, Extra-pyramidal symptoms
SAS-G	Self Assessment – Geriatric	High	Global assessment
	Snodgrass Picture Naming Task		Specific cognitive test
SBI	Spontaneous Behavior Interview		Behavior
SCAG	Sandoz Clinical Assessment – Geriatric	High	Global assessment
SCB	Screen for Caregiver Burden	NR	Caregiver burden
SCWIT	Stroop Color Word Interference Test		Specific cognitive test
	Set Test		Specific cognitive test
SF-36	Medical Outcomes Study Short-Form 36-Item Health Survey	Low	Function
SGRS	Stockton Geriatric Rating Scale	High	Global assessment
SHGRT	Stuard Hospital Geriatric Rating Scale		Behavior

Acronym/ Abbreviation	Test name	Level of Impairment with higher score	Domain
SKT	Syndrome Kurztest; Syndrome Short Test	High	Specific cognitive test
SIB	Severe Impairment Battery	Low	General cognitive function
SIP	Sickness Impact Profile		Function
SMQ	Squire's Memory Questionnaire		General cognitive function
SMST	Sternberg Memory Scanning	Low	General cognitive function
SRT-DR	Selective Reminding Procedure – delayed recall		Specific cognitive test
SRT SRT-Anxiety SRT- Depression	Kellner and Sheffield Rating Test		Behavior
SWFT	Semantic Word Fluency Test	---	Specific cognitive test
	Time to functional decline		Function
TK	Token Test	Low	Specific cognitive test
TP TPAT	Toulouse Piéron Toulouse Piéron Attention Test	Low	General cognitive function
TMT	Trail Making Test		Specific cognitive test
TSI	Test for Severe Impairment		Global assessment
UPDRS	Unified Parkinson's Disease Rating Scale	High	Adverse effects, Extrapyramidal symptoms
VHB	Video-recorder home-behavioral assessment		Behavior
VRGI	Video Rating of Global Impression		Global assessment
WAIS	Wechsler Adult Intelligence Scale (Verbal and Memory Performance scales)	Low	General cognitive function
WMS (MQ)	Memory Learning Restauration		Specific cognitive test
WMS-RR	Wechsley Memory Scale – Russel Revised		Specific cognitive test
ZVT ZVTG	Zahlen-Verbindungs Test – Trail Making Test	High	Specific cognitive test

Appendix F

List of excluded studies

Appendix F. List of excluded studies

Aarsland D, Larsen JP, Lim NG, et al. Olanzapine for psychosis in patients with Parkinson's disease with and without dementia. *J Neuropsychiatry Clin Neurosci* 1999; 11(3):392-4.

Status: Not included because dementia population not randomized to treatment

Aarsland D, Laake K, Larsen JP, et al. Donepezil for cognitive impairment in Parkinson's disease: A randomised controlled study. *J Neurol Neurosurg Psychiatry* 2002; 72(6):708-12.

Status: Cross-over trial;

Aarsland D. Erratum: Donepezil for cognitive impairment in Parkinson's disease. A randomised controlled study. *J Neurol Neurosurg Psychiatry* 2002; 73(3):354.

Status: Not included because not a full article

Abalan F, Manciet G, Dartigues JF, et al. Nutrition and SDAT. *Biol Psychiatry* 1992 Jan 1; 31(1):103-5.

Status: Not included because not a full article

Abuzzahab FS, Sr., Merwin GE, Zimmermann RL, et al. A double-blind investigation of piracetam (nootropil) versus placebo in the memory of geriatric inpatients. *Psychopharmacol Bull* 1978 Jan; 14(1):23-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Abyad A. Prevalence of vitamin B12 deficiency among demented patients and cognitive recovery with cobalamin replacement. *J Nutr Health Aging* 2002; 6(4):254-60.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Adler LA, Peselow E, Rosenthal M, et al. A controlled comparison of the effects of propranolol, benztropine, and placebo on akathisia: An interim analysis. *Psychopharmacol Bull* 1993; 29(2):283-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Aerssens J, Raeymaekers P, Lilienfeld S, et al. APOE genotype: no influence on galantamine treatment efficacy nor on rate of decline in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2001 Mar; 12(2):69-77.

Status: Not included because does not meet criteria for treatment for dementia patients

Agnoli A, Martucci N, Manna V, et al. Effect of cholinergic and anticholinergic drugs on short-term memory in Alzheimer's dementia: a neuropsychological and computerized electroencephalographic study. *Clin Neuropharmacol* 1983; 6(4):311-23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Agnoli A, Martucci N, Manna V. Quantitative EEG as a tool in neuropharmacological studies: The effect of naftidrofuryl in chronic cerebrovascular diseases (C.C.V.D.). *Curr Ther Res Clin Exp* 1985; 37(3):387-97.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Agnoli A, Manna V, Martucci N, et al. Randomized double-blind study of flunarizine versus placebo in patients with chronic cerebrovascular disorders. *Int J Clin Pharmacol Res* 1988; 8(3):189-97.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Aguglia E, Caraceni T, Genitrini S, et al. Comparison of teniloxazine and piracetam in Alzheimer-type or vascular dementia. *Curr Ther Res Clin Exp* 1995; 56(3):250-7.

Status: Not included because Jadad Quality Scale score less than three

Ahlin A, Nyback H, Junthe T, et al. THA in Alzheimer's dementia clinical biochemical and pharmacokinetic findings. *Alzheimer's disease basic mechanisms diagnosis and therapeutic strategies* 1990; 621-5.

Status: Not included because not a full article

Ahlin A, Nyback H, Junthe T, et al. Tetrahydroaminoacridine in Alzheimer's dementia: Clinical and biochemical results of a double-blind crossover trial. *Hum Psychopharmacol* 1991; (2):109-18.

Status: Cross-over trial;

Ahlin A, Hassan M, Junthe T, et al. Tacrine in Alzheimer's disease: Pharmacokinetic and clinical comparison of oral and rectal administration. *Int*

Clin Psychopharmacol 1994; 9(4):263-70.
Status: Cross-over trial;

Aisen PS, Marin D, Davis KL. Anti-inflammatory drug studies in Alzheimer's disease. Biol Psychiatry 1996; 39(7):563
Status: Not included because not a full article

Aisen PS, Marin DB, Brickman AM, et al. Pilot tolerability studies of hydroxychloroquine and colchicine in Alzheimer disease. Alzheimer Dis Assoc Disord 2001 Apr; 15(2):96-101.
Status: Not included because dementia population not randomized to treatment

Aisen PS, Berg JD, Craft S, et al. Steroid-induced elevation of glucose in Alzheimer's disease: Relationship to gender, apolipoprotein E genotype and cognition. Psychoneuroendocrinology 2003; 28(1):113-20.
Status: Not included because dementia population not randomized to treatment

Alafuzoff I, Helisalmi S, Heinonen EH, et al. Selegiline treatment and the extent of degenerative changes in brain tissue of patients with Alzheimer's disease. Eur J Clin Pharmacol 2000 Feb; 55(11-12):815-12.
Status: Not included because no extractable data relevant to review

Albizzati MG, Bassi S, Calloni E, et al. Cycandelate versus flunarizine. A double-blind study in a selected group of patients with dementia. Drugs 1987; 33(Suppl 2):90-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Aldenkamp AP, van Wieringen A, Alpherts WC, et al. Double-blind placebo-controlled, neuropsychological and neurophysiological investigations with oxiracetam (CGP 21690E) in memory-impaired patients with epilepsy. Neuropsychobiology 1990 Sep; 24(2):90-101.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Alexopoulos GS, Meyers BS, Young RC, et al. Executive dysfunction and long-term outcomes of geriatric depression. Arch Gen Psychiatry 2000 Mar; 57(3):285-90.
Status: Not included because does not meet criteria for treatment for dementia patients

Allain H, Denmat J, Bentue-Ferrer D, et al. Randomized, double-blind trial of exifone versus

cognitive problems in Parkinson's disease. Fundam Clin Pharmacol 1988; 2(1):1-12.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Allain H, Raoul P, Lieury A, et al. Effect of two doses of Gingko biloba extract (EGb 761) on the dual-coding test in elderly subjects. Clin Ther 1993; 15(3):549-58.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Allain H, Neuman E, Malbezin M, et al. Bridging study of S12024 in 53 in-patients with Alzheimer's disease. J Am Geriatr Soc 1997; 45(1):125-6.
Status: Not included because not a full article

Almkvist O, Jelic V, Amberla K, et al. Responder characteristics to a single oral dose of cholinesterase inhibitor: A double-blind placebo-controlled study with tacrine in Alzheimer patients. Dement Geriatr Cogn Disord 2001 Jan; 12(1):22-32.
Status: Cross-over trial;

Altman H, Mehta D, Evenson RC, et al. Behavioral effects of drug therapy on psychogeriatric inpatients. II. Multivitamin supplement. J Am Geriatr Soc 1973 Jun; 21(6):249-52.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Altman H, Mehta D, Evenson RC, et al. Behavioral effects of drug therapy on psychogeriatric inpatients. I. Chlorpromazine and thioridazine. J Am Geriatr Soc 1973 Jun; 21(6):241-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Alvarez XA, Laredo M, Corzo D, et al. Citicoline improves memory performance in elderly subjects. Methods & Findings in Experimental & Clinical Pharmacology 1997 Apr; 19(3):201-10.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Alvarez XA, Mouzo R, Pichel V, et al. Double-blind placebo-controlled study with citicoline in APOE genotyped Alzheimer's disease patients. Effects on cognitive performance, brain bioelectrical activity and cerebral perfusion. Methods & Findings in Experimental & Clinical Pharmacology 1999 Nov; 21(9):633-44.

Status: Not included because Jadad Quality Scale score less than three

Amaducci L, Maurer K, Winblad B, et al. A long-term, double-blind, placebo-controlled efficacy and safety study of nicergoline in patients with mild to moderate Alzheimer's disease. *J Eur Coll Neuropsychopharmacol* 1999; (Supl 5):S323.
Status: Not included because not a full article

Amar K, Wilcock GK, Scot M, et al. The presence of leuko-araiosis in patients with Alzheimer's disease predicts poor tolerance to tacrine, but does not discriminate responders from non-responders. *Age Ageing* 1997 Jan; 26(1):25-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ambrozi L, Danielczyk W. Treatment of impaired cerebral function in psychogeriatric patients with memantine: Results of a phase II double-blind study. *Pharmacopsychiatry* 1988 May; 21(3):144-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anand R, Gharabawi G, Enz A. Efficacy and safety results of the early phase studies with Exelon(tm) (ENA-713) in Alzheimer's disease: An overview. *J Drug Dev Clin Pract* 1996; 1-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ananth JV, Deutsch M, Ban TA. Senilex in the treatment of geriatric patients. *Curr Ther Res Clin Exp* ; 13(5):316-21.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ancoli-Israel S, Martin JL, Kripke DF, et al. Effect of light treatment on sleep and circadian rhythms in demented nursing home patients. *J Am Geriatr Soc* 2002; 50(2):282-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anderer P, Barbanoj MJ, Saletu B, et al. Restriction to a limited set of EEG-target variables may lead to misinterpretation of pharmaco-EEG results. *Neuropsychobiology* 1993; 27(2):112-6.
Status: Not included because no extractable data relevant to review

Anderson J, Arens K, Johnson R, et al. Spaced retrieval vs. memory tape therapy in memory rehabilitation for dementia of the Alzheimer's type.

Clin Gerontol 2001; (1-2):123-39.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Darvon and Darvon-N. *Med Lett Drugs Ther* 1972 May; 14(11):37-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Severely demented patients beyond help of drugs. *Modern Geriatrics* 1976; (10):36.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Double-blind, placebo-controlled evaluation of cinromide in patients with the Lennox-Gastaut Syndrome. The Group for the Evaluation of Cinromide in the Lennox-Gastaut Syndrome. *Epilepsia* 1989 Jul; 30(4):422-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Safety and tolerability of the antioxidant OPC-14117 in HIV-associated cognitive impairment. The Dana Consortium on the Therapy of HIV Dementia and Related Cognitive Disorders. *Neurology* 1997 Jul; 49(1):142-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Study suggests antioxidants slow decline in Alzheimer's disease. *Am J Health Syst Pharm* 1997; 54(13):1478.
Status: Not included because not a full article

Anonymous. Selegiline hydrochloride: Antiparkinsonian cognition enhancer. *Drugs of the Future* 1998; 23(2):240-1.
Status: Not included because not a full article

Anonymous. Benefits of new Alzheimer disease therapies. *J Pharm Technol* 1998; 14(3):125.
Status: Not included because not a full article

Anonymous. Eptastigmine tartrate: Cognition enhancer acetylcholinesterase inhibitor. *Drugs of the Future* 1998; 23(2):217-8.
Status: Not included because not a full article

Anonymous. Phosphatidylserine. *Altern Med Rev* 1999; 4(2):115-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. The alternative to tube-feeding patients with advanced dementia. *Volunt Leader* 1999; 40(4):13-4.

Status: Not included because not a full article

Anonymous. Tacrine and Alzheimer disease. *WHO Drug Information* 1999; 13(1):7-8.

Status: Not included because not a full article

Anonymous. New hope for early Alzheimer's disease. *Harv Womens Health Watch* 2000 Apr; 7(8):7

Status: Not included because not a full article

Anonymous. Rivastigmine for Alzheimer's disease. *Drug Ther Bull* 2000; 38(2):15-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Erratum: Estrogen replacement therapy for treatment of mild to moderate Alzheimer disease: A randomized controlled trial (*Journal of the American Medical Association* (February 23,2000) 283 (1007-1015)). *JAMA* 2000; 284(20):2597

Status: Not included because not a full article

Anonymous. Lead success for Nuero-gen. *Manuf Chem* 2001; 72(4):11

Status: Not included because not a full article

Anonymous. New Alzheimer's drug is first therapy to show efficacy in vascular dementia. *Hosp Formul* 2001; 36(8):569.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Idebenone. *Altern Med Rev* 2001; 6(1):83-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Double-blind trial will compare two anti-Alzheimer's drugs. *Journal of Dementia Care* 2001; 9(5):6

Status: Not included because not a full article

Anonymous. Colostrinin. *Journal of Dementia Care* 2001; 9(6):37.

Status: Not included because not a full article

Anonymous. Erratum: A 24-week, randomized, double-blind study of donepezil in moderate to severe alzheimer's disease (*Neurology* (2001) 57 (613-620)). *Neurology* 2001; 57(11):2153.

Status: Not included because not a full article

Anonymous. Galantamine (Reminyl) for Alzheimer's disease. *Med Lett Drugs Ther* 2001; 43(1107):53-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Caregiver experience relates to clinical trial involvement: research looks at how caregivers of Alzheimer's patients make decisions regarding care. *Caremanagement* 2001 Jun; 7(3):55.

Status: Not included because not a full article

Anonymous. Aromatherapy trial. *Journal of Dementia Care* 2001; 9(6):38.

Status: Not included because not a full article

Anonymous. Galantamine: New preparation. The fourth cholinesterase inhibitor for Alzheimer's disease. *Prescrire Int* 2001 Dec; 10(56):180-1.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Galantamine effective in treating dementia in patients with cerebrovascular disease. *Pharm J* 2001; 266(7153):842.

Status: Not included because not a full article

Anonymous. Trial of new immunotherapeutic agent for Alzheimer's suspended. *Pharm J* 2002; 268(7187):279

Status: Not included because not a full article

Anonymous. Drug that modulates glutamate levels promising for Alzheimer's disease. *Pharm J* 2002; 269(7209):152.

Status : Not included because not a full article

Anonymous. Drugs to treat dementia and psychosis: Management of Parkinson's disease. *Mov Disord* 2002; 17(Suppl 4):S120-S127.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. NSAID use could reduce Alzheimer's risk. *Pharm J* 2002; 269(7217):428.

Status: Not included because not a full article

Anonymous. Memantine launched for treatment of Alzheimer's. *Pharm J* 2002; 269(7219):516.

Status: Not included because not a full article

Anonymous. Greater satisfaction, ease of use reported with donepezil versus galantamine. *Hosp Formul* 2002; 37(8):383-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Anonymous. Reminyl of benefit in vascular dementia. *Pharm J* 2002; 268(7194):526.
Status: Not included because not a full article

Anonymous. Perindopril protects against dementia. *Pharm J* 2002; 268(7204):899.
Status: Not included because not a full article

Anonymous. NSAIDs: Protection against Alzheimer's? *Med Today* 2002; 3(2):9
Status: Not included because not a full article

Arendt G, von Giesen HJ, Heffer H, et al. Therapeutic effects of nucleoside analogues on psychomotor slowing in HIV infection. *AIDS* 2001 Mar 9; 15(4):493-500.
Status: Not included because dementia population not randomized to treatment

Arkin SM. Alzheimer memory training: Quizzes beat repetition, especially with more impaired. *Am J Alzheimers Dis* 1997; (4):147-58.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Arkin SM. Alzheimer memory training: Students replicate learning successes. *Am J Alzheimers Dis* 2000 May; 15(3):152-62.
Status: Not included because dementia population not randomized to treatment

Arrigo A, Moglia A, Borsotti L. A double-blind, placebo-controlled, crossover trial with nicergoline in patients with senile dementia. *Int J Clin Pharmacol Res* 1982; 2(4 Suppl 1):33-41.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Arrigo A, Casale R, Giorgi I, et al. Effects of intravenous high dose c-dergocrine mesylate ('Hydergine' (R)) in elderly patients with severe multi-infarct dementia: A double-blind, placebo-controlled trial. *Curr Med Res Opin* 1989; 11(8):491-500.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ashford JW, Soldinger S, Schaeffer J, et al. Physostigmine and its effect on six patients with dementia. *Am J Psychiatry* 1981; 138(6):829-30.
Status: Not included because dementia population not randomized to treatment

Asthana S, Raffaele KC, Berardi A, et al. Treatment of Alzheimer's disease by continuous intravenous infusion of physostigmine. *Alzheimer Dis Assoc Disord* 1995; 9(4):223-32.
Status: Cross-over trial;

Asthana S, Greig NH, Holloway HW, et al. Clinical pharmacokinetics of arecoline in subjects with Alzheimer's disease. *Clin Pharmacol Ther* 1996 Sep; 60(3):276-82.
Status: Not included because no extractable data relevant to review

Asthana S, Raffaele KC, Greig NH, et al. Neuroendocrine responses to intravenous infusion of physostigmine in patients with Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1999 Apr; 13(2):102-8.
Status: Cross-over trial;

Asthana S, Craft S, Baker LD, et al. Cognitive and neuroendocrine response to transdermal estrogen in postmenopausal women with Alzheimer's disease: Results of a placebo-controlled, double-blind, pilot study. *Psychoneuroendocrinology* 1999 Aug; 24(6):657-77.
Status: Not included because Jadad Quality Scale score less than three

Ather SA, Shaw SH, Stoker MJ. A comparison of chlormethiazole and thioridazine in agitated confusional states of the elderly. *Acta Psychiatr Scand* 1986; 73(Suppl 329):81-91.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Auer SR, Monteiro IM, Reisberg B. Behavioral symptoms in dementia: community-based research. *Int Psychogeriatr* 1996; 8(Suppl 3):363-6, 381, 382.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Avorn J, Soumerai SB, Everitt DE, et al. A randomized trial of a program to reduce the use of psychoactive drugs in nursing homes. *N Engl J Med* 1992 Jul 16; 327(3):168-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Avorn J, Benner J, Ford I, et al. Measuring the cost-effectiveness of lipid-lowering drugs in the elderly: The outcomes research and economic analysis components of the PROSPER trial. *Control Clin Trials* 2002; 23(6):757-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Azuma T, Nagai Y, Saito T, et al. The effect of dehydroepiandrosterone sulfate administration to patients with multi-infarct dementia. JNS 1999 Jan 1; 162(1):69-73.

Status: Not included because dementia population not randomized to treatment

Bach D, Bach M, Bohmer F, et al. Reactivating occupational therapy: A method to improve cognitive performance in geriatric patients. Age Ageing 1995 May; 24(3):222-6.

Status: Not included because does not meet criteria for treatment for dementia patients

Bachynsky J, McCracken P, Lier D, et al. Propentofylline treatment for Alzheimer disease and vascular dementia: An economic evaluation based on functional abilities. Alzheimer Dis Assoc Disord 2000 Apr; 14(2):102-11.

Status: Not included because Jadad Quality Scale score less than three

Backonja M, Beydoun A, Edwards KR, et al. Gabapentin for the symptomatic treatment of painful neuropathy in patients with diabetes mellitus: A randomized controlled trial. JAMA 1998 Dec 2; 280(21):1831-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baines S, Saxby P, Ehler K. Reality orientation and reminiscence therapy. A controlled cross-over study of elderly confused people. Br J Psychiatry 1987; Vol 151:222-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baker R, Dowling Z, Wareing LA, et al. Snoezelen: Its long-term and short-term effects on older people with dementia. Br J Occup Ther 1997; (5):213-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baker R, Bell S, Baker E, et al. A randomized controlled trial of the effects of multi-sensory stimulation (MSS) for people with dementia. Br J Clin Psychol 2001 Mar; 40(Pt 1):1-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baladi JF, Bailey PA, Black S, et al. Rivastigmine for Alzheimer's disease: Canadian interpretation

of intermediate outcome measures and cost implications. Clin Ther 2000 Dec; 22(12):1549-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Baldereschi M, Di Carlo A, Lepore V, et al. Estrogen-replacement therapy and Alzheimer's disease in the Italian Longitudinal Study on Aging. Neurology 1998 Apr; 50(4):996-1002.

Status: Not included because dementia population not randomized to treatment

Balestreri R, Bompani R, Cerrato G. Comparative study of sulotidil and dihydroergotoxine in chronic cerebrovascular insufficiency. Results of a double blind double dummy multicentric trial. Acta Ther 1984; 10(2):163-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Balestreri R, Fontana L, Astengo F. A double-blind placebo controlled evaluation of the safety and efficacy of vinpocetine in the treatment of patients with chronic vascular senile cerebral dysfunction. J Am Geriatr Soc 1987 May; 35(5):425-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ball JA, Taylor AR. Effect of cyclandelate on mental function and cerebral blood flow in elderly patients. BMJ 1967 Aug 26; 3(564):525-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ballard C, O'Brien J, James I, et al. Quality of life for people with dementia living in residential and nursing home care: The impact of performance on activities of daily living, behavioral and psychological symptoms, language skills, and psychotropic drugs. Int Psychogeriatr 2001 Mar; 13(1):93-106.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ballard C, Powell I, James I, et al. Can psychiatric liaison reduce neuroleptic use and reduce health service utilization for dementia patients residing in care facilities? Int J Geriatr Psychiatry 2002 Feb; 17(2):140-5.

Status: Not included because dementia population not randomized to treatment

Ballard CG, O'Brien JT, Reichelt K, et al. Aromatherapy as a safe and effective treatment

for the management of agitation in severe dementia: The results of a double-blind, placebo-controlled trial with Melissa. *J Clin Psychiatry* 2002; 63(7):553-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Balldin J, Gottries CG, Karlson I, et al. Relationship between DST and the serotonergic system. Results from treatments with two 5-HT reuptake blockers in dementia disorders. *Int J Geriatr Psychiatry* 1988; 3(1):17-26.

Status: Not included because no extractable data relevant to review

Bambasova E, Bilkova J, Budinska K. Papaverin in the treatment of geriatric patients. *Act Nerv Super (Praha)* 1974 Aug; 16(3):192-3.

Status: Not included because dementia population not randomized to treatment

Ban TA, Modafferi A, Morey L. Global changes with glycosaminoglycan polysulfate in primary degenerative and multi-infarct dementia. *Curr Ther Res Clin Exp* 1987; 41(5):631-6.

Status: Not included because Jadad Quality Scale score less than three

Ban TA, Morey LC, Fjetland OK, et al. Early manifestations of dementing illness: Treatment with glycosaminoglycan polysulfate. *Prog Neuropsychopharmacol Biol Psychiatry* 1992 Sep; 16(5):661-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Banerjee S. Randomized controlled trials. *Int Rev Psychiatry* 1998; 10(4):291-303.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Barak Y, Levine J, Glasman A, et al. Inositol treatment of Alzheimer's disease: A double blind, cross-over placebo controlled trial. *Prog Neuropsychopharmacol Biol Psychiatry* 1996 May; 20(4):729-35.

Status: Cross-over trial;

Baro F, Malfroid M, Waegemans T, et al. Double-blind trial of suloctidil versus placebo in moderate to severe mental deterioration. *Pharmatherapeutica* 1985; 4(6):399-404.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bass DM, McClendon MJ, Brennan PF, et al. The buffering effect of a computer support network on caregiver strain. *J Aging Health* 1998; 10(1):20-43.

Status: Not included because does not meet criteria for treatment for dementia patients

Bassi S, Albizzati MG, Corsini GU, et al. Therapeutic experience with transdihydrolisuride in Huntington's disease. *Neurology* 1986; 36(7):984-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Battaglia A, Bruni G, Ardia A, et al. Nicergoline in mild to moderate dementia. A multicenter, double-blind, placebo-controlled study. *J Am Geriatr Soc* 1989 Apr; 37(4):295-302.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Battaglia A, Bruni G, Sacchetti G, et al. A double-blind randomized study of two ergot derivatives in mild to moderate dementia. *Curr Ther Res Clin Exp* 1990; 48(4):597-612.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Battistin L, Pizzolato G, Dam M, et al. Effects of acetyl-L-carnitine (ALC) treatment in dementia: A multicentric, randomized, double-blind study. *New Trends in Clinical Neuropharmacology* 1989; (2):131-2.

Status: Not included because Jadad Quality Scale score less than three

Bavazzano A, Guarducci R, Gestri G, et al. Clinical trial with amantadine and hydergine in elderly patients. *J Clin Exp Gerontol* 1980; (4):289-99.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bazo AJ. An ergot alkaloid preparation (Hydergine) versus papaverine in treating common complaints of the aged: Double-blind study. *J Am Geriatr Soc* 1973 Feb; 21(2):63-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beck C, Heacock P, Mercer SO, et al. Improving dressing behavior in cognitively impaired nursing home residents. *Nurs Res* 1997 May; 46(3):126-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beck CK, Vogelpohl TS, Rasin JH, et al. Effects of behavioral interventions on disruptive behavior and affect in demented nursing home residents. *Nurs Res* 2002; 51(4):219-28.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Beck C, Heacock P, Mercer S, et al. The impact of cognitive skills remediation training on persons with Alzheimer's disease or mixed dementia. *J Geriatr Psychiatry* 1988; 21(1):73-88.

Status: Not included because dementia population not randomized to treatment

Beckers T, Wagemans J, Boucart M, et al. Different effects of lorazepam and diazepam on perceptual integration. *Vision Res* 2001 Aug; 41(17):2297-303.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bedard M, Molloy DW, Standish T, et al. Clinical trials in cognitively impaired older adults: Home versus clinic assessments. *J Am Geriatr Soc* 1995 Oct; 43(10):1127-30.

Status: Not included because does not meet criteria for treatment for dementia patients

Bedard MA, Pilon B, Dubois B, et al. Acute and long-term administration of anticholinergics in Parkinson's disease: Specific effects on the subcortico-frontal syndrome. *Brain Cogn* 1999 Jul; 40(2):289-313.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Belanoff JK, Jurik J, Schatzberg LD, et al. Slowing the progression of cognitive decline in Alzheimer's disease using mifepristone. *J Mol Neurosci* 2002; 19(1-2):201-6.

Status: Not included because no extractable data relevant to review

Belfiore G, Di Maio L, Napolitano G, et al. Long-term effect of a single dose of flunarizine in Huntington's disease. *Eur J Neurol* 1998; 5(3):249-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bell IR, Edman JS, Morrow FD, et al. Brief communication. Vitamin B1, B2, and B6 augmentation of tricyclic antidepressant treatment in geriatric depression with cognitive dysfunction. *J Am Coll Nutr* 1992 Apr; 11(2):159-63.

Status: Not included because does not meet criteria for treatment for dementia patients

Beller SA, Overall JE, Swann AC. Efficacy of oral physostigmine in primary degenerative dementia. A double-blind study of response to different dose level. *Psychopharmacologia* 1985; 87(2):147-51.

Status: Cross-over trial;

Bellus SB, Vergo JG, Kost PP, et al. Behavioral rehabilitation and the reduction of aggressive and self-injurious behaviors with cognitively impaired, chronic psychiatric inpatients. *Psychiatr Q* 1999; 70(1):27-37.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ben Shlomo Y, Churchyard A, Head J, et al. Investigation by Parkinson's Disease Research Group of United Kingdom into excess mortality seen with combined levodopa and selegiline treatment in patients with early, mild Parkinson's disease: Further results of randomised trial and confidential inquiry. *BMJ* 1998 Apr 18; 316(7139):1191-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Benedict RH, Shapiro A, Priore R, et al. Neuropsychological counseling improves social behavior in cognitively-impaired multiple sclerosis patients. *Mult Scler* 2000 Dec; 6(6):391-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bentham PW. A double-blind placebo-controlled trial of L-tryptophan to assess the degree of cognitive and behavioural improvement in patients with Alzheimer-type dementia and to compare differential response in clinical sub-groups. *Int Clin Psychopharmacol* 1990; 5(4):261-72.

Status: Not included because Jadad Quality Scale score less than three

Bergamasco B, Villardita C, Coppi R. Idebenone in the treatment of multi-infarct dementia: A randomised, double-blind, placebo controlled multicentre trial. *Arch Gerontol Geriatr* 1992; 15(3):271-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bergamasco B, Villardita C, Coppi R. Effects of idebenone in elderly subjects with cognitive decline. Results of a multicentre clinical trial. *Arch Gerontol Geriatr* 1992; 15(3):279-86.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bernardi F, Lanzone A, Cento RM, et al. Allopregnanolone and dehydroepiandrosterone response to corticotropin-releasing factor in patients suffering from Alzheimer's disease and vascular dementia. *Eur J Endocrinol* 2000; 142(5):466-71.

Status: Not included because does not meet criteria for treatment for dementia patients

Besson JAO, Palin AN, Ebmeier KP, et al. Calcium antagonists and multi-infarct dementia: A trial involving sequential NMR and psychometric assessment. *Int J Geriatr Psychiatry* 1988; 3(2):99-105.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bierer LM, Aisen PS, Davidson M, et al. A pilot study of oral physostigmine plus yohimbine in patients with Alzheimer disease. *Alzheimer Dis Assoc Disord* 1993; 7(2):98-104.

Status: Not included because dementia population not randomized to treatment

Bierer LM, Aisen PS, Davidson M, et al. A pilot study of clonidine plus physostigmine in Alzheimer's disease. *Dementia* 1994 Sep; 5(5):243-6.

Status: Not included because Jadad Quality Scale score less than three

Binder EF, Schechtman KB, Birge SJ, et al. Effects of hormone replacement therapy on cognitive performance in elderly women. *Maturitas* 2001 Apr; 38(2):137-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Birge SJ. The role of estrogen in the treatment of Alzheimer's disease. *Neurology* 1997; 48(5 Suppl 7):S36-S41.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Birkett DP, Boltuch B. Chlorpromazine in geriatric psychiatry. *J Am Geriatr Soc* 1972 Aug; 20(8):403-6.

Status: Not included because no extractable data relevant to review

Bjorkman T, Hansson L, Sandlund M. Outcome of case management based on the strengths model compared to standard care. A randomised

controlled trial. *Soc Psychiatry Psychiatr Epidemiol* 2002 Apr; 37(4):147-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blahe L, Erzigkeit H, Adamczyk A, et al. Clinical evidence of the effectiveness of vinpocetine in the treatment of organic psychosyndrome. *Hum Psychopharmacol* 1989; (2):103-11.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blass JP, Gleason P, Brush D, et al. Thiamine and Alzheimer's disease. A pilot study. *Arch Neurol* 1988 Aug; 45(8):833-5.

Status: Cross-over trial;

Blass JP, Cyrus PA, Bieber F, et al. Randomized, double-blind, placebo-controlled, multicenter study to evaluate the safety and tolerability of metrifonate in patients with probable Alzheimer's disease. *Alzheimer Dis Assoc Disord* 2000; 14(1):39-45.

Status: Not included because no extractable data relevant to review

Blazer DG, Landerman LR, Hays JC, et al. Symptoms of depression among community-dwelling elderly African-American and white older adults. *Psychol Med* 1998 Nov; 28(6):1311-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blin J, Piercey MF, Giuffra ME, et al. Metabolic effects of scopolamine and physostigmine in human brain measured by positron emission tomography. *JNS* 1994 May; 123(1-2):44-2.

Status: Not included because does not meet criteria for treatment for dementia patients

Blin J, Ivanoiu A, Coppens A, et al. Cholinergic neurotransmission has different effects on cerebral glucose consumption and blood flow in young normals, aged normals, and Alzheimer's disease patients. *Neuroimage* 1997 Nov; 6(4):335-43.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Blin J, Ivanoiu A, De Volder A, et al. Physostigmine results in an increased decrement in brain glucose consumption in Alzheimer's disease. *Psychopharmacologia* 1998 Apr; 136(3):256-63.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Block RI, DeVoe M, Stanley B, et al. Memory performance in individuals with primary degenerative dementia: Its similarity to diazepam-induced impairments. *Exp Aging Res* 1985; 11(3-4):151-4.

Status: Not included because does not meet criteria for treatment for dementia patients

Blume J, Ruhlmann KU, de la Haye R, et al. Treatment of chronic cerebrovascular disease in elderly patients with pentoxifylline. *J Med* 1992; 23(6):417-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bodick NC, Offen WW, Shannon HE, et al. The selective muscarinic agonist xanomeline improves both the cognitive deficits and behavioral symptoms of Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1997; 11(Suppl 4):S16-S22.

Status: Not included because Jadad Quality Scale score less than three

Boelhouwer C, Henry CE, Glueck BC, Jr. Positive spiking: A double-blind control study on its significance in behavior disorders, both diagnostically and therapeutically. *Am J Psychiatry* 1968 Oct; 125(4):473-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bollen ELEM, Gaw A, Buckley BM, et al. Statin therapy and the prevention of dementia. *Arch Neurol* 2001; 58(6):1023-4.

Status: Not included because not a full article

Bompani R, Scali G. Fipexide, an effective cognition activator in the elderly: A placebo-controlled, double-blind clinical trial. *Curr Med Res Opin* 1986; 10(2):99-106.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bonavita E. Study of the efficacy and tolerability of L-acetylcarnitine therapy in the senile brain. *Int J Clin Pharmacol Ther Toxicol* ; 24(9):511-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bonura ML, Grigoletto F. A 6-month, multicentre, double-blind trial of nicergoline in the treatment of mild to moderate Alzheimer's disease and its 12-month follow-up: Preliminary results. *J Neural Transm Gen Sect* 2000; XVIII.

Status: Not included because not a full article

Boon AJ, Tans JT, Delwel EJ, et al. Does CSF outflow resistance predict the response to shunting in patients with normal pressure hydrocephalus? *Acta Neurochir Suppl* 1998; 71:331-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Borjesson A, Karlsson T, Adolfsson R, et al. Linopirdine (DUP 996): Cholinergic treatment of older adults using successive and non-successive tests. *Neuropsychobiology* 1999; 40(2):78-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Borromei A, Gaggi R, Giancola LC. Involuntional dementias: New perspectives. *Ital J Neurol Sci* 1985; 6(2):167-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Borroni B, Colciaghi F, Pastorino L, et al. Amyloid precursor protein in platelets of patients with Alzheimer's disease: Effect of acetylcholinesterase inhibitor treatment. *Arch Neurol* 2001; 58(3):442-6.

Status: Not included because dementia population not randomized to treatment

Bottiglieri T, Godfrey P, Flynn T, et al. Cerebrospinal fluid S-adenosylmethionine in depression and dementia: Effects of treatment with parenteral and oral S-adenosylmethionine. *J Neurol Neurosurg Psychiatry* 1990; 53(12):1096-8.

Status: Not included because dementia population not randomized to treatment

Bourgeois MS, Burgio LD, Schulz R, et al. Modifying repetitive verbalizations of community-dwelling patients with AD. *Gerontologist* 1997 Feb; 37(1):30-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Bourgeois MS, Dijkstra K, Burgio L, et al. Memory aids as an augmentative and alternative communication strategy for nursing home residents with dementia. *Aac: Augmentative & Alternative Communication* 2001 Sep; 17(3):196-210.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bower HM, McDonald C. A controlled trial of A.N.P. 235 ("Lucidril") in senile dementia. Med J Aust 1966 Aug 6; 2(6):270-1.

Status: Not included because dementia population not randomized to treatment

Bowles EJ, Griffiths DM, Quirk L, et al. Effects of essential oils and touch on resistance to nursing care procedures and other dementia-related behaviours in a residential care facility. Int J Aromather 2002; 12(1):22-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bowling A, Formby J, Grant K, et al. A randomized controlled trial of nursing home and long-stay geriatric ward care for elderly people. Age Ageing 1991 Sep; 20(5):316-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO. A memory assessment technique for use in geriatric psychopharmacology: Drug efficacy trial with naftidrofuryl. J Am Geriatr Soc 1977 Apr; 25(4):186-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO. The impairment index as a symptom-independent parameter of drug efficacy in geriatric psychopharmacology. J Gerontol 1978 Mar; 33(2):217-23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO, Gardos G. ACTH 4-10 in the amelioration of neuropsychological symptomatology associated with senile organic brain syndrome. Psychopharmacologia 1979 Mar; 61(2):161-5.

Status : Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO, Ghazvinian S, et al. Treating the depressed elderly patient: The comparative behavioral pharmacology of mianserin and amitriptyline. Adv Biochem Psychopharmacol 1982; Vol 32:195-212.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Cole JO, Dessain EC, et al. The therapeutic efficacy of pramiracetam in Alzheimer's disease: Preliminary observations. Psychopharmacol Bull 1983; 19(4):726-30.

Status: Not included because dementia population not randomized to treatment

Branconnier RJ, Harto NE, Dessain EC, et al. Speech blockage, memory impairment, and age: a prospective comparison of amitriptyline and maprotiline. Psychopharmacol Bull 1987; 23(1):230-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Branconnier RJ, Harto NE, Dessain EC, et al. Palliation of the progressive memory impairment of Alzheimer's disease by nimodipine. Psychopharmacologia 1988; 96(Suppl):242.

Status: Not included because not a full article

Branconnier RJ, Cole JO. Effects of chronic papaverine administration on mild senile organic brain syndrome. J Am Geriatr Soc 1977; 25(10):458-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brandimonte MA, Passolunghi MC. The effect of cue-familiarity, cue-distinctiveness, and retention interval on prospective remembering. Q J Exp Psychol A 1994 Aug; 47(3):565-87.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brass EP, Polinsky R, Sramek JJ, et al. Effects of the cholinomimetic SDZ ENS-163 on scopolamine-induced cognitive impairment in humans. J Clin Psychopharmacol 1995 Feb; 15(1):58-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brautigam MRH, Blommaert FA, Verleye G, et al. Treatment of age-related memory complaints with ginkgo biloba extract: A randomized double blind placebo-controlled study. Phytomedicine 1998; 5(6):425-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brauzer B, Goldstein BJ. The differential response to parenteral chlorpromazine and mesoridazine in psychotic patients. J Clin Pharmacol New Drugs 1970 Mar; 10(2):126-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Braverman AM, Naylor CD. Vasoactive substances in the management of elderly patients

suffering from dementia. *Modern Geriatrics* 1975; (5):20-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brazzelli M, Capitani E, Della SS, et al. A neuropsychological instrument adding to the description of patients with suspected cortical dementia: The Milan overall dementia assessment. *J Neurol Neurosurg Psychiatry* 1994; 57(12):1510-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Breeze RW, Cox S, Rodgers CJ. Changes in P-300 latency as a result of co-dergocrine mesylate therapy in patients with senile dementia. *Int J Geriatr Psychiatry* 1988; 3(4):263-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Breitbart W, Marotta R, Platt MM, et al. A double-blind trial of haloperidol, chlorpromazine, and lorazepam in the treatment of delirium in hospitalized AIDS patients. *Am J Psychiatry* 1996 Feb; 153(2):231-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Breitner JC, Gau BA, Welsh KA, et al. Inverse association of anti-inflammatory treatments and Alzheimer's disease: Initial results of a co-twin control study. *Neurology* 1994 Feb; 44(2):227-32.

Status: Not included because dementia population not randomized to treatment

Breitner JC. The role of anti-inflammatory drugs in the prevention and treatment of Alzheimer's disease. *Annu Rev Med* 1996; Vol 47:401-11.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Breitner JCS, Zandi PP, In't Veld BA. Long-term use of non-steroidal anti-inflammatory drugs (NSAIDs) reduce the relative risk of Alzheimer's disease to 0.2. *J Neurol* 2002; 249(3):355-6.

Status: Not included because not a full article

Breuil V, de Rotrou J, Forette F, et al. Cognitive stimulation of patients with dementia: Preliminary results. *Int J Geriatr Psychiatry* 1994; 9(3):211-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Bridges-Parlet S, Knopman D, Steffes S. Withdrawal of neuroleptic medications from

institutionalized dementia patients: Results of a double-blind, baseline-treatment-controlled pilot study. *J Geriatr Psychiatry Neurol* 1997 Jul; 10(3):119-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brinkman SD, Smith RC, Meyer JS. Lecithin and memory training in suspected Alzheimer's disease. *J Gerontol* 1982; 37(1):4-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brinkman SD, Pomara N, Goodnick PJ, et al. A dose-ranging study of lecithin in the treatment of primary degenerative dementia (Alzheimer's disease). *J Clin Psychopharmacol* 1982 Aug; 2(4):281-5.

Status: Cross-over trial;

Brodsky H, Roberts K, Peters K. Quasi-experimental evaluation of an educational model for dementia caregivers. *Int J Geriatr Psychiatry* 1994; 9(3):195-204.

Status: Not included because dementia population not randomized to treatment

Brodsky H, Gresham M, Luscombe G. The Prince Henry Hospital dementia caregivers' training programme. *Int J Geriatr Psychiatry* 1997 Feb; 12(2):183-92.

Status: Not included because does not meet criteria for treatment for dementia patients

Broderick JP, Gaskill M, Dhawan A, et al. Temporal changes in brain volume and cognition in a randomized treatment trial of vascular dementia. *J Neuroimaging* 2001 Jan; 11(1):6-12.

Status: Not included because Jadad Quality Scale score less than three

Brodersen P, Philbert A, Gulliksen G, et al. The effect of L-Deprenyl on on-off phenomena in Parkinson's disease. *Acta Neurol Scand* 1985 Jun; 71(6):494-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brodie NH. A double-blind trial of naftidrofuryl in treating confused elderly patients in general practice. *Practitioner* 1977 Feb; 218(1304):274-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brody EM, Kleban MH, Lawton MP, et al. A longitudinal look at excess disabilities in the

mentally impaired aged. J Gerontol 1974; 29(1):79-84.

Status: Not included because dementia population not randomized to treatment

Brooker D, Duce L. Wellbeing and activity in dementia: A comparison of group reminiscence therapy, structured goal-directed group activity and unstructured time. Aging Ment Health 2000; 4(4):354-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brooker DJ, Snape M, Johnson E, et al. Single case evaluation of the effects of aromatherapy and massage on disturbed behaviour in severe dementia. Br J Clin Psychol 1997 May; 36(Pt 2):287-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brouwers P, Hendricks M, Lietzau JA, et al. Effect of combination therapy with zidovudine and didanosine on neuropsychological functioning in patients with symptomatic HIV disease: A comparison of simultaneous and alternating regimens. AIDS 1997 Jan; 11(1):59-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Brown S, Gotell E, Ekman SL. Singing as a therapeutic intervention in dementia care. Journal of Dementia Care 2001; (4):33-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Brun A, Gustafson L. The lund longitudinal dementia study a 25 year prespective on neuropathology differential diagnosis and treatment. In: Corian B, Iqbal K, Nicolini M, Winblad B, Wisniewski HM, Zatta PF, editors. Alzheimer's Disease: Advances in Clinical and Basic Research, Chichester: John Wiley & Sons; 1993. Chapter 1. p. 3-18.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bruno G, Mohr E, Gillespie M, et al. Muscarinic agonist therapy of Alzheimer's disease. A clinical trial of RS-86. Arch Neurol 1986 Jul; 43(7):659-61.

Status: Cross-over trial;

Buettner LL, Lundegren H, Lago D, et al. Therapeutic recreation as an intervention for persons with dementia and agitation: An efficacy

study. Am J Alzheimers Dis 1996; (5):4-10.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Buettner LL. Focus on caregiving. Falls prevention in dementia populations: Following a trial program of recreation therapy, falls were reduced by 164 percent. Provider 2002 Feb; 28(2):41-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Buettner LL, Fitzsimmons S. AD-venture program: Therapeutic biking for the treatment of depression in long-term care residents with dementia. Am J Alzheimers Dis Other Dement 2002 Mar; 17(2):121-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Bukatina EE, Grigor'eva IV, Sokol'chik EI. The effectiveness of amiridin in senile dementia of the Alzheimer's type. Neurosci Behav Physiol 1993 Jan; 23(1):83-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Burgener SC, Bakas T, Murray C, et al. Effective caregiving approaches for patients with Alzheimer's disease. Geriatr Nurs (Minneapolis) 1998 May; 19(3):121-6.

Status: Not included because does not meet criteria for treatment for dementia patients

Burgio LD, Reynolds CFI, Janosky JE, et al. A behavioral microanalysis of the effects of haloperidol and oxazepam in demented psychogeriatric inpatients. Int J Geriatr Psychiatry 1992; 7(4):253-62.

Status: Not included because Jadad Quality Scale score less than three

Burns A, Marsh A, Bender DA. A trial of vitamin supplementation in senile dementia. Int J Geriatr Psychiatry 1989; 4(6):333-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cahn LA, Diesfeldt HF. The use of neuroleptics in the treatment of dementia in old age. A critical analysis with reference to an experiment with a long-acting oral neuroleptic (penfluridol Janssen). Psychiatr Neurol Neurochir 1973; 76(6):411-20.

Status: Not included because dementia population not randomized to treatment

Caligiuri MP, Lacro JP, Jeste DV. Incidence and predictors of drug-induced parkinsonism in older psychiatric patients treated with very low doses of neuroleptics. *J Clin Psychopharmacol* 1999 Aug; 19(4):322-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Camberg L, Woods P, Ooi WL, et al. Evaluation of Simulated Presence: A personalized approach to enhance well-being in persons with Alzheimer's disease. *J Am Geriatr Soc* 1999 Apr; 47(4):446-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Campi N, Todeschini GP, Scarzella L. Selegiline versus L-acetylcarnitine in the treatment of Alzheimer-type dementia. *Clin Ther* 1990 Jul; 12(4):306-14.

Status: Not included because Jadad Quality Scale score less than three

Campion D, Brice A, Hannequin D, et al. A large pedigree with early-onset Alzheimer's disease: Clinical, neuropathologic, and genetic characterization. *Neurology* 1995 Jan; 45(1):80-5.

Status: Not included because does not meet criteria for treatment for dementia patients

Canal N, Imbimbo N. A 25-week double-blind randomized placebo-controlled trial of Eptastigmine in patients with diagnosis of probable Alzheimer's disease. *J Neural Transm Gen Sect* 1996; 103:XXIV.

Status: Not included because not a full article

Cantillon M, Brunswick R, Molina D, et al. Buspirone vs. haloperidol: A double-blind trial for agitation in a nursing home population with Alzheimer's disease. *Am J Geriatr Psychiatry* 1996; 4(3):263-7.

Status: Not included because Jadad Quality Scale score less than three

Capote B, Parikh N. Cyclandelate in the treatment of senility: a controlled study. *J Am Geriatr Soc* 1978 Aug; 26(8):360-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Capurso A, Capurso S, Panza F, et al. Efficacy of cytidine diphosphate choline in patients affected by chronic cerebrovascular disease. *Clin Drug Investig* 1996; (1):26-38.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Caraceni TA, Girotti F, Celano I, et al. 2-dimethylaminoethanol (Deanol) in Huntington's chorea. *J Neurol Neurosurg Psychiatry* 1978; 41(12):1114-8.

Status: Not included because no extractable data relevant to review

Carbonin PU, Greco A, Pisanti P, et al. Efficacy of almitrine-raubasine in cognitive disorders of aging: A double-blind, placebo-controlled, clinical and psychometric study. *Clin Neuropharmacol* 1990; Vol 13(Suppl 3):S92-S99

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cardebat D, Demonet J-F, Puel M, et al. Brain correlates of memory processes in patients with dementia of Alzheimer's type: A SPECT activation study. *J Cereb Blood Flow Metab* 1998; 18(4):457-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Carman JS, Shoulson I, Chase TN. Huntington's chorea treated with lithium carbonate. *Lancet* 1974; 1(7861):811.

Status: Not included because not a full article

Carmel R. Mild cobalamin deficiency. *West J Med* 1998 Jun; 168(6):522-3.

Status: Not included because not a full article

Caro AJ, Caro S. Vitamin E in treatment of Huntington's chorea. *BMJ* 1978 Jan 21; 1(6106):153.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Caro JJ, Salas M, Ward A, et al. Economic analysis of galantamine, a cholinesterase inhibitor, in the treatment of patients with mild to moderate Alzheimer's disease in the Netherlands. *Dement Geriatr Cogn Disord* 2002; 14(2):84-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Carter LT, Howard BE, O'Neil WA. Effectiveness of cognitive skill remediation in acute stroke patients. *Am J Occup Ther* 1983 May; 37(5):320-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Carver A, Dobson AM. Effects of dietary supplementation on demented elderly hospital residents. *Age Ageing* 1993; 22(Suppl 3):37.
Status: Not included because not a full article

Carver AD, Dobson AM. Effects of dietary supplementation of elderly demented hospital residents. *J Hum Nutr Diet* 1995; 8(6):389-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Casale R, Giorgi I, Guarnaschelli C. Evaluation of the effect of vincamine teprosilate on behavioural performances of patients affected with chronic cerebrovascular disease. *Int J Clin Pharmacol Res* 1984; 4(4):313-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cebul RD. Aspirin and MID Notes of caution. *J Am Geriatr Soc* 1989; 37(6):573-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cenacchi T, Bertoldin T, Farina C, et al. Cognitive decline in the elderly: A double-blind, placebo-controlled multicenter study on efficacy of phosphatidylserine administration. *Aging (Milano)* 1993 Apr; 5(Milano):123-33.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chabriat H, Pappata S, Ostergaard L, et al. Cerebral hemodynamics in CADASIL before and after acetazolamide challenge assessed with MRI bolus tracking. *Stroke* 2000 Aug; 31(8):1904-12.
Status: Not included because dementia population not randomized to treatment

Challis D, Von Abendorff R, Brown P, et al. Care management, dementia care and specialist mental health services: An evaluation. *Int J Geriatr Psychiatry* 2002; 17(4):315-25.
Status: Not included because dementia population not randomized to treatment

Chandra B. Treatment of multi-infarct dementia with citicholine. *Journal of Stroke and Cerebrovascular Diseases* 1992; 232-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chandra B. Treatment of vascular dementia with CDP choline. *Journal of Stroke and Cerebrovascular Diseases* 2000; 9(Suppl 2):128-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Chang BL. Cognitive-behavioral intervention for homebound caregivers of persons with dementia. *Nurs Res* 1999 May; 48(3):173-82.
Status: Not included because does not meet criteria for treatment for dementia patients

Chatellier G, Lacomblez L, and Group. Tacrine (tetrahydroaminoacridine; THA) and lecithin in senile dementia of the Alzheimer type: A multicentre trial. *BMJ* 1990; 300(6723):495-9.
Status: Cross-over trial;

Christe C, Janssens JP, Armenian B, et al. Midazolam sedation for upper gastrointestinal endoscopy in older persons: A randomized, double-blind, placebo-controlled study. *J Am Geriatr Soc* 2000 Nov; 48(11):1398-403.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Christensen DB, Benfield WR. Alprazolam as an alternative to low-dose haloperidol in older, cognitively impaired nursing facility patients. *J Am Geriatr Soc* 1998 May; 46(5):620-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Christie JE, Shering A, Ferguson J, et al. Physostigmine and arecoline: Effects of intravenous infusions in Alzheimer presenile dementia. *Br J Psychiatry* 1981; Vol 138:46-50.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Churchill M, Safaoui J, McCabe BW, et al. Using a therapy dog to alleviate the agitation and desocialization of people with Alzheimer's disease. *J Psychosoc Nurs Ment Health Serv* 1999 Apr; 37(4):16-22.
Status: Not included because dementia population not randomized to treatment

Citrin RS, Dixon DN. Reality Orientation: A Milieu Therapy Used in an Institution for the Aged. *Gerontologist* 1977; 17(1):39-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clair AA, Bernstein B. Effect of no music, stimulative background music and sedative background music on agitated behaviors in persons with severe dementia. *Activities Adaptation Aging* 1994; (1):61-70.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clair AA. Effect of singing on alert responses in persons with late stage dementia. *J Music Ther* 1996; (No. 4):234-47.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clark LR, Fraaza V, Schroeder S, et al. Alternative nursing environments: Do they affect hospital outcomes? *J Gerontol Nurs* 1995 Nov; 21(11):32-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clark ME, Lipe AW, Bilbrey M. Use of music to decrease aggressive behaviors in people with dementia. *J Gerontol Nurs* 1998 Jul; 24(7):10-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Claus JJ, Ludwig C, Mohr E, et al. Nootropic drugs in Alzheimer's disease: Symptomatic treatment with pramiracetam. *Neurology* 1991 Apr; 41(4):570-4.

Status: Cross-over trial;

Claus JJ, Mohr E, Chase TN. Clinical trials in dementia: Learning effects with repeated testing. *J Psychiatry Neurosci* 1991 Mar; 16(1):1-4.
Status: Not included because does not meet criteria for treatment for dementia patients

Claus JJ, van Harksamp F, de K, I, et al. Serotonergic mechanisms in Alzheimer's disease: Preliminary results of a controlled clinical trial with lisuride. *Can J Neurol Sci* 1993; Vol 20:126.
Status : Not included because not a full article

Clifford DB, McArthur JC, Schifitto G, et al. A randomized clinical trial of CPI-1189 for HIV-associated cognitive-motor impairment. *Neurology* 2002 Nov 26; 59(10):1568-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Clipp EC, Moore MJ. Caregiver time use: an outcome measure in clinical trial research on Alzheimer's disease. *Clin Pharmacol Ther* 1995 Aug; 58(2):228-36.
Status: Not included because Jadad Quality Scale score less than three

Cole MG, McCusker J, Bellavance F, et al. Systematic detection and multidisciplinary care of

delirium in older medical inpatients: a randomized trial. *CMAJ* 2002 Oct 1; 167(7):753-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Colling KB, Buettner LL. Simple pleasures. Interventions from the need-driven Dementia-Compromised Behavior model. *J Gerontol Nurs* 2002 Oct; 28(10):16-20.

Status: Not included because does not meet criteria for treatment for dementia patients

Comelli M, Lucca U, Spagnoli A. Statistical analysis of the clinical trial of a therapy for Alzheimer's disease. Univariate tests and logistic regression. *Acta Neurol (Napoli)* 1990 Jun; 12(3):222-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Commissaris K, Verhey FR, Jolles J. A controlled study into the effects of psychoeducation for patients with cognitive disturbances. *J Neuropsychiatry Clin Neurosci* 1996; 8(4):429-35.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Conti L, Re F, Lazzerini F, et al. Glycosaminoglycan polysulfate (Ateroid) in old-age dementias: Effects upon depressive symptomatology in geriatric patients. *Prog Neuropsychopharmacol Biol Psychiatry* 1989; 13(6):977.
Status: Not included because Jadad Quality Scale score less than three

Convit A, de Asis J, de Leon MJ, et al. Atrophy of the medial occipitotemporal, inferior, and middle temporal gyri in non-demented elderly predict decline to Alzheimer's disease. *Neurobiol Aging* 2000 Jan; 21(1):19-26.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cook WA. Methylperiodol: Clinical trials of a new tranquilizer. *Med J Aust* 1966 Jul 16; 2(3):117-9.
Status: Not included because dementia population not randomized to treatment

Cools R, Barker RA, Sahakian BJ, et al. Enhanced or impaired cognitive function in Parkinson's disease as a function of dopaminergic medication and task demands. *Cereb Cortex* 2001 Dec; 11(12):1136-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cooney C, Mortimer A, Smith A, et al. Carbamazepine use in aggressive behaviour associated with senile dementia. *Int J Geriatr Psychiatry* 1996; 11(10):901-5.

Status: Cross-over trial;

Cooper AJ, Wong YT, Packer H. A controlled trial of cosaldon in arteriosclerotic dementia: Penicillin versus penicillin-malaria in the treatment of dementia paralytica. *British Journal Of Psychiatry: British Journal of Venereal Diseases* 1964; 25(1):415-8, 439.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cooper AJ, Magnus RV. A placebo-controlled study of pyritinol ('Encephabol') in dementia. *Pharmatherapeutica* 1980; 2(5):317-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cooper JA, Sagar HJ, Doherty SM, et al. Different effects of dopaminergic and anticholinergic therapies on cognitive and motor function in Parkinson's disease. *Brain* 1992; 115(6):1701-25.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Corbeil RR, Quayhagen MP, Quayhagen M. Intervention effects on dementia caregiving interaction: A stress-adaptation modeling approach. *J Aging Health* 1999 Feb; 11(1):79-95.

Status: Not included because does not meet criteria for treatment for dementia patients

Corcoran MA, Gitlin LN. Family caregiver acceptance and use of environmental strategies provided in an occupational therapy intervention. *Phys Occup Ther Geriatr* 2001; 19(1):1-20.

Status: Not included because does not meet criteria for treatment for dementia patients

Corona GL, Cucchi ML, Frattini P, et al. Clinical and biochemical responses to therapy in Alzheimer's disease and multi-infarct dementia. *Eur Arch Psychiatry Neurol Sci* 1989; 239(2):79-86.

Status: Not included because Jadad Quality Scale score less than three

Corrigan FM, Van Rhijn A, Horrobin DF. Essential fatty acids in Alzheimer's disease. *Ann N Y Acad Sci* 1991; Vol 640:250-2.

Status: Not included because no extractable data relevant to review

Cott CA, Dawson P, Sidani S, et al. The effects of a walking/talking program on communication, ambulation, and functional status in residents with Alzheimer's disease. *Alzheimer Dis Assoc Disord* 2002; 16(2):81-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Coull JT, Sahakian BJ, Hodges JR. The alpha(2) antagonist idazoxan remediates certain attentional and executive dysfunction in patients with dementia of frontal type. *Psychopharmacologia* 1996 Feb; 123(3):239-49.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Covington JS. Alleviating agitation, apprehension, and related symptoms in geriatric patients: A double-blind comparison of a phenothiazine and a benzodiazepien. *South Med Assoc J* 1975 Jun; 68(6):719-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cowan P. "Special care" for dementia patients. *Can Nurse* 1999 Jun; 95(6):49-50.

Status: Not included because dementia population not randomized to treatment

Cowley LM, Glen RS. Double-blind study of thioridazine and haloperidol in geriatric patients with a psychosis associated with organic brain syndrome. *J Clin Psychiatry* 1979 Oct; 40(10):411-9.

Status : Not included because dementia population not defined by DSM, NINCDS or ICD

Cox JR. Double-blind evaluation of naftidrofuryl in treating elderly confused hospitalised patients. *Gerontol Clin (Basel)* 1975; 17(3):160-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cox JR, Shaw AM. Controlled trial of naftidrofuryl in dementia in old age. *J Clin Exp Gerontol* 1981; (4):339-43.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Coyle J, Kershaw P. Galantamine, a cholinesterase inhibitor that allosterically modulates nicotinic receptors: Effects on the course of Alzheimer's disease. *Biol Psychiatry*

2001; 49(3):289-99.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Coyne AC, Potenza M, Broken-Nose MA. Caregiving and dementia: The impact of telephone helpline services. *Am J Alzheimers Dis* 1995 Jul; 4(4):27-32.

Status: Not included because does not meet criteria for treatment for dementia patients

Coyne ML, Hoskins L. Improving eating behaviors in dementia using behavioral strategies. *Clin Nurs Res* 1997 Aug; 6(3):275-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Craft S, Asthana S, Newcomer JW, et al. Enhancement of memory in Alzheimer disease with insulin and somatostatin, but not glucose. *Arch Gen Psychiatry* 1999 Dec; 56(12):1135-40.

Status: Not included because dementia population not randomized to treatment

Crook, T.H. A 6-month, double-blind, placebo-controlled trial of nicergoline in patients with mild to moderate probable Alzheimer's disease. *J Neural Transm Gen Sect* 2000; 107:XVIII.

Status: Not included because not a full article

Crook T, Ferris S, Sathananthan G, et al. The effect of methylphenidate on test performance in the cognitively impaired aged. *Psychopharmacologia* 1977 May 9; 52(3):251-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Crook TH, Tinklenberg J, Yesavage J, et al. Effects of phosphatidylserine in age-associated memory impairment. *Neurology* 1991 May; 41(5):644-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Crook TH. Nicergoline in the treatment of probable Alzheimer's disease: Preliminary results of a double-blind, randomized, placebo-controlled study. *JNS* 1997; 150(Suppl 1):S18.

Status: Not included because not a full article

Cucinotta D, Passeri M, Ventura S, et al. Multicenter clinical placebo-controlled study with acetyl-L-carnitine (LAC) in the treatment of mildly demented elderly patients. *Drug Dev Res* 1988; 14(3-4):213-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cui Y. Anti-senility potential of Zusanli. *Int J Clin Acupunct* 1995; 6(1):1-4.

Status: Not included because dementia population not randomized to treatment

Culebras A. Effect of papaverine on cerebral electrogenesis. *Neurology* 1976 Jul; 26(7):673-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Cummings JL, Gorman DG, Shapira J. Physostigmine ameliorates the delusions of Alzheimer's disease. *Biol Psychiatry* 1993 Apr 1; 33(7):536-41.

Status: Not included because dementia population not randomized to treatment

Cummings JL, Knopman D. Advances in the treatment of behavioral disturbances in Alzheimer's disease. *Neurology* 2000; 53(5):899

Status: Not included because not a full article

Cummings JL, Nadel A, Masterman D, et al. Efficacy of metrifonate in improving the psychiatric and behavioral disturbances of patients with Alzheimer's disease. *J Geriatr Psychiatry Neurol* 2001; 14(2):101-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cummings JL, Street J, Masterman D, et al. Efficacy of olanzapine in the treatment of psychosis in dementia with lewy bodies. *Dement Geriatr Cogn Disord* 2002; 13(2):67-73.

Status: Not included because Jadad Quality Scale score less than three

Curless R, James OFW, McKeith I, et al. Effects of propranolol on aggressive behaviour in elderly patients with dementia. *Age Ageing* 1994; 23:P19.

Status: Not included because not a full article

Curtis-Prior P, Vere D, Fray P. Therapeutic value of Ginkgo biloba in reducing symptoms of decline in mental function. *J Pharm Pharmacol* 1999 May; 51(5):535-41.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cutler NR, Haxby J, Kay AD, et al. Evaluation of zimeldine in Alzheimer's disease. Cognitive and biochemical measures. *Arch Neurol* 1985 Aug;

42(8):744-8.

Status: Cross-over trial;

Cutler NR, Haxby JV, Narang PK, et al. Evaluation of an analogue of somatostatin (L363,586) in Alzheimer's disease. *N Engl J Med* 1985 Mar; 312(11):725.

Status: Not included because not a full article

Cutler NR, Murphy MF, Nash RJ, et al. Clinical safety, tolerance, and plasma levels of the oral anticholinesterase 1,2,3,4-tetrahydro-9-aminoacridin-1-ol-maleate (HP 029) in Alzheimer's disease: Preliminary findings. *J Clin Pharmacol* 1990 Jun; 30(6):556-61.

Status: Not included because no extractable data relevant to review

Cutler NR, Sramek JJ, Murphy MF, et al. Implications of the study population in the early evaluation of anticholinesterase inhibitors for Alzheimer's disease. *Ann Pharmacother* 1992 Sep; 26(9):1118-22.
Status: Not included because no extractable data relevant to review

Cutler NR, Sramek JJ, Murphy MF, et al. Alzheimer's patients should be included in phase I clinical trials to evaluate compounds for Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1992 Oct; 5(4):192-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Cutler NR, Fakouhi TD, Smith WT, et al. Evaluation of multiple doses of milacemide in the treatment of senile dementia of the Alzheimer's type. *J Geriatr Psychiatry Neurol* 1993 Apr; 6(2):115-9.
Status: Not included because Jadad Quality Scale score less than three

Cutler NR, Sramek JJ, Anand R. Safety and tolerance of ENA 713 in patients with Alzheimer's disease. *Biol Psychiatry* 1995; 37(9):643.
Status: Not included because not a full article

Cutler NR, Sramek JJ, Viereck C, et al. Tolerability and pharmacodynamics of besipirdine in Alzheimer's disease. *Biol Psychiatry* 1995; 37(9):643.
Status: Not included because not a full article

Cutler NR, Polinsky RJ, Sramek JJ, et al. Dose-dependent CSF acetylcholinesterase inhibition by SDZ ENA 713 in Alzheimer's disease. *Acta*

Neurol Scand 1998 Apr; 97(4):244-50.

Status: Not included because dementia population not randomized to treatment

Cutler NR, Jhee SS, Cyrus P, et al. Safety and tolerability of metrifonate in patients with Alzheimer's disease: Results of a maximum tolerated dose study. *Life Sci* 1998; 62(16):1433-41.

Status: Not included because dementia population not randomized to treatment

Czerwinski AW, Clark ML, Serafetinides EA, et al. Safety and efficacy of zinc sulfate in geriatric patients. *Clin Pharmacol Ther* 1974 Apr; 15(4):436-41.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Daniele A, Moro E, Bentivoglio AR. Zolpidem in progressive supranuclear palsy. *N Engl J Med* 1999 Aug 12; 341(7):543-4.
Status: Not included because not a full article

Daniels L. A group cognitive-behavioural and process-oriented approach to treating the social impairment and negative symptoms associated with chronic mental illness. *J Psychother Pract Res* 1998; 7:167-76.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dansky KH, Dellasega C, Shellenbarger T, et al. After hospitalization: Home health care for elderly persons. *Clin Nurs Res* 1996 May; 5(2):185-98.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Danysz W. CX-516 Cortex Pharmaceuticals Inc. *Idrugs* 1999; 2(8):814-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Darreh-Shori T, Almkvist O, Guan ZZ, et al. Sustained cholinesterase inhibition in AD patients receiving rivastigmine for 12 months. *Neurology* 2002 Aug 27; 59(4):563-72.
Status: Not included because dementia population not randomized to treatment

Davidson M, Mohs RC, Hollander E, et al. Lecithin and piracetam in Alzheimer's disease. *Biol Psychiatry* 1987; 22(1):112-4.
Status: Not included because not a full article

Davidson M, Zemishlany Z, Mohs RC, et al. 4-Aminopyridine in the treatment of Alzheimer's disease. *Biol Psychiatry* 1988 Mar 1; 23(5):485-90.

Status: Cross-over trial;

Davies AE. A pilot study to measure aluminium levels in hair samples of patients with dementia and the influence of aluminium 30c compared with placebo. *Communications of the British Homoeopathy Research Group Issue* 18, 1988; 42-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davies B, Andrewes D, Stargatt R, et al. Tacrine in Alzheimer's disease. *Lancet* 1989 Jul 15; 2(8655):163-4.

Status: Not included because not a full article

Davies B, Andrewes D, Stargatt R, et al. Tetrahydroaminoacridine in Alzheimer's disease. *Int J Geriatr Psychiatry* 1990; 5(5):317-21.

Status: Cross-over trial;

Davies G, Hamilton S, Hendrickson E, et al. The effect of cylandelate in depressed and demented patients: A controlled study in psychogeriatric patients. *Age Ageing* 1977 Aug; 6(3):156-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davies G. Drugs for dementia. *Modern Geriatrics* 1978; 8(1):56

Status: Not included because not a full article

Davis KJ, Sloane PD, Mitchell CM, et al. Specialized dementia programs in residential care settings. *Gerontologist* 2000 Feb; 40(1):32-42.

Status: Not included because does not meet criteria for treatment for dementia patients

Davis KL, Mohs RC. Enhancement of memory processes in Alzheimer's disease with multiple-dose intravenous physostigmine. *Am J Psychiatry* 1982; 139(11):1421-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davis KL, Mohs RC, Davis BM, et al. Oral physostigmine in Alzheimer's disease. *Psychopharmacol Bull* 1983; 19(3):451-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Davis KL, Thal LJ, Gamzu ER, et al. A double-blind, placebo-controlled multicenter study of tacrine for Alzheimer's disease. *N Engl J Med* 1992; 327(18):1253-9.

Status: Not included because Jadad Quality Scale score less than three

Davis KL, Yang RK, Davidson M, et al. Alzheimer's disease: Tacrine and tacrine metabolite concentrations in plasma and cognitive change. *Drug Dev Res* 1995; 34(1):55-65.

Status: Not included because Jadad Quality Scale score less than three

Davis RN, Massman PJ, Doody RS. Cognitive intervention in Alzheimer disease: A randomized placebo-controlled study. *Alzheimer Dis Assoc Disord* 2001 Jan; 15(1):1-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Dawson P, Kontos P. Nursing assistants reduce aggressive behaviour during bathing cognitively impaired nursing home residents. *Perspectives (Montclair)* 1998; 22(2):20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dawson P. Bright light treatment for people with Alzheimer's disease. *Perspectives (Montclair)* 1999; 23(1):25-6.

Status: Not included because dementia population not randomized to treatment

Day JJ, Grant I, Atkinson JH, et al. Incidence of AIDS dementia in a two-year follow-up of AIDS and ARC patients on an initial phase II AZT placebo-controlled study: San Diego cohort. *J Neuropsychiatry Clin Neurosci* 1992; 4(1):15-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

de Boer JB, van Dam FS, Sprangers MA, et al. Longitudinal study on the Quality of Life of symptomatic HIV-infected patients in a trial of zidovudine versus zidovudine and interferon-alpha. *AIDS* 1993 Jul; 7(7):947-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

De Deyn PP, Scheltens P, Kittner B. A double-blind placebo-controlled trial assessing the effects of propentofylline in patient's with Alzheimer's disease and vascular dementia: safety, efficacy, and impact on disease progression. *Alzheimers*

Rep 1999; (2):51.

Status: Not included because not a full article

De Deyn PP, Rabheru K, Rasmussen A. Neuroleptic for behavioral symptoms of dementia. J Fam Pract 2000; 49(1):28-9.

Status: Not included because not a full article

DeLuca J, Johnson SK, Ellis SP, et al. Cognitive functioning is impaired in patients with chronic fatigue syndrome devoid of psychiatric disease. J Neurol Neurosurg Psychiatry 1997; 62(2):151-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Delwaide PJ, Hurlet A. Bromocriptine and buccolinguofacial dyskinesias in patients with senile dementia. A quantitative study. Arch Neurol 1980 Jul; 37(7):441-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Delwaide PJ, Devoitille JM, Ylief M. Acute effect of drugs upon memory of patients with senile dementia. Acta Psychiatr Belg 1980; 748-54.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Delwaide PJ, Gyselynck-Mambourg AM, Hurlet A, et al. Double-blind randomized controlled study of phosphatidylserine in senile demented patients. Acta Neurol Scand 1986 Feb; 73(2):136-40.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dencker SJ, Lindberg D. A controlled double blind study of piracetam in the treatment of senile dementia. Nord J Psychiatry 1977; (1):48-52.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Denney A. Quiet music. An intervention for mealtime agitation? J Gerontol Nurs 1997 Jul; 23(7):16-23.
Status: Not included because dementia population not randomized to treatment

Denolle T, Sassano P, Allain H, et al. Effects of nicardipine and clonidine on cognitive functions and electroencephalography in hypertensive patients. Fundam Clin Pharmacol 2002; 16(6):527-35.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Derouesne C, Renault B, Gueguen B, et al. Neuropsychophysiological evaluation of three doses of S 12024-2 in mild-to-moderate Alzheimer's disease. Clin Drug Investig 1997; (4):301-6.

Status: Cross-over trial;

Desai A, Grossberg G. Review of rivastigmine and its clinical applications in Alzheimer's disease and related disorders. Expert Opin Pharmacother 2001; 2(4):653-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

desRosiers G, Hodges JR, Berrios G. The neuropsychological differentiation of patients with very mild Alzheimer's disease and/or major depression. J Am Geriatr Soc 1995 Nov; 43(11):1256-63.

Status: Not included because does not meet criteria for treatment for dementia patients

Devanand DP, Sackeim HA, Brown RP, et al. A pilot study of haloperidol treatment of psychosis and behavioral disturbance in Alzheimer's disease. Arch Neurol 1989; 46(8):854-7.
Status: Not included because dementia population not randomized to treatment

Devanand DP, Cooper T, Sackeim HA, et al. Low dose oral haloperidol and blood levels in Alzheimer's disease: A preliminary study. Psychopharmacol Bull 1992; 28(2):169-73.
Status: Not included because Jadad Quality Scale score less than three

Devanand DP, Marder K, Michaels KS, et al. A randomized, placebo-controlled dose-comparison trial of haloperidol for psychosis and disruptive behaviors in Alzheimer's disease. Am J Psychiatry 1998 Nov; 155(11):1512-20.
Status: Cross-over trial;

di Bari M, Pahor M, Franse LV, et al. Dementia and disability outcomes in large hypertension trials: Lessons learned from the systolic hypertension in the elderly program (SHEP) trial. Am J Epidemiol 2001; 153(1):72-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Di Perri R, Coppola G, Ambrosio LA, et al. A multicentre trial to evaluate the efficacy and tolerability of alpha-glycerylphosphorylcholine versus cytosine diphosphocholine in patients with vascular dementia. J Int Med Res 1991 Jul;

19(4):330-41.

Status: Not included because Jadad Quality Scale score less than three

Dick MB, Nielson KA, Beth RE, et al. Acquisition and long-term retention of a fine motor skill in Alzheimer's disease. *Brain Cogn* 1995 Dec; 29(3):294-306.

Status: Not included because does not meet criteria for treatment for dementia patients

Dierks T, Maurer K, Ihl R. Influence of tenilsetam on AEP-P300 in Alzheimer's disease. *J Neural Transm Gen Sect* 1989; 1:49.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ditzler K. Efficacy and tolerability of memantine in patients with dementia syndrome. A double-blind, placebo controlled trial. *Arzneimittelforschung* 1991 Aug; 41(8):773-418.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Diwan S, Phillips VL. Agitation and dementia-related problem behaviors and case management in long-term care. *Int Psychogeriatr* 2001; 13(1):5-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Djernes JK, Gulmann NC, Abelskov KE, et al. Psychopathologic and functional outcome in the treatment of elderly inpatients with depressive disorders, dementia, delirium and psychoses. *Int Psychogeriatr* 1998 Mar; 10(1):71-83.

Status: Not included because does not meet criteria for treatment for dementia patients

Dominguez D, De CCL, Gomensoro J, et al. Modification of psychometric, practical and intellectual parameters in patients with diffuse cerebrovascular insufficiency during prolonged treatment with pentoxifylline: A double blind, placebo controlled trial. *Pharmatherapeutica* 1977; 1(8):498-506.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Doody RS, Dunn JK, Clark CM, et al. Chronic donepezil treatment is associated with slowed cognitive decline in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2001 Jul; 12(4):295-300.

Status: Not included because dementia population not randomized to treatment

Doraiswamy PM, Kaiser L, Bieber F, et al. The Alzheimer's Disease Assessment Scale: Evaluation of psychometric properties and patterns of cognitive decline in multicenter clinical trials of mild to moderate Alzheimer's disease. *Alzheimer Dis Assoc Disord* 2001 Oct; 15(4):174-83.

Status: Not included because does not meet criteria for treatment for dementia patients

Doraiswamy PM, Krishen A, Stallone F, et al. NSAIDs and cognition in Alzheimer's disease. *Neurology* 1996; 46(4):1194.

Status: Not included because not a full article

Droes RM, Breebaart E, Ettema TP, et al. Effect of integrated family support versus day care only on behavior and mood of patients with dementia. *Int Psychogeriatr* 2000; 12(1):99-115.

Status: Not included because does not meet criteria for treatment for dementia patients

Duffy FH, McAnulty G, Albert M, et al. Lecithin: Absence of neurophysiologic effect in Alzheimer's disease by EEG topography. *Neurology* 1987 Jun; 37(6):1015-9.

Status: Not included because no extractable data relevant to review

Dunn JC, Thiru-Chelvam B, Beck CH. Bathing. Pleasure or pain? *J Gerontol Nurs* 2002 Nov; 28(11):6-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Duret M, Goldman S, Messina D, et al. Effect of L-dopa on dementia-related rigidity. *Acta Neurol Scand* 1989 Jul; 80(1):64-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Durso R, Fedio P, Brouwers P. Lysine vasopressin in Alzheimer's disease. *Neurology* 1982; 32(6):674-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dykierek P, Stadtmuller G, Schramm P, et al. The value of REM sleep parameters in differentiating Alzheimer's disease from old-age depression and normal aging. *J Psychiatr Res* 1998 Jan; 32(1):1-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Dysken M, Kuskowski M, Love S, et al. Ondansetron in the treatment of cognitive decline in Alzheimer's dementia. *Am J Geriatr Psychiatry* 2002 Mar; 10(2):212-5.

Status: Not included because Jadad Quality Scale score less than three

Dysken MW, Anton JS, Klein L, et al. CI-911: A placebo-controlled study in patients with primary degenerative dementia. *Drug Dev Res* 1988; 12(3-4):267-4.

Status: Cross-over trial;

Dysken MW, Mendels J, Lewitt P, et al. Milacemide: a placebo-controlled study in senile dementia of the Alzheimer type. *J Am Geriatr Soc* 1992 May; 40(5):503-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Dysken MW, Johnson SB, Holden L, et al. Haloperidol concentrations in patients with Alzheimer's dementia. *Am J Geriatr Psychiatry* 1994; 2(2):124-33.

Status: Not included because Jadad Quality Scale score less than three

Eagger S, Levi I. Serum levels of tacrine in relation to clinical response in Alzheimer's disease. *Int J Geriatr Psychiatry* 1992a; 7(2):115-9.

Status: Cross-over trial;

Eagger S, Morant N, Levy R, et al. Tacrine in Alzheimer's disease. Time course of changes in cognitive function and practice effects. *Br J Psychiatry* 1992b; 160:36-40.

Status: Cross-over trial;

Eagger S. Searching for a treatment for Alzheimer's disease - tales from the cutting-room floor. *Int J Geriatr Psychiatry* 1996; 11(4):337-42.

Status: Not included because dementia population not randomized to treatment

Eagger SA, Levy R, Sahakian BJ. Tacrine in Alzheimer's disease. *Lancet* 1991 Apr 27; 337(8748):989-92.

Status: Cross-over trial;

Eagger SA, Levy R, Sahakian BJ. Tacrine in Alzheimer's disease. *Acta Neurol Scand Suppl* 1992; 139:75-80.

Status: Cross-over trial;

Eaton M, Mitchell-Bonair IL, Friedmann E. The effect of touch on nutritional intake of chronic organic brain syndrome patients. *J Gerontol* 1986 Sep; 41(5):611-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ebmeier KP, Hunter R, Curran SM, et al. Effects of a single dose of the acetylcholinesterase inhibitor velnacrine on recognition memory and regional cerebral blood flow in Alzheimer's disease. *Psychopharmacologia* 1992; 108(1-2):103-2.

Status: Cross-over trial;

Edberg A, Hallberg IR. Actions seen as demanding in patients with severe dementia during one year of intervention. Comparison with controls. *Int J Nurs Stud* 2001 Jun; 38(3):271-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Edberg AK, Hallberg IR. Effects of clinical supervision on nurse-patient cooperation quality: A controlled study in dementia care. *Clin Nurs Res* 1996 May; 5(2):127-46.

Status: Not included because does not meet criteria for treatment for dementia patients

Edberg AK, Norberg A, Hallberg I. Mood and general behavior of patients with severe dementia during one year of supervised, individualized planned care and systematic clinical supervision: Comparison with a similar control group. *Aging Clin Exp Res* 1999; (6):395-403.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Edwards NE, Beck AM. Animal-assisted therapy and nutrition in Alzheimer's disease. *West J Nurs Res* 2002 Oct; 24(6):697-712.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ellingrod VL, Schultz SK, Ekstam-Smith K, et al. Comparison of risperidone with olanzapine in elderly patients with dementia and psychosis. *Pharmacotherapy* 2002 Jan; 22(1):1-5.

Status: Not included because dementia population not randomized to treatment

Eloniemi-Sulkava U, Notkola IL, Hentinen M, et al. Effects of supporting community-living demented patients and their caregivers: A randomized trial. *J Am Geriatr Soc* 2001 Oct; 49(10):1282-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Engel RR, Satzger W, Gunther W, et al. Double-blind cross-over study of phosphatidylserine vs. placebo in patients with early dementia of the Alzheimer type. *Eur Neuropsychopharmacol* 1992 Jun; 2(2):149-55.

Status: Cross-over trial;

Engelberts NH, Klein M, Ader HJ, et al. The effectiveness of cognitive rehabilitation for attention deficits in focal seizures: A randomized controlled study. *Epilepsia* 2002; 43(6):587-95.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Etienne P, Dastoor D, Gauthier S. Alzheimer's disease: Lack of effect of lecithin treatment for 3 months. *Neurology* 1981; 31(12):1552-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

European Propentofylline Study Group. Propentofylline in dementia (vascular dementia and Alzheimer's disease). *Cardiovasc Dis* 1994; 4:258.
Status: Not included because not a full article

Evans M, Hammond M, Wilson K, et al. Treatment of depression in the elderly: Effect of physical illness on response. *Int J Geriatr Psychiatry* 1997 Dec; 12(12):1189-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Evans M, Ellis A, Watson D, et al. Sustained cognitive improvement following treatment of Alzheimer's disease with donepezil. *Int J Geriatr Psychiatry* 2000 Jan; 15(1):50-3.
Status: Not included because dementia population not randomized to treatment

Evers S, Grotemeyer KH, Reichelt D, et al. Impact of antiretroviral treatment on AIDS dementia: A longitudinal prospective event-related potential study. *J Acquir Immune Defic Syndr Hum Retrovirol* 1998 Feb 1; 17(2):143-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fakouhi TD, Jhee SS, Sramek JJ, et al. Evaluation of cycloserine in the treatment of Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1995 Oct; 8(4):226-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Falsaperla A, Monici Preti PA, Oliani C. Selegiline versus oxiracetam in patients with Alzheimer-type dementia. *Clin Ther* 1990 Sep; 12(5):376-84.
Status: Not included because Jadad Quality Scale score less than three

Farina E, Fioravanti R, Chiavari L, et al. Comparing two programs of cognitive training in Alzheimer's disease: A pilot study. *Acta Neurol Scand* 2002 May; 105(5):365-71.
Status: Not included because does not meet criteria for treatment for dementia patients

Farlow M, Gracon SI, Hershey LA, et al. A controlled trial of tacrine in Alzheimer's disease. The Tacrine Study Group. *JAMA* 1992 Nov 11; 268(18):2523-9.
Status: Not included because Jadad Quality Scale score less than three

Farlow M, Brashear A, Hiu S, et al. The effects of tacrine in patients with mild versus moderate stage Alzheimer's disease. In: Iqbal K, editors. *Research Advances in Alzheimer's disease and related disorders*, Chichester: John Wiley & Sons; 1995. p. 283-92.
Status: Not included because not a full article

Farlow MR, Lahiri DK, Poirier J, et al. Apolipoprotein E genotype and gender influence response to tacrine therapy. *Ann N Y Acad Sci* 1996; 802(Dec 16):101-10.
Status: Not included because Jadad Quality Scale score less than three

Farlow MR, Cyrus PA, Nadel A, et al. Metrifonate treatment of AD: Influence of APOE genotype. *Neurology* 1999 Dec 10; 53(9):2010-6.
Status: Not included because dementia population not randomized to treatment

Farlow MR, Cyrus PA. Metrifonate therapy in Alzheimer's disease: A pooled analysis of four randomized, double-blind, placebo-controlled trials. *Dement Geriatr Cogn Disord* 2000 Jul; 11(4):202-11.
Status: Not included because dementia population not randomized to treatment

Faxen-Irving G, Andren-Olsson B, af GA, et al. The effect of nutritional intervention in elderly subjects residing in group-living for the demented. *Eur J Clin Nutr* 2002 Mar; 56(3):221-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Feldman H, Gauthier S, Hecker J, et al. Benefits of Donepezil on global function, behavior, cognition and ADLs in patients with moderate to severe Alzheimer's disease. *Neurology* 2000; 54(Suppl 3):A469.

Status: Not included because not a full article

Feldman H, Sauter A, Donald A, et al. The disability assessment for dementia scale: A 12-month study of functional ability in mild to moderate severity Alzheimer's disease. *Alzheimer Dis Assoc Disord* 2001 Apr; 15(2):89-95.

Status: Not included because does not meet criteria for treatment for dementia patients

Fenn P, Gray A. Estimating long-term cost savings from treatment of Alzheimer's disease. A modelling approach. *Pharmacoeconomics* 1999 Aug; 16(2):165-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fernandez HH, Trieschmann ME, Burke MA, et al. Quetiapine for psychosis in Parkinson's disease versus dementia with Lewy bodies. *J Clin Psychiatry* 2002 Jun; 63(6):513-5.

Status: Not included because dementia population not randomized to treatment

Ferris SH, Sathananthan G, Gershon S, et al. Cognitive effects of ACTH 4-10 in the elderly. *Pharmacol Biochem Behav* 1976; 5(Suppl 1):73-8. *Status: Not included because dementia population not defined by DSM, NINCDS or ICD*

Ferris SH, Mittelman MS. Behavioral treatment of Alzheimer's disease. *Int Psychogeriatr* 1996; 8(Suppl 1):87-90.

Status: Not included because no extractable data relevant to review

Fichter MM, Bruce ML, Schroppel H, et al. Cognitive impairment and depression in the oldest old in a German and in U.S. communities. *Eur Arch Psychiatry Clin Neurosci* 1995; 245(6):319-25.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Filip V, David I, Klatschka J, et al. Selegiline slows down the decline of cognitive and electrophysiological functions in Alzheimer's

disease. *Basic and clinical science of mental and additive disorders. Bibl Psychiatr* 1997; 238-40.

Status: Not included because not a full article

Finali G, Piccirilli M, Oliani C, et al. L-deprenyl therapy improves verbal memory in amnesic Alzheimer patients. *Clin Neuropharmacol* 1991 Dec; 14(6):523-36.

Status: Cross-over trial;

Finali G, Piccirilli M, Oliani C, et al. Alzheimer-type dementia and verbal memory performances: Influence of selegiline therapy. *Ital J Neurol Sci* 1992 Mar; 13(2):141-8.

Status: Cross-over trial;

Findlay DJ, Sharma J, McEwen J, et al. Double-blind controlled withdrawal of thioridazine treatment in elderly female inpatients with senile dementia. *Int J Geriatr Psychiatry* 1989; 4(2):115-20.

Status: Not included because Jadad Quality Scale score less than three

Finkel SI, Lyons JS, Anderson RL, et al. A randomized, placebo-controlled trial of thiothixene in agitated, demented nursing home patients. *Int J Geriatr Psychiatry* 1995; 10(2):129-36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Finkel SI, Lyons J. Nursing home research from investigators' perspective. *Int Psychogeriatr* 1996; 8(Suppl 3):371-3, 381-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fioravanti M, Bergamasco B, Bocola V, et al. A multicentre, double-blind, controlled study of piracetam vs placebo in geriatric patients with nonvascular mild-moderate impairment in cognition. *New Trends in Clinical Neuropharmacology* 1991; (1):27-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fioravanti M, Di Cesare F. Memory improvements and pharmacological treatment: A method to distinguish direct effects on memory from secondary effects due to attention improvement. *Int Psychogeriatr* 1992; 4(1):119-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fioravanti M, Ferrario E, Massaia M, et al. Low folate levels in the cognitive decline of elderly

patients and the efficacy of folate as a treatment for improving memory deficits. Arch Gerontol Geriatr 1997; 26(1):1-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fischer, Gotz P. Blood transferrin and ferritin in Alzheimer's disease. Life Sci 1997; 60(25):2273-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Fischhof PK, Saletu B, Ruther E, et al. Therapeutic efficacy of pyritinol in patients with senile dementia of the Alzheimer type (SDAT) and multi-infarct dementia (MID). Neuropsychobiology 1992; 26(1-2):65-2.

Status: Not included because Jadad Quality Scale score less than three

Fischhof PK. Divergent neuroprotective effects of nimodipine in PDD and MID provide indirect evidence of disturbances in Ca²⁺ homeostasis in dementia. Methods & Findings in Experimental & Clinical Pharmacology 1993 Oct; 15(8):549-55.
Status: Not included because Jadad Quality Scale score less than three

Fisman M, Merksey H, Helmes E. Double blind study of lecithin in patients with Alzheimer's disease. Can J Psychiatry 1981; (6):426-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fisman M, Merskey H, Helmes E. Double-blind trial of 2-dimethylaminoethanol in Alzheimer's disease. Am J Psychiatry 1981; 138(7):970-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fitten LJ, Perryman KM, Gross PL, et al. Treatment of Alzheimer's disease with short- and long-term oral THA and lecithin: A double-blind study. Am J Psychiatry 1990 Feb; 147(2):239-42.
Status: Cross-over trial;

Fitten LJ, Ganzell S. Spouses' assessments of Alzheimer patients' response to THA and lecithin. Am J Psychiatry 1992; 149(4):575.
Status: Not included because not a full article

Fitzsimmons S, Buettner LL. Therapeutic recreation interventions for need-driven dementia-compromised behaviors in community-dwelling elders. Am J Alzheimers Dis Other Dement 2002; 17(6):367-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fleischhacker WW, Buchgeher A, Schubert H. Memantine in the treatment of senile dementia of the Alzheimer type. Prog Neuropsychopharmacol Biol Psychiatry 1986; 10(1):87-93.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Flicker C, Ferris SH, Kalkstein D, et al. A double-blind, placebo-controlled crossover study of ganglioside GM1 treatment for Alzheimer's disease. Am J Psychiatry 1994 Jan; 151(1):126-9.
Status: Cross-over trial;

Flicker L, Grimley EG. Piracetam for dementia or cognitive impairment. In: The Cochrane Library, 2001. Issue 2. Oxford: Update Software.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Folnegovic Smalc V, Knezevic S, Bokonic R, et al. European pentoxifylline multi-infarct dementia trial: The epmid study. J Neurol 1994; 2(41):158.
Status: Not included because not a full article

Fontana RJ, Turgeon DK, Woolf TF, et al. The caffeine breath test does not identify patients susceptible to tacrine hepatotoxicity. Hepatology 1996 Jun; 23(6):1429-35.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fontana RJ, deVries TM, Woolf TF, et al. Caffeine based measures of CYP1A2 activity correlate with oral clearance of tacrine in patients with Alzheimer's disease. Br J Clin Pharmacol 1998 Sep; 46(3):221-8.
Status: Cross-over trial;

Food and Drug Administration. Tacrine as a treatment for Alzheimer's disease: Editor's note. An interim report from the FDA. A response from Summers et al. N Engl J Med 1991 Jan 31; 324(5):349-52.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ford JM, Truman CA, Wilcock GK, et al. Serum concentrations of tacrine hydrochloride predict its adverse effects in Alzheimer's disease. Clin Pharmacol Ther 1993; 53(6):691-5.
Status: Not included because dementia population not randomized to treatment

Forette F, Amery A, Staessen J, et al. Is prevention of vascular dementia possible? The Syst-Eur Vascular Dementia Project. Aging (Milano) 1991; 3(4):373-82.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Forette F, Hoover T, Gracon S, et al. A double-blind, placebo-controlled, enriched population study of tacrine in patients with Alzheimer's disease. Eur J Neurol 1995; 2:229-38.

Status: Not included because Jadad Quality Scale score less than three

Forette F, Seux ML, Staessen JA, et al. Prevention of dementia in randomised double-blind placebo-controlled Systolic Hypertension in Europe (Syst-Eur) trial. Lancet 1998 Oct 24; 352(9137):1347-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Forssell LG, Sjkovist B, Winblad B. Early stages of late onset Alzheimer's disease. III. Double blind treatment with choline chloride and lecithin with and without L-dopa and L-tryptophan, alternatively placebo. Acta Neurol Scand Suppl 1989; 79(121):43-66.

Status: Not included because dementia population not randomized to treatment

Forster DP, Newens AJ, Kay DW, et al. Risk factors in clinically diagnosed presenile dementia of the Alzheimer type: A case-control study in northern England. J Epidemiol Commun Health 1995 Jun; 49(3):253-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Foster HG, Hillbrand M, Chi CC. Efficacy of carbamazepine in assaultive patients with frontal lobe dysfunction. Prog Neuropsychopharmacol Biol Psychiatry 1989; 13(6):865

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Foster NL, Aldrich MS, Bluemlein L, et al. Failure of cholinergic agonist RS-86 to improve cognition and movement in PSP despite effects on sleep. Neurology 1989 Feb; 39(2 Pt 1):257-61.

Status: Cross-over trial;

Foster NL, Petersen RC, Gracon SI, et al. An enriched-population, double-blind, placebo-controlled, crossover study of tacrine and lecithin

in Alzheimer's disease. Dementia 1996; 7(5):260-6.

Status : Cross-over trial;

Foster NA, Valentine ER. Effect of auditory stimulation on autobiographical recall in dementia. Exp Aging Res 2001 Jul; (3):215-28.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fovall P, Dysken MW, Lazarus LW. Choline bitartrate treatment of Alzheimer-type dementias. Commun Psychopharmacol 1980; 4(2):141-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Francesse T. Research corner. The effects of regular exercise on muscle strength and functional abilities of late stage Alzheimer's residents. Va Nurse Today 1995; (4):25-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Francesse T, Sorrell J, Butler FR. The effects of regular exercise on muscle strength and functional abilities of late stage Alzheimer's residents. Am J Alzheimers Dis 1997; (3):122-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Franciosi A, Zavattini G. Dihydroergocristine in the treatment of elderly patients with cognitive deterioration: A double-blind, placebo-controlled, dose- response study. Curr Ther Res Clin Exp 1994; 55(11):1391-401.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Frattola L, Piolti R, Bassi S, et al. Multicenter clinical comparison of the effects of choline alfoscerate and cytidine diphosphocholine in the treatment of multi-infarct dementia. Curr Ther Res Clin Exp 1991; 49(4):683-93.

Status: Not included because Jadad Quality Scale score less than three

Frattola L, Trabucchi H, Cucinotta D, et al. Dopamine agonist and free-radicals scavenger activities of dihydroergokryptine in dementia of alzheimer type (DAT): Multicentre, long-term double-blind clinical study versus placebo. J Neurol 1994; 241:159.

Status: Not included because not a full article

Frederick B, Satlin A, Wald LL, et al. Brain proton magnetic resonance spectroscopy in Alzheimer's

disease: Changes after treatment with xanomeline. *Am J Geriatr Psychiatry* 2002 Jan; 10(1):81-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fric M, Horn R, Hasse SI, et al. Effects of nimodipine-treatment in primary degenerative dementia. Results of a clinical and psychometric study. *Pharmacopsychiatry* 1995; 28:177.

Status: Not included because not a full article

Friedman JI, Adler DN, Howanitz E, et al. Erratum: A double blind placebo controlled trial of donepezil adjunctive treatment to risperidone for the cognitive impairment of schizophrenia. *Biol Psychiatry* 2002; 51(12):1014.

Status: Not included because not a full article

Friedman JI, Adler DN, Howanitz E, et al. A double blind placebo controlled trial of donepezil adjunctive treatment to risperidone for the cognitive impairment of schizophrenia. *Biol Psychiatry* 2002 Mar 1; 51(5):349-57.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Friedman R, Tappen RM. The effect of planned walking on communication in Alzheimer's disease. *J Am Geriatr Soc* 1991 Jul; 39(7):650-4.

Status: Not included because does not meet criteria for treatment for dementia patients

Frisoni GB, Gozzetti A, Bignamini V, et al. Special care units for dementia in nursing homes: A controlled study of effectiveness. *Arch Gerontol Geriatr* 1998; 27(Suppl 6):215-24.

Status: Not included because dementia population not randomized to treatment

Frith CD, Stevens M, Johnstone EC, et al. Effects of ECT and depression on various aspects of memory. *Br J Psychiatry* 1983; Vol 142(Jun):610-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fuglum E, Schillinger A, Andersen JB, et al. Zuclopenthixol and haloperidol/levomepromazine in the treatment of elderly patients with symptoms of aggressiveness and agitation: A double-blind, multi-centre study. *Pharmatherapeutica* 1989; 5(5):285-91.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fulop TJ, Worum I, Csongor J, et al. Effects of centrophenoxine on body composition and some biochemical parameters of demented elderly people as revealed in a double-blind clinical trial. *Arch Gerontol Geriatr* 1990; 10(3):239-51.

Status: Not included because no extractable data relevant to review

Funfgeld EW, Baggen M, Nedwidek P, et al. Double blind study with phosphatidylserine (PS) in parkinsonian patients with senile dementia of Alzheimer's type (SDAT). *Alzheimers Dis Relat Disord* 1989; 1235-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Fuschillo C, La Pia S, Campana F, et al. Cognitive deficits in Alzheimer's disease: Treatment with acetylcholinesterase inhibitor agents. *Arch Gerontol Geriatr* 2001; 33(suppl 1):151-8.

Status: Not included because Jadad Quality Scale score less than three

Fusgen I, Bressel H-U, De Mey C. A randomised, placebo-controlled, double-blind study of the efficacy and tolerability of dimenhydrinate in multi-morbid patients with senile dizziness. *Eur J Geriatr* 2002; 4(2):92-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gaber S, Ronzoni S, Bruni A, et al. Sertaline versus small doses of haloperidol in the treatment of agitated behaviour in patients with dementia. *Arch Gerontol Geriatr Psychiatry* 2001; (Suppl 1):159-62.

Status: Not included because Jadad Quality Scale score less than three

Gabrynowicz JW, Dumbrell M. A clinical trial of leptazole with nicotinic acid in the management of psycho-geriatric patients. *Med J Aust* 1968 May 11; 1(19):799-802.

Status: Not included because no extractable data relevant to review

Gainotti G, Nocentini U, Sena E. Can the pattern of neuropsychological improvement obtained with cholinergic drugs be used to infer a cholinergic mechanism in other nootropic drugs? *Prog Neuropsychopharmacol Biol Psychiatry* 1989; 13(Suppl):S47-S59

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gaitz CM, Varner RV, Overall JE. Pharmacotherapy for organic brain syndrome in late life. Evaluation of an ergot derivative vs placebo. Arch Gen Psychiatry 1977 Jul; 34(7):839-45.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gallai V, Mazzotta G, Firenze C, et al. Study of the P300 and cerebral maps in subjects with multi-infarct dementia treated with cytidine. Psychopharmacologia 1991; 103(1):1-5.

Status: Not included because no extractable data relevant to review

Gallai V, Mazzotta G, Del Gatto F, et al. A clinical and neurophysiological trial on nootropic drugs in patients with mental decline. Acta Neurol (Napoli) 1991 Feb; 13(1):1-12.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gao H, Yan L, Liu B, et al. Clinical study on treatment of senile vascular dementia by acupuncture. J Tradit Chin Med 2001 Jun; 21(2):103-9.

Status: Not included because Jadad Quality Scale score less than three

Garetz FK, Baron JJ, Barron PB, et al. Efficacy of nylidrin hydrochloride in the treatment of cognitive impairment in the elderly. J Am Geriatr Soc 1979 May; 27(5):235-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gasnault J, Gueguen B, Bourdel MC, et al. Oral tacrine effects on computerised EEG activity during a double blind cross-over study in dementia of the Alzheimer type. J Neurol 1990; 237:33.

Status: Not included because not a full article

Gauthier S, Leblanc R, Robitaille Y, et al. Transmitter-replacement therapy in Alzheimer's disease using intracerebroventricular infusions of receptor agonists. Can J Neurol Sci 1986 Nov; 13(4 Suppl):394-402.

Status: Not included because dementia population not randomized to treatment

Gauthier S, Bouchard R, Lamontagne A, et al. Tetrahydroaminoacridine-lecithin combination treatment in patients with intermediate-stage Alzheimer's disease. N Engl J Med 1990;

322(18):1272-6.

Status: Cross-over trial;

Gedye JL, Exton-Smith AN, Wedgwood J. A method for measuring mental performance in the elderly and its use in a pilot clinical trial of meclofenoxate in organic dementia (preliminary communication). Age Ageing 1972 May; 1(2):74-80.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gedye JL, Ibrahim GS, McDonald C. A double blind controlled trial of piracetam (2-pyrrolidone acetamide) on two groups of psychogeriatric patients. IRCS Med Sci Clin Med 1978; 6(5):202.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gelinas I, Gauthier S, Cyrus PA. Metrifonate enhances the ability of Alzheimer's disease patients to initiate, organize, and execute instrumental and basic activities of daily living. J Geriatr Psychiatry Neurol 2000; 13(1):9-16.

Status: Not included because dementia population not randomized to treatment

Geng J. Treatment of 50 cases of senile dementia by acupuncture combined with inhalation of herbal drugs and oxygen. J Tradit Chin Med 1999 Dec; 19(4):287-9.

Status: Not included because Jadad Quality Scale score less than three

George TP, Vessicchio JC, Termine A, et al. Effects of smoking abstinence on visuospatial working memory function in schizophrenia. Neuropsychopharmacology 2002; 26(1):75-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gerber GJ, Prince PN, Snider HG, et al. Group activity and cognitive improvement among patients with Alzheimer's disease. Hospital & Community Psychiatry 1991; 42(8):843-5.

Status: Not included because does not meet criteria for treatment for dementia patients

Gerdner LA. Effects of individualized versus classical "relaxation" music on the frequency of agitation in elderly persons with Alzheimer's disease and related disorders. Int Psychogeriatr 2000 Mar; 12(1):49-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gerdner LA, Buckwalter KC, Reed D. Impact of a psychoeducational intervention on caregiver response to behavioral problems. *Nurs Res* 2002 Nov; 51(6):363-74.

Status: Not included because does not meet criteria for treatment for dementia patients

Gessner B, Voelp A, Klasser M. Study of the long-term action of a Ginkgo biloba extract on vigilance and mental performance as determined by means of quantitative pharmaco-EEG and psychometric measurements. *Arzneimittelforschung* 1985; 35(9):1459-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Getsios D, Caro JJ, Caro G, et al. Assessment of health economics in Alzheimer's disease (AHEAD): Galantamine treatment in Canada. *Neurology* 2001 Sep 25; 57(6):972-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ghatak R. Effects of an intervention program on dementia patients and their caregivers. *Caring* 1994 Aug; (No. 8):34-9.

Status: Not included because dementia population not randomized to treatment

Ghose K. Plasma drug levels of oxpentifylline in patients with dementia. An assessment of compliance. *Brain Dysfunct* 1989; (2):105-10.

Status: Not included because no extractable data relevant to review

Giacobini E, Spiegel R, Enz A, et al. Inhibition of acetyl- and butyryl-cholinesterase in the cerebrospinal fluid of patients with Alzheimer's disease by rivastigmine: Correlation with cognitive benefit. *J Neural Transm Gen Sect* 2002 Jul; 109(7-8):1053-65.

Status: Not included because dementia population not randomized to treatment

Ginsburg R, Weintraub M. Caffeine in the "sundown syndrome." Report of negative results. *J Gerontol* 1976 Jul; 31(4):419-20.

Status: Not included because dementia population not randomized to treatment

Gitlin LN, Corcoran M, Winter L, et al. A randomized, controlled trial of a home environmental intervention: Effect on efficacy and upset in caregivers and on daily function of persons with dementia. *Gerontologist* 2001 Feb; 41(1):4-14.

Status: Not included because does not meet criteria for treatment for dementia patients

Giuffra M, Mouradian MM, Bammert J, et al. Prolonged intravenous infusion of physostigmine in Alzheimer's disease. *Neurology* 1990; 40(Suppl 1):229, 1990.

Status: Not included because not a full article

Gleason RP, Schneider LS. Carbamazepine treatment of agitation in Alzheimer's outpatients refractory to neuroleptics. *J Clin Psychiatry* 1990; 51(3):115-8.

Status: Not included because dementia population not randomized to treatment

Goad DL, Davis CM, Liem P, et al. The use of selegiline in Alzheimer's patients with behavior problems. *J Clin Psychiatry* 1991; 52(8):342-5.

Status: Not included because dementia population not randomized to treatment

Gobburu JV, Tammara V, Lesko L, et al. Pharmacokinetic-pharmacodynamic modeling of rivastigmine, a cholinesterase inhibitor, in patients with Alzheimer's disease. *J Clin Pharmacol* 2001 Oct; 41(10):1082-90.

Status: Not included because dementia population not randomized to treatment

Goety CG, Tanner CM, Cohen JA, et al. L-acetylcarnitine in Huntington's disease: Double-blind placebo controlled crossover study of drug effects on movement disorder and dementia. *Mov Disord* 1990; 5(3):263-5.

Status: Not included because not a full article

Goldstein SE, Birnbom F. Nylidrin HCL in the treatment of symptoms of the aged: A double-blind placebo controlled study. *J Clin Psychiatry* 1979 Dec; 40(12):520-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Goldwasser AN, Auerbach SM, Harkins SW. Cognitive, affective, and behavioral effects of reminiscence group therapy on demented elderly. *Int J Aging Hum Dev* 1987; 25(3):209-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Golomb BA. Statins and dementia. *Arch Neurol* 2001; 58(7):1169-70.

Status: Not included because not a full article

Goodwin GM, Conway SC, Peyro-Saint-Paul H, et al. Executive function and uptake of 99mTc-exametazime shown by single photon emission tomography after oral idazoxan in probable Alzheimer-type dementia. *Psychopharmacologia* 1997 Jun; 131(4):371-8.

Status: Cross-over trial;

Gori G, Pientini S, Vespa A. The selection of meaningful activities as a treatment for day-care in dementia. *Arch Gerontol Geriatr* 2001; 33(Suppl):207-12.

Status: Not included because dementia population not randomized to treatment

Gorman DG, Read S, Cummings JL. Cholinergic therapy of behavioral disturbances in Alzheimer's disease. *Neuropsychiatry Neuropsychol Behav Neurol* ; 6(4):229-34.

Status: Cross-over trial;

Gormley N, Lyons D, Howard R. Behavioural management of aggression in dementia: A randomized controlled trial. *Age Ageing* 2001 Mar; 30(2):141-5.

Status: Not included because does not meet criteria for treatment for dementia patients

Gortelmeyer R, Erbler H. Memantine in the treatment of mild to moderate dementia syndrome. A double-blind placebo-controlled study. *Arzneimittelforschung* 1992 Jul; 42(7):904-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gotestam KG, Ljunghall S, Olsson B. A double-blind comparison of the effects of haloperidol and cis-clophenxol in senile dementia. *Acta Psychiatr Scand* 1981; 294(Suppl):46-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gotestam KG, Melin L. The effect of prompting and reinforcement of activity in elderly demented inpatients. *Scand J Psychol* 1990; 31(1):2-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gottfries CG, Karlsson I, Nyth AL. Treatment of depression in elderly patients with and without dementia disorders. *Int Clin Psychopharmacol* 1992; 6(Suppl 5):55-64.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gracon SI, Hoover TM, Lewis KW, et al. Tacrine in Alzheimer's disease efficacy and safety in a parallel group study. *Alzheimer's Disease Advances in Clinical and Basic Research* 1993; 549-57.

Status : Not included because not a full article

Gracon SI, Knapp MJ, Berghoff WG, et al. Safety of tacrine: Clinical trials, treatment IND, and postmarketing experience. *Alzheimer Dis Assoc Disord* 1998 Jun; 12(2):93-101.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Graf A, Wallner C, Schubert V, et al. The effects of light therapy on mini-mental state examination scores in demented patients. *Biol Psychiatry* 2001 Nov 1; 50(9):725-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grasser A, Gotthardt U, Heuser-Link M, et al. Behavioral effects of an ACTH4-9-fragment in a mixed sample of depressed patients and patients with Alzheimer's disease. *Pharmacopsychiatry* 1992; Vol 25:102

Status: Not included because not a full article

Green J, McDonald WM, Vitek JL, et al. Neuropsychological and psychiatric sequelae of pallidotomy for PD: Clinical trial findings. *Neurology* 2002; 58(6):858-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Green MF, Marder SR, Glynn SM, et al. The neurocognitive effects of low-dose haloperidol: A two-year comparison with risperidone. *Biol Psychiatry* 2002; 51(12):972-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Green RC, Goldstein FC, Auchus AP, et al. Treatment trial of oxiracetam in Alzheimer's disease. *Arch Neurol* 1992 Nov; 49(11):1135-6.

Status: Not included because dementia population not randomized to treatment

Greenberg SM, Tennis MK, Brown LB, et al. Donepezil therapy in clinical practice: A randomized crossover study. *Arch Neurol* 2000 Jan; 57(1):94-9.

Status: Cross-over trial;

Greendyke RM, Kanter DR. Therapeutic effects of pindolol on behavioral disturbances associated

with organic brain disease: A double-blind study. *J Clin Psychiatry* 1986 Aug; 47(8):423-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Greendyke RM, Kanter DR, Schuster DB, et al. Propranolol treatment of assaultive patients with organic brain disease. A double-blind crossover, placebo-controlled study. *J Nerv Ment Dis* 1986 May; 174(5):290-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Greendyke RM, Berkner JP, Webster JC, et al. Treatment of behavioral problems with pindolol. *Psychosomatics* 1989; 30(2):161-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Greer KL, Pustay KA, Zaun TC, et al. Comparison of the effects of toys versus live animals on the communication of patients with dementia of the Alzheimer's type. *Clin Gerontol* 2001; (3-4):157-82.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grioli S, Lomeo C, Quattropiani MC, et al. Pyroglutamic acid improves the age associated memory impairment. *Fundam Clin Pharmacol* 1990; 4(2):169-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gross RJ, Eisdorfer CE, Schiller HS, et al. Effect of ergot alkaloids on serum prolactin in non-psychotic organic brain syndrome of the elderly. *Exp Aging Res* 1979 Aug; 5(4):293-302.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grossberg GT, Stahelin HB, Messina JC, et al. Lack of adverse pharmacodynamic drug interactions with rivastigmine and twenty-two classes of medications. *Int J Geriatr Psychiatry* 2000 Mar; 15(3):242-7.
Status: Not included because dementia population not randomized to treatment

Grossman M, Mickanin J, Onishi K, et al. An aspect of sentence processing in Alzheimer's disease: Quantifier-noun agreement. *Neurology* 1995 Jan; 45(1):85-91.
Status: Not included because does not meet criteria for treatment for dementia patients

Grossmann WM, Standl A, May U, et al. Naftidrofuryl in the treatment of mild senile dementia. A double-blind study. *Pharmacopsychiatry* 1990 Nov; 23(6):265-73.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Grothe DR, Piscitelli SC, Dukoff R, et al. Penetration of tacrine into cerebrospinal fluid in patients with Alzheimer's disease. *J Clin Psychopharmacol* 1998; 18(1):78-81.
Status: Not included because dementia population not randomized to treatment

Group for the Advancement of Psychiatry and Committee on Aging. Impact of tacrine in the care of patients with Alzheimer's disease: What we know one year after FDA approval. *Am J Geriatr Psychiatry* 1994; 2(4):285-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Growdon JH, Corkin S, Huff FJ, et al. Piracetam combined with lecithin in the treatment of Alzheimer's disease. *Neurobiol Aging* 1986 Jul; 7(4):269-76.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Guerzoni A, Santambrogio S. Efficacy of dihydroergocristine 20 mg once daily in patients with organic brain psychosyndrome. A 3-month randomised, double-blind, placebo-controlled study. *Clin Drug Investig* 1995; 10(1):1-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Guimon J, Blanco J, Caso C. L-Dopa carbidopa treatment of senile dementia: A control study. *Eur J Psychiatry* 1995; (1):29-36.
Status: Not included because Jadad Quality Scale score less than three

Gustafson L, Risberg J, Johanson M, et al. Effects of piracetam on regional cerebral blood flow and mental functions in patients with organic dementia. *Psychopharmacologia* 1978 Mar 1; 56(2):115-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gustafson L, Edvinsson L, Dahlgren N, et al. Intravenous physostigmine treatment of Alzheimer's disease evaluated by psychometric testing, regional cerebral blood flow (rCBF) measurement, and EEG. *Psychopharmacologia*

1987; 93(1):31-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Gustafson L. Physostigmine and tetrahydroaminoacridine treatment of Alzheimer's disease. *Acta Neurol Scand Suppl* 1993; 149(Rand):39-41.

Status: Cross-over trial;

Haaland KY, Harrington DL, O'Brien S, et al. Cognitive-motor learning in Parkinson's disease. *Neuropsychology* 1997; 11(2):180-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hackman BW, Galbraith. Replacement therapy with piperazine oestrone sulphate (harmogen) and its effect on memory. *Curr Med Res Opin* 1976; 4(4):303-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hackman BW, Galbraith. Six-month pilot study of oestrogen replacement therapy and piperazine oestrone sulphate ('Harmogen') and its effect on memory. *Curr Med Res Opin* 1977; 4(3):21-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hadas-Lidor N, Katz N, Tyano S, et al. Effectiveness of dynamic cognitive intervention in rehabilitation of clients with schizophrenia. *Clin Rehabil* 2001; 15(4):349-59.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Haffmans PM, Sival RC, Lucius SA, et al. Bright light therapy and melatonin in motor restless behaviour in dementia: A placebo-controlled study. *Int J Geriatr Psychiatry* 2001 Jan; 16(1):106-10.

Status: Cross-over trial;

Hagstadius S, Gustafson L, Risberg J. The effects of bromvincamine and vincamine on regional cerebral blood flow and mental functions in patients with multi-infarct dementia. *Psychopharmacologia* 1984; 83(4):321-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Haley WE, Brown SL, Levine EG. Experimental evaluation of the effectiveness of group intervention for dementia caregivers. *Gerontologist* 1987; 27(3):376-82.

Status: Not included because does not meet criteria for treatment for dementia patients

Hall P, Harcup M. A trial of lipotropic enzymes in atheromatous ("arteriosclerotic") dementia. *Angiology* 1969 May; 20(5):287-300.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Halliday GM, Shepherd CE, McCann H, et al. Effect of anti-inflammatory medications on neuropathological findings in Alzheimer's disease. *Arch Neurol* 2000 Jun; 57(6):831-6.

Status: Not included because dementia population not randomized to treatment

Hammeke TA, Haughton VM, Grogan JP, et al. A preliminary study of cognitive and affective alterations following intrathecal administration of iopamidol or metrizamide. *Invest Radiol* 1984; 19(Suppl 5):S268-S271.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hammen CL, Jacobs M, Mayol A, et al. Dysfunctional cognitions and the effectiveness of skills and cognitive-behavioral assertion training. *J Consult Clin Psychol* 1980 Dec; 48(6):685-95.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hanley IG, McGuire RJ, Boyd WD. Reality orientation and dementia: A controlled trial of two approaches. *Br J Psychiatry* 1981; 138:10-4.

Status: Not included because dementia population not randomized to treatment

Hansson L. Antihypertensive treatment and the prevention of dementia: Further insights from the Syst-Eur trial. *J Hypertens* 1999; 17(3):307-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hanyu H, Tanaka Y, Sakurai H, et al. Atrophy of the substantia innominata on magnetic resonance imaging and response to donepezil treatment in Alzheimer's disease. *Neurosci Lett* 2002 Feb 8; 319(1):33-6.

Status: Not included because dementia population not randomized to treatment

Harbaugh RE, Roberts DW, Coombs DW, et al. Preliminary report: Intracranial cholinergic drug infusion in patients with Alzheimer's disease. *Neurosurgery* 1984 Oct; 15(4):514-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harbaugh RE. Intracerebroventricular cholinergic drug administration in Alzheimer's disease: Preliminary results of a double-blind study. *J Neural Transm Suppl* 1987; 24:271-7.
Status: Cross-over trial;

Harbaugh RE, Reeder TM, Senter HJ, et al. Intracerebroventricular bethanechol chloride infusion in Alzheimer's disease. Results of a collaborative double-blind study. *J Neurosurg* 1989 Oct; 71(4):481-6.
Status: Cross-over trial;

Harding GF, Hall P, Young J, et al. Multifocal infarct dementia treated by cyclandelate and monitored by quantitative EEG. *Angiology* 1978 Feb; 29(2):139-40.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harenko A. A comparison between chlormethiazole and nitrazepam as hypnotics in psycho-geriatric patients. *Curr Med Res Opin* 1974 Jul; 2(10):657-63.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harkins SW, Taylor JR, Mattay VS. Response to tacrine in patients with dementia of the Alzheimer's type: cerebral perfusion change is related to change in mental status. *Int J Neurosci* 1996 Feb; 84(1-4):149-4.
Status: Not included because dementia population not randomized to treatment

Harrell LE, Callaway R, Morere D, et al. The effect of long-term physostigmine administration in Alzheimer's disease. *Neurology* 1990; 40(9):1350-4.
Status: Cross-over trial;

Harrell LE, Jope RS, Falgout J, et al. Biological and neuropsychological characterization of physostigmine responders and nonresponders in Alzheimer's disease. *J Am Geriatr Soc* 1990 Feb; 38(2):113-22.
Status: Cross-over trial;

Harrell TH, Ryon NB. Cognitive-behavioral assessment of depression: Clinical validation of the automatic thoughts questionnaire. *J Consult Clin Psychol* 1983 Oct; 51(5):721-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harris S, Dowson JH. The effects of Meclofenoxate on cognitive performance in elderly individuals with memory impairment: A placebo-controlled study. *Int J Geriatr Psychiatry* 1986; 1:93-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hart BD, Wells DL. The effects of language used by caregivers on agitation in residents with dementia. *Clin Nurse Spec* 1997 Jan; 11(1):20-3.
Status: Not included because dementia population not randomized to treatment

Hart S, Smith CM, Swash M, et al. Word fluency in patients with early dementia of Alzheimer's type. *Br J Clin Psychol* 1988; 46(Pt 2):115-24, 1592-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hartmann A. Cerebral blood flow in patients with cerebrovascular disorders: study with pentoxifylline. *Ric Clin Lab* 1981; 11(Suppl 1):243-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hartmann A. Comparative randomized study of cerebral blood flow after long-term administration of pentoxifylline and co-dergocrine mesylate in patients with chronic cerebrovascular disease. *Curr Med Res Opin* 1985; 9(7):475-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hartmann A, Tsuda Y. A controlled study on the effect of pentoxifylline and an ergot alkaloid derivative on regional cerebral blood flow in patients with chronic cerebrovascular disease. *Angiology* 1988 May; 39(5):449-57.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harvey PD, Moriarty PJ, Serper MR, et al. Practice-related improvement in information processing with novel antipsychotic treatment. *Schizophr Res* 2000; 46(2-3):139-48.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Harwart D. The treatment of chronic cerebrovascular insufficiency. A double-blind

study with pentoxifylline ('Trental' 400). *Curr Med Res Opin* 1979; 6(2):73-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hauber AB, Gnanasakthy A, Mauskopf JA. Savings in the cost of caring for patients with Alzheimer's disease in Canada: An analysis of treatment with rivastigmine. *Clin Ther* 2000 Apr; 22(4):439-51.

Status: Not included because does not meet criteria for treatment for dementia patients

Hauer K, Marburger C, Oster P. Motor performance deteriorates with simultaneously performed cognitive tasks in geriatric patients. *Arch Phys Med Rehabil* 2002 Feb; 83(2):217-23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Haupt M, Karger A, Janner M. Improvement of agitation and anxiety in demented patients after psychoeducative group intervention with their caregivers. *Int J Geriatr Psychiatry* 2000 Dec; 15(12):1125-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Hebert R, Leclerc G, Bravo G, et al. Efficacy of a support group programme for care-givers of demented patients in the community: A randomized controlled trial. *Arch Gerontol Geriatr* 1994; 18(1):1-14.

Status: Not included because does not meet criteria for treatment for dementia patients

Hebert R, Girouard D, Leclerc G, et al. The impact of a support group programme for care-givers on the institutionalisation of demented patients. *Arch Gerontol Geriatr* 1995; 20(2):129-34.

Status: Not included because does not meet criteria for treatment for dementia patients

Heinonen EH, Savijarvi M, Kotila M, et al. Effects of monoamine oxidase inhibition by selegiline on concentrations of noradrenaline and monoamine metabolites in CSF of patients with Alzheimer's disease. *J Neural Transm Park Dis Dement Sect* 1993; 5(3):193-202.

Status: Cross-over trial;

Heinze B, Karrass W, Peters T. Pharmacopsychological effects of flunarizine in geriatric patients with light brainorganic psychosyndrome. Preliminary communication.

Eur Neurol 1986; 25(Suppl 1):115-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heiss WD, Kessler J, Slansky I, et al. Activation PET as an instrument to determine therapeutic efficacy in Alzheimer's disease. *Ann N Y Acad Sci* 1993; 695(Sep 24):327-13.

Status: Not included because Jadad Quality Scale score less than three

Heiss WD, Kessler J, Mielke R, et al. Long-term effects of phosphatidylserine, pyritinol, and cognitive training in Alzheimer's disease. A neuropsychological, EEG, and PET investigation. *Dementia* 1994 Mar; 5(2):88-98.

Status: Not included because Jadad Quality Scale score less than three

Helkala EL, Koivisto K, Hanninen T, et al. Stability of age-associated memory impairment during a longitudinal population-based study. *J Am Geriatr Soc* 1997 Jan; 45(1):120-2.

Status: Not included because not a full article

Henderson VW, Roberts E, Wimer C, et al. Multicenter trial of naloxone in Alzheimer's disease. *Ann Neurol* 1989 Apr; 25(4):404-6.

Status: Cross-over trial;

Henderson VW, Paganini-Hill A, Miller BL, et al. A randomized controlled trial of estrogen for the treatment of Alzheimer's disease in women. *Neurology* 2000; 54(Suppl 3):A470.

Status: Not included because not a full article

Herrmann WM, Kern U, Rohmel J. On the effects of pyritinol on functional deficits of patients with organic mental disorders. *Pharmacopsychiatry* 1986 Sep; 19(5):378-85.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Kern U, Rohmel J. Contribution to the search for vigilance-indicative EEG variables. Results of a controlled, double-blind study with pyritinol in elderly patients with symptoms of mental dysfunction. *Pharmacopsychiatry* 1986 Mar; 19(2):75-83.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Dietrich B, Hiersemenzel R. Pharmacoelectroencephalographic and clinical effects of the cholinergic substance--acetyl-L-carnitine--in patients with organic brain syndrome.

Int J Clin Pharmacol Res 1990; 10(1-2):81-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herrmann WM, Stephan K. Moving from the question of efficacy to the question of therapeutic relevance: an exploratory reanalysis of a controlled clinical study of 130 inpatients with dementia syndrome taking piracetam. Int Psychogeriatr 1992; 4(1):25-44.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Herz LR, Volicer L, Ross V, et al. A single-case-study method for treating resistiveness in patients with Alzheimer's disease. Hospital & Community Psychiatry 1992 Jul; 43(7):720-4.
Status: Cross-over trial;

Heseker H, Kubler W, Pudel V, et al. Interaction of vitamins with mental performance. Bibl Nutr Dieta 1995; (52):43-55.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heseltine PN, Goodkin K, Atkinson JH, et al. Randomized double-blind placebo-controlled trial of peptide T for HIV-associated cognitive impairment. Arch Neurol 1998 Jan; 55(1):41-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heuser I, Heuser-Link M, Gotthardt U, et al. Behavioral effects of a synthetic corticotropin 4-9 analog in patients with depression and patients with Alzheimer's disease. J Clin Psychopharmacol 1993 Jun; 13(3):171-4.
Status: Not included because dementia population not randomized to treatment

Hewawasam L. Floor patterns limit wandering of people with Alzheimer's. Nurs Times 1996 May 29; 92(22):41-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Heyman A, Schmechel D, Wilkinson W, et al. Failure of long term high-dose lecithin to retard progression of early-onset Alzheimer's disease. J Neural Transm Suppl 1987; Vol 24:279-86.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hinchliffe AC, Katona C, Livingston G. The assessment and management of behavioural manifestations of dementia: A review and results

of a controlled trial. Int J Psychiatry Clin Pract 1997; 1(3):157-68.
Status: Not included because does not meet criteria for treatment for dementia patients

Hincliffe AC, Hyman IL, Blizard B, et al. Behavioural complications of dementia - Can they be treated? Int J Geriatr Psychiatry 1995; 10(10):839-47.
Status: Not included because does not meet criteria for treatment for dementia patients

Hindle JV, Meara RJ, Sharma JC, et al. Prescribing pergolide in the elderly - An open label study of pergolide in elderly patients with Parkinson's disease. Int J Geriatr Psychopharmacol 1998; (2):78-81.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hindmarch I, Fuchs HH, Erzigkeit H. Efficacy and tolerance of vinpocetine in ambulant patients suffering from mild to moderate organic psychosyndromes. Int Clin Psychopharmacol 1991; 6(1):31-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hinkin CH, Castellon SA, Hardy DJ, et al. Methylphenidate improves HIV-1-associated cognitive slowing. J Neuropsychiatry Clin Neurosci 2001; 13(2):248-54.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hirohata S, Suda H, Hashimoto T. Low-dose weekly methotrexate for progressive neuropsychiatric manifestations in Behcet's disease. JNS 1998 Aug 14; 159(2):181-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hjorthor A, Browne E, Jakobsen K, et al. Organic brain syndrome treated with oxiracetam. A double-blind randomized controlled trial. Acta Neurol Scand 1987 Apr; 75(4):271-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hock C, Maddalena A, Heuser I, et al. Treatment with the selective muscarinic agonist talsaclidine decreases cerebrospinal fluid levels of total amyloid beta-peptide in patients with Alzheimer's disease. Ann N Y Acad Sci 2000; 920:285-91.
Status: Not included because no extractable data relevant to review

Hodges, Graham JR, K.S. A reversal of the temporal gradient for famous person knowledge in semantic dementia: implications for the neural organisation of long-term memory. *Neuropsychologia* 1998; 36(8):803-25.

Status: Not included because does not meet criteria for treatment for dementia patients

Hoeffler B, Rader J, McKenzie D, et al. Reducing aggressive behavior during bathing cognitively impaired nursing home residents. *J Gerontol Nurs* 1997 May; 23(5):16-23.

Status: Not included because dementia population not randomized to treatment

Hofferberth B. The efficacy of EGb 761 in patients with senile dementia of the Alzheimer type, a double-blind, placebo-controlled study on different levels of investigation. *Hum Psychopharmacol* 1994; (3):215-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Holden M, Kelly C. Use of cholinesterase inhibitors in dementia. *Adv Psychiatr Treat* 2002; 8(2):89-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hollander E, Davidson M, Mohs RC, et al. RS 86 in the treatment of Alzheimer's disease: cognitive and biological effects. *Biol Psychiatry* 1987 Sep; 22(9):1067-78.

Status: Cross-over trial;

Holliman DC, Orgassa UC, Forney JP. Developing an interactive physical activity group in a geriatric psychiatry facility. *Activities Adaptation Aging* 2001; 26(1):57-69.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Holm A, Michel M, Stern GA, et al. The outcomes of an inpatient treatment program for geriatric patients with dementia and dysfunctional behaviors. *Gerontologist* 1999 Dec; 39(6):668-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Holmes C, Hopkins V, Hensford C, et al. Lavender oil as a treatment for agitated behaviour in severe dementia: A placebo controlled study. *Int J Geriatr Psychiatry* 2002 Apr; 17(4):305-8.

Status: Not included because dementia population not randomized to treatment

Holzman D. Ginkgo biloba for Alzheimer's disease. *Altern Complement Ther* 1998; (5):361-3.

Status: Not included because not a full article

Homma A, Takeda M, Imai Y, et al. Clinical efficacy and safety of donepezil on cognitive and global function in patients with Alzheimer's disease. A 24-week, multicenter, double-blind, placebo-controlled study in Japan. *E2020 Study Group. Dement Geriatr Cogn Disord* 2000 Nov; 11(6):299-313.

Status: Not included because Jadad Quality Scale score less than three

Honjo H, Ogino Y, Naitoh K, et al. In vivo effects by estrone sulfate on the central nervous system-senile dementia (Alzheimer's type). *J Steroid Biochem* 1989; 34(1-6):521-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Honjo H, Tanaka T, Urabe M, et al. Senile-dementia Alzheimer's type and estrogen. *Hormone & Metabolic Research* 1995; 27(4):204-7.

Status: Not included because dementia population not randomized to treatment

Hopman-Rock M, Staats PG, Tak EC, et al. The effects of a psychomotor activation programme for use in groups of cognitively impaired people in homes for the elderly. *Int J Geriatr Psychiatry* 1999 Aug; 14(8):633-42.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hornig P. Commentary on A 30-week randomized controlled trial of high-dose tacrine in patients with Alzheimer's disease. *Nursing Scan in Research* 1994; 7(4):6

Status: Not included because not a full article

Hossain M, Jhee SS, Shiovitz T, et al. Estimation of the absolute bioavailability of rivastigmine in patients with mild to moderate dementia of the Alzheimer's type. *Clin Pharmacokinet* 2002; 41(3):225-34.

Status: Not included because no extractable data relevant to review

Hoyer S, Oesterreich K, Stoll KD. Effects of Pyritinol HCl on blood flow and oxidative metabolism of the brain in patients with dementia. *Arzneimittelforschung* 1977; 27(3):671-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hozumi S, Hori H, Okawa M, et al. Favorable effect of transcranial electrostimulation on behavior disorders in elderly patients with dementia: a double-blind study. *Int J Neurosci* 1996 Nov; 88(1-2):1-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Huff FJ, Shipley J, Somerville NJ, et al. Besipirdine (HP 749) treatment trial in Alzheimer's disease. *Neurology* 1995; 45(Suppl 4):A288.

Status: Not included because not a full article

Huff FJ. Preliminary evaluation of besipirdine for the treatment of Alzheimer's disease. *Ann N Y Acad Sci* 1996; 777:410-4.

Status: Not included because Jadad Quality Scale score less than three

Huff FJ, Antuono PG, Delagandara JE, et al. A treatment and withdrawal trial of besipirdine in Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1996; 10(2):93-102.

Status: Not included because Jadad Quality Scale score less than three

Huusko TM, Karppi P, Avikainen V, et al. Randomised, clinically controlled trial of intensive geriatric rehabilitation in patients with hip fracture: Subgroup analysis of patients with dementia. *BMJ* 2000 Nov 4; 321(7269):1107-11.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Hyman BT, Eslinger PJ, Damasio AR. Effect of naltrexone on senile dementia of the Alzheimer type. *J Neurol Neurosurg Psychiatry* 1985; 48(11):1169-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ikedo S, Yamada Y, Ikegami N. Economic evaluation of donepezil treatment for Alzheimer's disease in Japan. *Dement Geriatr Cogn Disord* 2002; 13(1):33-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Imbimbo BP, Nicoli M, Martini C, et al. Acetylcholinesterase assay may predict cognitive response of Alzheimer patients to eptastigmine treatment. *Eur J Clin Pharmacol* 1998 Nov; 54(9-10):809-10.

Status: Not included because not a full article

Imbimbo BP, Lucca U, Lucchelli F, et al. A 25-week placebo-controlled study of eptastigmine in patients with Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1998 Dec; 12(4):313-22.

Status: Not included because Jadad Quality Scale score less than three

Imbimbo BP, Verdelli G, Martelli P, et al. Two-year treatment of Alzheimer's disease with eptastigmine. *Dement Geriatr Cogn Disord* 1999; 10(2):139-47.

Status: Not included because dementia population not randomized to treatment

Imbimbo BP, Troetel WM, Martelli P, et al. A 6-month, double-blind, placebo-controlled trial of eptastigmine in Alzheimer's disease. *Dement Geriatr Cogn Disord* 2000 Jan; 11(1):17-24.

Status: Not included because Jadad Quality Scale score less than three

In't Veld BA, Ruitenberg A, Hofman A, et al. Antihypertensive drugs and incidence of dementia: The Rotterdam Study. *Neurobiol Aging* 2001; 22(3):407-12.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ingersoll-Dayton B, Schroeffer T, Pryce J. The effectiveness of a solution-focused approach for problem behaviors among nursing home residents. *J Gerontol Soc Work* 1999; 32(3):49-64.

Status: Not included because does not meet criteria for treatment for dementia patients

Innes EH. Efficacy and tolerance of flurbiprofen in the elderly using liquid and tablet formulations. *Curr Med Res Opin* 1977; 5(1):122-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Inzitari D, Pantoni L. Subcortical vascular dementia: therapeutic aspects. *Arch Gerontol Geriatr* 1998; 263-8.

Status: Not included because not a full article

Iribar MC, Montes J, Gonzalez MR, et al. Alanyl-aminopeptidase activity decrease in cerebrospinal fluid of Alzheimer patients. *Dement Geriatr Cogn Disord* 1998 Jan; 9(1):44-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Ishizaki J, Meguro K, Ohe K, et al. Therapeutic psychosocial intervention for elderly subjects with very mild Alzheimer disease in a community: The tajiri project. *Alzheimer Dis Assoc Disord* 2002; 16(4):261-9.

Status: Not included because dementia population not randomized to treatment

Israel L, Melac M, Milinkevitch D, et al. Drug therapy and memory training programs: a double-blind randomized trial of general practice patients with age-associated memory impairment. *Int Psychogeriatr* 1994; 6(2):155-70.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Israel L, Myslinski M, Kozarevic D. Nootropic treatment and combined therapy in age-associated memory impairment. *Arch Gerontol Geriatr* 1998; 27(Suppl 6):269-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Itil T, Martorano D. Natural substances in psychiatry (Ginkgo biloba in dementia). *Psychopharmacol Bull* 1995; 31(1):147-58.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Itil TM, Mukherjee S, Michael ST. Clinical and electrophysiological effects of suloctidil in elderly patients with multi-infarct dementia (A double-blind placebo-controlled study). *Psychopharmacol Bull* 1983; 19(4):730-3.

Status: Not included because Jadad Quality Scale score less than three

Itil TM, Eralp E, Ahmed I, et al. The pharmacological effects of Ginkgo biloba, a plant extract, on the brain of dementia patients in comparison with tacrine. *Psychopharmacol Bull* 1998; 34(3):391-7.

Status: Not included because dementia population not randomized to treatment

Jacobs EA, Winter PM, Alvis HJ, et al. Hyperoxygenation effect on cognitive functioning in the aged. *N Engl J Med* 1969 Oct 2; 281(14):753-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jani J, Prettyman R. Use of a prescribing protocol in routine clinical practice: Experience following the introduction of donepezil. *Psychiatr Bull* 2001;

25(5):174-7.

Status: Not included because dementia population not randomized to treatment

Jann MW. Rivastigmine, a new-generation cholinesterase inhibitor for the treatment of Alzheimer's disease. *Pharmacotherapy* 2000; 20(11):1-12.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jansen W, Bruckner GW, Jansen P. The treatment of senile dementia associated with cerebrovascular insufficiency: a comparative study of buflomedil and dihydrogenated ergot alkaloids. *J Int Med Res* 1985; 13(1):48-53.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jansen W, O'Connolly M, Lehmann E, et al. Experimental clinical studies on the effect of eburnamonine in cerebrovascular disorders. *Pharmacopsychiatry* 1986 Sep; 19(5):389-94.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jean-Louis G, von Gizycki H, Zizi F. Melatonin effects on sleep, mood, and cognition in elderly with mild cognitive impairment. *J Pineal Res* 1998 Oct; 25(3):177-83.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jellinger K, Flament H, Riederer P. Levodopa in the treatment of (pre) senile dementia. *Mech Ageing Dev* 1980; 14(1-2):253-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jenike MA, Albert MS, Heller H, et al. Combination therapy with lecithin and ergoloid mesylates for Alzheimer's disease. *J Clin Psychiatry* 1986 May; 47(5):249-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jenike MA, Albert MS, Heller H, et al. Oral physostigmine treatment for patients with presenile and senile dementia of the Alzheimer's type: a double-blind placebo-controlled trial. *J Clin Psychiatry* 1990 Jan; 51(1):3-7.

Status: Not included because dementia population not randomized to treatment

Jenike MA, Albert M, Baer L, et al. Ergot mesylates for Alzheimer's disease: A year-long

double-blind trial of 3 mg vs 12 mg daily. *Int J Geriatr Psychiatry* 1990; 5(6):375-80.
Status: Not included because Jadad Quality Scale score less than three

Jennekens-Schinkel A, Wintzen AR, Lanser JB. A clinical trial with desglycinamide arginine vasopressin for the treatment of memory disorders in man. *Prog Neuropsychopharmacol Biol Psychiatry* 1985; 9(3):273-84.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jeste DV, Lacro JP, Bailey A, et al. Lower incidence of tardive dyskinesia with risperidone compared with haloperidol in older patients. *J Am Geriatr Soc* 1999 Jun; 47(6):716-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jhee SS, Shiovitz T, Hartman RD, et al. Centrally acting antiemetics mitigate nausea and vomiting in patients with Alzheimer's disease who receive rivastigmine. *Clin Neuropharmacol* 2002 Mar; 25(2):122-3.
Status: Not included because not a full article

Jobe JB, Smith DM, Ball K, et al. ACTIVE: a cognitive intervention trial to promote independence in older adults. *Control Clin Trials* 2001 Aug; 22(4):453-79.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Johnson SA, Simmon VF. Randomized, double-blind, placebo-controlled international clinical trial of the AMPAKINE CX516 in elderly participants with mild cognitive impairment. A progress report. *J Mol Neurosci* 2002; 19(1-2):197-200.
Status: Not included because no extractable data relevant to review

Johnstone EC, Owens DG, Crow TJ, et al. Does a four-week delay in the introduction of medication alter the course of functional psychosis? *J Psychopharmacol (Oxf)* 1999; 13(3):238-44.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jolkkonen JT, Soininen HS, Riekkinen PJ. The effect of ACTH4-9 analog (Org2766) on some cerebrospinal fluid parameters in patients with Alzheimer's disease. *Life Sci* 1985 Aug 19; 37(7):585-90.
Status: Not included because does not meet criteria for treatment for dementia patients

Jones GM, Sahakian BJ, Levy R, et al. Effects of acute subcutaneous nicotine on attention, information processing and short-term memory in Alzheimer's disease. *Psychopharmacologia* 1992; 108(4):485-94.
Status: Not included because dementia population not randomized to treatment

Jonsson A, Korfitzen EM, Heltberg A, et al. Effects of neuropsychological treatment in patients with multiple sclerosis. *Acta Neurol Scand* 1993 Dec; 88(6):394-400.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jonsson L, Lindgren P, Wimo A, et al. The cost-effectiveness of donepezil therapy in Swedish patients with Alzheimer's disease: A Markov model. *Clin Ther* 1999; 21(7):1230-40.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jonsson L, Gerth W, Fastbom J. The potential economic consequences of cognitive improvement with losartan. *Blood Press* 2002; 11(1):46-52.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jorgenson S, Bille A, Andersen J, et al. Fluvoxamine treatment of dementia: Tryptophan levels. *Biol Psychiatry* 1993; 34(8):587-8.
Status: Not included because not a full article

Jorissen BL, Brouns F, Van Boxtel MP, et al. The influence of soy-derived phosphatidylserine on cognition in age-associated memory impairment. *Nutr Neurosci* 2001; 4(2):121-34.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jorissen BL, Brouns F, Van Boxtel MPJ, et al. Safety of soy-derived phosphatidylserine in elderly people. *Nutr Neurosci* 2002; 5(5):337-43.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Jotkowitz S. Lack of clinical efficacy of chronic oral physostigmine in Alzheimer's disease. *Ann Neurol* 1983 Dec; 14(6):690-1.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Judge KS, Camp CJ, Orsulic-Jeras S. Use of Montessori-based activities for clients with

dementia in adult day care: Effects on engagement. *Am J Alzheimers Dis* 2000; 15(1):42-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Judge TG, Urquhart A. Naftidrofuryl: a double blind cross-over study in the elderly. *Curr Med Res Opin* 1972; 1(3):166-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Juvars KA, McDonald C. A controlled trial of butyrylperazine ("Randolectil") in senile dementia. *Med J Aust* 1967 Feb 18; 1(7):334-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kahana E, Kahana B. Therapeutic potential of age integration. Effects of age-integrated hospital environments on elderly psychiatric patients. *Arch Gen Psychiatry* 1970 Jul; 23(1):20-9.

Status: Not included because no extractable data relevant to review

Kanamori M, Suzuki M, Yamamoto K, et al. A day care program and evaluation of animal-assisted therapy (AAT) for the elderly with senile dementia. *Am J Alzheimers Dis* 2001; 16(4):234-9.

Status: Not included because dementia population not randomized to treatment

Kanowski S, Kinzler E, Lehmann E, et al. Confirmed clinical efficacy of Actovegin in elderly patients with organic brain syndrome. *Pharmacopsychiatry* 1995 Jul; 28(4):125-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kapur N, Ironside J, Abbott P, et al. A neuropsychological-neuropathological case study of variant Creutzfeldt-Jakob disease. *Neurocase* 2001; 7(3):261-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kario K, Matsuo T, Hoshida S, et al. Effect of thrombin inhibition in vascular dementia and silent cerebrovascular disease. An MR spectroscopy study. *Stroke* 1999 May; 30(5):1033-7.

Status: Not included because dementia population not randomized to treatment

Karlsson I, Brane G, Melin E, et al. Effects of environmental stimulation on biochemical and psychological variables in dementia. *Acta*

Psychiatr Scand 1988; 77(2):207-13.

Status: Not included because dementia population not randomized to treatment

Karlsson J, Hallgren P, Kral J, et al. Predictors and effects of long-term dieting on mental well-being and weight loss in obese women. *Appetite* 1994 Aug; 23(1):15-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Karoutas G, Milonas I, Artemis N, et al. The pharmacological effects of buflomedil in patients with multi-cerebral infarcts dementia: An open, preliminary assessment. *Curr Med Res Opin* 1986; Vol 10:380-9.

Status: Not included because dementia population not randomized to treatment

Kasckow JW, McElroy SL, Cameron RI. A pilot study on the use of divalproex sodium in the treatment of behavioral agitation in elderly patients with dementia: assessment with the BEHAVE-AD and CGI rating scales. *Current therapeutic research, clinical and experimental* 1997; 58(981):989.

Status: Not included because dementia population not randomized to treatment

Katzman R, Zhang M, YChen PJ, et al. Effects of apolipoprotein E on dementia and aging in the Shanghai Survey of Dementia. *Neurology* 1997; 49(3):779-85.

Status: Not included because does not meet criteria for treatment for dementia patients

Kaufer DI. Pharmacologic therapy of dementia with Lewy bodies. *J Geriatr Psychiatry Neurol* 2002; 15(4):224-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kaye WH, Sitaram N, Weingartner H, et al. Modest facilitation of memory in dementia with combined lecithin and anticholinesterase treatment. *Biol Psychiatry* 1982; 17(2):275-80.

Status: Cross-over trial;

Kazui H, Mori E, Hashimoto M, et al. Impact of emotion on memory. Controlled study of the influence of emotionally charged material on declarative memory in Alzheimer's disease. *Br J Psychiatry* 2000; 177:343-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Kempenaar L, McNamara C, Creaney B. Sensory stimulation work with carers in the community. *Journal of Dementia Care* 2001; 9(1):16.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kertesz A. Efficacy of galantamine in probable vascular dementia and Alzheimer's disease combined with cerebrovascular disease: A randomized trial. *Curr Neurol Neurosci Rep* 2002 Nov; 2(6):503-4.
Status: Not included because not a full article

Khan A, Mirolo MH, Claypoole K, et al. Low-dose thyrotropin-releasing hormone effects in cognitively impaired alcoholics. *Alcohol Clin Exp Res* 1993 Aug; 17(4):791-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kiebertz K, Schifitto G, McDermott M, et al. Safety and tolerability of the antioxidant OPC-14117 in HIV-associated cognitive impairment. *Neurology* 1997; 49(1):142-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kiebertz K, Schifitto G, McDermott M, et al. A randomized, double-blind, placebo-controlled trial of deprenyl and thioctic acid in human immunodeficiency virus-associated cognitive impairment. *Neurology* 1998; 50(3):645-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kim JM, Shin IS, Yoon JS. Correlates of dropout, efficacy, and adverse events in treatment with acetylcholinesterase inhibitors in Korean patients with Alzheimer's disease. *Int Psychogeriatr* 2002; 14(2):187-95.
Status: Not included because Jadad Quality Scale score less than three

Kimura M, Robinson RG, Kosier JT. Treatment of cognitive impairment after poststroke depression: A double-blind treatment trial. *Stroke* 2000 Jul; 31(7):1482-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

King AC, Brassington G. Enhancing physical and psychological functioning in older family caregivers: The role of regular physical activity. *Ann Behav Med* 1997; (No. 2):91-100.
Status: Not included because does not meet criteria for treatment for dementia patients

Kirby M, Denihan A, Bruce I, et al. Benzodiazepine use among the elderly in the community. *Int J Geriatr Psychiatry* 1999; 14(4):280-4.
Status: Not included because dementia population not randomized to treatment

Kirrane RM, Mitropoulou V, Nunn M, et al. Physostigmine and cognition in schizotypal personality disorder. *Schizophr Res* 2001 Mar 1; 48(1):1-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kishimoto T, Hiraoka Y, Oribe H, et al. Auditory P300 event-related potentials and mini mental state examination performance in dementia; effects of l-debenone and vinpocetine. *J Nara Med Assoc* 1995; 46(3):259-66.
Status: Not included because dementia population not randomized to treatment

Kittner B. Using a combined randomized start/withdrawal design to assess propentofylline's effect on disease progression in Alzheimer's disease and vascular dementia. *J Neural Transm Gen Sect* 2000; 107:XV.
Status: Not included because does not meet criteria for treatment for dementia patients

Kittner B. Propentofylline for the treatment of vascular dementia: A 48-week, placebo-controlled study examining safety, efficacy, and impact on disease progression. *J Cereb Blood Flow Metab* 1999; 19(Suppl 1):16.
Status: Not included because not a full article

Kittner B. Using a combined randomized start/withdrawal design to assess propentofylline's effects on disease progression in Alzheimer's disease and vascular dementia: Results of clinical studies. *J Eur Coll Neuropsychopharmacol* 1999; 9(Suppl 5):S320
Status: Not included because not a full article

Kittner B. Clinical trials of propentofylline in vascular dementia. European/Canadian Propentofylline Study Group. *Alzheimer Dis Assoc Disord* 1999; 13(Suppl 3):S166-S171.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kittner B, De Deyn PP, Erkinjuntti T. Investigating the natural course and treatment of vascular dementia and Alzheimer's disease. Parallel study

populations in two randomized, placebo-controlled trials. *Ann N Y Acad Sci* 2000 Apr; 903:535-41.
Status: Not included because Jadad Quality Scale score less than three

Knezevic S, Mubrin Z, Risberg J, et al. Pyritinol treatment of SDAT patients: evaluation by psychiatric and neurological examination, psychometric testing and rCBF measurements. *Int Clin Psychopharmacol* 1989 Jan; 4(1):25-38.
Status: Cross-over trial;

Knopman D. Long-term retention of implicitly acquired learning in patients with Alzheimer's disease. *J Clin Exp Neuropsychol* 1991; 13(6):880-94.
Status: Not included because does not meet criteria for treatment for dementia patients

Knopman D, Schneider L, Davis K, et al. Long-term tacrine treatment effects. *Neurology* 1998 Feb; 50(2):567-8.
Status: Not included because dementia population not randomized to treatment

Knopman DS. Metrifonate for Alzheimer's disease: Is the next cholinesterase inhibitor better? *Neurology* 1998; 50(5):1203-5.
Status: Not included because dementia population not randomized to treatment

Knopman D, Gracon S. Observations on the short-term "natural history" of probable Alzheimer's disease in a controlled clinical trial. *Neurology* 1994; 44(2):260-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Knott V, Mohr E, Mahoney C, et al. Pharmacologic EEG test dose response predicts cholinesterase inhibitor treatment outcome in Alzheimer's disease. *Methods & Findings in Experimental & Clinical Pharmacology* 2000 Mar; 22(2):115-22.
Status: Not included because dementia population not randomized to treatment

Knott V, Engeland C, Mohr E, et al. Acute nicotine administration in Alzheimer's disease: An exploratory EEG study. *Neuropsychobiology* 2000; 41(4):210-20.
Status: Not included because dementia population not randomized to treatment

Knox J, Hindmarch I, Wallace M. Effects of twice standard dosage of neuroactive drugs in dementia. A preliminary report. *Br J Clin Pract*

1984 Sep; 38(9):313-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kodjian A, Barriaga C, Turcot G. Double-blind study of pimozide in senile dementia patients. *Curr Ther Res Clin Exp* 1986; 40(4):694-701.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kofler B, Erhart C, Erhart P, et al. A multidimensional approach in testing nootropic drug effects (Cerebrolysin (R)). *Arch Gerontol Geriatr* 1990; 10(2):129-40.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kogan EA, Korczyn AD, Virchovsky RG, et al. EEG changes during long-term treatment with donepezil in Alzheimer's disease patients. *J Neural Transm Gen Sect* 2001; 108(10):1167-73.
Status: Not included because dementia population not randomized to treatment

Koh K, Ray R, Lee J, et al. Dementia in elderly patients: Can the 3R mental stimulation programme improve mental status? *Age Ageing* 1994; 23(3):195-9.
Status: Not included because dementia population not randomized to treatment

Koivisto, Portin M, Seinela R, et al. Automatic influences of memory in Alzheimer's disease. *Cortex* 1998; 34(2):209-19.
Status: Not included because does not meet criteria for treatment for dementia patients

Koivisto K, Helkala EL, Hanninen T, et al. Long-term selegiline treatment reduces the progression of Alzheimer's disease: Results after three years' follow-up. *Alzheimers Res* 1995; 1(Suppl 1):28.
Status: Not included because not a full article

Koltai DC, Welsh-Bohmer KA, Schmechel DE. Influence of anosognosia on treatment outcome among dementia patients. *Neuropsychol Rehab* 2001; 11(3-4):455-75.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Koltringer P, Langsteger W, Eber O. Dose-dependent hemorheological effects and microcirculatory modifications following intravenous administration of Ginkgo biloba special extract EGb 761. *Clin Hemorheol* 1995; 15(4):649-56.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kompoliti K, Goetz CG, Boeve BF, et al. Clinical presentation and pharmacological therapy in corticobasal degeneration. *Arch Neurol* 1998 Jul; 55(7):957-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kontush A, Mann U, Arlt S, et al. Influence of vitamin E and C supplementation on lipoprotein oxidation in patients with Alzheimer's disease. *Free Radic Biol Med* 2001 Aug 1; 31(3):345-54.
Status: Not included because dementia population not randomized to treatment

Kosten TR, Rosen MI, McMahon TL, et al. Treatment of early AIDS dementia in intravenous drug users: High versus low dose peptide T. *Am J Drug Alcohol Abuse* 1997 Nov; 23(4):543-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kountouris D. Therapeutic effects of piracetam combined with intravenous immunoglobulin premature of Alzheimer type. *J Neural Transm Gen Sect* 2000; 107:XVIII.
Status: Not included because not a full article

Krebs Roubicek EM. Group therapy with demented elderly. *Alzheimers Dis Relat Disord* 1989; 1261-72.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kristensen V, Olsen M, Theilgaard A. Levodopa treatment of presenile dementia. *Acta Psychiatr Scand* 1977 Jan; 55(1):41-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kruck TPA, Fisher EA, McLachlan DRC. Prediction of side effects in Alzheimer's disease patients on longterm deferoxamine-mesylate treatment. *Alzheimers Dis Relat Disord Adv Biosci* 1993; 87:257-8.
Status: Not included because no extractable data relevant to review

Kruck TPA, Fisher EA, McLachlan DRC. A predictor for side effects in patients with Alzheimer's disease treated with deferoxamine mesylate. *Clin Pharmacol Ther* 1993; 53(1):30-7.
Status: Not included because dementia population not randomized to treatment

Kuhl DE, Minoshima S, Frey KA, et al. Limited donepezil inhibition of acetylcholinesterase measured with positron emission tomography in living Alzheimer cerebral cortex. *Ann Neurol* 2000 Sep; 48(3):391-5.

Status: Not included because dementia population not randomized to treatment

Kuhn DR, de Leon CFM. Evaluating an educational intervention with relatives of persons in the early stages of Alzheimer's disease. *Res Soc Work Pract* 2001 Sep; 11(5):531-48.
Status: Not included because does not meet criteria for treatment for dementia patients

Kulisevsky J, Garcia-Sanchez C, Berthier ML, et al. Chronic effects of dopaminergic replacement on cognitive function in Parkinson's disease: A two-year follow-up study of previously untreated patients. *Mov Disord* 2000 Jul; 15(4):613-26.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kumar V, Smith RC, Sherman KA, et al. Cortisol responses to cholinergic drugs in Alzheimer's disease. *Int J Clin Pharmacol Ther Toxicol* ; 26(10):471-6.
Status: Not included because dementia population not randomized to treatment

Kumar V, Brecher M. Psychopharmacology of atypical antipsychotics and clinical outcomes in elderly patients. *J Clin Psychiatry* 1999; 60(Suppl 13):5-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kuskowski MA, Morley G, Malone SM, et al. Hydergine treatment and psychophysiological measures in primary degenerative dementia. *J Geriatr Psychiatry Neurol* 1990 Jan; 3(1):41-7.
Status: Not included because dementia population not randomized to treatment

Kwiecinski H, Lusakowska A, Mieszkowski J. Improvement in concentration following treatment with Ginseng/Ginkgo biloba combination in patients with chronic cerebrovascular disorders: A double-blind, placebo-controlled study. *Eur J Clin Res* 1997; 9:59-67.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Kwok T, Tang C, Woo J, et al. Randomized trial of the effect of supplementation on the cognitive

function of older people with subnormal cobalamin levels. *Int J Geriatr Psychiatry* 1998 Sep; 13(9):611-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LaBarge E, Rosenman LS, Leavitt K, et al. Counseling Clients with Mild Senile Dementia of the Alzheimer's Type: A Pilot Study. *J Neurol Rehabil* 1988; 2(4):167-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LaBrecque DC, Goldberg RI. A double-blind study of pentylentetrazol combined with niacin in senile patients. *Curr Ther Res Clin Exp* ; 9(12):611-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lacomblez L, Chatellier L. Tetrahydroaminoacridine (THA) in Alzheimer's disease (AD): A double-blind pilot study. *Fundam Clin Pharmacol* 1989; 3(2):172.

Status: Not included because not a full article

Lacomblez L, Chatellier L. A multicenter trial of tetrahydroaminoacridine in senile dementia of the Alzheimer type. *Fundam Clin Pharmacol* 1990; 4(4):456.

Status: Not included because not a full article

Lamb HM, Faulds D. Mefenolate. *Drugs Aging* 1997; 11(6):490-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lanctot KL, Herrmann N, Van Reekum R, et al. Gender, aggression and serotonergic function are associated with response to sertraline for behavioral disturbances in Alzheimer's disease. *Int J Geriatr Psychiatry* 2002; 17(6):531-41.

Status: Cross-over trial;

Landi F, Bernabei R, Russo A, et al. Predictors of rehabilitation outcomes in frail patients treated in a geriatric hospital. *J Am Geriatr Soc* 2002; 50(4):679-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lange KW, Robbins TW, Marsden CD, et al. L-dopa withdrawal in Parkinson's disease selectively impairs cognitive performance in tests sensitive to frontal lobe dysfunction. *Psychopharmacologia* 1992; 107(2-3):394-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lange KW, Paul GM, Naumann M, et al. Dopaminergic effects on cognitive performance in patients with Parkinson's disease. *J Neural Transm Suppl* 1995; 46:423-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Langlais PJ, Mair RG, Whalen PJ, et al. Memory effect of DL-threo-3,4-dihydroxyphenylserine (DOPS) in human Korsakoff's disease. *Psychopharmacologia* 1988; 95(2):250-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LaPorte DJ, Lahti AC, Koffel B, et al. Absence of ketamine effects on memory and other cognitive functions in schizophrenia patients. *J Psychiatr Res* 1996 Sep; 30(5):321-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lawlor BA, Mellow AM, Sunderland T, et al. A pilot study of serotonergic system responsivity in Alzheimer's disease. *Psychopharmacol Bull* 1988; 24(1):127-9.

Status: Not included because Jadad Quality Scale score less than three

Lawlor BA, Sunderland T, Mellow AM, et al. Hyperresponsivity to the serotonin agonist m-chlorophenylpiperazine in Alzheimer's disease. A controlled study. *Arch Gen Psychiatry* 1989 Jun; 46(6):542-9.

Status: Cross-over trial;

Lawlor BA, Sunderland T, Mellow AM, et al. A pilot placebo-controlled study of chronic m-CPP administration in Alzheimer's disease. *Biol Psychiatry* 1991 Jul 15; 30(2):140-4.

Status: Cross-over trial;

Lawlor BA, Radcliffe J, Molchan SE, et al. A pilot placebo-controlled study of trazodone and buspirone in Alzheimer's disease. *Int J Geriatr Psychiatry* 1994; 9(1):55-9.

Status: Cross-over trial;

Lawlor BA, Aisen PS, Green C, et al. Selegiline in the treatment of behavioural disturbance in Alzheimer's disease. *Int J Geriatr Psychiatry* 1997 Mar; 12(3):319-22.

Status: Cross-over trial;

Laws SM, Clarnette RM, Taddei K, et al. APOE-epsilon4 and APOE-491A polymorphisms in individuals with subjective memory loss. *Mol Psychiatry* 2002; 7(7):768-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lawton MP, Van Haitsma K, Klapper J, et al. A stimulation-retreat special care unit for elders with dementing illness. *Int Psychogeriatr* 1998 Dec; 10(4):379-95.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lawton MP, Brody EM, Saperstein AB, et al. Respite services for caregivers: Research findings for service planning. *Home Health Care Serv Q* 1989; 10(1-2):5-32.

Status: Not included because does not meet criteria for treatment for dementia patients

Lawton MP, Brody EM, Saperstein AR. Controlled study of respite service for caregivers of Alzheimer's patients. *Gerontologist* 1989 Feb; (1):8-16.

Status: Not included because does not meet criteria for treatment for dementia patients

Lazar PA. Ginkgo is not a smart pill. *J Fam Pract* 2002 Nov; 51(11):912.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Leblhuber F, Neubauer C, Peichl M, et al. Age and sex differences of dehydroepiandrosterone sulfate (DHEAS) and cortisol (CRT) plasma levels in normal controls and Alzheimer's disease (AD). *Psychopharmacologia* 1993; 111(1):23-6.

Status: Not included because dementia population not randomized to treatment

Leblhuber F, Walli J, Widner B, et al. Homocysteine and B vitamins in dementia. *Am J Clin Nutr* 2001; 73(1):127-8.

Status: Not included because not a full article

Lebowitz BD, Pollock BG, Schneider LS. Estrogen in geriatric psychopharmacology. *Psychopharmacol Bull* 1997; 33(2):287-8.

Status: Not included because not a full article

Lechner H, Walzl M, Walzl B, et al. HELP application in multi-infarct dementia. *Journal of Stroke and Cerebrovascular Diseases* 1992; 2:228-31.

Status: Not included because Jadad Quality Scale score less than three

Leckman J, Ananth JV, Ban TA, et al. Pentylentetrazol in the treatment of geriatric patients with disturbed memory function. *J Clin Pharmacol New Drugs* 1971 Jul; 11(4):301-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lehmann HE, Ban TA, Saxena BM. Nicotinic acid, thioridazine, fluoxymesterone and their combinations in hospitalized geriatric patients: A systematic clinical study. *Can Psychiatr Assoc J* 1972 Aug; 17(4):315-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Leighman L, Stotsky BA, Cole JO. A controlled study of drugs in long-term geriatric psychiatric patients: A double-blind comparison of pentylentetrazol, papaverine, and niacin. *Arch Gen Psychiatry* 1971 Sep; 25:284-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lerner V, Miodownik C, Kaptsan A, et al. Vitamin B6 as add-on treatment in chronic schizophrenic and schizoaffective patients: a double-blind, placebo-controlled study. *J Clin Psychiatry* 2002 Jan; 63(1):54-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Leszek J, Inglot AD, Janusz M, et al. Colostrinin proline-rich polypeptide complex from ovine colostrum - A long-term study of its efficacy in Alzheimer's disease. *Med Sci Monit* 2002; 8(10):I93-I96

Status: Not included because dementia population not randomized to treatment

Levin HS. Cognitive rehabilitation. Unproved but promising. *Arch Neurol* 1990 Feb; 47(2):223-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Levine FM, Spitalnik R, Dobos C. Caudate nucleus effects on geriatric senility: Effect of belladonna on learning and memory of geriatric patients. *Percept Mot Skills* 1973 Dec; 37(3):1003-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

LeVine H, III, Scholten JD. Screening for pharmacologic inhibitors of amyloid fibril formation. *Methods Enzymol* 1999; 309:467-76.

Status: Not included because dementia population not randomized to treatment

Levinson B, Wright P, Barklem S. Effect of buflomedil on behaviour, memory, and intellectual capacity in patients with dementia. A placebo-controlled study. *S Afr Med J* 1985 Aug 31; 68(5):302-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Levkovitz Y, Bloch Y, Kaplan D, et al. Fluvoxamine for psychosis in Alzheimer's disease. *J Nerv Ment Dis* 2001 Feb; 189(2):126-9.

Status: Cross-over trial;

Levy A, Brandeis R, Treves TA, et al. Transdermal physostigmine in the treatment of Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1994; 8(1):15-21.

Status: Not included because dementia population not randomized to treatment

Levy MA, Burgio LD, Sweet R, et al. A trial of buspirone for the control of disruptive behaviors in community-dwelling patients with dementia. *Int J Geriatr Psychiatry* 1994; 9(10):841-8.

Status: Not included because dementia population not randomized to treatment

Levy R, Little A, Chuaqui P, et al. Early results from double-blind, placebo controlled trial of high dose phosphatidylcholine in Alzheimer's disease. *Lancet* 1983 Apr 30; 1(8331):987-8.

Status: Not included because not a full article

Lewis C, Ballinger BR, Presly AS. Trial of levodopa in senile dementia. *BMJ* 1978 Mar 4; 1(6112):550.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Li C, Jiang Z, Wu D, et al. Clinical study on treating vascular dementia by muskiness injection in points. *Int J Clin Acupunct* 2002; 13(1):1-7.

Status: Not included because Jadad Quality Scale score less than three

Lincoln NB, Dent A, Harding J, et al. Evaluation of cognitive assessment and cognitive intervention for people with multiple sclerosis. *J Neurol Neurosurg Psychiatry* 2002; 72(1):93-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lindner K, Bedard M, William MD, et al. Changes in medication use and functional status of community-dwelling Alzheimer's patients after consultation at a memory clinic. *Clin Gerontol* 2001; 22(3-4):13-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Linn MW, Gurel L, Williford WO, et al. Nursing home care as an alternative to psychiatric hospitalization. A Veterans Administration cooperative study. *Arch Gen Psychiatry* 1985 Jun; 42(6):544-51.

Status: Not included because does not meet criteria for treatment for dementia patients

Lipper S, Tuchman MM. Treatment of chronic post-traumatic organic brain syndrome with dextroamphetamine: First reported case. *J Nerv Ment Dis* 1976 May; 162(5):366-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lipton SA, Dafni U, Simpson D, et al. Double blind randomised placebo-controlled trial of the calcium channel antagonist Nimodipine for the neurological manifestations of acquired immunodeficiency syndrome, including dementia and painful neuropathy. *Ann Neurol* 1995; 38:347

Status: Not included because not a full article

Little A, Levy R, Chuaqui-Kidd P, et al. A double-blind, placebo controlled trial of high-dose lecithin in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 1985; 48(8):736-42.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Litvan I, Gomez C, Atack JR, et al. Physostigmine treatment of progressive supranuclear palsy. *Ann Neurol* 1989 Sep; 26(3):404-7.

Status: Cross-over trial;

Litvan I, Blesa R, Clark K, et al. Pharmacological evaluation of the cholinergic system in progressive supranuclear palsy. *Ann Neurol* 1994 Jul; 36(1):55-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Litvan I, Sirigu A, Toothman J, et al. What can preservation of autobiographic memory after

muscarinic blockade tell us about the scopolamine model of dementia? *Neurology* 1995 Feb; 45(2):387-9.

Status: Not included because dementia population not randomized to treatment

Litvan I, FitzGibbon EJ. Can tropicamide eye drop response differentiate patients with progressive supranuclear palsy and Alzheimer's disease from healthy control subjects? *Neurology* 1996; 47(5):1324-6.

Status: Not included because does not meet criteria for treatment for dementia patients

Litvan I, Phipps M, Pharr VL, et al. Randomized placebo-controlled trial of donepezil in patients with progressive supranuclear palsy. *Neurology* 2001 Aug 14; 57(3):467-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Llorente AM, van Gorp WG, Stern MJ, et al. Long-term effects of high-dose zidovudine treatment on neuropsychological performance in mildly symptomatic HIV-positive patients: Results of a randomized, double-blind, placebo-controlled investigation. *J Int Neuropsychol Soc* 2001 Jan; 7(1):27-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lloyd-Evans S, Brocklehurst JC, Palmer MK. Piracetam in chronic brain failure. *Curr Med Res Opin* 1979; 6(5):351-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Loher TJ, Krauss JK, Wielepp JP, et al. Pallidal deep brain stimulation in a parkinsonian patient with late-life dementia: Sustained benefit in motor symptoms but not in functional disability. *Eur Neurol* 2002; 47(2):122-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Londos E, Passant U, Gustafson L. Blood pressure and drug treatment in clinically diagnosed Lewy body dementia and Alzheimer's disease. *Arch Gerontol Geriatr* 2000; 30(1):35-46.

Status: Not included because does not meet criteria for treatment for dementia patients

Lopez OL, Becker JT, Wisniewski S, et al. Cholinesterase inhibitor treatment alters the natural history of Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2002 Mar; 72(3):310-4.

Status: Not included because dementia population not randomized to treatment

Lord TR, Garner JE. Effects of music on Alzheimer patients. *Percept Mot Skills* 1993 Apr; 76(2):451-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lott IT, Osann K, Doran E, et al. Down syndrome and Alzheimer disease: Response to donepezil. *Arch Neurol* 2002; 59(7):1133-6.

Status: Not included because dementia population not randomized to treatment

Lovett WC, Stokes DK, Taylor LB, et al. Management of behavioral symptoms in disturbed elderly patients: Comparison of trifluoperazine and haloperidol. *J Clin Psychiatry* 1987 Jun; 48(6):234-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lucca U, Tettamanti M, Forloni G, et al. Nonsteroidal antiinflammatory drug use in Alzheimer's disease. *Biol Psychiatry* 1994 Dec; 36(12):854-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Lucca U, Lucchelli F, Alberoni M, et al. Reliability and correlation measures of cognitive function and behavioural scales in controlled clinical trial of eptastigmine in Alzheimer's disease patients. *J Neurol* 1995; 242:S106.

Status: Not included because not a full article

Lupien SJ, Wilkinson CW, Briere S, et al. Acute modulation of aged human memory by pharmacological manipulation of glucocorticoids. *J Clin Endocrinol Metab* 2002; 87(8):3798-807.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Luqman WA, Riesenber D, Oakley N, et al. Tacrine for Alzheimer's disease. *JAMA* 1994; (24):1896-8.

Status: Not included because not a full article

Lyford J. Statins reduce dementia risk by 70%. *Cur Control Trials Cardiovasc Med* 2000; 1(3):172.

Status: Not included because not a full article

Lyketsos CG, Lindell VL, Baker A, et al. A randomized, controlled trial of bright light therapy

for agitated behaviors in dementia patients residing in long-term care. *Int J Geriatr Psychiatry* 1999 Jul; 14(7):520-5.

Status: Cross-over trial;

MacGowan SH, Wilcock GK, Scott M. Effect of gender and apolipoprotein E genotype on response to anticholinesterase therapy in Alzheimer's disease. *Int J Geriatr Psychiatry* 1998 Sep; 13(9):625-30.

Status: Not included because dementia population not randomized to treatment

MacMahon S, Kermode S. A clinical trial of the effect of aromatherapy on motivational behaviour in a dementia care setting using a single subject design. *Aust J Holist Nurs* 1998 Oct; 5(2):47-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Madhusoodanan S, Brenner R, Cohen CI. Role of atypical antipsychotics in the treatment of psychosis and agitation associated with dementia. *CNS Drugs* 1999; 12(2):135-50.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Maestri NE, Brusilow SW, Clissold DB, et al. Long-term treatment of girls with ornithine transcarbamylase deficiency. *N Engl J Med* 1996 Sep 19; 335(12):855-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Magai C, Cohen CI, Gomberg D. Impact of training dementia caregivers in sensitivity to nonverbal emotion signals. *Int Psychogeriatr* 2002 Mar; 14(1):25-38.

Status: Not included because does not meet criteria for treatment for dementia patients

Magnus RV. A controlled trial of chlormethiazole in the management of symptoms of the organic dementias in the elderly. *Clin Ther* 1978; 1(6):387-96.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Maidment R, Livingston G, Katona C. 'Just keep taking the tablets': Adherence to antidepressant treatment in older people in primary care. *Int J Geriatr Psychiatry* 2002; 17(8):752-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Mair RG, McEntee WJ. Cognitive enhancement in Korsakoff's psychosis by clonidine: A comparison with L-dopa and ephedrine.

Psychopharmacologia 1986; 88(3):374-80.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Malaguarnera M, Pistone G, Vinci M, et al. Tacrine treatment of Alzheimer's disease: Many expectations, few certainties.

Neuropsychobiology 1998 Nov; 38(4):226-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Malaret B. Use of music to decrease aggressive behaviours in people with dementia. *J Am Geriatr Soc* 1998; 46(12):1586.

Status: Not included because not a full article

Manes F, Jorge R, Morcuende M, et al. A controlled study of repetitive transcranial magnetic stimulation as a treatment of depression in the elderly. *Int Psychogeriatr* 2001; 13(2):225-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mann AH, Schneider J, Mozley CG, et al. Depression and the response of residential homes to physical health needs. *Int J Geriatr Psychiatry* 2000; 15(12):1105-12.

Status: Not included because does not meet criteria for treatment for dementia patients

Mann K, Gunther A, Stetter F, et al. Rapid recovery from cognitive deficits in abstinent alcoholics: A controlled test-retest study. *Alcohol Alcohol* 1999 Jul; 34(4):567-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mantero MA, Barbero M, Giannini R, et al. Acetyl-L-carnitine as a therapeutic agent for mental deterioration in geriatric patients. (Double-blind placebo controlled study). *New Trends in Clinical Neuropharmacology* 1989; (1):17-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marder K, Tang MX, Alfaro B, et al. Postmenopausal estrogen use and Parkinson's disease with and without dementia. *Neurology* 1998 Apr; 50(4):1141-3.

Status: Not included because does not meet criteria for treatment for dementia patients

Marin DB, Bierer LM, Lawlor BA, et al. L-deprenyl and physostigmine for the treatment of Alzheimer's disease. *Psychiatry Res* 1995 Oct 16; 58(3):181-9.

Status: Cross-over trial;

Marini G, Caratti C, Peluffo F, et al. Placebo-controlled double-blind study of pramiracetam (CI-879) in the treatment of elderly subjects with memory impairment. *Adv Ther* 1992; 9(3):136-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marra CM, Boutin P, McArthur JC, et al. A pilot study evaluating ceftriaxone and penicillin G as treatment agents for neurosyphilis in human immunodeficiency virus-infected individuals. *Clin Infect Dis* 2000 Mar; 30(3):540-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marriott A, Donaldson C, Tarrier N, et al. Effectiveness of cognitive-behavioural family intervention in reducing the burden of care in carers of patients with Alzheimer's disease. *Br J Psychiatry* 2000 Jun; 176:557-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Marsh L, Lyketsos C, Reich SG. Olanzapine for the treatment of psychosis in patients with Parkinson's disease and dementia. *Psychosomatics* 2001 Nov; 42(6):477-81.

Status: Not included because dementia population not randomized to treatment

Martignoni E, Bono G, Blandini F, et al. Monoamines and related metabolite levels in the cerebrospinal fluid of patients with dementia of Alzheimer type. Influence of treatment with L-deprenyl. *J Neural Transm Park Dis Dement Sect* 1991; 3(1):15-25.

Status: Not included because dementia population not randomized to treatment

Martin JC. Effect of a synthetic peptide, ORG 2766, on inpatients with severe senile dementia. *Acta Psychiatr Scand* 1983; 67(3):205-7.

Status: Cross-over trial;

Martin PR, Adinoff B, Eckardt MJ, et al. Effective pharmacotherapy of alcoholic amnestic disorder with fluvoxamine. Preliminary findings. *Arch Gen Psychiatry* 1989 Jul; 46(7):617-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Martin PR, Adinoff B, Lane E, et al. Fluvoxamine treatment of alcoholic amnestic disorder. *Eur Neuropsychopharmacol* 1995 Mar; 5(1):27-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Martin RM, Rink E, Wilkinson DG, et al. Did knowledge, opinions, background, and health authority advice influence early prescribing of the novel Alzheimer's disease drug donepezil in general practice? - National postal survey. *Pharmacoepidemiol Drug Saf* 1999; 8(6):413-22.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Martinez Perez JA, Chavida GF, Sanchez-Seco HP, et al. Epidemiology of cognitive impairment in Spain. *Eur J Gen Pract* 2000; 6(2):52-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Marx J. Alzheimer's congress. Drug shows promise for advanced disease. *Science* 2000 Jul 21; 289(5478):375-7.

Status: Not included because not a full article

Marx JL. Alzheimer's drug trial put on hold. *Science* 1987; 238(4830):1041-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Masaki KH, Losonczy KG, Izmirlian G, et al. Association of vitamin E and C supplement use with cognitive function and dementia in elderly men. *Neurology* 2000; 54(6):1265-72.

Status: Not included because dementia population not randomized to treatment

Masuda Y, Akagawa Y, Hishikawa Y. Effect of serotonin 1A agonist tandospirone on depression symptoms in senile patients with dementia. *Hum Psychopharmacol* 2002 Jun; 17(4):191-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Matthews HP, Korbey J, Wilkinson DG, et al. Donepezil in Alzheimer's disease: Eighteen month results from Southampton Memory Clinic. *Int J Geriatr Psychiatry* 2000 Aug; 15(8):713-20.

Status: Not included because dementia population not randomized to treatment

Maurer I, Moller HJ, Saletu B. Treatment with propentofylline in dementia. *Pharmacopsychiatry*

1993; 26:179.

Status: Not included because not a full article

McCaffrey RJ, Steckler RA, Gansler DA, et al. An experimental evaluation of the efficacy of suloctidil in the treatment of primary degenerative dementia. *Arch Clin Neuropsychol* 1987; (2):155-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McCallion P, Toseland RW, Freeman K. An evaluation of a family visit education program. *J Am Geriatr Soc* 1999 Feb; 47(2):203-14.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McCarten JR, Kovera C, Maddox MK, et al. Triazolam in Alzheimer's disease: Pilot study on sleep and memory effects. *Pharmacol Biochem Behav* ; 52(2):447-52.

Status: Not included because dementia population not randomized to treatment

McClendon MJ, Bass DM, Brennan PF, et al. A computer network for Alzheimer's caregivers and use of support group services. *J Ment Health Aging* 1998; (4):403-20.

Status: Not included because dementia population not randomized to treatment

McConnachie RW. A clinical trial comparing 'Hydergine' with placebo in the treatment of cerebrovascular insufficiency in elderly patients. *Curr Med Res Opin* 1973; 1(8):463-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McConnachie RW. The clinical assessment of brain failure in the elderly. *Pharmacology* 1978; Vol 16(Suppl 1):27-35.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McCusker J, Verdon J, Tousignant P, et al. Rapid emergency department intervention for older people reduces risk of functional decline: Results of a multicenter randomized trial. *J Am Geriatr Soc* 2001; 49(10):1272-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McDonald C, Mowbray RM, Wilson JM. A sequential trial of amylobarbitone sodium used as sedation for confused female psychogeriatric patients. *Gerontol Clin (Basel)* 1970; 12(6):335-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McEntee WJ, Crook TH, Jenkyn LR, et al. Treatment of age-associated memory impairment with guanfacine. *Psychopharmacol Bull* 1991; 27(1):41-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McGeer PL, McGeer EG. Anti-inflammatory drugs in the fight against Alzheimer's disease. *Ann N Y Acad Sci* 1996; 777:213-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McKeith I, Del Ser T, Anand Rao. Rivastigmine provides symptomatic benefit in dementia with lewy bodies: Findings from a placebo-controlled international multicenter study. *Neurology* 2000; 54(Suppl 3):A450.

Status: Not included because not a full article

McKeith I, Del Ser T, Anand R. Erratum: Rivastigmine provides symptomatic benefit in dementia with lewy bodies: Findings from a placebo-controlled international multicenter study. *Expert Opin Pharmacother* 2001; 2(5):907.

Status: Not included because not a full article

McKeith IG, Grace JB, Walker Z, et al. Rivastigmine in the treatment of dementia with Lewy bodies: Preliminary findings from an open trial. *Int J Geriatr Psychiatry* 2000 May; 15(5):387-92.

Status: Not included because dementia population not randomized to treatment

McLachlan DR, Smith WL, Kruck TP. Desferrioxamine and Alzheimer's disease: Video home behavior assessment of clinical course and measures of brain aluminum. *Ther Drug Monit* 1993 Dec; 15(6):602-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McLean A, Jr., Stanton KM, Cardenas DD, et al. Memory training combined with the use of oral physostigmine. *Brain Inj* 1987 Oct; 1(2):145-59.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McLean A, Jr., Cardenas DD, Burgess D, et al. Placebo-controlled study of pramiracetam in young males with memory and cognitive problems resulting from head injury and anoxia. *Brain Inj*

1991 Oct; 5(4):375-80.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McLennan J, Findlay DJ, Sharma J, et al. Prolactin response to withdrawal of thioridazine in dementia. *Int J Geriatr Psychiatry* 1992; 7(10):739-42.

Status: Not included because Jadad Quality Scale score less than three

McNamara MJ, Gomez-Isla T, Hyman BT. Apolipoprotein E genotype and deposits of Abeta40 and Abeta42 in Alzheimer disease. *Arch Neurol* 1998 Jul; 55(7):1001-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

McNeil JK. Neuropsychological characteristics of the dementia syndrome of depression: Onset, resolution, and three-year follow-up. *Clin Neuropsychol* 1999; 13(2):136-46.

Status: Not included because does not meet criteria for treatment for dementia patients

Mcpherson A, Furniss FG, Sdogati C, et al. Effects of individualized memory aids on the conversation of persons with severe dementia a pilot study. *Aging Ment Health* 2001; 5(3):289-94.

Status: Not included because dementia population not randomized to treatment

McRae T, Griesing T, Whalen E. Donepezil and sertraline for the management of behavioral symptoms in patients with Alzheimer's disease. *Neurology* 2000; 54(Suppl 3):A416.

Status: Not included because not a full article

Mead MG, Castleden CM. Confusion and hypnotics in demented patients. *J R Coll Gen Pract* 1982 Dec; 32(245):763-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meador K, Loring D, Nichols M, et al. Preliminary findings of high-dose thiamine in dementia of Alzheimer's type. *J Geriatr Psychiatry Neurol* 1993 Oct; 6(4):222-9.

Status: Cross-over trial;

Mecocci P, Grossi E, Buscema M, et al. Use of artificial networks in clinical trials: A pilot study to predict responsiveness to donepezil in Alzheimer's disease. *J Am Geriatr Soc* 2002 Nov; 50(11):1857-60.

Status: Not included because dementia population not randomized to treatment

Medina A, Bodick N, Goldberger AL, et al. Effects of central muscarinic-1 receptor stimulation on blood pressure regulation. *Hypertension* 1997 Mar; 29(3):828-34.

Status: Not included because dementia population not randomized to treatment

Mega MS, Dinov ID, Lee L, et al. Orbital and dorsolateral frontal perfusion defect associated with behavioral response to cholinesterase inhibitor therapy in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 2000; 12(2):209-18.

Status: Not included because dementia population not randomized to treatment

Mega MS, Cummings JL, O'Connor SM, et al. Cognitive and metabolic responses to metrifonate therapy in Alzheimer disease. *Neuropsychiatry Neuropsychol Behav Neurol* 2001 Jan; 14(1):63-8.

Status: Not included because dementia population not randomized to treatment

Meier DE, Ahronheim JC, Morris J, et al. High short-term mortality in hospitalized patients with advanced dementia: Lack of benefit of tube feeding. *Arch Intern Med* 2001; 161(4):594-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meier U, Kintzel D. Clinical experiences with different valve systems in patients with normal-pressure hydrocephalus: Evaluation of the Miethke dual-switch valve. *Childs Nerv Syst* 2002 Jul; 18(6-7):288-94.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Melin L, Gotestam KG. Effects of rearranging Ward routines on communication and eating behaviors of psychogeriatric patients. *J Appl Behav Anal* 1981; (No. 1):47-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mellow AM, Sunderland T, Cohen RM, et al. Acute effects of high-dose thyrotropin releasing hormone infusions in Alzheimer's disease. *Psychopharmacologia* 1989; 98(3):403-7.

Status: Cross-over trial;

Mendez MF, Perryman KM. Neuropsychiatric features of frontotemporal dementia: Evaluation of

consensus criteria and review. J Neuropsychiatry Clin Neurosci 2002; 14(4):424-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Merrick BJ. A geriatric care management approach to a treatment plan for dementia. J Geriatr Psychiatry 2001; 34(2):233-45.
Status: Not included because dementia population not randomized to treatment

Mervis RJ, Ganzell S, Fitten LJ, et al. Comparison of Carbamazepine and Trazodone in the control of aggression / agitation in demented institutionalized patients. J Am Geriatr Soc 1991; 39(8):A75
Status: Not included because not a full article

Meszaros Z, Borcsiczky D, Mate M, et al. Platelet MAO-B activity and serotonin content in patients with dementia: Effect of age, medication, and disease. Neurochem Res 1998 Jun; 23(6):863-8.
Status: Not included because does not meet criteria for treatment for dementia patients

Meyer JS, Rogers RL, McClintic K, et al. Controlled clinical trial of daily aspirin therapy in multi-infarct dementia. Stroke 1988; 19(1):148
Status: Not included because not a full article

Meyer JS, Rogers RL, McClintic K, et al. Randomized clinical trial of daily aspirin therapy in multi-infarct dementia. A pilot study. J Am Geriatr Soc 1989 Jun; 37(6):549-55.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meyer JS, Lotfi J, Martinez G, et al. Effects of medical and surgical treatment on cerebral perfusion and cognition in patients with chronic cerebral ischemia. Surg Neurol 1990 Nov; 34(5):301-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Meyer JS, Li Y, Xu G, et al. Feasibility of treating mild cognitive impairment with cholinesterase inhibitors. Int J Geriatr Psychiatry 2002 Jun; 17(6):586-8.
Status: Not included because dementia population not randomized to treatment

Meythaler JM, Depalma L, Devivo MJ, et al. Sertraline to improve arousal and alertness in severe traumatic brain injury secondary to motor vehicle crashes. Brain Inj 2001 Apr; 15(4):321-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Micca JL, Sky AJ, Uhrig-Hitchcock LG. Quality care: A practical guide to managing behavioral symptoms of dementia. Journal of the American Medical Directors Association 2002; 3(Suppl 4):H21-H25
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miceli G, Caltagirone C, Gainotti G. Gangliosides in the treatment of mental deterioration. A double-blind comparison with placebo. Acta Psychiatr Scand 1977 Feb; 55(2):102-10.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mielke R, Moller HJ, Erkinjuntti T, et al. Propentofylline in the Treatment of Vascular Dementia and Alzheimer-Type Dementia: Overview of Phase I and Phase II Clinical Trials. Alzheimer Dis Assoc Disord 1998; 12:S29-S35.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miguel-Hidalgo JJ. Rivastigmine Novartis AG. Curr Opin Cent Peripher Nerv Syst Invest Drugs 2000; 2(4):438-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Milders M, Deelman B, Berg I. Rehabilitation of memory for people's names. Memory 1998 Jan; 6(1):21-36.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miller E. On the nature of the memory disorder in presenile dementia. Neuropsychologia 1971; 9(1):75-81.
Status: Not included because does not meet criteria for treatment for dementia patients

Miller E. Efficiency of coding and the short-term memory defect in presenile dementia. Neuropsychologia 1972; 10(1):133-6.
Status: Not included because does not meet criteria for treatment for dementia patients

Miller IW, Norman WH, Keitner GI. Treatment response of high cognitive dysfunction depressed inpatients. Compr Psychiatry 1990 Jan; 31(1):62-71.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miller PA, Butin D. The role of occupational therapy in dementia - C.O.P.E. (Caregiver Options for Practical Experiences). *Int J Geriatr Psychiatry* 2000; 15(1):86-9.

Status: Not included because dementia population not randomized to treatment

Miller R, Newcomer R, Fox P. Effects of the Medicare Alzheimer's disease demonstration on nursing home entry. *Health Serv Res* 1999 Aug; 34(3):691-714.

Status: Not included because does not meet criteria for treatment for dementia patients

Miller RG. Leptazol and meso-inositol hexanicotinate in the treatment of chronic cerebrovascular degenerative disorders. A double blind study. *Gerontol Clin (Basel)* 1995 Oct; 5:95-102.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Miller TP, Fong K, Tinklenberg JR. An ACTH 4-9 analog (Org 2766) and cognitive performance: High-dose efficacy and safety in dementia of the Alzheimer's type. *Biol Psychiatry* 1993 Feb 15; 33(4):307-9.

Status: Not included because no extractable data relevant to review

Minger SL, Esiri MM, McDonald B, et al. Cholinergic deficits contribute to behavioral disturbance in patients with dementia. *Neurology* 2000; 55(10):1460-7.

Status: Not included because does not meet criteria for treatment for dementia patients

Minthon L, Gustafson L, Dalfelt G, et al. Oral tetrahydroaminoacridine treatment of Alzheimer's disease evaluated clinically and by regional cerebral blood flow and EEG. *Dementia* 1993 Jan; 4(1):32-42.

Status: Cross-over trial;

Minthon L, Edvinsson L, Gustafson L. Tacrine treatment modifies cerebrospinal fluid neuropeptide levels in Alzheimer's disease. *Dementia* 1994 Nov; 5(6):295-301.

Status: Not included because no extractable data relevant to review

Minthon L, Nilsson K, Edvinsson L, et al. Long-term effects of tacrine on regional cerebral blood flow changes in Alzheimer's disease. *Dementia* 1995 Sep; 6(5):245-51.

Status: Not included because dementia population not randomized to treatment

Mintzer JE, Brawman MO, Mirski DF, et al. Anxiety in the behavioral and psychological symptoms of dementia. *Int Psychogeriatr* 2000; 12(Suppl 1):139-42.

Status: Not included because dementia population not randomized to treatment

Mintzer JE, Madhusoodanan S, Brenner Ronald. Risperidone in dementia. *Psychiatr Ann* 2000; 30(3):181-7.

Status: Not included because dementia population not randomized to treatment

Mishima K, Okawa M, Hishikawa Y, et al. Morning bright light therapy for sleep and behavior disorders in elderly patients with dementia. *Acta Psychiatr Scand* 1994; 89(1):1-7.

Status: Not included because dementia population not randomized to treatment

Mishima K, Hishikawa Y, Okawa M. Randomized, dim light controlled, crossover test of morning bright light therapy for rest-activity rhythm disorders in patients with vascular dementia and dementia of Alzheimer's type. *Chronobiol Int* 1998 Nov; 15(6):647-54.

Status: Cross-over trial;

Mitchell A, Drachman DA, O'Connell B, et al. Oral physostigmine in Alzheimer's disease. *Neurology* 1986; 36(Suppl 1):295.

Status: Not included because not a full article

Mitchell A, Maercklein LA. The effect of individualized special instruction on the behaviors of nursing home residents diagnosed with dementia. *Am J Alzheimers Dis* 1996; (1):23

Status: Not included because does not meet criteria for treatment for dementia patients

Mitchell S. Aromatherapy's Effectiveness in Disorders Associated With Dementia. *Int J Aromatherapy* 1993; (2):20-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mittelman MS, Ferris SH, Steinberg G, et al. An intervention that delays institutionalization of Alzheimer's disease patients: Treatment of spouse-caregivers. *Gerontologist* 1993 Dec; 33(6):730-40.

Status: Not included because does not meet criteria for treatment for dementia patients

Mittelman MS, Ferris SH, Shulman E, et al. A family intervention to delay nursing home placement of patients with Alzheimer disease. A randomized controlled trial. *JAMA* 1996 Dec 4; 276(21):1725-31.

Status: Not included because does not meet criteria for treatment for dementia patients

Miyamoto Y, Ito H, Otsuka T, et al. Caregiver burden in mobile and non-mobile demented patients: A comparative study. *Int J Geriatr Psychiatry* 2002 Aug; 17(8):765-73.

Status: Not included because does not meet criteria for treatment for dementia patients

Moffoot A, O'Carroll RE, Murray C, et al. Clonidine infusion increases uptake of 99mTc-Exametazime in anterior cingulate cortex in Korsakoff's psychosis. *Psychol Med* 1994 Feb; 24(1):53-61.

Status: Not included because dementia population not randomized to treatment

Moglia A, Arrigo A, Bono G. Citicoline in patients with chronic cerebrovascular diseases (CCVD): Quantitative EEG study. *Curr Ther Res Clin Exp* 1984; 36(2):309-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohr E, Bruno G, Foster N, et al. GABA-agonist therapy for Alzheimer's disease. *Clin Neuropharmacol* 1986; 9(3):257-63.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohr E, Schlegel J, Fabbrini G, et al. Clonidine treatment of Alzheimer's disease. *Arch Neurol* 1989 Apr; 46(4):376-8.

Status: Cross-over trial;

Mohr E, Knott V, Sampson M, et al. Cognitive and quantified electroencephalographic correlates of cycloserine treatment in Alzheimer's disease. *Clin Neuropharmacol* 1995 Feb; 18(1):28-38.

Status: Not included because Jadad Quality Scale score less than three

Mohr E, Walker D, Randolph C, et al. Utility of clinical trial batteries in the measurement of Alzheimer's and Huntington's dementia. *Int Psychogeriatr* 1996; 8(3):397-411.

Status: Not included because does not meet criteria for treatment for dementia patients

Mohs R, Doody R, Morris J, et al. Donepezil preserves functional status in Alzheimer's disease patients: Results from a 1-year prospective placebo-controlled attrition study. *J Eur Coll Neuropsychopharmacol* 1999; 9(Suppl 5):S328
Status: Not included because not a full article

Mohs R, Doody R, Morris J, et al. Donepezil preserves activities of daily living in Alzheimer's disease patients results from a one-year placebo-controlled functional survival study. *Neurology* 2000 Apr; 54(Suppl 3):1

Status: Not included because not a full article

Mohs RC, Davis KL. A signal detectability analysis of the effect of physostigmine on memory in patients with Alzheimer's disease. *Neurobiol Aging* 1982; 3(2):105-10.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Davis BM, Johns CA, et al. Oral physostigmine treatment of patients with Alzheimer's disease. *Am J Psychiatry* 1985 Jan; 142(1):28-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Davis BM, Greenwald BS. Clinical studies of the cholinergic deficit in Alzheimer's disease. II. Psychopharmacologic studies. *J Am Geriatr Soc* 1985; 33(11):749-57.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mohs RC, Ferris SH. Measuring response to treatment in Alzheimer's disease: What constitutes meaningful change? *Int J Geriatr Psychopharmacol* 1998; 1(Suppl 1):S7-S14.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Molchan SE, Mellow AM, Hill JL, et al. The effects of thyrotropin-releasing hormone and scopolamine in Alzheimer's disease and normal volunteers. *J Psychopharmacol (Oxf)* 1992; 6(4):489-500.

Status: Cross-over trial;

Molchan SE, Manji H, Chen G, et al. Effects of chronic lithium treatment on platelet PKC isozymes in Alzheimer's and elderly control subjects. *Neurosci Lett* 1993; 162(1-2):187-2.

Status: Not included because dementia population not randomized to treatment

Molchan SE, Hill JL, Minichiello M, et al. Scopolamine effects on the pressor response to thyrotropin-releasing hormone in humans. *Life Sci* 1994; 54(13):933-8.

Status: Not included because no extractable data relevant to review

Molloy DW, Cape RD. Acute effects of oral pyridostigmine on memory and cognitive function in SDAT. *Neurobiol Aging* 1989 Mar; 10(2):199-204.

Status: Cross-over trial;

Molloy DW, Guyatt GH, Wilson DB, et al. Effect of tetrahydroaminoacridine on cognition, function and behaviour in Alzheimer's disease. *CMAJ* 1991 Jan 1; 144(1):29-34.

Status: Cross-over trial;

Molloy DW, Guyatt G, Brown G. Effects of oxiracetam on cognition, behaviour and activities of daily living in dementia. *J Clin Exp Gerontol* 1992; (3-4):217-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Molloy DW, Guyatt GH, Standish T, et al. Effect of a new nootropic agent, CGS 5649B, on cognition, function, and behavior in dementia. *J Gen Intern Med* 1993 Aug; 8(8):444-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Molloy DW, Standish TI. Clinical experience with Cerebrolysin. *J Neural Transm Suppl* 2000; 59:293-300.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Moniz-Cook E, Agar S, Gibson G, et al. A preliminary study of the effects of early intervention with people with dementia and their families in a memory clinic. *Aging Ment Health* 1998 Aug; 2(3):199-211.

Status: Not included because dementia population not randomized to treatment

Monreal M, Lafoz E, Olive A, et al. Comparison of subcutaneous unfractionated heparin with a low molecular weight heparin (Fragmin (R)) in patients with venous thromboembolism and contraindications to coumarin. *Thromb Haemost* 1994; 71(1):7-11.

Status: Not included because does not meet criteria for treatment for dementia patients

Monteverde A, Gnemmi P, Rossi F, et al. Selegiline in the treatment of mild to moderate Alzheimer-type dementia. *Clin Ther* 1990 Jul; 12(4):315-22.

Status: Not included because Jadad Quality Scale score less than three

Montgomery P, Erickson GK. Neuropsychological perspectives in amyotrophic lateral sclerosis. *Neurol Clin* 1987; 5(1):61-81.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Moore MJ, Clipp EC. Alzheimer's disease and caregiver time. *Lancet* 1994 Jan 22; 343(8891):239-40.

Status: Not included because not a full article

Moore S, Sandman CA, McGrady K, et al. Memory training improves cognitive ability in patients with dementia. *Neuropsychol Rehab* 2001; 11(3-4):245-61.

Status: Not included because does not meet criteria for treatment for dementia patients

Moretti R, Torre P, Antonello RM, et al. Rivastigmine in subcortical vascular dementia: A comparison trial on efficacy and tolerability for 12 months follow-up. *Eur J Neurol* 2001 Jul; 8(4):361-2.

Status: Not included because not a full article

Moretti R, Torre P, Antonello RM, et al. Depression and Alzheimer's disease: Symptom or comorbidity? *Am J Alzheimers Dis Other Dement* 2002; 17(6):338-44.

Status: Not included because Jadad Quality Scale score less than three

Moretti R, Torre P, Antonello RM, et al. An open-label pilot study comparing rivastigmine and low-dose aspirin for the treatment of symptoms specific to patients with subcortical vascular dementia. *Curr Ther Res Clin Exp* 2002; 63(7):443-58.

Status: Not included because dementia population not randomized to treatment

Moretti R, Torre P, Antonello RM, et al. Frontotemporal dementia: Paroxetine as a possible treatment of behavior symptoms: A randomized, controlled, open 14-month study. *Eur Neurol* 2003; 49(1):13-9.

Status: Not included because Jadad Quality Scale score less than three

Morey LC, Ban TA, Cassano G, et al. Glycosaminoglycan polysulfate in old-age dementias: A factor-analytic study of change in psychopathologic symptoms. *Neuropsychobiology* 1988; 19(3):135-8.

Status: Not included because dementia population not randomized to treatment

Morganroth J, Graham S, Hartman R, et al. Electrocardiographic effects of rivastigmine. *J Clin Pharmacol* 2002 May; 42(5):558-68.

Status: Not included because does not meet criteria for treatment for dementia patients

Mori T, Inoue D, Kosugi S, et al. Effects of low dose L-triiodothyronine administration on mental, behavioural and thyroid states in elderly subjects. *Endocrinol Jpn* 1988 Aug; 35(4):585-92.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Morich FJ, Bieber F, Lewis JM, et al. Nimodipine in the treatment of probable Alzheimer's disease: Results of two multicentre trials. *Clin Drug Investig* 1996; (4):185-95.

Status: Not included because dementia population not randomized to treatment

Moroney JT, Tseng C-L, Paik MC, et al. Treatment for the secondary prevention of stroke in older patients: The influence of dementia status. *J Am Geriatr Soc* 1999; 47(7):824-9.

Status: Not included because dementia population not randomized to treatment

Moroney JT, Tang MX, Berglund L, et al. Low-density lipoprotein cholesterol and the risk of dementia with stroke. *JAMA* 1999; 282(3):254-60.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Morrison RS, Siu AL. A comparison of pain and its treatment in advanced dementia and cognitively intact patients with hip fracture. *J Pain Symptom Manage* 2000 Apr; 19(4):240-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Moss DE, Berlanga P, Hagan MM, et al. Methanesulfonyl fluoride (MSF): A double-blind, placebo-controlled study of safety and efficacy in the treatment of senile dementia of the Alzheimer type. *Alzheimer Dis Assoc Disord* 1999 Jan; 13(1):20-5.

Status: Not included because Jadad Quality Scale score less than three

Mouradian MM, Mohr E, Williams JA, et al. No response to high-dose muscarinic agonist therapy in Alzheimer's disease. *Neurology* 1988 Apr; 38(4):606-8.

Status: Not included because dementia population not randomized to treatment

Mouradian MM, Blin J, Giuffra M, et al. Somatostatin replacement therapy for Alzheimer dementia. *Ann Neurol* 1991 Oct; 30(4):610-3.

Status: Not included because dementia population not randomized to treatment

Mubrin Z, Knezevic S, Spilich G, et al. Normalization of rCBF pattern in senile dementia of the Alzheimer's type. *Psychiatry Res* 1989 Sep; 29(3):303-6.

Status: Not included because no extractable data relevant to review

Mucke HAM. Metrifonate. Treatment of Alzheimer's disease, acetylcholinesterase inhibitor. *Drugs of the Future* 1998; 23(5):491-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Mulsant BH, Mazumdar S, Pollock BG, et al. Methodological issues in characterizing treatment response in demented patients with behavioral disturbances. *Int J Geriatr Psychiatry* 1997 May; (No. 5):537-47.

Status: Not included because dementia population not randomized to treatment

Munch-Petersen S, Pakkenberg H, Kornerup H, et al. RNA treatment of dementia. A double-blind study. *Acta Neurol Scand* 1974; 50(5):553-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Murali DP, Kaiser L. Variability of the mini-mental state examination in dementia. *Neurology* 2000 Apr 11; 54(7):1538-9.

Status: Not included because not a full article

Muramoto O, Sugishita M, Sugita H, et al. Effect of physostigmine on constructional and memory tasks in Alzheimer's disease. *Arch Neurol* 1979 Aug; 36(8):501-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Muramoto O, Sugishita M, Ando K. Cholinergic system and constructional praxis: A further study of physostigmine in Alzheimer's disease. *J Neurol*

Neurosurg Psychiatry 1984; 47(5):485-91.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Muratorio A, Bonuccelli U, Nuti A, et al. A neurotropic approach to the treatment of multi-infarct dementia using L-alpha-glycerylphosphorylchlorine. Curr Ther Res Clin Exp 1992; 52(5):741-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Muresanu DF, Rainer M, Moessler H. Improved global function and activities of daily living in patients with AD: a placebo-controlled clinical study with the neurotrophic agent Cerebrolysin. J Neural Transm Suppl 2002; (62):277-85.

Status: Not included because Jadad Quality Scale score less than three

Murialdo G, Barreca A, Nobili F, et al. Dexamethasone effects on cortisol secretion in Alzheimer's disease: Some clinical and hormonal features in suppressor and nonsuppressor patients. J Endocrinol Invest 2000 Mar; 23(3):178-86.

Status: Not included because does not meet criteria for treatment for dementia patients

Murri L, Bardi C, Arena R. A comparison between lorazepam and flunitrazepam in insomniac patients affected by chronic cerebrovascular disorders. Curr Ther Res Clin Exp 1984; 35(1):113-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nadeau SE, Malloy PF, Andrew ME. A crossover trial of bromocriptine in the treatment of vascular dementia. Ann Neurol 1988 Aug; 24(2):270-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nagaraja D, Jayashree S. Randomized study of the dopamine receptor agonist piribedil in the treatment of mild cognitive impairment. Am J Psychiatry 2001 Sep; 158(9):1517-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nair NP, Ban TA, Hontela S, et al. Trazodone in the treatment of organic brain syndromes, with special reference to psychogeriatrics. Curr Ther Res Clin Exp ; 15(10):769-75.

Status: Not included because dementia population not randomized to treatment

Nakamura H, Nakanishi M, Hamanaka T, et al. Semantic priming in patients with Alzheimer and semantic dementia. Cortex 2000 Apr; 36(2):151-62.

Status: Not included because does not meet criteria for treatment for dementia patients

Nakamura K. Aniracetam: Its novel therapeutic potential in cerebral dysfunctional disorders based on recent pharmacological discoveries. CNS Drug Rev 2002; 8(1):70-89.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nakano S, Asada T, Matsuda H, et al. Donepezil hydrochloride preserves regional cerebral blood flow in patients with Alzheimer's disease. J Nucl Med 2001 Oct; 42(10):1441-5.

Status: Not included because Jadad Quality Scale score less than three

Namazi KH, Haynes SR. Sensory stimuli reminiscence for patients with Alzheimer's disease: Relevance and implications. Clin Gerontol 1994; 29-46.

Status: Not included because dementia population not randomized to treatment

Namazi KH, Johnson BD. Environmental effects on incontinence problems in Alzheimer's disease patients. Am J Alzheimers Care Relat Disord 1991 Nov; (6):16-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Naritomi H, Murata S, Shimizu T, et al. Long-term effects of bifemelane hydrochloride on post-stroke deterioration of cognitive function and cerebral blood flow. Curr Ther Res Clin Exp 1995; Vol 56(231):238

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nasrallah HA, Varney N, Coffman JA, et al. Effects of naloxone on cognitive deficits following electroconvulsive therapy. Psychopharmacol Bull 1985; 21(1):89-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Navia BA, Dafni U, Simpson D, et al. A phase I/II trial of nimodipine for HIV-related neurologic complications. Neurology 1998 Jul; 51(1):221-8.

Status: Not included because dementia population not randomized to treatment

Nebes RD, Pollock BG, Mulsant BH, et al. Cognitive effects of paroxetine in older depressed patients. *J Clin Psychiatry* 1999; 60(Suppl):9
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Netz Y, Yaretzki A, Salganik I, et al. The effect of supervised physical activity on cognitive and affective state of geriatric and psychogeriatric in-patients. *Clin Gerontol* 1994; 15(1):47-56.
Status: Not included because does not meet criteria for treatment for dementia patients

Neumeister A, Graf A, Willeit M, et al. Effects of Light Therapy in dementia. *Biol Psychiatry* 2000; Vol 47:S536
Status: Not included because not a full article

Newcomer R, Spitalny M, Fox P, et al. Effects of the Medicare Alzheimer's disease demonstration on the use of community-based services. *Health Serv Res* 1999 Aug; 34(3):645-67.
Status: Not included because does not meet criteria for treatment for dementia patients

Newhouse PA, Sunderland T, Tariot PN, et al. Intravenous nicotine in Alzheimer's disease: A pilot study. *Psychopharmacologia* 1988; 95(2):171-5.
Status: Not included because dementia population not randomized to treatment

Newhouse PA, Sunderland T, Narang PK, et al. Neuroendocrine, physiologic, and behavioral responses following intravenous nicotine in nonsmoking healthy volunteers and in patients with Alzheimer's disease. *Psychoneuroendocrinology* 1990; 15(5-6):471-6.
Status: Not included because dementia population not randomized to treatment

Niemann H, Ruff RM, Baser CA. Computer-assisted attention retraining in head-injured individuals: A controlled efficacy study of an outpatient program. *J Consult Clin Psychol* 1990 Dec; 58(6):811-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nijhuis E, Hinloopen B, van Duijn C, et al. Decreased sensitivity to dexamethasone in lymphocytes from patients with Alzheimer's disease. *Clin Immunol Immunopathol* 1994 Oct; 73(1):45-52.

Status: Not included because does not meet criteria for treatment for dementia patients

Nikolova G, Traykov L. Efficacy of donepezil in patients with Alzheimer's disease - Results of 12-week open clinical trial. *Acta Med Bulg* 2001; 28:70-5.
Status: Not included because dementia population not randomized to treatment

Nimodipine Clinical Study Group. Effects of Nimodipine on vascular dementia. *Cardiovasc Dis* 1996; 6(Suppl 2):70.
Status: Not included because not a full article

Nirenberg TD. Relocation of institutionalized elderly. *J Consult Clin Psychol* 1983 Oct; 51(5):693-701.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nobili F, Vitali P, Canfora M, et al. Effects of long-term Donepezil therapy on rCBF of Alzheimer's patients. *Clin Neurophysiol* 2002; 113(8):1241-8.
Status: Not included because dementia population not randomized to treatment

Nobili F, Koulibaly M, Vitali P, et al. Brain perfusion follow-up in Alzheimer's patients during treatment with acetylcholinesterase inhibitors. *J Nucl Med* 2002 Aug; 43(8):983-90.
Status: Not included because dementia population not randomized to treatment

Noel G, Jeanmart M, Reinhardt B. Treatment of the organic brain syndrome in the elderly. A double-blind comparison on the effects of a neurotropic drug and placebo. *Neuropsychobiology* 1983; 10(2-3):90-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Norbergh KG, Hellzen O, Sandman PO, et al. The relationship between organizational climate and the content of daily life for people with dementia living in a group-dwelling. *J Clin Nurs* 2002 Mar; 11(2):237-46.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Novo FP, Ryan RP, Frazier EL. Dihydroergotoxine mesylate in treatment of symptoms of idiopathic cerebral dysfunction in geriatric patients. *Clin Ther* 1978; 1(5):359-69.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nyback H, Hassan M, Junthe T, et al. Clinical experiences and biochemical findings with tacrine (THA). *Acta Neurol Scand Suppl* 1993; 88(149):36-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nygaard H, Bakke K, Brudvik E, et al. Zuclopenthixol and melperon in the treatment of elderly patients: A double-blind, controlled, multi-centre study. *Pharmatherapeutica* 1987; 5(3):152-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nygaard HA, Bakke K, Brudvik E, et al. Dosing of neuroleptics in elderly demented patients with aggressive and agitated behaviour: A double-blind study with zuclopenthixol. *Curr Med Res Opin* 1994; 13(4):222-32.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Nyth AL, Gottfries CG, Lyby K, et al. A controlled multicenter clinical study of citalopram and placebo in elderly depressed patients with and without concomitant dementia. *Acta Psychiatr Scand* 1992 Aug; 86(2):138-45.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Brien BJ, Goeree R, Hux M, et al. Economic evaluation of donepezil for the treatment of Alzheimer's disease in Canada. *J Am Geriatr Soc* 1999; 47(5):570-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Brien JT, Schweitzer I, Ames D, et al. The function of the hypothalamic-pituitary-adrenal axis in Alzheimer's disease. Response to insulin hypoglycaemia. *Br J Psychiatry* 1994 Nov; 165(5):650-7.

Status: Not included because dementia population not randomized to treatment

O'Carroll RE, Moffoot A, Ebmeier KP, et al. Korsakoff's syndrome, cognition and clonidine. *Psychol Med* 1993 May; 23(2):341-7.

Status: Not included because dementia population not randomized to treatment

O'Carroll RE, Moffoot AP, Ebmeier KP, et al. Effects of fluvoxamine treatment on cognitive functioning in the alcoholic Korsakoff syndrome.

Psychopharmacologia 1994 Sep; 116(1):85-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Connolly M, Dierdorf D, Greb WH, et al. Efficacy of denbufylline in patients with multi-infarct dementia. *Drug Dev Res* 1988; 14(3-4):195-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

O'Connor DW, Pollitt PA, Brook CP, et al. Does early intervention reduce the number of elderly people with dementia admitted to institutions for long term care? *BMJ* 1991; 302(6781):871-5.

Status: Not included because dementia population not randomized to treatment

O'Keeffe ST, Lavan JN. Subcutaneous fluids in elderly hospital patients with cognitive impairment. *Gerontology* 1996; 42(1):36-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Oakley F, Sunderland T. Assessment of motor and process skills as a measure of IADL functioning in pharmacologic studies of people with Alzheimer's disease: A pilot study. *Int Psychogeriatr* 1997 Jun; 9(2):197-206.

Status: Cross-over trial;

Obonsawin MC, Robertson A, Crawford JR, et al. Non-mnemonic cognitive function in the scopolamine model of Alzheimer's disease. *Hum Psychopharmacol* 1998; 13:439-49.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ohno Y. The effects of magnetized mineral water on memory loss delay in Alzheimer's disease. *The centre for frontier sciences* 1997; (6):38-43.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Oishi M, Mochizuki Y, Takasu T, et al. Effectiveness of traditional Chinese medicine in Alzheimer disease. *Alzheimer Dis Assoc Disord* 1998 Sep; 12(3):247-50.

Status: Not included because dementia population not randomized to treatment

Okajima Y. Quantitative pharmaco-EEG study of indeloxazine hydrochloride in psychogeriatric patients. *Integr Psychiatry* 1993; 9(1):25-33.

Status: Not included because dementia population not randomized to treatment

Olin JT, Schneider LS. Assessing response to tacrine using the factor analytic structure of the Alzheimer's Disease Assessment Scale (ADAS) - Cognitive subscale. *Int J Geriatr Psychiatry* 1995; 10(9):753-6.

Status: Not included because Jadad Quality Scale score less than three

Onofrj M, Thomas A, Luciano AL, et al. Donepezil versus vitamin E in Alzheimer's disease: Part 2: Mild versus moderate-severe Alzheimer's disease. *Clin Neuropharmacol* 2002 Jul; 25(4):207-15.

Status: Not included because Jadad Quality Scale score less than three

Opie J, Doyle C, O'Connor DW. Challenging behaviours in nursing home residents with dementia: A randomized controlled trial of multidisciplinary interventions. *Int J Geriatr Psychiatry* 2002 Jan; 17(1):6-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Orengo CA, Kidwell K, Kunik ME, et al. The effect of risperidone on cognitive performance in elderly psychotic and aggressive patients with dementia; a pilot study. *Int J Geriatr Psychopharmacol* 1998; 1(4):193-6.

Status: Not included because dementia population not randomized to treatment

Orten JD, Allen M, Cook J. Reminiscence Groups with Confused Nursing Center Residents: An Experimental Study. *Soc Work Health Care* 1989; 14(1):73-86.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ostwald SK, Hepburn KW, Caron W, et al. Reducing caregiver burden: A randomized psychoeducational intervention for caregivers of persons with dementia. *Gerontologist* 1999 Jun; 39(3):299-309.

Status: Not included because does not meet criteria for treatment for dementia patients

Ousset PJ, Viallard G, Puel M, et al. Lexical Therapy and episodic word learning in dementia of the Alzheimer type. *Brain Lang* 2002; 80(1):14-20.

Status: Not included because does not meet criteria for treatment for dementia patients

Owensworth TL, McFarland K. Memory remediation in long-term acquired brain injury: Two

approaches in diary training. *Brain Inj* 1999 Aug; 13(8):605-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Paire JA, Karney RJ. The effectiveness of sensory stimulation for geropsychiatric inpatients. *Am J Occup Ther* 1984 Aug; 38(8):505-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Palmer GC. Neuroprotection by NMDA receptor antagonists in a variety of neuropathologies. *Curr Drug Targets* 2001; 2(3):241-71.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Palmieri G, Palmieri R, Inzoli MR. Double-blind controlled trial of phosphatidylserine in patients with senile mental deterioration. *Clin Trials J* 1987; 24(1):73-83.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pantev M, Ritter R, Gortelmeyer R. Therapy responder analysis of dementia study with memantine. *Pharmacopsychiatry* 1993; 26:185.

Status: Not included because not a full article

Parkes JD, Marsden CD, Rees JE. Parkinson's disease, cerebral arteriosclerosis, and senile dementia. Clinical features and response to levodopa. *Q J Med* 1974; 43(169):49-61.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Ciuffetti G, Mercuri M, et al. Relationship between haemorheological factors and initial mental deterioration in the elderly. A preliminary study. *Clin Hemorheol* 1985; 5(4):361-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Ciuffetti G, Mercuri M, et al. Haemorheological pattern in initial mental deterioration: Results of a long-term study using piracetam and pentoxifylline. *Arch Gerontol Geriatr* 1985 Jul; 4(2):141-55.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Ciuffetti G, Mercuri M, et al. The role of haemorheological factors in the ageing brain: Long-term therapy with pentoxifylline ('Trental' 400) in elderly patients with initial mental

deterioration. *Pharmatherapeutica* 1986; 4(10):617-27.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Parnetti L, Senin U, Carosi M, et al. Mental deterioration in old age: results of two multicenter, clinical trials with nimodipine. The Nimodipine Study Group. *Clin Ther* 1993 Mar; 15(2):394-406.
Status: Not included because dementia population not randomized to treatment

Parnetti L, Abate G, Bartorelli L, et al. Multicentre study of l-alpha-glyceryl-phosphorylcholine vs ST200 among patients with probable senile dementia of Alzheimer's type. *Drugs Aging* 1993 Mar; 3(2):159-64.
Status: Not included because Jadad Quality Scale score less than three

Parnetti L, Amici S, Lanari A, et al. Cerebrospinal fluid levels of biomarkers and activity of acetylcholinesterase (AChE) and butyrylcholinesterase in AD patients before and after treatment with different AChE inhibitors. *Neurol Sci* 2002; 23(Suppl 2):S95-S96.
Status: Not included because not a full article

Partanen J, Wolters EC, van Duijn H. Clinical neurophysical correlates of vasopressin derivative therapy in Alzheimer's disease. *Clin Neurol Neurosurg* 1987; 2:34.
Status: Not included because not a full article

Passeri M, Cucinotta D, de Mello M, et al. Minaprine for senile dementia. *Lancet* 1985 Apr 6; 1(8432):824.
Status: Not included because not a full article

Passeri M, Cucinotta D. Ateroid in the clinical treatment of multi-infarct dementia. *Mod Probl Pharmacopsychiatry* 1989; 23:85-94.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Passeri M, Cucinotta D, Bonati PA, et al. Acetyl-L-carnitine in the treatment of mildly demented elderly patients. *Int J Clin Pharmacol Res* 1990; 10(1-2):75-82.
Status: Not included because Jadad Quality Scale score less than three

Paterson J, Hamilton MM, Grant H. The effectiveness of the Hierarchic Dementia Scale in tailoring interventions to reduce problem behaviours in people with Alzheimer's disease.

Aust Occup Ther J 2000; 47(3):134-40.
Status: Not included because does not meet criteria for treatment for dementia patients

Pathy J, Menon G, Reynolds A, et al. Betahistine hydrochloride (Serc) in cerebrovascular disease: A placebo-controlled study. *Age Ageing* 1977 Aug; 6(3):179-84.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Peabody CA, Thiemann S, Pigache R, et al. Desglycinamide-9-arginine-8-vasopressin (DGAVP, Organon 5667) in patients with dementia. *Neurobiol Aging* 1985; 6(2):95-100.
Status: Not included because dementia population not randomized to treatment

Peabody CA, Deblois TE, Tinklenberg JR. Thyrotropin-releasing hormone (TRH) and Alzheimer's disease. *Am J Psychiatry* 1986; 143(2):262-3.
Status: Not included because not a full article

Penn RD, Martin EM, Wilson RS, et al. Intraventricular bethanechol infusion for Alzheimer's disease: Results of double-blind and escalating-dose trials. *Neurology* 1988 Feb; 38(2):219-22.
Status: Cross-over trial;

Pepping J. Phosphatidylserine. *Am J Health Syst Pharm* 1999; 56(20):2038-44.
Status: Not included because not a full article

Pepping J. Alternative therapies. Huperzine A: A potent and selective acetylcholinesterase inhibitor. *Am J Health Syst Pharm* 2000 Mar 15; 57(6):530-4.
Status: Not included because dementia population not randomized to treatment

Perini M, Montanini R, Casucci R. Effects of eptastigmine: A new cholinesterase inhibitor on regional cerebral blood flow in Alzheimer patients. *J Neurol* 1995; 242:S57.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Perryman KM, Fitten LJ. Quantitative EEG during a double-blind trial of THA and lecithin in patients with Alzheimer's disease. *J Geriatr Psychiatry Neurol* 1991 Jul; 4(3):127-33.
Status: Not included because no extractable data relevant to review

Peskind ER, Wingerson D, Murray S, et al. Effects of Alzheimer's disease and normal aging on cerebrospinal fluid norepinephrine responses to yohimbine and clonidine. *Arch Gen Psychiatry* 1995 Sep; 52(9):774-82.

Status: Not included because does not meet criteria for treatment for dementia patients

Peters BH, Levin HS. Memory enhancement after physostigmine treatment in the amnesic syndrome. *Arch Neurol* 1977 Apr; 34(4):215-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Peters BH, Levin HS. Effects of physostigmine and lecithin on memory in Alzheimer's disease. *Ann Neurol* 1979 Sep; 6(3):219-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Peterson LG, Bongar B. Navane versus Haldol. Treatment of acute organic mental syndromes in the general hospital. *Gen Hosp Psychiatry* 1989 Nov; 11(6):412-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Petit D, Montplaisir J, Lorrain D, et al. THA does not affect sleep or EEG spectral power in Alzheimer's disease. *Biol Psychiatry* 1993; 33(10):753-4.

Status: Not included because dementia population not randomized to treatment

Petracca G, Teson A, Chemerinski E, et al. A double-blind placebo-controlled study of clomipramine in depressed patients with Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1996; 8(3):270-5.

Status: Cross-over trial;

Pettegrew JW, Klunk WE, Panchalingam K, et al. Clinical and neurochemical effects of acetyl-L-carnitine in Alzheimer's disease. *Neurobiol Aging* 1995 Jan; 16(1):1-4.

Status: Not included because dementia population not randomized to treatment

Pettegrew JW, Levine J, McClure RJ. Acetyl-L-carnitine physical-chemical, metabolic, and therapeutic properties: Relevance for its mode of action in Alzheimer's disease and geriatric depression. *Mol Psychiatry* 2000; 5(6):616-32.

Status: Not included because does not meet criteria for treatment for dementia patients

Pettegrew JW, McClure RJ. Acetyl-L-carnitine as a possible therapy for Alzheimer's disease. *Expert Rev Neurother* 2002; 2(5):647-54.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Petursson H. A controlled study of Moclobemide in elderly depression with cognitive decline. *Psychopharmacologia* 1993; Vol 111:B5

Status: Not included because not a full article

Pérodeau G, Lauzon S, Lévesque L, et al. Mental health stress correlates and psychotropic drug use or non user among aged caregivers to elders with dementia. *Aging Ment Health* 2001; 5(3):225-34.

Status: Not included because does not meet criteria for treatment for dementia patients

Pfefferbaum A, Davis KL, Coulter CL, et al. EEG effects of physostigmine and choline chloride in humans. *Psychopharmacologia* 1979 Apr 25; 62(3):225-33.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Phanjoo AL, Link C. Remoxipride versus thioridazine in elderly psychotic patients. *Acta Psychiatr Scand Suppl* 1990; Vol 358:181-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Phillips CD, Spry KM, Sloane PD, et al. Use of physical restraints and psychotropic medications in Alzheimer special care units in nursing homes. *Am J Public Health* 2000; 90(1):92-6.

Status: Not included because dementia population not randomized to treatment

Piccinin GL, Finali G, Piccirilli M. Neuropsychological effects of L-deprenyl in Alzheimer's type dementia. *Clin Neuropharmacol* 1990 Apr; 13(2):147-63.

Status: Cross-over trial;

Piccoli F, Battistini N, Carbonin P, et al. CDP-choline in the treatment of chronic cerebrovasculopathies. *Arch Gerontol Geriatr* 1994; 18(3):161-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pillemer K, Jill SJ. Peer support for Alzheimer's caregivers: Is it enough to make a difference? *Res Aging* 2002; 24(2):171-92.

Status: Not included because does not meet criteria for treatment for dementia patients

Pincus MM, Kilander L, Ohrvall M. Alpha-tocopherol and Alzheimer's disease. *N Engl J Med* 1997; 337(8):572-3.

Status: Not included because not a full article

Pisvejc J, Hyrman V, Sikora J, et al. A comparison of brief and ultrabrief pulse stimuli in unilateral ECT. *J ECT* 1998 Jun; 14(2):68-75.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pittera A, Ciancitto S. Effect of oral nimodipine on cerebral blood flow in patients with chronic cerebrovascular disorders. A supra-aortic Doppler ultra-sound open study. *Curr Ther Res Clin Exp* 1990; 48(4):716-29.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Poitrenaud J, Piette F, Malbezin M, et al. Almitrine-raubasine and cognitive impairment in the elderly: Results of a 6-month controlled multicenter study. *Clin Neuropharmacol* 1990; Vol 13(Suppl 3):S100-S108.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Polinsky RJ. Clinical pharmacology of rivastigmine: A new-generation acetylcholinesterase inhibitor for the treatment of Alzheimer's disease. *Clin Ther* 1998; 20(4):634-47.
Status: Not included because dementia population not randomized to treatment

Pomara N, Block R, Abraham J. Combined cholinergic precursor treatment and dihydroergotoxine mesylate in Alzheimer's disease. *IRCS Med Sci* 1983; (12):1048-9.
Status: Not included because dementia population not randomized to treatment

Pomara N, Block R, Moore N, et al. Combined Piracetam and cholinergic precursor treatment for primary degenerative dementia. *IRCS Med Sci Psychol Psychiatry* 1984; (5-6):388-6.
Status: Cross-over trial;

Pomara N, Roberts R, Rhiew HB, et al. Multiple, single-dose naltrexone administrations fail to effect overall cognitive functioning and plasma cortisol in individuals with probable Alzheimer's

disease. *Neurobiol Aging* 1985; 6(3):233-6.
Status: Cross-over trial;

Pomara N, Deptula D, Singh R. Pretreatment postural blood pressure drop as a possible predictor of response to the cholinesterase inhibitor velnacrine (HP 029) in Alzheimer's disease. *Psychopharmacol Bull* 1991; 27(3):301-7.
Status: Not included because dementia population not randomized to treatment

Pomara N, Doraiswamy PM, Tun H, et al. Mifepristone (RU 486) for Alzheimer's disease. *Neurology* 2002; 58(9):1436.
Status: Not included because Jadad Quality Scale score less than three

Pomeroy VM. The effect of physiotherapy input on mobility skills of elderly people with severe dementing illness. *Clin Rehabil* 1993; 7(2):163-70.
Status: Not included because does not meet criteria for treatment for dementia patients

Pomeroy VM. Immobility and severe dementia: When is physiotherapy treatment appropriate? *Clin Rehabil* 1994; 8:226-32.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Pomeroy VM, Warren CM, Honeycombe C, et al. Mobility and dementia: Is physiotherapy treatment during respite care effective? *Int J Geriatr Psychiatry* 1999 May; 14(5):389-97.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Portegies P, Enting RH, de Jong MD, et al. AIDS dementia complex and didanosine. *Lancet* 1994 Sep 10; 344(8924):759
Status: Not included because not a full article

Porter RJ, Lunn BS, Walker LL, et al. Cognitive deficit induced by acute tryptophan depletion in patients with Alzheimer's disease. *Am J Psychiatry* 2000 Apr; 157(4):638-40.
Status: Not included because dementia population not randomized to treatment

Porter RJ, Marshall EF, O'Brien JT. Effects of rapid tryptophan depletion on salivary and plasma cortisol in Alzheimer's disease and the healthy elderly. *J Psychopharmacol (Oxf)* 2002 Mar; 16(1):73-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Postiglione A, Soricelli A, Cicerano U, et al. Effect of acute administration of L-acetyl carnitine on cerebral blood flow in patients with chronic cerebral infarct. *Pharmacol Res* 1991 Apr; 23(3):241-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Potkin SG, Fleming K, Jin Y, et al. Clozapine enhances neurocognition and clinical symptomatology more than standard neuroleptics. *J Clin Psychopharmacol* 2001; 21(5):479-83.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Potkin SG, Anand R, Hartman R, et al. Impact of Alzheimer's disease and rivastigmine treatment on activities of daily living over the course of mild to moderately severe disease. *Prog Neuropsychopharmacol Biol Psychiatry* 2002; 26(4):713-20.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Potkin SG, Alva G, Keator D, et al. Brain metabolic effects of Neotrofin in patients with Alzheimer's disease. *Brain Res* 2002; 951(1):87-95.

Status: Not included because dementia population not randomized to treatment

Potter A, Corwin J, Lang J, et al. Acute effects of the selective cholinergic channel activator (nicotinic agonist) ABT-418 in Alzheimer's disease. *Psychopharmacologia* 1999 Mar; 142(4):334-42.

Status: Cross-over trial;

Pratt RD. Patient populations in clinical trials of the efficacy and tolerability of donepezil in patients with vascular dementia. *J Neurol Sci* 2002 Nov 15; 203-204:57-65.:57-65.

Status: Not included because no extractable data relevant to review

Pratt RD, Perdomo CA, Surick IW, et al. Donepezil: Tolerability and safety in Alzheimer's disease. *Int J Clin Pract* 2002 Nov; 56(9):710-7. *Status: Not included because dementia population not randomized to treatment*

Predescu V, Riga D, Riga S, et al. Antagonic-stress. A new treatment in gerontopsychiatry and

for a healthy productive life. *Ann N Y Acad Sci* 1994; 717(Jun 30):315-31.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Preston GC, Brazell C, Ward C, et al. The scopolamine model of dementia: Determination of central cholinomimetic effects of physostigmine on cognition and biochemical markers in man. *J Psychopharmacol (Oxf)* 1988; 2(2):67-79.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Preston GC, Ward C, Lines CR, et al. Scopolamine and benzodiazepine models of dementia: Cross-reversals by Ro 15-1788 and physostigmine. *Psychopharmacologia* 1989; 98(4):487-94.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Proctor R, Burns A, Powell HS, et al. Behavioural management in nursing and residential homes: a randomised controlled trial. *Lancet* 1999 Jul 3; 354(9172):26-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Puchler K, Schaffler K, Plenker A. The comparative effects of single and multiple doses of RS-8359, moclobemide and placebo on psychomotor function in healthy subjects. *Int Clin Psychopharmacol* 1997; Vol 12(Suppl 5):S17-S23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Purandare N, Bloom C, Page S, et al. The effect of anticholinesterases on personality changes in Alzheimer's disease. *Aging Ment Health* 2002; 6(4):350-4.

Status: Not included because dementia population not randomized to treatment

Puri BK, Bydder GM, Counsell SJ, et al. MRI and neuropsychological improvement in Huntington disease following ethyl-EPA treatment. *NeuroReport* 2002; 13(1):123-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Putt ME, Ravina B. Randomized, placebo-controlled, parallel group versus crossover study designs for the study of dementia in Parkinson's disease. *Control Clin Trials* 2002 Apr; 23(2):111-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Quayhagen MP, Quayhagen M, Corbeil RR, et al. A dyadic remediation program for care recipients with dementia. *Nurs Res* 1995 May; 44(3):153-9.
Status: Not included because does not meet criteria for treatment for dementia patients

Quayhagen MP, Quayhagen M, Corbeil RR, et al. Coping with dementia: Evaluation of four nonpharmacologic interventions. *Int Psychogeriatr* 2000 Jun; 12(2):249-65.
Status: Not included because does not meet criteria for treatment for dementia patients

Quayhagen MP, Quayhagen M. Testing of a cognitive stimulation intervention for dementia caregiving dyads. *Neuropsychol Rehab* 2001; 11(Special issue (3-4)):319-22.
Status: Not included because does not meet criteria for treatment for dementia patients

Quinlivan R, Hough R, Crowell A, et al. Service utilization and costs of care for severely mentally ill clients in an intensive case management program. *Psychiatr Serv* 1995 Apr; 46(4):365-71.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rabey JM, Neufeld MY, Treves TA, et al. Cognitive effects of scopolamine in dementia. *J Neural Transm Gen Sect* 1996; 103(7):873-81.
Status: Cross-over trial;

Rada RT, Kellner R. Thiothixene in the treatment of geriatric patients with chronic organic brain syndrome. *J Am Geriatr Soc* 1976 Mar; 24(3):105-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rader MA, Alston JB, Ellis DW. Sensory stimulation of severely brain-injured patients. *Brain Inj* 1989 Apr; 3(2):141-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Raffaele KC, Berardi A, Asthana S, et al. Effects of long-term continuous infusion of the muscarinic cholinergic agonist arecoline on verbal memory in dementia of the Alzheimer type. *Psychopharmacol Bull* 1991; 27(3):315-9.
Status: Not included because dementia population not randomized to treatment

Raffaele KC, Asthana S, Berardi A, et al. Differential response to the cholinergic agonist arecoline among different cognitive modalities in Alzheimer's disease. *Neuropsychopharmacology* 1996; 15(2):163-70.
Status: Not included because dementia population not randomized to treatment

Ragneskog H, Brane G, Karlsson I, et al. Influence of dinner music on food intake and symptoms common in dementia. *Scand J Caring Sci* 1996; 10(1):11-7.
Status: Not included because dementia population not randomized to treatment

Ragneskog H, Kihlgren M, Karlsson I, et al. Dinner music for demented patients: Analysis of video-recorded observations. *Clin Nurs Res* 1996 Aug; 5(3):262-77, 278-82.
Status: Not included because dementia population not randomized to treatment

Ragneskog H, Asplund K, Kihlgren M, et al. Individualized music played for agitated patients with dementia: Analysis of video-recorded sessions. *Int J Nurs Pract* 2001 Jun; 7(3):146-55.
Status: Not included because dementia population not randomized to treatment

Rai GS, Shovlin C, Wesnes KA. A double-blind, placebo controlled study of Ginkgo biloba extract ('tanakan') in elderly outpatients with mild to moderate memory impairment. *Curr Med Res Opin* 1991; 12(6):350-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rained M, Mucke HAM. Long-term cognitive benefit from galanthamine in Alzheimer's disease. *Int J Geriatr Psychopharmacol* 1998; 1(4):197-201.
Status: Not included because dementia population not randomized to treatment

Rainer M, Mucke HA, Chwatal K, et al. Alcohol-induced organic cerebral psychosyndromes: Partial reversal of cognitive impairments assisted by dihydroergocristine. *Psychopharmacologia* 1996 Oct; 127(4):365-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Randolph C, Roberts JW, Tierney MC, et al. D-cycloserine treatment of Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1994; 8(3):198-205.
Status: Cross-over trial;

Ransmayr G, Plorer S, Gerstenbrand F, et al. Double-blind placebo-controlled trial of phosphatidylserine in elderly patients with arteriosclerotic encephalopathy. *Clin Trials J* 1987; 24(1):62-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rao DB, Georgiev EL, Paul PD, et al. Cyclandelate in the treatment of senile mental changes: A double-blind evaluation. *J Am Geriatr Soc* 1977 Dec; 25(12):548-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rapp S, Brenes G, Marsh AP. Memory enhancement training for older adults with mild cognitive impairment: A preliminary study. *Aging Ment Health* 2002 Feb; 6(1):5-11.

Status: Not included because does not meet criteria for treatment for dementia patients

Rascol O, Sieradzan K, Peyro-Saint-Paul H, et al. Efaroxan, an alpha-2 antagonist, in the treatment of progressive supranuclear palsy. *Mov Disord* 1998 Jul; 13(4):673-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rasmusen L, Yan B, Robillard A, et al. Effects of washout and dose-escalation periods on the efficacy, safety, and tolerability of galantamine in patients previously treated with donepezil: Ongoing clinical trials. *Clin Ther* 2001; 23(Suppl):A25-A30.

Status: Not included because dementia population not randomized to treatment

Ray PG, Meador KJ, Loring DW, et al. Effects of scopolamine on visual evoked potentials in aging and dementia. *Electroencephalogr Clin Neurophysiol* 1991 Sep; 80(5):347-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ray WA, Taylor JA, Meador KG, et al. Reducing antipsychotic drug use in nursing homes. A controlled trial of provider education. *Arch Intern Med* 1993; 153(6):713-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Raz N, Torres IJ, Briggs SD, et al. Selective neuroanatomic abnormalities in Down's syndrome and their cognitive correlates: Evidence from MRI

morphometry. *Neurology* 1995 Feb; 45(2):356-66.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Reddy H, De Stefano N, Mortilla M, et al. Functional reorganization of motor cortex increases with greater axonal injury from CADASIL. *Stroke* 2002 Feb; 33(2):502-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Reeve W. Use of environmental manipulation and classroom and modified informal reality orientation with institutionalized, confused elderly patients. *Age Ageing* 1985; 14(2):119-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Regland B, Lehmann W, Abedini I, et al. Treatment of Alzheimer's disease with clioquinol. *Dement Geriatr Cogn Disord* 2001 Nov; 12(6):408-14.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rehman SA. Two trials comparing 'Hydergine' with placebo in the treatment of patients suffering from cerebrovascular insufficiency. *Curr Med Res Opin* 1973; 1(8):456-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Reisberg B, Ferris SH, Schneck MK. Piracetam in the treatment of cognitive impairment in the elderly. *Drug Dev Res* 1982; 2(5):475-80.

Status: Cross-over trial;

Reisberg B, Ferris SH, Anand R. Effects of naloxone in senile dementia: A double-blind trial. *N Engl J Med* 1983; 308(12):721-2.

Status: Not included because not a full article

Remington R. Calming music and hand massage with agitated elderly. *Nurs Res* 2002 Sep; 51(5):317-23.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Richard F, Helbecque N, Neuman E, et al. APOE genotyping and response to drug treatment in Alzheimer's disease. *Lancet* 1997 Feb 22; 349(9051):539

Status: Not included because Jadad Quality Scale score less than three

Riekkinen M, Laakso MP, Jakala P, et al. Clonidine impairs sustained attention and memory in Alzheimer's disease. *Neuroscience* 1999; 92(3):975-82.

Status: Not included because dementia population not randomized to treatment

Riekkinen P, Jr., Kuikka J, Soininen H, et al. Tetrahydroaminoacridine modulates technetium-99m labelled ethylene dicysteinate retention in Alzheimer's disease measured with single photon emission computed tomography imaging. *Neurosci Lett* 1995 Jul 28; 195(1):53-6.

Status: Not included because no extractable data relevant to review

Riekkinen P, Jr., Riekkinen M, Soininen H, et al. Frontal dysfunction blocks the therapeutic effect of THA on attention in Alzheimer's disease. *NeuroReport* 1997 May 27; 8(8):1845-9.

Status: Not included because dementia population not randomized to treatment

Riekkinen PJ, Koivisto K, Helkala EL, et al. Long-term, double-blind trial of selegiline in Alzheimer's disease. *Neurobiol Aging* 1994; 15(Suppl 1):67.

Status: Not included because not a full article

Riekkinen PJ, Soininen H, Helkala E-L, et al. Hippocampal atrophy, acute THA treatment and memory in Alzheimer's disease. *NeuroReport* 1995; 6(9):1297-300.

Status: Not included because dementia population not randomized to treatment

Riekkinen PJr, Soininen H, Partanen J, et al. The ability of THA treatment to increase cortical alpha waves is related to apolipoprotein E genotype of Alzheimer disease patients. *Psychopharmacologia* 1997 Feb; 129(3):285-8.

Status: Not included because dementia population not randomized to treatment

Riekkinen PJr, Paakkonen A, Karhu J, et al. THA disrupts mismatch negativity in Alzheimer's disease. *Psychopharmacologia* 1997 Sep; 133(2):203-6.

Status: Not included because no extractable data relevant to review

Riekkinen PJr, Riekkinen M. THA improves word priming and clonidine enhances fluency and working memory in Alzheimer's disease. *Neuropsychopharmacology* 1999 Apr; 20(4):357-64.

Status: Not included because dementia population not randomized to treatment

Rigaud A-S, Traykov L, Latour F, et al. Presence or absence of at least one epsilon4 allele and gender are not predictive for the response to donepezil treatment in Alzheimer's disease. *Pharmacogenetics* 2002; 12(5):415-20.

Status: Not included because dementia population not randomized to treatment

Rigaud AS, Andre G, Vellas B, et al. No additional benefit of HRT on response to rivastigmine in menopausal women with AD. *Neurology* 2003; 60(1):148-50.

Status: Not included because Jadad Quality Scale score less than three

Rimon R, Rakkolainen V. Lithium iodide in the treatment of confusional states. *Br J Psychiatry* 1968 Jan; 114(506):109-10.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rinne JO, Portin R, Ruottinen H, et al. Cognitive impairment and the brain dopaminergic system in Parkinson's disease: Fluorodopa positron emission tomographic study. *Arch Neurol* 2000 Apr; 57(4):470-5.

Status: Not included because dementia population not randomized to treatment

Rinsky JR, Wikler A, Way JG, et al. 'Mental set' in controls, postalcoholics, chronic schizophrenics, and 'organics'. *Biol Psychiatry* 1979; 14(6):881-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Riordan JM, Bennet AV. An evaluation of an augmented domiciliary service to older people with dementia and their carers. *Aging Ment Health* 1998; 2(2):137-43.

Status: Not included because dementia population not randomized to treatment

Rivera VM, Meyer JS, Baer PE, et al. Vertebrobasilar arterial insufficiency with dementia. Controlled trials of treatment with betahistine hydrochloride. *J Am Geriatr Soc* 1974 Sep; 22(9):397-406.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Riviere S, Gillette-Guyonnet S, Voisin T, et al. A nutritional education program could prevent

weight loss and slow cognitive decline in Alzheimer's disease. *J Nutr Health Aging* ; 5(4):295-9.

Status: Not included because dementia population not randomized to treatment

Rizzo JA, Bogardus ST, Jr., Leo-Summers L, et al. Multicomponent targeted intervention to prevent delirium in hospitalized older patients: What is the economic value? *Med Care* 2001 Jul; 39(7):740-52.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rizzo M, Ficola U, Marozzi P, et al. Effects of aniracetam on clinical pattern and cerebral blood flow of old patients with degenerative dementia (SDAT). *J Neurol* 1994; 241:S126.

Status: Not included because not a full article

Robbins TW, Semple J, Kumar R, et al. Effects of scopolamine on delayed-matching-to-sample and paired associates tests of visual memory and learning in human subjects: Comparison with diazepam and implications for dementia. *Psychopharmacologia* 1997 Nov; 134(1):95-106.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Robert PH, Allain H. Clinical management of agitation in the elderly with tiapride. *Eur Psychiatry* 2001 Jan; 16(Suppl):47S.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Roberts CJ, Ford JM, Truman CA, et al. Assessment of the value of therapeutic monitoring of tacrine in Alzheimer's disease. *Eur J Clin Pharmacol* 1998 Nov; 54(9-10):721-10.

Status: Not included because dementia population not randomized to treatment

Robichaud L, Hebert R, Desrosiers J. Efficacy of a sensory integration program on behaviors of inpatients with dementia. *Am J Occup Ther* 1994 Apr; 48(4):355-60.

Status: Not included because does not meet criteria for treatment for dementia patients

Rochtchina I, Gavrilova S, Kolykhalov I, et al. Neuropsychological assessment of modification of efficacy of cholinergic therapy by preceding cerebrolysin treatment in Alzheimer's disease. *J Eur Coll Neuropsychopharmacol* 1999; (Suppl 5):S330.

Status: Not included because not a full article

Rockwood K, Stolee P, Howard K, et al. Use of Goal Attainment Scaling to measure treatment effects in an anti-dementia drug trial. *Neuroepidemiology* 1996; 15(6):330-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rockwood K, Graham JE, Fay S, et al. Goal setting and attainment in Alzheimer's disease patients treated with donepezil. *J Neurol Neurosurg Psychiatry* 2002 Nov; 73(5):500-7.

Status: Not included because dementia population not randomized to treatment

Rogers JD, Sanchez-Saffon A, Frol AB, et al. Elevated plasma homocysteine levels in patients treated with levodopa: Association with vascular disease. *Arch Neurol* 2003; 60(1):59-64.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rogers JF, Morrison AL, Nafziger AN, et al. Flumazenil reduces midazolam-induced cognitive impairment without altering pharmacokinetics. *Clin Pharmacol Ther* 2002; 72(6):711-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rogers SL, Doody R, Mohs R, et al. E2020 produces both clinical, global and cognitive test improvement in patients with mild to moderately severe Alzheimer's disease: Results of a 30-week phase III trial. *Neurology* 1996; 46(Suppl):217

Status: Not included because not a full article

Rombouts SA, Barkhof F, Van Meel CS, et al. Alterations in brain activation during cholinergic enhancement with rivastigmine in Alzheimer's disease. *J Neurol Neurosurg Psychiatry* 2002 Dec; 73(6):665-71.

Status: Not included because dementia population not randomized to treatment

Ronnberg L. Quality of life in nursing-home residents: An intervention study of the effect of mental stimulation through an audiovisual programme. *Age Ageing* 1998; 27(3):393-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosen HJ. Mental decline in the elderly: Pharmacotherapy (ergot alkaloids versus papaverine). *J Am Geriatr Soc* 1975 Apr; 23(4):169-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosenberg SJ, Ryan JJ, Prifitera A. Rey Auditory-Verbal Learning Test performance of patients with and without memory impairment. *J Clin Psychol* 1984; (3):785-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosenzweig P, Patat A, Zieleniuk I, et al. Cognitive performance in elderly subjects after a single dose of befloxacitane, a new reversible selective monoamine oxidase A inhibitor. *Clin Pharmacol Ther* 1998 Aug; 64(2):211-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rosewarne R, Bruce A, McKenna M. Dementia programme effectiveness in long-term care. *Int J Geriatr Psychiatry* 1997 Feb; (2):173-82.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rossi R, Inzitari D, Pantoni L, et al. Nimodipine in subcortical vascular dementia trial. *Alzheimer Dis Assoc Disord* 1999; 13(Suppl 3):S159-S165.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rostow CD. The effect of self- vs. external-monitoring and locus of control upon the pacing and general adjustment of psychiatric inpatients. *Behav Res Ther* 1980; 18(6):541-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rother M, Kittner B, Rudolphi K, et al. HWA 285 (propentofylline) - A new compound for the treatment of both vascular dementia and dementia of the Alzheimer type. *Ann N Y Acad Sci* 1996; 777:404-9.
Status: Not included because dementia population not randomized to treatment

Rother M. Long-term effects of propentofylline in patients with Alzheimer's disease: A 72-week, placebo-controlled study assessing safety, efficacy, and impact on disease progression. *J Cereb Blood Flow Metab* 1999; 19(Suppl 1):18.
Status: Not included because not a full article

Rouy JM, Douillon AM, Compan B, et al. Ergoloid mesylates ('Hydergine') in the treatment of mental deterioration in the elderly: A 6-month double-blind, placebo-controlled trial. *Curr Med Res Opin*

1989; 11(6):380-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Rovner BW, Steele CD, Shmueli Y, et al. A randomized trial of dementia care in nursing homes. *J Am Geriatr Soc* 1996 Jan; 44(1):7-13.
Status: Not included because does not meet criteria for treatment for dementia patients

Rozzini R, Ferrucci L, Losonczy K, et al. Protective effect of chronic NSAID use on cognitive decline in older persons. *J Am Geriatr Soc* 1996; 44(9):1025-9.
Status: Not included because dementia population not randomized to treatment

Rudman D, Racette D, Rudman IW, et al. Hyponatremia in tube-fed elderly men. *J Chronic Dis* 1986; 39(2):73-80.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Ruether E, Husmann R, Kinzler E, et al. A 28-week, double-blind, placebo-controlled study with cerebrolysin in patients with mild to moderate Alzheimer's disease. *Int Clin Psychopharmacol* 2001; 16(6):372
Status: Not included because not a full article

Ruggiero, Ovallesco U. Clozapine and validation therapy in the treatment of dementia. *Funct Neurol* 1997; 3-4(12):240-412.
Status: Not included because dementia population not randomized to treatment

Ryden MB, Snyder M, Gross CR, et al. Value-added outcomes: The use of advanced practice nurses in long-term care facilities. *Gerontologist* 2000 Dec; 40(6):654-62.
Status: Not included because does not meet criteria for treatment for dementia patients

Ryynanen OP, Myllykangas M, Kinnunen J, et al. Doctors' willingness to refer elderly patients for elective surgery. *Fam Pract* 1997 Jun; 14(3):216-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saarela T, Kiviharju U. Evaluating the usefulness of training in psychogeriatrics. *Int J Geriatr Psychiatry* 1995 Dec; (12):1019-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sabe L, Kuzis G, Garcia CA, et al. A randomized, double-blind, placebo-controlled study of idebenone in Alzheimer's disease (AD). *JNS* 1997; 150(Suppl):S296.

Status: Not included because not a full article

Sacktor N, Schifitto G, McDermott MP, et al. Transdermal selegiline in HIV-associated cognitive impairment: Pilot, placebo-controlled study. *Neurology* 2000 Jan 11; 54(1):233-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sacktor N, McDermott MP, Marder K, et al. HIV-associated cognitive impairment before and after the advent of combination therapy. *J Neurovirol* 2002; 8(2):136-42.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sahakian BJ, Coull JT. Tetrahydroaminoacridine (THA) in Alzheimer's disease: An assessment of attentional and mnemonic function using CANTAB. *Acta Neurol Scand Suppl* 1993; Vol 149:29-35.
Status: Cross-over trial;

Sahakian BJ, Owen AM, Morant NJ, et al. Further analysis of the cognitive effects of tetrahydroaminoacridine (THA) in Alzheimer's disease: Assessment of attentional and mnemonic function using CANTAB. *Psychopharmacologia* 1993; 110(4):395-401.
Status: Cross-over trial;

Sahakian BJ, Coull JT. Nicotine and tetrahydroaminoacridine: Evidence for improved attention in patients with dementia of the Alzheimer type. *Drug Dev Res* 1994; 31(1):80-8.
Status: Not included because dementia population not randomized to treatment

Sahin HA, Gurvit IH, Bilgic B, et al. Therapeutic effects of an acetylcholinesterase inhibitor (donepezil) on memory in Wernicke-Korsakoff's disease. *Clin Neuropharmacol* 2002 Jan; 25(1):16-20.
Status: Not included because dementia population not randomized to treatment

Saine K, Cullum CM, Martin-Cook K, et al. Comparison of functional and cognitive donepezil effects in Alzheimer's disease. *Int Psychogeriatr* 2002; 14(2):181-5.
Status: Not included because dementia population not randomized to treatment

Sajatovic M, Mullen JA, Sweitzer DE. Efficacy of quetiapine and risperidone against depressive symptoms in outpatients with psychosis. *J Clin Psychiatry* 2002; 63(12):1156-63.

Status: Not included because no extractable data relevant to review

Saldmann F, Funel A, Jacquet P. Efficacy of naftidrofuryl in patients with moderate senile dementia. *Curr Med Res Opin* 1991; 12(6):379-89.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saletu B, Linzmayer L, Grunberger J, et al. Double-blind, placebo-controlled, clinical, psychometric and neurophysiological investigations with oxiracetam in the organic brain syndrome of late life. *Neuropsychobiology* 1985; 13(1-2):44-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saletu B, Gruenberger J, Linzmayer L, et al. EEG brain mapping and psychometry in age-associated memory impairment after acute and 2-week infusions with the hemoderivative Actovegin®: Double-blind, placebo-controlled trials. *Neuropsychobiology* 1990; 24(3):135-48.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Saletu B, Anderer P, Fischhof PK, et al. EEG mapping and psychopharmacological studies with denbufylline in SDAT and MID. *Biol Psychiatry* 1992 Oct 15; 32(8):668-81.
Status: Not included because Jadad Quality Scale score less than three

Saletu B, Anderer P, Semlitsch HV, et al. Amantadine infusions in mild dementia: Acute double-blind placebo-controlled EEG mapping and psychometric studies. *Arch Gerontol Geriatr* 1992; 15(1):43-58.
Status: Cross-over trial;

Saletu M, Grunberger J, Saletu B, et al. Accelerated remission of the alcoholic organic brain syndrome with EMD 21657. Double-blind clinical and psychometric trials. *Arzneimittelforschung* 1978; 28(9):1525-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Salib E, Sheridan T, Allington M. Ventricular measurements in computed tomography of responders and non-responders to donepezil in the treatment of Alzheimer's disease. *Int J Psychiatry Clin Pract* 2001; 5(3):189-94.

Status: Not included because dementia population not randomized to treatment

Salvioli G, Neri G. L-acetylcarnitine treatment of mental decline in the elderly. *Drugs Exp Clin Res* 1994; 20(4):169-76.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Samorajski T, Vroulis GA, Smith RC. Piracetam plus lecithin trials in senile dementia of the Alzheimer type. *Ann N Y Acad Sci* 1985; 444:478-81.

Status: Not included because dementia population not randomized to treatment

Samuel W, Caligiuri M, Galasko D, et al. Better cognitive and psychopathologic response to donepezil in patients prospectively diagnosed as dementia with Lewy bodies: A preliminary study. *Int J Geriatr Psychiatry* 2000 Sep; 15(9):794-802.

Status: Not included because dementia population not randomized to treatment

Sanders DS, Carter MJ, D'Silva J, et al. Survival analysis in percutaneous endoscopic gastrostomy feeding: A worse outcome in patients with dementia. *Am J Gastroenterol* 2000; 95(6):1472-5.

Status: Not included because dementia population not randomized to treatment

Sano M, Stern Y, Marder K, et al. A controlled trial of piracetam in intellectually impaired patients with Parkinson's disease. *Mov Disord* 1990; 5(3):230-4.

Status: Not included because no extractable data relevant to review

Sano M, Bell K, Marder K, et al. Safety and efficacy of oral physostigmine in the treatment of Alzheimer's disease. *Clin Neuropharmacol* 1993 Feb; 16(1):61-9.

Status: Cross-over trial;

Sano M, Growdon J, Klauber M, et al. Expanding the severity range of patients in clinical trials for Alzheimer's disease: A multicentre clinical trial of Selegiline and alpha-tocopherol. *Neurology* 1995; 45(Suppl 4):A289.

Status: Not included because not a full article

Sano M, Ernesto C, Thomas RG, et al. Effects of Selegiline and alpha-Tocopherol on cognitive and functional outcome measures in moderately impaired patients with Alzheimer's disease. *Neurology* 1997; 48(Suppl):A377-A378

Status: Not included because not a full article

Sano M, Bell K, Jacobs D. Cognitive effects of estrogens in women with cardiac disease: What we do not know. *Am J Med* 2002; 113(7):612-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sansom G. Comparing multisensory stimulation with (1) tactile stimulation and (2) themed reminiscence. *Journal of Dementia Care* 2002; 10(4):38

Status: Not included because not a full article

Satlin A, Volicer L, Ross V, et al. Bright light treatment of behavioral and sleep disturbances in patients with Alzheimer's disease. *Am J Psychiatry* 1992; 149(8):1028-32.

Status: Not included because dementia population not randomized to treatment

Sato K, Kamiya S, Okawa M, et al. Effect of transcranial electrostimulation on EEG component waves of elderly patients with dementia. *J Brain Sci* 1998; 24(1-2):65-72.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Savoldi F, Nappi G, Martignoni E, et al. Brain phospholipids in the treatment of chronic cerebrovascular insufficiency. *Curr Ther Res Clin Exp* 1978; 24(2):209.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Scarzella L, Bono G, Bergamasco B. Dihydroergocryptine in the management of senile psycho-organic syndrome. *Int J Clin Pharmacol Res* 1992; 12(1):37-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Scherder E, Bouma A, Steen L. Effects of simultaneously applied short-term transcutaneous electrical nerve stimulation and tactile stimulation on memory and affective behaviour of patients with probable Alzheimer's disease. *Behav Neurol* 1995; (1):3-13.

Status: Not included because dementia population not randomized to treatment

Scherder EJ, Bouma A, Steen L. Influence of transcutaneous electrical nerve stimulation on memory in patients with dementia of the Alzheimer type. *J Clin Exp Neuropsychol* 1992 Nov; 14(6):951-60.

Status: Not included because dementia population not randomized to treatment

Scherder EJ, Bouma A, Steen AM. Effects of short-term transcutaneous electrical nerve stimulation on memory and affective behaviour in patients with probable Alzheimer's disease. *Behav Brain Res* 1995; 67(2):211-9.

Status: Not included because dementia population not randomized to treatment

Scherder EJ, Bouma A, Steen LM. Effects of "isolated" transcutaneous electrical nerve stimulation on memory and affective behavior in patients with probable Alzheimer's disease. *Biol Psychiatry* 1998 Mar 15; 43(6):417-24.

Status: Not included because does not meet criteria for treatment for dementia patients

Scherder EJ, Van Someren EJ, Swaab DF. Transcutaneous electrical nerve stimulation (TENS) improves the rest-activity rhythm in midstage Alzheimer's disease. *Behav Brain Res* 1999 May; 101(1):105-7.

Status: Not included because no extractable data relevant to review

Scherder EJ, Bouma A. Effects of transcutaneous electrical nerve stimulation on memory and behavior in Alzheimer's disease may be stage-dependent. *Biol Psychiatry* 1999 Mar 15; 45(6):743-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Scherder EJ, Van Someren EJ, Bouma A, et al. Effects of transcutaneous electrical nerve stimulation (TENS) on cognition and behaviour in aging. *Behav Brain Res* 2000 Jun 15; 111(1-2):223-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Scherder EJ, Deijen JB, Vreeswijk SH, et al. Cranial electrostimulation (CES) in patients with probable Alzheimer's disease. *Behav Brain Res* 2002 Jan 22; 128(2):215-7.

Status: Not included because Jadad Quality Scale score less than three

Scherder E, Bouma A, Steen L. Effects of peripheral tactile stimulation on memory in patients with probable Alzheimer's disease. *Am J Alzheimers Dis* 1995 May; (3):15-21.

Status: Not included because dementia population not randomized to treatment

Scherder E, Bouma A, Steen L. Effects of peripheral tactile nerve stimulation on affective behavior of patients with probable Alzheimer's disease. *Am J Alzheimers Dis* 1998 Mar; (2):61-9.

Status: Not included because does not meet criteria for treatment for dementia patients

Schiffmann R, Heyes MP, Aerts JM, et al. Prospective study of neurological responses to treatment with macrophage-targeted glucocerebrosidase in patients with type 3 Gaucher's disease. *Ann Neurol* 1997 Oct; 42(4):613-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schifitto G, Sacktor N, Marder K, et al. Randomized trial of the platelet-activating factor antagonist lexipafant in HIV-associated cognitive impairment. *Neurological AIDS Research Consortium. Neurology* 1999 Jul 22; 53(2):391-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schlegel J, Mohr E, Williams J, et al. Guanfacine treatment of Alzheimer's disease. *Clin Neuropharmacol* 1989; 12(2):124-8.

Status: Cross-over trial;

Schmechel DE, Schmitt F, Horner J, et al. Lack of effect of oral physostigmine and lecithin in patients with probable Alzheimer's disease. *Neurology* 1984; 34(Suppl 1):280.

Status: Not included because not a full article

Schmidt IK, Fastbom J. Quality of drug use in Swedish nursing homes. A follow-up study. *Clin Drug Investig* 2000; 20(6):433-46.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schmitt R, Capo T, Frazier H, et al. Cranial electrotherapy stimulation treatment of cognitive brain dysfunction in chemical dependence. *J Clin Psychiatry* 1962 Mar; 45(2):60-1.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schneider LS, Gleason RP. Carbamazepine in behaviorally disturbed primary dementia patients. *J Clin Psychiatry* 1990; 51(12):524
Status: Not included because not a full article

Schneider LS, Lyness SA, Pawluczyk S, et al. Do blood pressure and age predict response to tacrine (THA) in Alzheimer's disease? A preliminary report. *Psychopharmacol Bull* 1991; 27(3):309-14.
Status: Not included because no extractable data relevant to review

Schneider LS, Olin JT, Pawluczyk S. A double-blind crossover pilot study of l-deprenyl (selegiline) combined with cholinesterase inhibitor in Alzheimer's disease. *Am J Psychiatry* 1993 Feb; 150(2):321-3.
Status: Cross-over trial;

Schneider LS, Farlow MR, Henderson VW, et al. Estrogen replacement therapy may enhance response to tacrine in women with Alzheimer's disease. *Neurology* 1995; 45:288.
Status: Not included because not a full article

Schneider LS, Farlow MK. Severity of Alzheimer's disease and response to cholinergic therapy. *Eur J Neurol* 1996; 3:238.
Status: Not included because not a full article

Schneider LS, Farlow MR, Pogoda JM. Potential role for estrogen replacement in the treatment of Alzheimer's dementia. *Am J Med* 1997 Sep 22; 103(3A):46S-50S.
Status: Not included because Jadad Quality Scale score less than three

Schneider LS, Tariot PN, Lyketsos CG, et al. National Institute of Mental Health Clinical Antipsychotic Trials of Intervention Effectiveness (CATIE): Alzheimer's disease trial methodology. *Am J Geriatr Psychiatry* 2001; 9(4):346-60.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schredl M, Weber B, Braus D, et al. The effect of rivastigmine on sleep in elderly healthy subjects. *Exp Gerontol* 2000; 35(2):243-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schreiber M, Schweizer A, Lutz K, et al. Potential of an interactive computer-based training in the rehabilitation of dementia: An initial study. *Neuropsychol Rehab* 1999; (2):155-67.

Status: Not included because dementia population not randomized to treatment

Schrijnemaekers V, van Rossum E, Candel M, et al. Effects of emotion-oriented care on elderly people with cognitive impairment and behavioral problems. *Int J Geriatr Psychiatry* 2002 Oct; 17(10):926-37.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schuck S, Lebreton S. Aminotransferase levels and silymarine in de novo tacrine-treated Alzheimer's disease patients. *Fundam Clin Pharmacol* 1999; 13(3):422.
Status: Not included because not a full article

Schulz R, O'Brien A, Czaja S, et al. Dementia caregiver intervention research: in search of clinical significance. *Gerontologist* 2002 Oct; 42(5):589-602.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Schwartz AS, Kohlstaedt EV. Physostigmine effects in Alzheimer's disease: Relationship to dementia severity. *Life Sci* 1986 Mar 17; 38(11):1021-8.
Status: Not included because no extractable data relevant to review

Schwartz BL, Hashtroudi S, Herting RL, et al. D-Cycloserine enhances implicit memory in Alzheimer patients. *Neurology* 1996 Feb; 46(2):420-4.
Status: Not included because no extractable data relevant to review

Schweiger C. A 48-week, placebo-controlled study examining propentofylline's safety, efficacy, and impact on disease progression in patients with vascular dementia. *J Eur Coll Neuropsychopharmacol* 1999; (Suppl 5):S319.
Status: Not included because not a full article

Seipel JH, Fisher R, Floam JE, et al. Rheoencephalographic and other studies of betahistine in humans. III. Improved methods of diagnosis and selection in arteriosclerotic dementia. *J Clin Pharmacol* 1977 Jan; 17(1):63-75.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Seipel JH, Fisher R, Blatchley RJ, et al. Rheoencephalographic and other studies of

betahistine in humans. IV. Prolonged administration with improvement in arteriosclerotic dementia. *J Clin Pharmacol* 1977 Feb; 17(2-3):140-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Semlitsch HV, Anderer P, Saletu B, et al. Topographic mapping of cognitive event-related potentials in a double-blind, placebo-controlled study with the hemoderivative Actovegin in age-associated memory impairment. *Neuropsychobiology* 1990 Sep; 24(1):49-56.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Semlitsch HV, Anderer P, Saletu B. Topographic mapping of long latency "cognitive" event-related potentials (P 300): A double-blind, placebo-controlled study with amantadine in mild dementia. *J Neural Transm Park Dis Dement Sect* 1992; 4:319-36.
Status: Not included because no extractable data relevant to review

Semlitsch HV, Anderer P, Saletu B, et al. Cognitive psychophysiology in nootropic drug research: Effects of Ginkgo biloba on event-related potentials (P300) in age-associated memory impairment. *Pharmacopsychiatry* 1995 Jul; 28(4):134-42.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Senin U, Parnetti L, Barbagallo-Sangiorgi G, et al. Idebenone in senile dementia of Alzheimer type: A multicentre study. *Arch Gerontol Geriatr* 1992; 15(3):249-60.
Status: Not included because Jadad Quality Scale score less than three

Serafetinides EA, Willis D, Clark ML. The EEG effects of zinc in geriatric psychiatric patients. *Int Pharmacopsychiatry* 1974; 9(2):95-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Serby M, Angrist B, Corwin J, et al. Cholecystokinin octapeptide in dementia. *Psychopharmacol Bull* 1984; 20(3):546-7.
Status: Not included because dementia population not randomized to treatment

Serby M, Resnick R, Jordan B, et al. Naltrexone and Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 1986;

10(3-5):587-90.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Serfaty M, Kennell-Webb S, Warner J, et al. Double blind randomised placebo controlled trial of low dose melatonin for sleep disorders in dementia. *Int J Geriatr Psychiatry* 2002; 17(12):1120-7.
Status: Cross-over trial;

Seux M-L, Forette F, Staessen JA, et al. Treatment of isolated systolic hypertension and dementia prevention in older patients. Results of the Systolic Hypertension in Europe trial (SYST-EUR) vascular dementia project. *Eur Heart J Suppl* 1999; 1(M):M6-M12.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sevush S, Guterman A, Villalon AV. Improved verbal learning after outpatient oral physostigmine therapy in patients with dementia of the Alzheimer type. *J Clin Psychiatry* 1991 Jul; 52(7):300-3.
Status: Not included because dementia population not randomized to treatment

Shaw FE, Bond J, Richardson DA, et al. Multifactorial intervention after a fall in older people with cognitive impairment and dementia presenting to the accident and emergency department: Randomised controlled trial. *BMJ* 2003; 326(7380):73-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Shaw TG, Meyer JS. Double-blind trial of oral papaverine in chronic cerebrovascular ischemia. *Angiology* 1978 Nov; 29(11):839-51.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sheikh JI, Hill RD, Yesavage JA. Long-term efficacy of cognitive training for age-associated memory impairment: A six-month follow-up study. *Dev Neuropsychol* 1986; 2(4):413-21.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Shelton P, Schraeder C, Dworak D, et al. Caregivers' utilization of health services: results from the Medicare Alzheimer's Disease Demonstration, Illinois site. *J Am Geriatr Soc* 2001 Dec; 49(12):1600-5.
Status: Not included because does not meet criteria for treatment for dementia patients

Sherwin BB. Estrogen and cognitive functioning in men with Mild Cognitive Impairment. *J Mol Neurosci* 2002; 19(1-2):219-23.

Status: Not included because no extractable data relevant to review

Shua-Haim JR, Shua-Haim V, Sabo M, et al. Case report: Donepezil in the treatment of advanced Alzheimer's disease. *Ann Long Term Care* 1999; 7(2):67-71.

Status: Not included because dementia population not randomized to treatment

Shua-Haim JR, Shua-Haim V, Comsti E, et al. Donepezil (Aricept(TM)) treatment of multi infarct dementia: The caregivers and clinical impression. *Am J Alzheimers Dis* 2000; 15(4):201-11.

Status: Not included because dementia population not randomized to treatment

Shumaker SA, Reboussin BA, Espeland MA, et al. The Women's Health Initiative Memory Study (WHIMS): A trial of the effect of estrogen therapy in preventing and slowing the progression of dementia. *Control Clin Trials* 1998 Dec; 19(6):604-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sidtis JJ, Gatsonis C, Price RW, et al. Zidovudine treatment of the AIDS dementia complex: Results of a placebo-controlled trial. *AIDS Clinical Trials Group. Ann Neurol* 1993 Apr; 33(4):343-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Silverberg GD, Levinthal E, Sullivan EV, et al. Assessment of low-flow CSF drainage as a treatment for AD: Results of a randomized pilot study. *Neurology* 2002 Oct 22; 59(8):1139-45.

Status: Not included because does not meet criteria for treatment for dementia patients

Sinforiani E, Iannuccelli M, Mauri M, et al. Neuropsychological changes in demented patients treated with acetyl-L-carnitine. *Int J Clin Pharmacol Res* 1990; 10(1-2):69-74.

Status: Not included because Jadad Quality Scale score less than three

Sival RC, Haffmans PM, Jansen PA, et al. Sodium valproate in the treatment of aggressive behavior in patients with dementia: A randomized placebo controlled clinical trial. *Int J Geriatr*

Psychiatry 2002; 17(6):579-85.

Status: Cross-over trial;

Skelly J, Flint A, Brunt S, et al. Treatment of urinary incontinence in dementia using low dose oxybutynin chloride. *Age Ageing* 1995; 24(S):19.

Status: Not included because not a full article

Slooter AJC, Houwing-Duistermaat JJ, van Harskamp F, et al. Apolipoprotein E genotype and progression of Alzheimer's disease: The Rotterdam Study. *J Neurol* 1999; 246(4):304-8.

Status: Not included because dementia population not randomized to treatment

Smallwood J, Brown R, Coulter F, et al. Aromatherapy and behaviour disturbances in dementia: A randomized controlled trial. *Int J Geriatr Psychiatry* 2001 Oct; 16(10):1010-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smid HG, Trumper BG, Pottag G, et al. Differentiation of hypoglycaemia induced cognitive impairments. An electrophysiological approach. *Brain* 1997 Jun; 120(Pt 6):1041-56.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith CM, Swash M, Exton-Smith AN, et al. Choline therapy in Alzheimer's disease. *Lancet* 1978 Aug 5; 2(8084):318.

Status: Not included because not a full article

Smith CM, Swash M. Physostigmine in Alzheimer's disease. *Lancet* 1979 Jan 6; 1(8106):42.

Status: Not included because not a full article

Smith DF, Stromgren E, Petersen HN. Lack of effect of tryptophan treatment in demented gerontopsychiatric patients. A double-blind, crossover-controlled study. *Acta Psychiatr Scand* 1984; 70(5):470-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith DJ, Yuhknevich S. Adverse reactions to rivastigmine in three cases of dementia. *Aust N Z J Psychiatry* 2001; 35(5):694-5.

Status: Not included because not a full article

Smith F. Mixed-model analysis of incomplete longitudinal data from a high-dose trial of tacrine (Cognex) in Alzheimer's patients. *J Biopharm Stat* 1996 Mar; 6(1):59-67.

Status: Not included because Jadad Quality Scale score less than three

Smith F, Gracon S, Knopman D, et al. Survival analysis to evaluate the effect of long-term tacrine (Cognex) treatment on nursing home placement in Alzheimer's patients. *J Neurol* 1997; 244(Suppl 3):S88.

Status: Not included because not a full article

Smith F, Gracon S, Knopman D, et al. Tacrine treatment and nursing home placement: Application of the Cox proportional hazards model with time-dependent covariates. *Drug Inf J* 1998; 32(3):729-35.

Status: Not included because dementia population not randomized to treatment

Smith GR, Taylor CW, Linkous P. Haloperidol versus thioridazine for the treatment of psychogeriatric patients: A double-blind clinical trial. *Psychosomatics: Journal of Consultation Liaison Psychiatry* 1974; 15(3):134-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith MK. The effects of participatory music on the reality orientation and sociability of Alzheimer's residents in a long-term-care setting. *Activities Adaptation Aging* 1994; 18(2):41-55.
Status: Not included because dementia population not randomized to treatment

Smith RC, Vroulis G, Johnson R, et al. Comparison of therapeutic response to long-term treatment with lecithin versus piracetam plus lecithin in patients with Alzheimer's disease. *Psychopharmacol Bull* 1984; 20(3):542-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Smith WL, Lowrey JB, Davis JA. The effects of cyclandelate on psychological test performance in patients with cerebral vascular insufficiency. *Curr Ther Res Clin Exp* 1968; 10(12):613-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Snaedal J, Johannesson T, Jonsson JE, et al. The effects of nicotine in dermal plaster on cognitive functions in patients with Alzheimer's disease. *Dementia* 1996 Jan; 7(1):47-52.
Status: Not included because dementia population not randomized to treatment

Snyder M, Egan EC, Burns KR. Interventions for decreasing agitation behaviors in persons with dementia. *J Gerontol Nurs* 1995 Jul; 21(7):34-40.
Status: Not included because dementia population not randomized to treatment

Snyder M, Tseng Y, Brandt C, et al. A glider swing intervention for people with dementia. *Geriatr Nurs (Minneap)* 2001 Mar; 22(2):86-90.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sobel BP. Bingo vs. physical intervention in stimulating short-term cognition in Alzheimer's disease patients. *Am J Alzheimers Dis Other Demen* 2001 Mar; 16(2):115-20.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sobow TM, Maczkiewicz M, Kloszewska I. Tianeptine versus fluoxetine in the treatment of depression complicating Alzheimer's disease. *Int J Geriatr Psychiatry* 2001; 16(11):1108-9.
Status: Not included because not a full article

Solomon PR, Knapp MJ, Gracon SI, et al. Long-term tacrine treatment in patients with Alzheimer's disease. *Lancet* 1996 Jul 27; 348(9022):275-6.
Status: Not included because not a full article

Soncrant TT, Raffaele KC, Asthana S, et al. Memory improvement without toxicity during chronic, low dose intravenous arecoline in Alzheimer's disease. *Psychopharmacologia* 1993; 112(4):421-7.
Status: Cross-over trial;

Sourander LB, Portin R, Molsa P, et al. Senile dementia of the Alzheimer type treated with aniracetam: A new nootropic agent. *Psychopharmacologia* 1987; 91(1):90-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Spagnolo C, Dallasta D, Iannuccelli M. A controlled double-blind trial comparing etoperidone with thioridazine in the management of severe senile dementia. *Drugs Exp Clin Res* 1983; 9(12):873-80.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Spector A, Orrell M, Davies S, et al. Can reality orientation be rehabilitated? Development and piloting of an evidence-based programme of cognition-based therapies for people with

dementia. Neuropsychol Rehab 2001; 11(3-4):377-97.

Status: Not included because does not meet criteria for treatment for dementia patients

Sramek JJ, Cutler NR, Hurley DJ, et al. The utility of salivary amylase as an evaluation of M3 muscarinic agonist activity in Alzheimer's disease. Prog Neuropsychopharmacol Biol Psychiatry 1995 Jan; 19(1):85-91.

Status: Not included because does not meet criteria for treatment for dementia patients

Sramek JJ, Sedman AJ, Reece PA, et al. Safety and tolerability of CI-979 in patients with Alzheimer's disease. Life Sci 1995; 57(5):503-10.

Status: Not included because no extractable data relevant to review

Sramek JJ, Hurley DJ, Wardle TS, et al. The safety and tolerance of xanomeline tartrate in patients with Alzheimer's disease. J Clin Pharmacol 1995 Aug; 35(8):800-6.

Status: Not included because no extractable data relevant to review

Sramek JJ, Viereck C, Huff FJ, et al. A "bridging" (safety/tolerance) study of besipirdine hydrochloride in patients with Alzheimer's disease. Life Sci 1995; 57(12):1241-8.

Status: Not included because no extractable data relevant to review

Sramek JJ, Block GA, Reines SA, et al. A multiple-dose safety trial of eptastigmine in Alzheimer's disease, with pharmacodynamic observations of red blood cell cholinesterase. Life Sci 1995; 56(5):319-26.

Status: Not included because no extractable data relevant to review

Sramek JJ, Anand R, Wardle TS, et al. Safety/tolerability trial of SDZ ENA 713 in patients with probable Alzheimer's disease. Life Sci 1996; 58(15):1201-7.

Status: Not included because no extractable data relevant to review

Sramek JJ, Forrest M, Mengel H, et al. A bridging study of LU 25-109 in patients with probable Alzheimer's disease. Life Sci 1998; 62(3):195-202.

Status: Not included because dementia population not randomized to treatment

Sramek JJ, Hourani J, Jhee SS, et al. NXX-066 in patients with Alzheimer's disease: A bridging study. Life Sci 1999; 64(14):1215-21.

Status: Not included because no extractable data relevant to review

St Clair D, Norrman J, Perry R, et al. Apolipoprotein E epsilon 4 allele frequency in patients with Lewy body dementia, Alzheimer's disease and age-matched controls. Neurosci Lett 1994 Jul 18; 176(1):45-6.

Status: Not included because dementia population not randomized to treatment

Stapleton JM, Eckardt MJ, Martin P, et al. Treatment of alcoholic organic brain syndrome with the serotonin reuptake inhibitor fluvoxamine: A preliminary study. Adv Alcohol Subst Abuse 1988; 7(3-4):47-4.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Starr JM, Whalley LJ. Hypertensive Old People in Edinburgh (HOPE) Study: Electrocardiographic changes after captopril or bendrofluzide treatment. Age Ageing 1993 Sep; 22(5):343-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Starr JM, Whalley LJ, Inch S, et al. A double-blind trial of captopril or bendrofluzide in newly diagnosed senile hypertension. Curr Med Res Opin 1994; 13(4):214-21.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Starr JM, Whalley LJ, Deary IJ. The effects of antihypertensive treatment on cognitive function: Results from the HOPE study. J Am Geriatr Soc 1996 Apr; 44(4):411-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Steele C, Lucas MJ, Tune L. Haloperidol versus thioridazine in the treatment of behavioral symptoms in senile dementia of the Alzheimer's type: Preliminary findings. J Clin Psychiatry 1986 Jun; 47(6):310-2.

Status: Not included because dementia population not randomized to treatment

Stegink AJ. The clinical use of piracetam, a new nootropic drug. The treatment of symptoms of senile involution. Arzneimittelforschung 1972 Jun; 22(6):975-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Steiger William A. Effects of naloxone in treatment of senile dementia. *J Am Geriatr Soc* 1985; 33(2):155.

Status: Not included because not a full article

Stern FH. Management of chronic brain syndrome secondary to cerebral arteriosclerosis, with special reference to papaverine hydrochloride. *J Am Geriatr Soc* 1970 Jun; 18(6):507-12.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Stern Y, Sano M, Mayeux R. Effects of oral physostigmine in Alzheimer's disease. *Ann Neurol* 1987 Sep; 22(3):306-10.

Status: Not included because dementia population not randomized to treatment

Stern Y, Sano M, Mayeux R. Long-term administration of oral physostigmine in Alzheimer's disease. *Neurology* 1988 Dec; 38(12):1837-41.

Status: Cross-over trial;

Stevermer JJ, Lindbloom EJ. Ginkgo biloba for dementia. *J Fam Pract* 1998; 46(1):20.

Status: Not included because not a full article

Stewart A, Phillips R, Dempsey G. Pharmacotherapy for people with Alzheimer's disease: A Markov-cycle evaluation of five years' therapy using donepezil. *Int J Geriatr Psychiatry* 1998 Jul; 13(7):445-53.

Status: Not included because dementia population not randomized to treatment

Stewart WF, Kawas C, Corrada M, et al. Risk of Alzheimer's disease and duration of NSAID use. *Neurology* 1997 Mar; 48(3):626-32.

Status: Not included because does not meet criteria for treatment for dementia patients

Stoppe G, Sandholzer H, Staedt J, et al. Sleep disturbances in the demented elderly: Treatment in ambulatory care. *Sleep* 1995; 18(10):844-8.

Status: Not included because dementia population not randomized to treatment

Stotsky BA, Cole JO, Tang YT, et al. Sodium butabarbital (butisol sodium) as an hypnotic agent for aged psychiatric patients with sleep disorders. *J Am Geriatr Soc* 1971 Oct; 19(10):860-70.

Status: Not included because no extractable data relevant to review

Stotsky BA, Cole JO, Lu LM, et al. A controlled study of the efficacy of pentyleneetetrazol (Metrazol) with hard-core hospitalized psychogeriatric patients. *Am J Psychiatry* 1972 Oct; 129(4):387-91.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Stracciari A, Ciucci G, Loreta R, et al. Multi-infarct dementia: Clinical trial with acetyl-L-carnitine vs placebo. *J Neurol* 1988; 235(Suppl):92.

Status: Not included because not a full article

Street J, Clark WS, Gannon KS, et al. Reduction of psychotic symptoms in patients with Lewy Body-like symptoms treated with olanzapine. *J Eur Coll Neuropsychopharmacol* 1999; 9(Suppl 5):S331.

Status : Not included because not a full article

Streim JE, Oslin DW, Katz IR, et al. Drug treatment of depression in frail elderly nursing home residents. *Am J Geriatr Psychiatry* 2000; 8(2):150-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sudilovsky A, Cutler NR, Sramek JJ, et al. A pilot clinical trial of the angiotensin-converting enzyme inhibitor ceranapril in Alzheimer's disease. *Alzheimer Dis Assoc Disord* 1993; 7(2):105-11.

Status: Cross-over trial;

Sultzer DL, Gray KF, Gunay I, et al. A double-blind comparison of trazodone and haloperidol for treatment of agitation in patients with dementia. *Am J Geriatr Psychiatry* 1997; 5(1):60-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sultzer DL, Gray KF, Gunay I, et al. Does behavioral improvement with haloperidol or trazodone treatment depend on psychosis or mood symptoms in patients with dementia? *J Am Geriatr Soc* 2001 Oct; 49(10):1294-300.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sumiyoshi T, Matsui M, Nohara S, et al. Enhancement of cognitive performance in schizophrenia by addition of tandospirone to neuroleptic treatment. *Am J Psychiatry* 2001; 158(10):1722-5.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Summers WK, Majovski LV, Marsh GM, et al. Oral tetrahydroaminoacridine in long-term treatment of senile dementia, Alzheimer type. *N Engl J Med* 1986 Nov 13; 315(20):1241-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Sunderland T, Tariot P, Murphy DL, et al. Scopolamine challenges in Alzheimer's disease. *Psychopharmacologia* 1985; 87(2):247-9.
Status: Cross-over trial;

Sunderland T, Tariot PN, Cohen RM, et al. Anticholinergic sensitivity in patients with dementia of the Alzheimer type and age-matched. *Arch Gen Psychiatry* 1987; 44(5):418-26.
Status: Not included because Jadad Quality Scale score less than three

Sunderland T, Weingartner H, Cohen RM, et al. Low-dose oral lorazepam administration in Alzheimer subjects and age-matched controls. *Psychopharmacologia* 1989; 99(1):129-33.
Status: Cross-over trial;

Sunderland T, Molchan S, Lawlor B, et al. A strategy of "combination chemotherapy" in Alzheimer's disease: Rationale and preliminary results with physostigmine plus deprenyl. *Int Psychogeriatr* 1992; 4(Suppl 2):291-309.
Status: Not included because dementia population not randomized to treatment

Swanson EA, Maas ML, Buckwalter KC. Alzheimer's residents' cognitive and functional measures: Special and traditional care unit comparison. *Clin Nurs Res* 1994 Feb; 3(1):27-41.
Status: Not included because does not meet criteria for treatment for dementia patients

Swanson EA, Maas ML, Buckwalter KC. Catastrophic reactions and other behaviors of Alzheimer's residents: Special unit compared with traditional units. *Arch Psychiatr Nurs* 1993; 7(5):292-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Swartz JR, Miller BL, Lesser IM, et al. Frontotemporal dementia: Treatment response to serotonin selective reuptake inhibitors. *J Clin Psychiatry* 1997; 58(5):212-6.

Status: Not included because dementia population not randomized to treatment

Tabet N, Mantle D, Walker Z, et al. Dietary and endogenous antioxidants in dementia. *Int J Geriatr Psychiatry* 2001; 16(6):639-41.
Status: Not included because not a full article

Tabet N, Mantle D, Walker Z, et al. Endogenous antioxidant activities in relation to concurrent vitamins A, C, and E intake in dementia. *Int Psychogeriatr* 2002; 14(1):7-15.
Status: Not included because does not meet criteria for treatment for dementia patients

Tabourne CE. The effects of a life review program on disorientation, social interaction and self-esteem of nursing home residents. *Int J Aging Hum Dev* 1995; 41(3):251-66.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Talwalker S. The cardinal features of cognitive and noncognitive dysfunction and the differential efficacy of tacrine in Alzheimer's disease patients. *J Biopharm Stat* 1996 Nov; 6(4):443-56.
Status: Not included because no extractable data relevant to review

Talwalker S, Overall JE, Srirama MK, et al. Cardinal features of cognitive dysfunction in Alzheimer's disease: A factor-analytic study of the Alzheimer's Disease Assessment Scale. *J Geriatr Psychiatry Neurol* 1996 Jan; 9(1):39-46.
Status: Not included because does not meet criteria for treatment for dementia patients

Talwalker S. Analysis of repeated measurements with dropouts among Alzheimer's disease patients using summary measures and meta-analysis. *J Biopharm Stat* 1996; 6(1):49-58.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tappen RM. The effect of skill training on functional abilities of nursing home residents with dementia. *Res Nurs Health* 1994 Jun; 17(3):159-65.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tappen RM, Roach KE, Applegate EB, et al. Effect of a combined walking and conversation intervention on functional mobility of nursing home residents with Alzheimer disease. *Alzheimer Dis Assoc Disord* 2000 Oct; 14(4):196-201.

Status: Not included because does not meet criteria for treatment for dementia patients

Tappen RM, Williams CL, Barry C, et al. Conversation intervention with Alzheimer's patients: Increasing the relevance of communication. Clin Gerontol 2002; 24(3-4):63-75.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Targum SD, Wieland S, Glasky MS, et al. Evaluation of AIT-082 in patients with mild to moderate senile dementia of the Alzheimer's type. J Eur Coll Neuropsychopharmacol 1999; (Suppl 5):S320.

Status : Not included because not a full article

Tariot P, Parys W, Kershaw P. The efficacy and tolerability of galantamine in Alzheimer's disease a 5 month placebo controlled study with slow dose escalation. Neurology 2000; 54(Suppl 3):A415.

Status: Not included because not a full article

Tariot PN, Sunderland T, Weingartner H, et al. Low- and high-dose naloxone in dementia of the Alzheimer type. Psychopharmacol Bull 1985; 21(3):680-2.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Sunderland T, Murphy DL, et al. Design and interpretation of opiate antagonist trials in dementia. Prog Neuropsychopharmacol Biol Psychiatry 1986; 10(3-5):611-26.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Sunderland T, Weingartner H, et al. Naloxone and Alzheimer's disease. Cognitive and behavioral effects of a range of doses. Arch Gen Psychiatry 1986 Aug; 43(8):727-32.

Status: Cross-over trial;

Tariot PN, Cohen RM, Sunderland T, et al. L-deprenyl in Alzheimer's disease. Preliminary evidence for behavioral change with monoamine oxidase B inhibition. Arch Gen Psychiatry 1987 May; 44(5):427-33.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Sunderland T, Weingartner H, et al. Cognitive effects of L-deprenyl in Alzheimer's disease. Psychopharmacologia 1987; 91(4):489-95.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Cohen RM, Welkowitz JA, et al. Multiple-dose arecoline infusions in Alzheimer's disease. Arch Gen Psychiatry 1988 Oct; 45(10):901-5.

Status: Cross-over trial;

Tariot PN, Sunderland T, Cohen RM, et al. Tranylcypromine compared with L-deprenyl in Alzheimer's disease. J Clin Psychopharmacol 1988 Feb; 8(1):23-7.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Gross M, Sunderland T, et al. High-dose naloxone in older normal subjects: Implications for Alzheimer's disease. J Am Geriatr Soc 1988 Aug; 36(8):681-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tariot PN, Leibovici A, Erb R, et al. Carbamazepine therapy for agitation in dementia: Status report. Am J Geriatr Psychiatry 1994; 2(3):257-9.

Status: Not included because not a full article

Tariot PN, Erb R, Leibovici A, et al. Carbamazepine treatment of agitation in nursing home patients with dementia: A preliminary study. J Am Geriatr Soc 1994 Nov; 42(11):1160-6.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Frederiksen K, Erb R, et al. Lack of carbamazepine toxicity in frail nursing home patients: A controlled study. J Am Geriatr Soc 1995 Sep; 43(9):1026-9.

Status: Not included because dementia population not randomized to treatment

Tariot PN, Goldstein B, Podgorski CA, et al. Short-term administration of selegiline for mild-to-moderate dementia of the Alzheimer's type. Am J Geriatr Psychiatry 1998; 6(2):145-54.

Status: Cross-over trial;

Tariot PN, Upadhyaya A, Sunderland T, et al. Physiologic and neuroendocrine responses to intravenous naloxone in subjects with Alzheimer's disease and age-matched controls. Biol Psychiatry 1999 Aug 1; 46(3):412-9.

Status: Not included because no extractable data relevant to review

Tariot PN, Salzman C, Yeung PP, et al. Long-Term use of quetiapine in elderly patients with psychotic disorders. *Clin Ther* 2000 Sep; 22(9):1068-84.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tekin S, Aykut BC, Tanridag T, et al. Antiglutamatergic therapy in Alzheimer's disease: effects of Lamotrigine. *J Neural Transm Gen Sect* 1998; 105(2-3):295-303.

Status: Cross-over trial;

Templeton L, Barker A, Wesnes K, et al. A double-blind, placebo-controlled single dose trial of intravenous flumazenil in Alzheimer's disease. *Hum Psychopharmacol* 1999; 14(4):239-45.

Status: Cross-over trial;

Tennant FS. Preliminary observations on naltrexone for treatment of Alzheimer's type dementia. *J Am Geriatr Soc* 1987; 35(4):369-70.

Status: Not included because not a full article

ter Haar HW. A comparison of chlormethiazole and haloperidol in the treatment of elderly patients with confusion of organic and psychogenic origin: A double-blind crossover study.

Pharmatherapeutica 1977; 1(9):563-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Terano T, Fujishiro S, Ban T, et al. Docosahexaenoic acid supplementation improves the moderately severe dementia from thrombotic cerebrovascular diseases. *Lipids* 1999; 34(Suppl):S345-S346.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Teri L, Reifler BV, Veith RC, et al. Imipramine in the treatment of depressed Alzheimer's patients: Impact on cognition. *J Gerontol* 1991 Nov; 46(6):P372-P377.

Status: Not included because Jadad Quality Scale score less than three

Teri L. Behavioral treatment of depression patients with dementia. *Alzheimer Dis Assoc Disord* 1994; 8(Suppl 3):66-74.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Teri L, Logsdon RG, Uomoto J, et al. Behavioral treatment of depression in dementia patients: A

controlled clinical trial. *J Gerontol B Psychol Sci Soc Sci* 1997 Jul; 52(4):159-66.

Status: Not included because does not meet criteria for treatment for dementia patients

Teri L, Logsdon RG, Whall AL, et al. Treatment for agitation in dementia patients: A behavior management approach. *Psychotherapy: Theory, Research, Practice, Training* 1998; 35(4):354-443.

Status: Not included because dementia population not randomized to treatment

Tewfik GI, Jain VK, Harcup M, et al. Effectiveness of various tranquilisers in the management of senile restlessness. *Gerontol Clin (Basel)* 1970; 12(6):351-9.

Status: Not included because dementia population not randomized to treatment

Thal LJ, Rosen W, Sharpless NS, et al. Choline chloride fails to improve cognition in Alzheimer's disease. *Neurobiol Aging* 1981; (3):205-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thal LJ, Fuld PA, Masur DM, et al. Oral physostigmine and lecithin improve memory in Alzheimer disease. *Ann Neurol* 1983; 13(5):491-6.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thal LJ, Masur DM, Sharpless NS, et al. Acute and chronic effects of oral physostigmine and lecithin in Alzheimer's disease. *Prog Neuropsychopharmacol Biol Psychiatry* 1986; 10(3-5):627-36.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thibault A. A double-blind evaluation of 'Hydergine' and placebo in the treatment of patients with organic brain syndrome and cerebral arteriosclerosis in a nursing home. *Curr Med Res Opin* 1974; 2(8):482-7.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Thienhaus OJ, Wheeler BG, Simon S. A controlled double-blind study of high-dose dihydroergotamine mesylate (Hydergine(R)) in mild dementia. *J Am Geriatr Soc* 1987; 35(3):219-23.

Status: Not included because Jadad Quality Scale score less than three

Thorpe L, Middleton J, Russell G, et al. Bright light therapy for demented nursing home patients with behavioural disturbance. *Am J Alzheimers Dis* 2000; 15(1):18-26.

Status: Not included because dementia population not randomized to treatment

Tian J, Du H, Yang H, et al. A clinical study on compound Da Huang (Radix et Rhizoma Rhei) preparations for improvement of senile persons' memory ability. *J Tradit Chin Med* 1997 Sep; 17(3):168-73.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tian J, Yin J, Liu H, et al. Jian Nao Ning for treatment of memory impairment in patients with mild to moderate multi-infarct dementia. *J Tradit Chin Med* 2002 Dec; 22(4):247-51.

Status: Not included because Jadad Quality Scale score less than three

Tignol J, Pujol-Domenech J, Chartres JP, et al. Double-blind study of the efficacy and safety of milnacipran and imipramine in elderly patients with major depressive episode. *Acta Psychiatr Scand* 1998 Feb; 97(2):157-65.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tinetti ME, Baker D, Gallo WT, et al. Evaluation of restorative care vs usual care for older adults receiving an acute episode of home care. *JAMA* 2002 Apr 24; 287(16):2098-105.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tollefson GD. Short-term effects of the calcium channel blocker nimodipine (Bay-e-9736) in the management of primary degenerative dementia. *Biol Psychiatry* 1990 May 15; 27(10):1133-42.

Status: Not included because Jadad Quality Scale score less than three

Toseland RW, Diehl M, Freeman K, et al. The impact of validation group therapy on nursing home residents with dementia. *J Appl Gerontol* 1997; 16(1):31-50.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Trappler B, Cohen CI. Use of SSRIs in "very old" depressed nursing home residents. *Am J Geriatr Psychiatry* 1998; 6(1):83-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Trappler B, Cohen CI. Using fluoxetine in "very old" depressed nursing home residents. *Am J Geriatr Psychiatry* 1996; 4(3):258-62.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tsai GE, Falk WE, Gunther J. A preliminary study of D-cycloserine treatment in Alzheimer's disease. *J Neuropsychiatry Clin Neurosci* 1998; 10(2):224-6.

Status: Cross-over trial;

Tsai GE, Falk WE, Gunther J, et al. Improved cognition in Alzheimer's disease with short-term D-cycloserine treatment. *Am J Psychiatry* 1999 Mar; 156(3):467-9.

Status: Cross-over trial;

Tsiskaridze A, Vashadze A. Comparison study of Amiridine and Piracebral effects on cognition in patients with primary dementias. *J Neural Transm Gen Sect* 1996; 103:LII.

Status: Not included because not a full article

Tsuang MM, Lu LM, Stotsky BA, et al. Haloperidol versus thioridazine for hospitalized psychogeriatric patients: Double-blind study. *J Am Geriatr Soc* 1971 Jul; 19(7):593-600.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tune L, Brandt J, Frost JJ, et al. Physostigmine in Alzheimer's disease: Effects on cognitive functioning, cerebral glucose metabolism analyzed by positron emission tomography and cerebral blood flow analyzed by single photon emission tomography. *Acta Psychiatr Scand Suppl* 1991; 366:61-5.

Status: Cross-over trial;

Tune LE, Steele C, Cooper T. Neuroleptic drugs in the management of behavioral symptoms of Alzheimer's disease. *Psychiatr Clin North Am* 1991 Jun; 14(2):353-73.

Status: Not included because dementia population not randomized to treatment

Turek I, Kurland AA, Ota KY, et al. Effects of pipradrol hydrochloride on geriatric patients. *J Am Geriatr Soc* 1969 Apr; 17(4):408-13.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Tweedy JR, Garcia CA. Lecithin treatment of cognitively impaired Parkinson's patients. *Eur J*

- Clin Invest 1982 Feb; 12(1):87-90.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Udani JK. Extract of ginkgo biloba for treatment of dementia. Integr Med 1998; (1):43-4.
Status: Not included because not a full article
- Ulmar G, Weickelt G. Anticholinergic treatment, cognition and the acetylcholine hypothesis of Alzheimer's disease (AD). J Neural Transm Park Dis Dement Sect 1989; 1:144.
Status: Not included because not a full article
- Underhill JA. Exercise for older people with dementia. Age Ageing 1993; 22(Suppl 3):35.
Status: Not included because not a full article
- Uney JB, Jones GM, Rebeiro A, et al. The effect of long-term high dose lecithin on erythrocyte choline transport in Alzheimer patients. Biol Psychiatry 1992 Mar 15; 31(6):630-3.
Status: Not included because no extractable data relevant to review
- van Asselt DZ, Pasman JW, van Lier HJ, et al. Cobalamin supplementation improves cognitive and cerebral function in older, cobalamin-deficient persons. J Gerontol A Biol Sci Med Sci 2001 Dec; 56(12):M775-M779.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- van Diepen E, Baillon SF, Redman J, et al. A pilot study of the physiological and behavioural effects of snoezelen in dementia. Br J Occup Ther 2002; 65(2):61-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- van Dongen MC, van Rossum E, Kessels AG, et al. The efficacy of ginkgo for elderly people with dementia and age-associated memory impairment: New results of a randomized clinical trial. J Am Geriatr Soc 2000 Oct; 48(10):1183-94.
Status: Not included because no extractable data relevant to review
- van Duijn H, Wolters EC, Beckmann MKF. Desglycinamide arginine vasopressin (DGAVP) in Alzheimer's disease. Clin Neurol Neurosurg 1987; 2:21
Status: Not included because not a full article
- Van Dyck CH, McMahon TJ, Rosen MI, et al. Sustained-release methylphenidate for cognitive impairment in HIV-1-infected drug abusers: A pilot study. J Neuropsychiatry Clin Neurosci 1997; 9(1):29-36.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Van Gool WA. Erratum: effect of hydroxychloroquine on progression of dementia in early alzheimer's disease: An 18-month randomised, double-blind, placebo-controlled study. Lancet 2001; 358(9288):1188.
Status: Not included because not a full article
- Van Loveren-Huyben CM, Engelaar HFWJ, Hermans MBM. Double-blind clinical and psychologic study of ergoloid mesylates (Hydergine®) in subjects with senile mental deterioration. J Am Geriatr Soc 1984; 32(8):584-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Van Reekum R, Clarke D, Conn D, et al. A randomized, placebo-controlled trial of the discontinuation of long-term antipsychotics in dementia. Int Psychogeriatr 2002; 14(2):197-210.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Van Someren EJ, Scherder EJ, Swaab DF. Transcutaneous electrical nerve stimulation (TENS) improves circadian rhythm disturbances in Alzheimer's disease. Alzheimer Dis Assoc Disord 1998 Jun; 12(2):114-8.
Status: Not included because no extractable data relevant to review
- Vangtorp A, Simmelsgaard H, Mellegaard M. Experience with a new butyrophenone derivative (Buronil). Acta Psychiatr Scand Suppl 1968 Jan 1; 203:235-8.
Status: Not included because dementia population not randomized to treatment
- Vecchi GP, Chiari G, Cipolli C, et al. Acetyl-L-carnitine treatment of mental impairment in the elderly: Evidence from a multicentre study. Arch Gerontol Geriatr 1991; (Suppl 2):159-68.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Vellas B, Inglis F, Potkin S, et al. Interim results from an international clinical trial with rivastigmine evaluating a 2-week titration rate in mild to severe Alzheimer's disease patients. Int J Geriatr Psychopharmacol 1998; (3):140-4.

Status: Not included because dementia population not randomized to treatment

Velligan DI, Newcomer J, Pultz J, et al. Does cognitive function improve with quetiapine in comparison to haloperidol? Schizophr Res 2002 Jan 15; 53(3):239-48.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vennerica A, Shanks MF, Staff RT, et al. Cerebral blood flow and cognitive responses to rivastigmine treatment in Alzheimer's disease. NeuroReport 2002; 13(1):83-7.

Status: Not included because dementia population not randomized to treatment

Versiani M, da Silva JA, Mundim FD. Loxapine versus thioridazine in the treatment of organic psychosis. J Int Med Res 1980; 8(1):22-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vespa A, Gori G, Spazzafumo L. Evaluation of non-pharmacological intervention on antisocial behavior in patients suffering from Alzheimer's disease in a day care center. Arch Gerontol Geriatr 2002; 34(1):1-8.

Status: Not included because does not meet criteria for treatment for dementia patients

Vida S, Gauthier L, Gauthier S. Canadian collaborative study of tetrahydroaminoacridine (THA) and lecithin treatment of Alzheimer's disease: Effect on mood. Can J Psychiatry 1989; 34(3):165-70.

Status: Not included because dementia population not randomized to treatment

Villardita C, Parini J, Grioli S, et al. Clinical and neuropsychological study with oxiracetam versus placebo in patients with mild to moderate dementia. J Neural Transm Suppl 1987; 24:293-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Viukari M, Linnoila M. Effect of methyldopa on tardive dyskinesia in psychogeriatric patients. Curr Ther Res Clin Exp ; 18(3):417-24.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Viukari M, Salo H, Lamminsivu U, et al. Tolerance and serum levels of haloperidol during parenteral and oral haloperidol treatment in

geriatric patients. Acta Psychiatr Scand 1982 Apr; 65(4):301-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Viukari M, Jaatinen P, Kylmamaa T. Flunitrazepam, nitrazepam and psychomotor skills in psychogeriatric patients. Curr Ther Res Clin Exp 1983; 33(5):828-34.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Vojtechovsky M, Sobotkova J, Wodniakova J. Secatoxin Spofa-super(R) in the treatment of chronic organic psychosyndrome. Act Nerv Super (Praha) 1981; 23(3):232-3.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Volicer L, Stelly M, Morris J, et al. Effects of dronabinol on anorexia and disturbed behavior in patients with Alzheimer's disease. Int J Geriatr Psychiatry 1997 Sep; 12(9):913-9.

Status: Cross-over trial;

Vroulis GA, Smith RC, Brinkman S, et al. The effects of lecithin on memory in patients with senile dementia of the Alzheimer's type. Psychopharmacol Bull 1981; 17(1):127-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wakelin JS. Fluvoxamine in the treatment of the older depressed patient; Double-blind, placebo-controlled data. Int Clin Psychopharmacol 1986 Jul; 1(3):221-30.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Walker C. Ergoloid mesylates vs. Alzheimer's: The latest round. Geriatrics 1990; 45(12):22, 24.

Status: Not included because not a full article

Walshe TM. The use of nimodipine in vascular dementia. J Neurol 1985; Vol 232:74

Status: Not included because not a full article

Walzl M, Walzl B, Lechner H. Results of a two-month follow-up after single heparin-induced extracorporeal LDL precipitation in vascular dementia. National Stroke Association 1994; (3):179-83.

Status : Not included because no extractable data relevant to review

- Walzl M. A promising approach to the treatment of multi-infarct dementia. *Neurobiol Aging* 2000 Mar; 21(2):283-7.
Status: Not included because Jadad Quality Scale score less than three
- Wang D, Huang X, Du S. A clinical trial on yu cong tang in treatment of senile dementia. *J Tradit Chin Med* 1999 Mar; 19(1):32-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Wang X, Xie Z. A clinical study on the effect of reinforcement of kidney on senile brain functions. *J Tradit Chin Med* 1997 Jun; 17(2):92-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Ward CR, Los Kamp. The effects of participation in an intergenerational program on the behavior of residents with dementia. *Activities Adaptation Aging* 1996; 20(4):61-76.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Wardle J, Rogers P, Judd P, et al. Randomized trial of the effects of cholesterol-lowering dietary treatment on psychological function. *Am J Med* 2000 May; 108(7):547-53.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Waring SC, Rocca WA, Petersen RC, et al. Postmenopausal estrogen replacement therapy and risk of AD: A population-based study. *Neurology* 1999; 52(5):965-70.
Status: Not included because dementia population not randomized to treatment
- Warren PA, Dunn L, Jackson-Clark A. The Medicare Alzheimer's Project in Portland, Oregon. *Pride Inst J Long Term Home Health Care* 1991; 10(2):20-7.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Watkins PB, Zimmerman HJ, Knapp MJ, et al. Hepatotoxic effects of tacrine administration in patients with Alzheimer's disease. *JAMA* 1994 Apr 6; 271(13):992-8.
Status: Not included because dementia population not randomized to treatment
- Watson NM, Wells TJ, Cox C. Rocking chair therapy for dementia patients: Its effect on psychosocial well-being and balance. *Am J Alzheimers Dis* 1998; (6):296-308.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Weiner MF, Tractenberg RE, Sano M, et al. No long-term effect of behavioral treatment on psychotropic drug use for agitation in Alzheimer's disease patients. *J Geriatr Psychiatry Neurol* 2002; 15(2):95-8.
Status: Not included because does not meet criteria for treatment for dementia patients
- Weingartner H, Kaye W, Gold P, et al. Vasopressin treatment of cognitive dysfunction in progressive dementia. *Life Sci* 1981; 29(26):2721-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Weingartner H, Buchsbaum MS, Linnoila M. Zimelidine effects on memory impairments produced by ethanol. *Life Sci* 1983 Nov 28; 33(22):2159-63.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Weiser M, Rotmensch HH, Korczyn AD, et al. A pilot, randomized, open-label trial assessing safety and pharmacokinetic parameters of co-administration of rivastigmine with risperidone in dementia patients with behavioral disturbances. *Int J Geriatr Psychiatry* 2002 Apr; 17(4):343-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Wells D, Dawson P, Sidani S, et al. The benefits of abilities-focused morning care for residents with dementia and their caregivers. *Perspectives (Montclair)* 2000; 24(1):17.
Status: Not included because does not meet criteria for treatment for dementia patients
- Wesensten NJ, Balkin TJ, Davis HQ, et al. Reversal of triazolam- and zolpidem-induced memory impairment by flumazenil. *Psychopharmacologia* 1995 Sep; 121(2):242-9.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD
- Wesensten NJ, Balkin TJ, Belenky GL. Effects of daytime administration of zolpidem versus triazolam on memory. *Eur J Clin Pharmacol* 1995; 48(2):115-22.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wesnes K, Simmons D, Rook M, et al. A double-blind placebo-controlled trial of Tanakan in the treatment of idiopathic cognitive impairment in the elderly. *Hum Psychopharmacol* 1987; 2:159-69.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wesnes KA, McKeith IG, Ferrara R, et al. Effects of rivastigmine on cognitive function in dementia with lewy bodies: A randomised placebo-controlled international study using the cognitive drug research computerised assessment system. *Dement Geriatr Cogn Disord* 2002; 13(3):183-92.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Westreich G, Alter M, Lundgren S. Effect of cyclandelate on dementia. *Stroke* 1975 Sep; 6(5):535-8.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wettstein A. No effect from double-blind trial of physostigmine and lecithin in Alzheimer disease. *Ann Neurol* 1983; 13(2):210-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wettstein A, Spiegel R. Clinical trial with the cholinergic drug RS 86 in Alzheimer's disease (AD) and senile dementia of the Alzheimer type (SDAT). *Psychopharmacologia* 1984; 84(4):572-3.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

White CM, Dicks RS. Rivastigmine: An acetylcholinesterase inhibitor for patients with Alzheimer's disease. *Hosp Formul* 1999; 34(6):493-9.
Status: Not included because dementia population not randomized to treatment

White HK, Levin ED. Four-week nicotine skin patch treatment effects on cognitive performance in Alzheimer's disease. *Psychopharmacologia* 1999 Apr; 143(2):158-65.
Status: Cross-over trial;

White JC, Christensen JF, Singer CM. Methylphenidate as a treatment for depression in acquired immunodeficiency syndrome: an n-of-1 trial. *J Clin Psychiatry* 1992 May; 53(5):153-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

White L, Crovello JN, Rosenberg SN, et al. Evaluation of isobaric oxygenation for the aged with cognitive impairment: Pilot study. *J Am Geriatr Soc* 1975 Feb; 23(2):80-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Whitlatch CJ, Zarit SH, von Eye A. Efficacy of interventions with caregivers: A reanalysis. *Gerontologist* 1991 Feb; (1):9-14.
Status: Not included because does not meet criteria for treatment for dementia patients

Whitlatch CJ, Zarit SH, Goodwin PE, et al. Influence of the success of psychoeducational interventions on the course of family care. *Clin Gerontol* 1995; (1):17-30.
Status: Not included because does not meet criteria for treatment for dementia patients

Wiener PK, Kiosses DN, Klimstra S, et al. A short-term inpatient program for agitated demented nursing home residents. *Int J Geriatr Psychiatry* 2001; 16(9):866-72.
Status: Not included because dementia population not randomized to treatment

Wilcock GK, Scott M, Pearsall T, et al. Galanthamine and the treatment of Alzheimer's disease. *Int J Geriatr Psychiatry* 1993; 8(9):781-2.
Status: Not included because not a full article

Wilcock GK, Surmon DJ, Scott M, et al. An evaluation of the efficacy and safety of tetrahydroaminoacridine (THA) without lecithin in the treatment of Alzheimer's disease. *Age Ageing* 1993; 22(5):316-24.
Status: Cross-over trial;

Wilkinson D, Srikumar S, Shaw K, et al. Drama and movement therapy in dementia: A pilot study. *Arts in Psychotherapy* 1998; (3):195-201.
Status: Not included because dementia population not randomized to treatment

Wilkinson D. Clinical experience with Donepezil (Aricept) in the UK. *J Neural Transm Suppl* 1998; 54:311-5.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wilkinson DG, Hock C, Farlow M, et al. Galantamine provides broad benefits in patients with 'advanced moderate' Alzheimer's disease (MMSE < or = 12) for up to six months. *Int J Clin*

Pract 2002 Sep; 56(7):509-14.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wilkinson DG, Passmore AP, Bullock R, et al. A multinational, randomised, 12-week, comparative study of donepezil and rivastigmine in patients with mild to moderate Alzheimer's disease. *Int J Clin Pract* 2002 Jul; 56(6):441-6.

Status: Not included because Jadad Quality Scale score less than three

Williams R, Reeve W, Ivison D, et al. Use of environmental manipulation and modified informal reality orientation with institutionalized, confused elderly subjects: A replication. *Age Ageing* 1987; 16(5):315-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Willis SL, Schaie KW. Training the elderly on the ability factors of spatial orientation and inductive reasoning. *Psychol Aging* 1986 Sep; 1(3):239-47.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wilson AL, Langley LK, Monley J, et al. Nicotine patches in Alzheimer's disease: Pilot study on learning, memory, and safety. *Pharmacol Biochem Behav* ; 51(2-3):509-3.

Status: Not included because dementia population not randomized to treatment

Wilson RS, Martin EM. New intrathecal drugs in Alzheimer's disease and psychometric testing. *Ann N Y Acad Sci* 1988; 531:180-6.

Status: Not included because dementia population not randomized to treatment

Wincor MZ. Ginkgo biloba for dementia: A reasonable alternative? *J Am Pharm Assoc (Wash)* 1999 May; 39(3):415-6.

Status : Not included because dementia population not randomized to treatment

Winther K, Randlov C, Rein E, et al. Effects of Ginkgo biloba extract on cognitive function and blood pressure in elderly subjects. *Curr Ther Res Clin Exp* 1998; 59(12):881-8.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wishart L, Macerollo J, Loney P, et al. "Special steps": An effective visiting/walking program for persons with cognitive impairment. *Can J Nurs Res* 2000 Mar; 31(4):57-71.

Status: Not included because does not meet criteria for treatment for dementia patients

Wolfgang SA. Olanzapine in whole, not half, tablets for psychosis from Alzheimer's dementia. *Am J Health Syst Pharm* 1999 Nov 1; 56(21):2245-6.

Status: Not included because not a full article

Wolters EC, Riekkinen P, Lowenthal A, et al. DGAVP (Org 5667) in early Alzheimer's disease patients: An international double-blind, placebo-controlled, multicenter trial. *Neurology* 1990 Jul; 40(7):1099-101.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wong YT, Packer H. Penicillin versus penicillins in the treatment of dementia paralytica. *Br J Vener Dis* 2000; 25(1):39.

Status: Not included because not a full article

Woods B. Promoting well-being and independence for people with dementia. *Int J Geriatr Psychiatry* 1999; 14(2):97-109.

Status: Not included because dementia population not randomized to treatment

Wooltorton E. Risperidone (Risperdal): Increased rate of cerebrovascular events in dementia trials. *CMAJ* 2002 Nov 26; 167(11):1269-70.

Status: Not included because not a full article

Wouters-Wesseling W, Wouters AE, Kleijer CN, et al. Study of the effect of a liquid nutrition supplement on the nutritional status of psycho-geriatric nursing home patients. *Eur J Clin Nutr* 2002 Mar; 56(3):245-51.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Wright LK, Litaker M, Laraia MT, et al. Continuum of care for Alzheimer's disease: A nurse education and counseling program. *Issues Ment Health Nurs* 2001 Apr; 22(3):231-52.

Status: Not included because does not meet criteria for treatment for dementia patients

Wroblewski T, Silvestre J, Castaner J. JTP-4819. Cognition enhancer, prolyl endopeptidase inhibitor. *Drugs of the Future* 1998; 23(4):384-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Xu H, Shao N, Cui D, et al. A clinical study of yi zhi capsules in prevention of vascular dementia.

J Tradit Chin Med 2000 Mar; 20(1):10-3.
Status: Not included because Jadad Quality Scale score less than three

Yaffe K, Krueger K, Sarkar S, et al. Cognitive function in postmenopausal women treated with raloxifene. N Engl J Med 2001 Apr 19; 344(16):1207-13.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yamamoto Y, Akiguchi I, Oiwa K, et al. Twenty-four-hour blood pressure changes in the course of lacunar disease. Cardiovasc Dis 2001; 11(2):100-6.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yehuda S, Rabinovitz S, Carasso RL, et al. Essential fatty acids preparation (SR-3) improves Alzheimer's patients quality of life. Int J Neurosci 1996 Nov; 87(3-4):141-4.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yesavage JA, Hollister LE, Burian E. Dihydroergotoxine: 6-mg versus 3-mg dosage in the treatment of senile dementia. Preliminary report. J Am Geriatr Soc 1979 Feb; 27(2):80-2.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Yesavage JA, Westphal J, Rush L. Senile dementia: Combined pharmacologic and psychologic treatment. J Am Geriatr Soc 1981; 29(4):164-71.
Status: Not included because dementia population not randomized to treatment

Yesavage JA, Tinklenberg JR, Hollister LE, et al. Effect of nafrolyl on lactate and pyruvate in the cerebrospinal fluid of patients with senile dementia. J Am Geriatr Soc 1982; 30(2):105-8.
Status: Not included because dementia population not randomized to treatment

Zanetti O, Frisoni GB, De Leo D, et al. Reality orientation therapy in Alzheimer disease: Useful or not? A controlled study. Alzheimer Dis Assoc Disord 1995; 9(3):132-8.
Status: Not included because dementia population not randomized to treatment

Zanetti O, Bianchetti A, Trabucchi M. Cost effectiveness of non pharmacological interventions in Alzheimer's disease. J Am Geriatr

Soc 1998 Nov; 46(11):1481
Status: Not included because not a full article

Zanetti O, Zanieri G, Di Giovanni G, et al. Effectiveness of procedural memory stimulation in mild Alzheimer's disease patients: A controlled study. Neuropsychol Rehab 2001; (3-4):263-4.
Status: Not included because dementia population not randomized to treatment

Zank S, Schacke C. Evaluation of geriatric day care units: effects on patients and caregivers. J Gerontol B Psychol Sci Soc Sci 2002 Jul; 57(4):348-57.
Status: Not included because does not meet criteria for treatment for dementia patients

Zappoli R, Arnetoli G, Paganini M, et al. Contingent negative variation and reaction time in patients with presenile idiopathic cognitive decline and presenile Alzheimer-type dementia. Preliminary report on long-term nicergoline treatment. Neuropsychobiology 1987; 18(3):149-54.
Status: Not included because Jadad Quality Scale score less than three

Zappoli R, Arnetoli G, Paganini M, et al. Topographic bit-mapped event-related neurocognitive potentials and clinical status in patients with primary presenile mental decline chronically treated with nicergoline. Curr Ther Res Clin Exp 1991; 49(6):1078-97.
Status: Not included because Jadad Quality Scale score less than three

Zarit SH, Zarit JM, Reeve KE. Memory training for severe memory loss: Effects on senile dementia patients and their families. Gerontologist 1982; 22(4):373-7.
Status: Not included because does not meet criteria for treatment for dementia patients

Zemlan FP, Keys M, Richter RW, et al. Double-blind placebo-controlled study of velnacrine in Alzheimer's disease. Life Sci 1996; 58(21):1823-32.
Status: Not included because Jadad Quality Scale score less than three

Ziemba C, Foster G, Neufeld R, et al. Haloperidol holiday: Is it a beneficial vacation for some nursing home residents? Clin Gerontol 1997; 17(3):15-24.
Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Zill P, Burger K, Behrens S, et al. Polymorphisms in the alpha-2 macroglobulin gene in psychogeriatric patients. *Neurosci Lett* 2000 Nov 17; 294(2):69-72.

Status: Not included because does not meet criteria for treatment for dementia patients

Zimmer JG, Eggert GM, Chiverton P. Individual versus team case management in optimizing community care for chronically ill patients with dementia. *J Aging Health* 1990; 2(3):357-72.

Status: Not included because does not meet criteria for treatment for dementia patients

Zisselman MH, Rovner BW, Shmueli Y, et al. A pet therapy intervention with geriatric psychiatry inpatients. *Am J Occup Ther* 1996 Jan; 50(1):47-51.

Status: Not included because no extractable data relevant to review

Zissis NP, Alevizos V, Dontas AS. Flunarizine, an inhibitor of Casup +sup 2-induced vascular constriction in geriatric patients. *Curr Ther Res Clin Exp* 1991; 29(31):395-400.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Zivadinov R, Rudick RA, De Masi R, et al. Effects of IV methylprednisolone on brain atrophy in relapsing-remitting MS. *Neurology* 2001 Oct 9; 57(7):1239-47.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

Zwerling I, Plutchik R, Hotz M, et al. Effects of a procaine preparation (Gerovital H3) in hospitalized geriatric patients: A double-blind study. *J Am Geriatr Soc* 1975 Aug; 23(8):355-9.

Status: Not included because dementia population not defined by DSM, NINCDS or ICD

EvTable1. Key characteristics: Carnitine (ALCAR).

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Livingston 1991	IF IS	5	Placebo Acetyl-L-Carnitine	NINCDS	AD	Probable or Possible	71	40	72.8y (65-80y) 18%M	NR	24w	ADL CGI Drawing KOLT MMSE MNLT NART PADL Recognition memory for words and pictures Word Fluency	No
Rai 1990	IS	7	Placebo Acetyl-L-Carnitine	NINCDS	AD	Mild-Mod	36	20	79y (> 60y) 38%M	2 g/d (1 g bid)	24w	ADL CGA Computerized psychometric tests DCT Digit Span GDS GMS-A HMII NART NLT OLT P300 Reisberg GDS Word Fluency Test	No

EvTable1. Key characteristics: Carnitine (ALCAR) cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Sano 1992	PI IS	6	Placebo Acetyl-L-Carnitine	NINCDS	AD	Mild-Mod	30	27	Mean NR (60-80y) %M NR Community	12 w 2500 mg/d and 12 w 3000 mg/d	24w	Benton Visual Retention test Cancellations CGI Digit span mMMSE SIP SMQ BSRT Verbal Fluency Wechsler memory scale	MMSE
Spagnoli 1991	PI	7	Placebo Acetyl-L-Carnitine	DSM III	AD	Mild-Mod	130	108	75.2y (>40y) 29%M 60.8% Community 39.3% Institution	2 g/d	1y	Blessed Dementia Scale Blessed Information Memory Concentration test Block-tapping test Finger agnosia test Geometrical constructive apraxia test Ideomotor and buccofacial apraxia Prose memory test Raven's matrices SBI Supra-span verbal learning Token test Verbal judgement and mental calculation test Visual search on matrices of digits Word association test	No

EvTable1. Key characteristics: Carnitine (ALCAR) cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Thal 2000a	IF	6	Placebo Acetyl-L-Carnitine	NINCDS	AD	Probable	227	167	59y (46-65y) 53%M 90% White	1 g tid	1y	ADAS-Cog ADAS-Noncog ADL CDR CIBIC MMSE	No
Thal 1996a Auxiliary Brooks 1998	IF	6	Placebo Acetyl-L-Carnitine	NINCDS DSM-III-R	AD	Mild-Mod	431	355	72y (NR) 44%M 93.5% White	1 g tid	12m	ADAS –Noncog ADAS-Cog ADL CDR-S CGI-C CGI-S IADL MMSE	Age

EvTable2. Study results: Carnitine (ALCAR).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: 12w		Final: 24w	
Livingston, 1991	OC Analysis 1] Placebo 2] ALCAR (dose not specified) 3] Placebo Change from baseline 4] ALCAR change from baseline 5] Difference between placebo and ALCAR in change from baseline	ADL	1] 34.9 2] 33.0		1] 31.5 2] 33.6	3] < 0.01 4] NS 5] NS	1] 32.3 2] 34.5	3] < 0.05 4] NS 5] NS
		Clock Drawing	1] 5.66 2] 7.68		1] 5.63 2] 8.14	3] NS 4] NS 5] NS	1] 5.22 2] 8.74	3] NS 4] NS 5] NS
		Word fluency	1] 16.6 2] 18.2		1] 14.7 2] 15.7	3] <0.05 4] <0.05 5] NS	1] 15.3 2] 17.4	3] NS 4] NS 5] NS
		MMSE	1] 16.1 2] 15.8		1] 15.3 2] 16.0	3] NS 4] NS 5] NS	1] 15.1 2] 17.6	3] NS 4] NS 5] NS
		MNLT	1] 20.9 2] 18.4		1] 21.1 2] 21.6	3] NS 4] NS 5] NS	1] 21.9 2] 22.2	3] NS 4] NS 5] NS
		Object Learning	1] 12.3 2] 12.1		1] 11.0 2] 14.6	3] NS 4] NS 5] NS	1] 13.1 2] 14.1	3] NS 4] NS 5] NS
		RM – pictures	1] 14.5 2] 13.5		1] 14.6 2] 14.2	3] NS 4] NS 5] NS	1] 15.2 2] 15.1	3] NS 4] NS 5] NS
		RM – words	1] 15.2 2] 13.3		1] 14.2 2] 14.8	3] <0.05 4] NS 5] NS	1] 13.8 2] 15.5	3] <0.01 4] NS 5] < 0.01

EvTable3. Study results: Carnitine (ALCAR).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12w		Final: (specify) 24w	
Rai, 1990	OC Population 1] Placebo change from baseline 2] ALCAR 1g bid change from baseline 3] Placebo vs. ALCAR difference from baseline	GDS			1] 1.00 2] 0.25	1] NS 2] NS 3] NS	1] 2.00 2] 1.00	1] NS 2] NS 3] NS
		NLT			1] -3.38 2] 1.44	1] NS 2] NS 3] NS	1] -1.31 2] 0.57	1] NS 2] NS 3] NS
		Word Fluency Test			1] -1.00 2] -2.56	1] NS 2] NS 3] NS	1] -0.15 2] 0.57	1] NS 2] NS 3] NS
		ADL			1] 0.31 2] 0.22	1] NS 2] NS 3] NS	1] 0.15 2] 0	1] NS 2] NS 3] NS
		Digit Span			1] 0 2] 0.14	1] NS 2] NS 3] NS	1] -0.08 2] 0	1] NS 2] NS 3] NS
		Kendrick Battery Tests – Digit Copying Test			1] -4.25 2] 0.14	1] NS 2] NS 3] NS	1] -0.42 2] -1.00	1] NS 2] NS 3] NS
		Kendrick Battery Tests – Object Learning Test			1] 2.69 2] 1.00	1] NS 2] NS 3] NS	1] 1.77 2] -0.86	1] NS 2] NS 3] NS
		Clinical Global Improvement					1] 3.92 2] 3.6	3] NS
		Efficacy Index					1] 12.0 2] 10.29	3] NS

EvTable4. Study results: Carnitine (ALCAR).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24 w	
Sano 1992	OC Analysis 1] Placebo 2] ALCAR 2500 mg/d for 3m 3000mg/d for 6m 3] Difference between Placebo and ALCAR in change from baseline	SRT total recall	1] 21.4 (7.6) 2] 22.1 (7.3)				1] 16.0 (10.0) 2] 21.2 (8.8)	3] NS
		WMS	1] 2.9 (1.6) 2] 2.8 (1.8)				1] 2.2 (2.1) 2] 3.0 (1.6)	3] NS
		WMS Paired Associates	1] 6.3 (1.4) 2] 7.0 (2.0)				1] 6.4 (3.9) 2] 6.5 (1.9)	3] NS
		mMMSE	1] 35.3 (7.2) 2] 35.5 (5.4)				1] 32.4 (9.3) 2] 34.3 (6.3)	3] NS
		SIP	1] 27.3 (15.6) 2] 25.5 (12.5)				1] 24.17 (16.5) 2] 22.9 (12.5)	3] NS
		SMQ	1] 38.7 (8.5) 4] 38.5 (8.3)				1] 45.1 (10.8) 2] 45.0 (8.8)	3] NS
		Other Neuropsycholog ical Tests						3] NS

EvTable5. Study results: Carnitine (ALCAR).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: 12m	
Spagnoli, 1991	ITT Population 1] Placebo 2] ALCAR 250mg/bid 3] Difference between placebo and ALCAR 4] Difference between placebo and ALCAR in change from baseline	BDS	1] 9.5 (3.8) 2] 9.4 (4.3)	3] 0.84			1] 13.0 (4.6) 2] 11.0 (5.3)	3] 0.03 4] 0.01
		Blessed Information Memory	1] 17.1 (4.4) 2] 18.4 (4.8)	3] 0.10			1] 14.5 (6.7) 2] 16.8 (7.7)	3] 0.07 4] 0.33
		RPM	1] 5.6 (3.9) 2] 6.7 (4.8)	3] 0.32			1] 6.3 (4.7) 2] 3.8 (3.7)	3] 0.01 4] 0.03
		Supra-Span Verbal Learning	1] 2.1 (2.2) 2] 2.4 (2.7)	3] 0.76			1] 2.7 (3.4) 2] 1.5 (2.3)	3] 0.12 4] 0.24
		Block Tapping Task	1] 3.3 (4.0) 2] 3.7 (2.7)	3] 0.04			1] 2.8 (3.4) 2] 4.0 (3.8)	3] 0.03 4] 0.47
		Token Test	1] 21.9 (6.9) 2] 24.5 (7.1)	3] 0.04			1] 17.4 (9.9) 2] 21.8 (9.5)	3] 0.02 4] 0.41
		SBI*	1] 33.2 (8.5) 2] 29.7 (9.5)	3] 0.03			1] 41.7 (13.7) 2] 35.8 (13.6)	3] 0.02 4] 0.12

EvTable6. Study results: Carnitine (ALCAR).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline				FINAL 12m	
Thal, 2000a	MITT Analysis 1] Placebo 2] ALCAR 1g tid 3] Difference between ALCAR and placebo in change from baseline	<u>ADAS-cog</u> <u>CDR</u> ADAS-Non cog MMSE ADL	1] 22.9 (1.1) 2] 23.1 (1.2) 1] 5.1 (0.2) 2] 5.3 (0.3) 1] 3.2 (0.3) 2] 3.3 (0.3) 1] 20.6 (0.4) 2] 20.1 (0.5) 1] 7.1 (0.2) 2] 7.1 (0.2)				1] 30.4 (1.6) 2] 30.0 (1.7) 1] 6.8 (0.4) 2] 7.1 (0.4) 1] 5.3 (0.6) 2] 5.2 (0.5) 1] 17.3 (0.7) 2] 17.5 (0.8) 1] 8.3 (0.3) 2] 8.6 (0.4)	3] 0.58 3] 0.69 3] 0.89 3] 0.10 3] 0.43

→ Modified ITT sample

EvTable7. Study results: Carnitine (ALCAR).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12m	
Thal, 1996a	ITT Population	<u>ADAS-cog</u>	1] 26.0 (10.5) 2] 25.4 (9.8) 4] 25.8 (10.5) 5] 26.0 (10.5) 6] 27.8 (10.5) 7] 24.9 (9.5)				1] 33.0 (14.3) 2] 32.8 (14.8) 4] 35.1 (14.2) 5] 32.4 (14.4) 6] 34.7 (16.2) 7] 32.3 (14.5)	3] 0.434 8] 0.11 9] 0.085
Brooks, 1998	2] ALCAR 3g/d							
	3] ALCAR vs. Placebo							
	4] Placebo < 65 years	<u>CDR</u>	1] 6.6 (2.7) 2] 6.3 (2.4) 4] 6.2 (2.6) 5] 6.7 (2.7) 6] 6.1 (2.3) 7] 6.4 (2.4)				1] 8.8 (3.7) 2] 8.7 (3.8) 4] 9.5 (4.3) 5] 8.0 (3.7) 6] 8.5 (3.4) 7] 8.9 (3.8)	3] 0.562 8] 0.056 9] 0.047
	5] Placebo < 65 years							
	6] ALCAR < 65 years	MMSE	1] 9.6 (3.9) 2] 19.8 (3.9)				1] 15.8 (6.2) 2] 16.5 (6.4)	3] 0.818
	7] ALCAR > 65 years	ADAS-non cog	1] 4.7 (4.2) 2] 4.4 (3.8)				1] 7.6 (6.3) 2] 7.0 (6.3)	3] 0.466
	8] Difference between Placebo and ALCAR change from baseline for subgroup <65yrs	ADL	1] 7.6 (2.1) 2] 7.4 (2.1)				1] 10.1 (4.1) 2] 9.9 (4.2)	3] 0.733
		IADL	1] 16.8 (6.0) 2] 16.8 (5.7)				1] 20.3 (6.6) 2] 20.8 (6.0)	3] 0.62
		CGI-S	1] 3.6 (0.7) 2] 3.5 (0.6)				1] 4.1 (0.9) 2] 3.9 (0.9)	3] 0.172
	9] Difference between Placebo and ALCAR change from baseline for subgroup >65yrs.	CGI-C					1] 4.8 (0.9) 2] 4.9 (0.9)	3] 0.358

EvTable8. Adverse Events: Carnitine (ALCAR).

Adverse events (AE) identified in included studies	Livingston, 1991	Rai, 1989	Sano, 1992	Spagnoli, 1991	Thal, 1996a	Thal, 2000a
Withdrawn (%) due to AE	T: 0 C: 0	T: 44 C: 22	T: 0 C: 0	T: 0 C: 0	T: 3 C: 1	T: 1 C: 3
AE Checklist (Max 5)	3	3	3	3	1	1
None Reported						
Balance						
Accidental Injury						
Dizziness		X				
Falls						
Behavioral		X				
Agitation		X		NS		
Cardiovascular						
Arrhythmia						
Hypotension						
Hypertension						
Extrapyramidal						NS
Tremor						
Gastrointestinal	X					NS
Abdominal pain			X			
Constipation						
Diarrhea	X					
Dyspepsia						
Nausea, vomiting	X	X	X			
Metabolic/nutritional					NS	
Eating disorder						
Weight Change						
Neurological						
Asthenia						
Psychiatric		X				
Anxiety						
Confusion, delirium		X				
Depression		X				
Respiratory						NS
Cough, cold, infection						
Rhinitis						
Other					S	
Aberrant hematology						
Fatigue, weakness						
Fever, flu, pneumonia						
Headache						
Hepatic abnormality						
Muscle/joint disorder						
Pain						
Rash, skin disorder	X				NS	
Sleep disorder						
Urinary disorder						NS

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable9. Key characteristics: Donepezil (DPZ).

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Burns 1999	IF	6	Placebo Donepezil	DSM-III-R NINCDS	PDD	Mild-Mod	818	631	72y (50-93y) 42%M 99% white	5 mg/d for 7 d, then 10 mg/d	24w + 6w placebo washout period	ADAS-Cog CDR-SB CIBIC+ IDDD QOL	No
Feldman 2001 Auxiliary: Gauthier 2002	IF	8	Placebo Donepezil	NINCDS	AD	Moderate- Severe	290	247	73.7y (51-92y) 39%M Community	10 mg/d	24w	CAUST CIBIC CIBIC+ CSS DAD FRS IADL+ sMMSE NPI PSMS+ SF 36 SIB	MMSE Psychoactive drug use
Mohs 2001	IF	5	Placebo Donepezil	NINCDS DSM IV	AD	Probable	431	111	75.4y (50-93y) 37%M 92.15% white 2.75% black 5.1% other	10 mg/d	54w	ADL ADFACS CDR IADL IDDD MMSE	No

EvTable9. Key characteristics: Donepezil (DPZ) cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Prasher 2002	IF	6	Placebo Donepezil (DPZ)	ICD-10	AD	Mild–Mod	30	27	54y (43-66y) 50%M Community (99%) Institution (1%)	10 mg/d	24w	ABS DMR NPI SIB	Down Syndrome only
Pratt 2002	IF	5	Placebo Donepezil	NINCDS AIREN	VaD	Possible or Probable	893	707	74.0 (0.3)y range 41-95 Community	5 mg/d for 4w then either 5 or 10 mg/d for 20w	24w	ADAS-Cog CIBIC+ MMSE	No
Rogers 1996 Auxiliary: Rogers 2000 Neumann 1999 Rogers 1998	IF	6	Placebo Donepezil	DSM-III-R NINCDS	AD	Mild-Modly Sev	161	141	71.8y (54-85y) 40%M 99% white	5 mg/d	12w + 2w placebo washout period	ADAS-Cog ADL CDR-SB CGIC MMSE QoL	No
Rogers 1998a Auxiliary: Doody 2001 Steele 1999	IF	6	Placebo Donepezil	NINCDS DSM-III-R	AD	Mild-Modly severe	468	412	73.7y (50-94y) 36%M 96% white	5 mg/d for 7 d then 10 mg/d	15w	ADAS-Cog CIBIC+ MMSE CDR-SB QoL	No

EvTable9. Key characteristics: Donepezil (DPZ) cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Rogers 1998b Auxiliary: Doody 2001 Sparano 1998	IF	6	Placebo Donepezil	NINCDS DSM-III-R	AD	Mild-Mod	473	367	73.6 (51-94y) 38%M 95% white	10 mg/d	24w	ADAS-Cog CDR-SB CIBIC+ MMSE QoL	No
Tariot 2001a	IF	8	Placebo Donepezil	NINCDS	AD with CVD	Moderate- Severe	208	162	85.7y (65-100y) 18%M Institution (100%)	5 mg bid	24w	CDR-SB MMSE NPI-NH PSMS	MMSE Age

EvTable9. Key characteristics: Donepezil (DPZ) cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Thomas 2001	NR	7	Donepezil Vitamin E Rivastigmine (open label)	NINCDS	AD	Mild-Mod	60	54	66.5y (58-73y) 46%M	DPZ: 5 mg/d (month 1) 10 mg/d (until end) Vit E: 2000 IU (fixed) Rivastigmine: 1.5 mg/d (month 1) 3 mg/d (month 2) 6 mg/d (month 3) 9 mg/d (month 4) 12 mg/d (until end)	6m	ADAS-cog CT/MRI ERP scalp topography GBS GDS MMSE NPI WAIS	No
Winblad 2001b	IF	7	Placebo Donepezil	DSM IV NINCDS	AD	Mild-Mod	286	192	72.5y (50-87y) 36%M 100% white	10 mg/d	1y	ADL-PDS GBS GDS MMSE NPI PDS	APOE Genotype Gender

EvTable10. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12 w		Final: (specify) 24 w	
Burns 1999	ITT analysis	ADAS-cog	1] 69.84(1.68)* 2] 67.78 (1.61) 3] 69.85 (1.71)		4] 0.45	7] <0.0001	4] 1.5	7] 0.0315
	1] Placebo				5] -1.5	8] <0.0001	5] 0.5	8] <0.0001
	2] DPZ 5 mg/d				6] -1.8		6] -1.4	
	3] DPZ 10 mg/d	4] 4.25			7] 0.0545	4] 4.45	7] 0.0326	
	CIBIC+	5] 4.05			8] 0.0001	5] 4.25	8] 0.0009	
		6] 3.9				6] 4.1		
		4] Placebo change from baseline			4] 0.15	7] 0.0021	4] 0.375	7] 0.0387
	5] DPZ 5 change from baseline	5] -0.15			8] 0.0014	5] 0.075	9] <0.05	
	CDR-SB	6] -0.18				6] -0.13	8] 0.0020	
		6] DPZ 10 change from baseline			4] 69.5	8] 0.0085	4] 71.0	8] 0.0163
IDDD		5] 69.0		5] 70.8				
	6] 68.0		6] 69.0					
	7] DPZ 5 vs. placebo				4] NS			
8] DPZ 10 vs. placebo	QoL				5] NS			
9] DPZ both doses vs. placebo					6] NS			

*SEM

EvTable11. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12 w		Final: (specify) 24w LOCF	
Feldman 2001	ITT Analysis	<u>CIBIC+</u>			1] 4.1 2] 3.6 5] -0.11 6] -0.48	3] <0.0001 7] 0.0002	1] 4.65 2] 4.05 5] 0.24 6] -0.22	3] 0.0004 7] 0.0044
Gauthier 2002	1] Placebo mean change from baseline							
	2] DPZ 10 mg/d mean change from baseline	<u>CIBIC+</u> <u>% improved</u>					1] 42% 2] 63%	3] <0.0001
	3] Mean treatment difference DPZ vs. Placebo	DAD			1] -305 2] 1.25 5] -4 6] 2	3] 0.0037 7] 0.0037	1] -8.98 2] -0.74 5] -9 6] 0.1	3] <0.0001 7] <0.0001
	4] Mean treatment difference LOCF population	sMMSE			1] 0.2 2] 1.75 5] 0.0 6] 2.0	3] 0.0004 7] 0.0004	1] -0.5 2] 1.25 5] -0.5 6] 1.5	3] 0.0019 7] 0.0009
	5] Placebo change from baseline subgroup with MMSE of 10-17	SIB			1] -0.25 2] 4.75 5] -1.0 6] 3.5	3] <0.0001 7] 0.0004	1] -4.0 2] 2.0 5] -3.0 6] 2.5	3] <0.0001 7] 0.0012
	6] DPZ 10 mg d change from baseline subgroup with MMSE of 10-17	IADL + PSMS +					3] 6.83 3] 1.32	3] 0.0015 3] 0.0015
	7] Difference between Placebo and DPZ change from baseline subgroup with MMSE of 10-17	NPI FRS			5] -0.5 6] -3.8		1] -1.0 2] 4.6 5] 1.0 6] 5.0 1] -1.66 2] -0.38	7] 0.021 3] 0.0002 7] 0.0022

EvTable12. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 24 w		Final: (specify) 54 w	
Mohs 2001	ITT Endpoint Analysis	ADFACS					3] 4.0 4] 2.5	5] <0.001
	1] Placebo	ADFACS						5] 0.001
	2] DPZ 10 mg d	ADL – Instrumental						5] 0.007
	3] Placebo mean change from baseline	ADFACS ADL-basic						5] <0.001
	4] DPZ mean 10mg/d change from baseline	MMSE	1] 17.1 (0.2)* 2] 17.1 (0.2)*		3] 0.4 4] 1.8	5] <0.01	3] –0.7 4] 0.6	5] <0.001
	5] Mean change from baseline DPZ vs. Placebo	Time to functional decline (days)					3] 208 CI (165 to 252) 4] 356 CI (>280)	5] 0.0051

*SEM

EvTable 13. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24 w	
Prasher 2002	OC Analysis	<u>DMR</u>	1] 58.2 (16.9) 2] 54.3 (16.1)				1] 64.4 (14.2) 2] 55.1 (17.9)	5] 0.22
	1] Placebo							3] 0.002
	2] DPZ 10 mg d	SIB	1] 27.2 (13.6) 2] 36.8 (21.9)				1] 11.2 (8.7) 2] 31.6 (28.2)	4] 0.002 5] 0.06
	5] Change from baseline placebo vs. DPZ	NPI	1] 8.0 (7.6) 2] 7.9 (5.8)				1] 3.6 (5.0) 2] 5.7 (7.6)	3] 0.03
	3] Placebo change from baseline	ABS	1] 93.0 (19.2) 2] 121.4 (36.9)				1] 84.5 (22.4) 2] 120.5 (44.1)	3] 0.51
	4] DPZ change from baseline							
	5] Change from baseline placebo vs. DPZ							

EvTable14. Study results: Donepezil.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24 w	
Pratt 2002	ITT Analysis							
	1] Placebo change from baseline	<u>ADAS-cog</u>					1] 0.1 2] -1.1 3] -2.2	4] <0.001 5] <0.001
	2] Donepezil 5mg/d change from baseline	<u>CIBIC+</u>					1] 32% 2] 46% 3] 36%	4] 0.0006 5] 0.2096
	3] Donepezil 10mg/d change from baseline	MMSE					1] 0.5 2] 1.5 3] 1.6	4] <0.001 5] <0.001
	4] Donepezil 5mg/d vs Placebo							
	5] Donepezil 10mg/d vs Placebo							

EvTable15. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6w		Final: (specify) 12 w	
Rogers 1996	ITT Endpoint analysis	ADAS-Cog			1] 0.5 2] -2.0 3] -3.2 4] -3.8		1] 0.7 2] -0.9 3] -1.4 4] -2.5	5] 0.0359 6] 0.105 7] 0.036 8] 0.002
Rogers 2000	1] Placebo mean change from baseline	CGIC % success					1] 80% 2] 82% 3] 83% 4] 90%	8] 0.039
Newman 1999	2] DPZ 1mg/d mean change from baseline							
Rogers 1998	3] DPZ 3mg/d mean change from baseline	ADL					1] 1.5 2] 4.0 3] 0.6 4] -3.1	5] 0.0684
	4] DPZ 5 mg/d mean change from baseline	MMSE			1] 0.8 2] 1.15 3] 1.25 4] 1.85		4] 1.2 2] 0.6 3] 0.9 4] 2.0	5] 0.0275
	5] Dose response analysis							
	6] DPZ 1 mg/d difference from placebo	QoL-P					1] -1.3 2] 0.7 3] 2.6 4] 8.8	5] 0.0369
	7] DPZ 3 mg/d difference from placebo	QoL-C					1] 3.7 2] -5.3 3] 0.0 4] 0.3	5] 0.8860
	8] DPZ 5mg/d difference from placebo	CDR-SB			1] 0.10 2] -0.050 3] 0.0 4] -0.04		1] 0.10 2] 0.18 3] 0.23 4] -0.11	5] 0.3375

EvTable16. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6 w		Final: (specify) 12 w	
Rogers 1998a	ITT Analysis Endpoint	ADAS-cog			1] -0.02 2] -1.5 3] -2.9	4] 0 .011 5] <0.001	1] 0.4 (0.43)* 2] -2.1 (0.43)* 3] -2.7 (0.43)*	4] <0.001 5] <0.001
Doody 2001	1] Placebo change from baseline	CIBIC +			1] 3.99 2] 3.85 3] 3.93		1] 4.2 (0.07)* 2] 3.9 (0.08)* 3] 3.8 (0.08)*	4] 0.003 5] 0.008
Steele 1999	2] DPZ 5mg/d change from baseline	MMSE			1] 0.65 2] 0.95 3] 1.4	5] 0.03	1] 0.04 (0.25)* 2] 1.0 (0.25)* 3] 1.3 (0.24)*	4] <0.004 5] <0.001
	3] DPZ 10mg/d change from baseline	CDR-SB			1] -0.15 2] 0.0 3] -0.25	5] 0.008	1] -0.14 (0.11)* 2] -0.10 (0.11)* 3] -0.31 (0.11)*	4] 0.32
	4] DPZ 5mg/d vs placebo	QoL					1] 4.0 (2.7)* 2] 5.7 (2.7)* 3] -4.3 (2.7)*	4] 0.65 5] 0.02
	5] DPZ 10mg/d vs placebo							

*SEM

EvTable17. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12 w		Final: (specify) 24 w	
Rogers 1998b	ITT Population	<u>ADAS-cog</u>			4] 1.25 5] -1.25 6] 2.0	7] 0.0007 8] <0.0001	4] 1.82 (0.49)* 5] -0.67 (0.51)* 6] -1.06 (0.51)*	7] <0.0001 8] <0.0001
Doody 2001	1] Placebo	<u>CIBIC +</u>						
	2] DPZ 5 mg/d				4] 4.2 5] 3.95 6] 3.9	7] 0.0157 8] 0.009	4] 4.51 (0.08)* 5] 4.15 (0.09)* 6] 4.07 (0.07)*	7] <0.0047 8] <0.0001
Sparano 1998	3] DPZ 10 mg/d	MMSE						
	4] Placebo change from baseline				4] -0.5 5] 0.75 6] 1.0	7] 0.0002 8] <0.0001	4] -0.97 (0.28)* 5] 0.24 (0.29)* 6] 0.39 (0.29)*	7] 0.0007 8] 0.0002
	5] DPZSP5 change from baseline	CDR-SB			4] 0.1 5] -0.2 6] -0.025		4] 0.58 (0.14)* 5] -0.01 (0.14)* 6] -0.02 (0.14)*	7] 0.0008 8] 0.0007
	6] DPZ10 change from baseline	QoL						
	7] DPZ5 vs placebo							7] NS 8] NS
	8] DPZ10 vs. Placebo							

*SEM

EvTable18. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12 w		Final: (specify) 24 w	
Tariot 2001a	ITT Analysis							
	1] Placebo	<u>NPI-NH</u>	1] 20.5 (14.7) 2] 21.0 (14.5)				3] -4.9 (1.9)* 4] -2.3 (1.9)*	7] NS
	2] DPZ 10 mg /d	MMSE	1] 14.4 (5.8) 2] 14.4 (5.4)		3] -0.5 4] 0.35 5] -0.95 6] 0.6	6] <0.05 7] NS	3] -0.75 4] -0.1 5] -1.0 6] 0.0	7] NS 8] <0.05
	3] Placebo change from baseline all patients	CDR-SB						
	4] DPZ change from baseline all patients		1] 10.8 (3.7) 2] 11.2 (4.0)		3] 0.2 4] -0.15 5] 0.8 6] -0.3	4] 0.09 8] <0.05	3] 0.7 4] -0.1 5] 0.8 6] -0.2	4] <0.05 6] <0.05 7] <0.05 8] <0.05
	5] Placebo change from baseline for subgroup with baseline MMSE scores of 10-26	PSMS	1] 14.7 (5.0) 2] 15.1 (4.9)			7] 0.09	3] -1.0 4] -1.0	7] 0.31
	6] DPZ change from baseline for subgroup with baseline MMSE scores of 10-26							
	7] mean change DPZ over Placebo							
	8] subjects >85 DPZ over Placebo							

*SEM

EvTable19. Study results: Donepezil (DPZ), Vitamin E.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3 m		Final: (specify) 6 m	
Thomas 2001	OC Analysis							
	1] DPZ 10 mg d	WAIS	1] 72 (2.0)* 2] 72 (2.0)*		1] 74 (2.0)* 2] 72 (2.0)*		1] 75 (2.0)* 2] 71 (2.1)*	3] 0.15 4] 0.43
	2] Vitamin E 2,000 IU							
	3] change from baseline with DPZ	MMSE	1] 16 (0.5)* 2] 16 (0.5)*		1] 16 (0.6)* 2] 15 (0.5)*	5] <0.001 favors DZP	1] 16 (0.5)* 2] 15 (0.6)*	3] 0.06 4] 0.07 5] <0.001 favors DPZ
	4] change from baseline with Vitamin E	ADAS-cog	1] 33.34 (2.7)* 2] 33.45 (2.6)*		1] 31.55 (2.7)* 2] 36.09 (2.8)*		1] 31.84 (2.7)* 2] 39.07 (2.7)*	3] <0.001 4] <0.01
	5] DPZ vs Vitamin E change from baseline	NPI	1] 21.9 (0.5)* 2] 21.9 (0.5)*				1] 16.8 (0.2)* 2] 22.8 (1.2)*	

*SEM

EvTable20. Study results: Donepezil (DPZ).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 24 w		Final: (specify) 52 w	
Winblad 2001b	ITT Endpoint Analysis	<u>GBS total score</u>					3] 12.0 4] 9.0	5] 0.054 6] 0.532 7] 0.258
	1] Placebo							
	2] DPZ 10 mg/d	GBS						5] 0.012
	3] Placebo change from baseline	MMSE					3] -2.2 4] -0.4	5] <0.001 6] 0.712 7] 0.743
	4] DPZ change from baseline	ADL-PDS					3] -15 4] -11	5] <0.05
	5] DPZ vs. Placebo	GDS						5] 0.047
	6] Analysis for APOE Genotype Interaction	NPI				5] NS		5] NS
	7] Aanalysis for Gender Interaction							

EvTable21. Adverse Events: Donepezil (DPZ).

Adverse events (AE) identified in included studies	Burns, 1999	Feldman, 2001	Mohs, 2001	Prasher, 2002	Pratt, 2002	Rogers, 1996	Rogers, 1998a	Rogers, 1998b	Tariot, 2001a	Thomas, 2001	Winblad, 2001b
Withdrawn (%) due to AE	T:14 ⁺ C:10	T: 8 C: 6	T: 11 C: 7	T: 7 C: 0	T:15 ⁺ C: 9	T: 8 C: 5	T: 7 ⁺ C: 1	T:11 ⁺ C: 7	T: 18 C: 11	T: 0 C: 0	T: 7 C: 6
AE Checklist (Max 5)	3	2	3	4	1	2	3	3	2	3	4
None Reported										X	
Balance											S
Accidental Injury		X	NS			X	NS*		X		NS
Dizziness	NS*	X		NS		X	NS*	NS*	X		NS
Falls											
Behavioral		X									NS
Agitation			NS	NS		X	NS*		X		
Cardiovascular									X		
Arrhythmia							X				
Hypotension											
Hypertension											
Extrapyramidal											
Tremor									X		
Gastrointestinal						X	X				
Abdominal pain		X		NS			X		X		NS
Constipation						X					NS
Diarrhea	S*	X	S	NS		X	S*	S*	X		NS
Dyspepsia			S								
Nausea, vomiting	S*	X	S	NS		X	S*	S*	X		NS
Metabolic/nutritional									X		
Eating disorder	NS*		S	NS			X	NS*	X		
Weight Change		X	NS				X		X		
Neurological				NS							
Asthenia		X	NS						X		S
Psychiatric				X							
Anxiety											NS
Confusion, delirium	NS*	X							X		NS
Depression		X									NS
Respiratory		X				X	X				
Cough, cold, infection						X	NS*		X		
Rhinitis			NS				X	NS*	X		
Other						X	X		X		
Aberrant hematology								NS*	X		
Fatigue, weakness				S			NS*	S*			
Fever, flu, pneumonia									X		X
Headache		X	S			X	NS*		X		NS
Hepatic abnormality	NS*					NS	NS				
Muscle/joint disorder		X		NS			NS*	S*	X		
Pain		X				X	NS*		X		
Rash, skin disorder			NS						X		
Sleep disorder	NS*		NS	NS			S*				NS
Urinary disorder		X	S			X	S		X		NS

NR = Withdrawals due to AE Not Reported; + = Dose effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable22. Key characteristics: Galantamine.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Erkinjuntti 2002	PI	8	Placebo Galantamine	NINDS NINCDS	VaD AD	Probable	592	457	75.1y (40-90y) 53%M 99.9% white	4 mg/d on w 1 8 mg/d on w 2 12 mg/d on w 3 16 mg/d on w 4 20 mg/d on w 5 24 mg/d from w 6 to the end	6m	ADAS-Cog CIBIC+ DAD NPI	MID vs AD+vas cular
Raskind 2000	IF	8	Placebo Galantamine	NINCDS	AD	Mild-Mod	636	438	75.4y (NR) 38%M 91.5% white	Loading: 8 mg/d on w 1 16 mg/d on w 2 24 mg/d on w 3 Then one group stayed at 24 mg/d the other group 32 mg/d	6m	ADAS-Cog CIBIC+ DAD	APOE Genotyp e
Rockwood 2001	IF	7	Placebo Galantamine	NINCDS	AD	Probable Mild-Mod	386	288	74.9y (NR) 44%M	8 mg/d on w 1 16 mg/d in w 2 24 mg/d on w 3 32 mg/d on w 4 At the end of w 4 dose could be reduced to 24 mg/d	3m	ADAS-Cog CIBIC+ DAD NPI PSQI	No
Tariot 2000	NR	8	Placebo Galantamine	NINCDS	AD	Probable Mild-Mof	978	679	76.8y (NR) 36%M 93% white	Loading: 8 mg/d 16 mg/d Then 24 mg/d	5m	ADAS-Cog ADCS CIBIC+ IADL NPI	No

EvTable22. Key characteristics: Galantamine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Wilcock 2000 Auxiliary: Wilcock 2001	NI	8	Placebo Galantamine	NINCDS	AD	Mild-Mod	653	525	72.2y (NR) 37%M	Loading: 8 mg/d on w 1 16 mg/d on w 2 24 mg/d on w 3 one group stayed at 24 mg/d the other group 32 mg/d	6m	ADAS-Cog ADAS CIBIC+ DAD	MMSE APOE Genotyp e
Wilkinson 2001	IF	7	Placebo Galantamine	NINCDS DSM-III-R	AD	Mild-Mod	285	206	73.8y (>45y) 42%M 100% Community	Start at 4 mg/d and progressively increased every 2-3 d to reach the target doses 18mg/d 24 mg/d 36 mg/d	12w	ADAS-Cog CGIC PDS	No

EvTable23. Study results: Galantamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Erkinjuntti 2002	OC Analysis							
	1] Placebo from baseline	ADAS-cog/11 <u>change from baseline</u>					1] 1.0 (0.5)* 2] -1.7 (0.4)* 4] 1.8 (0.6)* 5] -1.0 (0.46)* 7] -0.4 (0.78)* 8] -2.4 (0.59)*	1] 0.045 2] <0.0001 3] <0.0001 6] 0.0005 9] 0.06
	2] Galantamine 24 mg/d from baseline							
	3] Placebo vs Galantamine							
	4] Placebo from baseline subgroup AD+vascular	CIBIC-plus % <u>improved or stable</u>					1] 59% 2] 74% 4] 54% 5] 75%	3] 0.0011 6] 0.001
	5] Galantamine from baseline subgroup AD+vascular	CIBIC-plus % improved					4] 19% 5] 32% 7] 23% 8] 31%	6] 0.019 9] 0.238
	6] Placebo vs Galantamine subgroup AD+vascular							
	7] Placebo from baseline subgroup VAD	ADAS-cog/13 Change from baseline					1] 0.9 (0.6)* 2] -2.4 (0.4)*	3] <0.0001
	8] Galantamine from baseline subgroup VAD	DAD					1] -4.4 (1.3)* 2] 0.2 0.9)*	3] 0.0017
	9] Placebo vs Galantamine subgroup VAD	NPI					1] 1.0 (0.9)* 2] -1.2 0.6)*	3] 0.0164

*SEM

EvTable24. Study results: Galantamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Raskind 2000	ITT Analysis							
	1] Placebo change from baseline	<u>ADAS-cog/11</u>					1] 2.0 (0.45)* 2] 1.9 (0.36)* 3] -1.4 (0.44)*	4] <0.001 5] <0.001
	2] Galantamine 24 mg/d change from baseline	<u>CIBIC-plus</u> <u>% improved</u>					1] 13.8% 2] 19.9% 3] 15.8%	4] <0.05 5] <0.05
	3] Galantamine 32 mg/d change from baseline	DAD (total)						4] NS 5] NS
	4] Placebo vs. Galantamine 24mg/d change from baseline							
	5] Placebo vs. Galantamine 32 mg/d change from baseline							

*SEM

EvTable25. Study results: Galantamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Rockwood 2001	ITT Analysis 1] Placebo change from baseline 2] Galantamine 24-32mg/d change from baseline 3] Difference between placebo and galantamine change from baseline	<u>ADAS-Cog-11</u>					1] 0.6 (0.45)* 2] -1.1 (0.33)*	3] <0.01
		<u>ADAS-Cog-13</u>					1] +0.7 (0.51)* 2] -1.2 (0.38)*	3] <0.01
		<u>CIBIC-+ % improved or stable</u>					1] 18.7% 2] 22.1%	3] <0.01
		NPI					1] 0.5 (0.65)* 2] -0.3 (0.7)*	1] NS 2] NS
		DAD					1] -5.2 (1.18)* 1] -0.4 (0.76)	3] <0.001

*SEM

EvTable26. Study results: Galantamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 5m	
Tariot 2000	ITT Analysis							
	1] Placebo change from baseline	ADAS-Cog					1] 1.7 (0.39)*	5] NS
	2] Galantamine 8mg/d change from baseline						2] +0.4 (0.52)*	6] <0.001
	3] Galantamine 16 mg/d change from baseline	CIBIC+ % improved					3] -1.4 (0.35)*	7] <0.001
	4] Galantamine 24mg/d change from baseline						4] -1.4 (0.39)*	8] <0.05
								9] <0.01
	5] Placebo vs. galantamine 8 mg/d in change from baseline	ADCS/ADL					1] 49%	6] <0.001
	6] Placebo vs. galantamine 16 mg/d in change from baseline						2] 53%	7] <0.001
	7] Placebo vs. galantamine 24 mg/d in change from baseline						3] 66%	8] <0.05
	8] Galantamine 8mg/d vs. 16mg/d in change from baseline						4] 64%	9] <0.05
	9] Galantamine 8mg/d vs. 24mg/d in change from baseline						1] -3.8 (0.6)*	6] <0.001
							2] -3.2 (0.8)*	7] <0.01
							3] -0.7 (0.5)*	8] <0.01
							4] -1.5 (0.6)*	
							1] 2.0 (0.7)*	6] <0.05
							2] 2.3 (1.0)*	7] <0.05
							3] -0.1 (0.7)*	
							4] 0.0 (0.8)*	

*SEM

EvTable27. Study results: Galantamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Wilcock 2000	ITT Analysis	<u>ADAS-cog</u>					1] 2.4 (0.41)* 2] -0.5 (0.38)* 3] -0.8 (0.43)*	1] <0.001 2] <0.001 3] <0.001
Wilcock 2001	1] Placebo change from baseline							4] <0.001 5] <0.001
	2] Galantamine 24mg/d change from baseline	<u>DAD</u>					1] -6.0 (1.08)* 2] -3.2 (1.02)* 3] -2.5 (1.07)*	4] 0.1 5] <0.05
	3] Galantamine 32 mg/d change from baseline	<u>CIBIC+</u>						
	4] Difference between placebo and Galantamine 24mg/d in change from baseline						1] 16.5% 2] 17% 3] 25%	4] <0.05 5] <0.001
	5] Difference between placebo and Galantamine 32 mg/d in change from baseline							

*SEM

EvTable28. Study results: Galantamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Wilkinson 2001	ITT Analysis	ADAS-Cog					1] 1.6 (0.7)* 2] -01 (0.7)* 3] -1.4 (0.9)* 4] -0.7 (0.7)*	5] NS 6] <0.01 7] 0.08
	1] Placebo change from baseline							
	2] Galantamine 18 mg/d Change from baseline	CGIC % improved					1] 31.3% 2] 36.7% 3] 28.3% 4] 31.9%	5] NS 6] NS 7] NS
	3] Galantamine 24 mg/d Change from baseline							
	4] Galantamine 36 mg/d Change from baseline	PDS-1 % improved					1] 9.2 % 2] 12.5 % 3] 14.3% 4] 7.4%	5] NS 6] NS 7] NS
	5] Placebo vs. Galantamine 18mg/d in change from baseline							
	6] Placebo vs. galantamine 24mg/d in change from baseline							
	7] Placebo vs. galantamine 36mg/d in change from baseline							

*SEM

EvTable29. Adverse Events: Galantamine.

Adverse events (AE) identified in included studies	Erkinjuntti, 2002	Raskind, 2000	Rockwood, 2001	Tariot, 2000	Wilcock, 2000	Wilkinson, 2001
Withdrawn (%) due to AE	T: 20 C: 8	T: 27 C: 8	T: 26 C: 4	T: 8 C: 7	T: 18 C: 9	T: 27 ⁺ C: 9
AE Checklist (Max 5)	3	3	4	3	3	3
None Reported						
Balance						
Accidental Injury	X					
Dizziness		X	X		X	X
Falls						
Behavioral						
Agitation			X	X		
Cardiovascular						
Arrhythmia						
Hypotension						
Hypertension						
Extrapyramidal						
Tremor		X				
Gastrointestinal	X			X		
Abdominal pain		X	X			
Constipation						
Diarrhea		X		X	X	X
Dyspepsia						
Nausea, vomiting	X	X	X	X	X	X
Metabolic/nutritional						
Eating disorder		X	X	X	X	X
Weight Change		S		X	S	
Neurological	X					
Asthenia						
Psychiatric						
Anxiety						
Confusion, delirium						
Depression						
Respiratory						
Cough, cold, infection						
Rhinitis						
Other						
Aberrant hematology		NS	NS	NS	NS	NS
Fatigue, weakness						
Fever, flu, pneumonia						
Headache					X	X
Hepatic abnormality						
Muscle/joint disorder				X		
Pain						
Rash, skin disorder						
Sleep disorder			X			
Urinary disorder						

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable30. Key characteristics. Metrifonate.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Becker 1996	NI	6	Placebo Metrifonate	NINCDS	AD	Probable	53	51	71.4y (57-85y) 40%M	Loading: 5.0 mg/kg for 2 w, 4.9 mg/kg for 1 w then 2.1 mg/kg weekly	3m	ADAS-Cog ADAS-Noncog ADAS-T ADLC Laboratory tests EEG GIS MMSE	No
Becker 1998	NI	6	Placebo Metrifonate	NINCDS	AD	Probable	47	46	73.0y (<90y) 51%M Community	Loading 2 mg/kg for 5 d, 0.95 mg/kg on d 6 then 2.9 mg/kg weekly	6m	ADAS-Cog ADAS-Noncog ADLC Laboratory tests GIS MMSE	No

EvTable30. Key characteristics: Metrifonate cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Cummings 1997	NR	5	Placebo Metrifonate	NINCDS	AD	Probable Mild-Mod	480	453	NR	Loading: Low dose group: 0.5 mg/kg Medium-dose group: 0.9 mg/kg High-dose group: 2.0 mg/kg Maintenance dose respectively: 0.2 mg/kg 0.3 mg/kg 0.65 mg/kg/d	12w	ADAS-Cog CIBIC+ MMSE	No
Cummings 1998 AUXILIARY Cummings 1998b	PI	8	Placebo Metrifonate	NINCDS	AD	Probable	480	443	73.5y (NR) 41%M	Loading for 2 w: low dose group 0.5 mg/kg ; mid-dose group: 0.9 mg/kg; high-dose group 2.0 mg/kg then respectively 0.2 mg/kg 0.3 mg/kg 0.65 mg/kg	12w	ADAS-Cog CIBIC+ CIBIS+ GERRI IADL MMSE PSMS	No

EvTable30. Key characteristics: Metrifonate cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Dubois 1999 AUXILIARY McKeith 1998	IF	6	Placebo Metrifonate	DSM IV	AD	Mild-Mod	605	516	72.1y (NR) 36%M	1 loading dose of 80 mg or 120 mg and then: 0.65 mg/Kg/d or 1.0 mg/Kg/d	26w	AChE ADAS-Cog ADAS-Noncog CIBIC+ CIBIS+ DAD ECG GDS MMSE NPI Laboratory tests	No
Jann 1999	PI	6	Placebo Metrifonate	NINCDS	AD	Mild-Mod	395	393	75.0y (45-90y) 42%M 91% White 3.8% Black 2.6% Hispanic 1.8% Asian	Loading dose group: 100 or 150 mg for 2 w Non loading dose group: 50 mg/d	6w	ADAS-Cog ADAS-Noncog CIBIC+ CIBIS+ EEG MMSE	No

EvTable30. Key characteristics: Metrifonate cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	#Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Morris 1998	PI	7	Placebo Metrifonate	NINCDS	AD	Mild-Mod	408	334	73.6y (NR) 39%M 93% White 3.3% Black 0.4% Asian 2.6% Hispanic	2 w loading (2.0mg/kg/d) 24w 0.65 mg/kg/d	26w	ADAS-Cog ADAS-NONCOG CIBIC+ CIBIS+ GDS MMSE NPI	No
Pettigrew 1998	PI	6	Placebo Metrifonate	NINCDS	AD	Probable	27	27	72.0y (55-85y) 59%M 100% community	Loading for 6 d: Panel 1: 1.5 mg/kg Panel 2: 2.5 mg/kg Panel 3: 4 .0 mg/kg Panel 4: 4.0 then respectively: 0.25 mg/kg; 4.0 mg/kg; 0.65 mg/kg 1.0 mg/kg	21d	ADAS-Cog ADAS-Noncog Blessed – DRS ECG MMSE MRS	No

EvTable30. Key characteristics: Metrifonate cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Raskind 1999	IF	7	Placebo Metrifonate	NINCDS	AD	Probable Mild-Mod	264	219	74.6y (NR) 36% M Community 90% White 3.7% Black 5.1% Hispanic 0.3% American Indian	50 mg/d	6m + 6 w post- treatment follow- up period	ADAS-Cog ADAS-Noncog CIBIC+ CIBIS+ DAD GDS MMSE NPI	No

EvTable31. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3 m	
Becker 1996	OC Analysis 1] Placebo 2] Metrifonate 5.0 mg/kg 2 w 4.9 mg/kg 1 w 2.1 mg/kg 9w 3] Baseline vs. Placebo 4] Baseline vs. Metrifonate 5] Metrifonate vs. placebo	ADAS-Cog	1] 26.39(10.20) 2] 25.64(11.86)				1] 27.49(11.04) 2] 24.89(11.80)	3] <0.02 4] 0.15 5] <0.01
		MMSE	1] 19.30 (5.50) 2] 19.47 (5.35)				1] 18.35 (5.77) 2] 19.36 (6.01)	3] <0.03 4] 0.76 5] 0.14
		ADAS-Noncog	1] 5.19 (5.38) 2] 5.91 (4.93)				1] 5.88 (5.74) 2] 5.94 (5.11)	3] 0.09 4] 0.88 5] 0.11
		ADAS-Total	1] 31.58(13.29) 2] 31.56(14.56)				1] 33.37(14.51) 2] 30.83(14.44)	3] <0.02 4] 0.21 5] <0.01
		GIS	1] 3.63 (0.46) 2] 4.08 (0.52)				1] 4.11 (0.59) 2] 4.10 (0.71)	3] <0.01 4] 0.90 5] 0.1
		ADLC	1] -2.51(13.26) 2] -7.80(17.21)				1] -2.23 (9.31) 2] -3.82(17.00)	3] 0.93 4] 0.17 5] 0.80

EvTable32. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Becker 1998	OC Analysis							
	1] Placebo	ADAS-Cog	1] 21.75 (8.40) 2] 20.61 (9.33)				1] 23.42 (9.53) 2] 20.61 (9.32)	3] 0.01 4] 1.00 5] <0.03
	2] Metrifonate 2.9 mg/kg w	MMSE	1] 19.78 (4.92) 2] 20.62 (3.85)				1] 18.60 (5.40) 2] 20.38 (4.42)	3] <0.01 4] 0.44 5] 0.09
	3] Placebo vs. baseline							
	4] Metrifonate vs. baseline	ADAS- Noncog	1] 3.75 (4.53) 2] 2.72 (3.57)				1] 4.07 (4.92) 2] 2.80 (3.40)	3] 0.31 4] 0.80 5] 0.53
	5] Metrifonate vs. Placebo	GIS	1] 3.85 (0.77) 2] 3.78 (0.45)				1] 4.23 (0.81) 2] 4.34 (0.80)	3] <0.02 4] <0.00 5] 0.42
		ADLC	1] 3.88 (8.66) 2] 4.93 (6.66)				1] 9.56 (10.93) 2] 8.94 (8.67)	3] 0.01 4] 0.00 5] 0.58

EvTable33. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12 w	
Cummings 1997	OC Population							
	1] Placebo	ADAS-Cog					5] 1.45 6] 3.17	5] <0.05 6] <0.001
	2] Low Dose Metrifonate 0.5 mg/kg loading 0.2 mg/kg maintenance	CIBIC+					5] 0.33 6] 0.40	5] <0.05 6] <0.001
	3] Medium Dose Metrifonate 0.9 mg/kg loading 0.3 mg/kg maintenance							
	4] High dose Metrifonate 2.0 mg/kg loading 0.65 mg/kg maintenance							
	5] Placebo vs. medium dose Metrifonate							
	6] Placebo vs. high dose Metrifonate							

EvTable34. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Cummings 1998a	ITT Analysis	<u>CIBIC+</u>					2] 0.04 CI (-0.16-0.24)	2] 0.735
	1] Placebo						3] 0.29 CI (0.09-0.48)	3] 0.005
Cummings 1998b	2] Metrifonate 20 mg qid difference from Placebo						4] 0.35 CI (0.15-0.54)	4] 0.0007
	3] Metrifonate 25mg qid difference from Placebo	<u>ADAS-Cog</u>					2] 1.5 CI (0.18-2.83)	2] 0.02
							3] 1.30 CI (-0.02-2.62)	3] 0.053
	4] Metrifonate 60mg qid difference from Placebo						4] 2.94 CI (1.61-4.27)	4] 0.0001
		MMSE					2] 1.11 C1(0.39 - 1.84)	2] 0.0029
							3] 0.63 C1(-0.10-1.35)	3] 0.0905
							4] 1.37 C1(0.64 - 2.10)	4] 0.0003
		PSMS						2] NS
								3] NS
								4] NS
		CIBS+						2] NS
								3] NS
								4] NS
		GERRI						2] NS
								3] NS
								4] NS
		IADL						2] NS
								3] NS
								4] NS

EvTable35. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 26 w	
Dubois 1999	ITT Analysis (LOCF)	<u>ADAS-cog</u>					3] 1.3 4] 3.24	3] 0.032 4] 0.0001
McKeith 1998	1] Placebo	<u>CIBIC+</u>					3] 0.21 4] 0.35	3] 0.052 4] 0.0014
	2] Metrifonate 40 to 50 mg/d variable by weight	DAD total					3] 3.00 4] 5.45	3] 0.0522 4] 0.0005
	3] Metrifonate 60 to 80 mg/d variable by weight	MMSE					3] 0.40 4] 1.19	3] 0.26 4] 0.0009
	4] mean change from baseline Placebo vs. 40/50mg dose Metrifonate	CIBIS+					3] 0.20 4] 0.23	3] 0.0002 4] 0.0001
		ADAS- Noncog Total					3] 0.59 4] 1.37	3] 0.14 4] 0.0008
	5] mean change from baseline Placebo vs. 60/80 mg dose Metrifonate	NPI total					3] 0.83 4] 1.44	3] 0.48 4] 0.23
		GDS					3] 0.08 4] 0.21	3] 0.22 4] 0.0026
		IADL						3] <0.05 4] <0.05
		ADL						3] NS 4] NS

EvTable36. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4w		Final: (specify) 6w	
Jann 1999	ITT Analysis							
	1] Placebo	ADAS-Cog			4] 0.0 5] -0.85 6] -0.1	7] NS	4] 0.55 5] -1.0 6] 0.0	7] 0.01 8] NS
	2] Metrifonate loading dose 100/150 mg d 2w by weight plus 50 mg d 4 w	CIBIC+			4] 4.02 5] 4.25 6] 3.85	7] <0.05	4] 4.1 5] 3.9 6] 3.72	7] <0.05 8] <0.05
	3] Metrifonate 50 mg d 6w	ADAS- Noncog						7] NS 8] NS
	4] Placebo change from baseline	CIBIS+						7] NS 8] NS
	5] Metifonate loading dose change from baseline	MMSE						7] NS 8] NS
	6] Metrifonate no loading dose change from baseline							
	7] Metrifonate loading dose vs. placebo change from baseline							
	8] Metrifonate no loading dose vs Placebo change from baseline							

EvTable37. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12w		Final: (specify) 26w	
Morris 1998	ITT Analysis							
	1] Placebo	<u>ADAS-Cog</u>			4] 1.5 5] -0.5	3] 0.0002	1] 2.7 2] -0.3 3] 2.86 CI (1.37-4.34) 4] 2.5 5] -0.3	3] 0.0001
	2] Metrifonate 180mg loading 60mg maintenance							
	3] Difference between Placebo and Metrifonate change from baseline	<u>CIBIC+</u>			1] 4.15 2] 3.95	3] 0.0152	1] 4.35 2] 4.05 3] 0.28 CI (0.06-0.50) 4] 4.38 5] 4.05	3] 0.0071
	4] Placebo change from baseline	DAD						3] 0.0860
	5] Metrifonate change from baseline	GDS						3] 0.0734
		ADAS-Noncog						3] 0.1221
		MMSE						3] 0.1788

EvTable38. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 21d	
Pettigrew 1998	OC Analysis	ADAS-Cog					1] 0.00 (3.65) 2] 1.67 (2.52) 3] 1.33 (4.04) 4] -1.33 (2.4) 5] -2.63 (6.7)	
	1] Placebo change from baseline							
	2] Metrifonate 135mg loading 25mg maintenance change from baseline	ADAS-Noncog					1] -1.86 (2.1) 2] -3.67 (1.5) 3] 1.33 (5.77) 4] -2.00 (2.1) 5] -1.75 (2.6)	
	3] Metrifonate 225mg loading 35mg maintenance change from baseline	MMSE					1] 0.71 (2.50) 2] -0.67 (3.5) 3] 1.33 (0.58) 4] 0.50 (1.87) 5] 2.63 (2.07)	
	4] Metrifonate 335mg loading 60mg maintenance change from baseline	Blessed-DRS					1] -2.64 (2.46) 2] -1.17 (2.52) 3] 0.50 (2.29) 4] 0.67 (3.40) 5] 0.56 (1.66)	
	5] Metrifonate 335mg loading 90mg maintenance change from baseline							

EvTable39. Study results: Metrifonate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12 w		Final: (specify) 26 w	
Raskind 1999	ITT analysis	MMSE			3] -0.60 4] 0.75	5] <0.05	3] -1.25 4] 0.60	6] 0.0001 5] <.05
	1] Placebo							
	2] Metrifonate 50 mg d	NPI					3] .075 4] 0.30	6] 0.013 5] <0.05
	3] Placebo mean change from baseline	CIBIC+			3] 4.2 4] 4.05		3] 4.4 4] 4.78	5] <0.05 6] 0.039
	4] Metrifonate mean change from baseline	GDS ADAS-Cog						6] 0.570 6] 0.012
	5] Metrifonate vs. placebo	ADAS- Noncog						6] 0.185
	6] Mean drug- placebo difference change from baseline	DAD CIBIS+						6] 0.036 6] 0.260

EvTable40. Adverse Events: Metrifonate.

Adverse events (AE) identified in included studies	Becker, 1996	Becker, 1998	Cummings, 1997	Cummings, 1998	Dubois, 1999	Jann, 1999	Morris, 1998	Pettigrew, 1998	Raskind, 1999
Withdrawn (%) due to AE	T: 0 C: 0	T: 0 C: 0	T: 6 C: 4	T: 6 C: 4	T: 6 C: 6	T: 6 C: 2	T: 12 C: 4	T: 0 C: 0	T: 11 C: 9
AE Checklist (Max 5)	5	2	4	3	3	4	4	5	3
None Reported		X							
Balance					X				
Accidental Injury					X				
Dizziness	X				X			X	
Falls									
Behavioral									
Agitation	X							X	X
Cardiovascular	X		X						
Arrhythmia				X	X	X		X	
Hypotension					X			X	
Hypertension									
Extrapyramidal									
Tremor									
Gastrointestinal	X		X	X				X	
Abdominal pain	X		X	X				X	X
Constipation									
Diarrhea	X		X	X	X	S*	X	X	
Dyspepsia						X			
Nausea, vomiting	X		X	X	X	S*	X	X	
Metabolic/nutritional									
Eating disorder									
Weight Change									
Neurological								X	
Asthenia					X				
Psychiatric									
Anxiety									
Confusion, delirium									
Depression									
Respiratory	X								
Cough, cold, infection									
Rhinitis							X	X	X
Other	X		X	X	X			X	
Aberrant hematology					X				
Fatigue, weakness					X				
Fever, flu, pneumonia									
Headache	X								
Hepatic abnormality				X					
Muscle/joint disorder			X	X	X	S*	X		X
Pain									
Rash, skin disorder									
Sleep disorder	X							X	
Urinary disorder									

NR = Withdrawals due to AE Not Reported

+ = Dose response effect on AE

x = Reported adverse event/side effect but not tested for significant differences between groups

S or NS = Reported and tested for statistical differences between placebo and treatment group

S* or NS* = Reported and tested for statistical differences between two (three) treatment groups

[] = Symptom NOT reported in the paper

EvTable41. Key characteristics: Nicergoline.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Hermann 1997	NR	6	Placebo Nicergoline	DSM III	MID	Mild-Mod	139	136	73.5y (55-85y) 48%M	30 mg bid	6m	BL-A CGI DSPT DST DTIC HIS MMSE Laboratory Tests SCAG WAIS	No
Nappi 1997	IF	6	Placebo Nicergoline	DSM-III-R	PDD VaD Mixed	Mild-Mod	108	101	69.3y (55-81y) 55%M	30 mg bid	12m	CGI HAM-D MMSE Neurological Exam Physician Global Impression Patient Global Impression SCAG	No

EvTable41. Key characteristic:. Nicergoline cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Saletu 1995 AUXILIARY Saletu 1997	NR	6	Placebo Nicergoline	DSM-III-R	SDAT MID	Mild-Mod	112	98	Mean NR (>60y) %M NR	30 mg bid	8w	AAMD CGI CT Scan EEG/ERG Mapping Ham-D Laboratory tests MMS NOSIE SCAG TESS-DOTES	SDAT vs MID
Winblad 2001a	IF	6	Placebo Nicergoline	NINCDS DSM-III-R	AD	Probable Mild-Mod	346	285	73.7y (NR) 38%M	30 mg bid	6m	ADAS-cog ADAS-noncog ADAS-Total IADL CGIC CT/MRI ECG MMSE PSMS	No
DRUG VS DRUG													
Schneider 1994	NR	5	Nicergoline Antagonic Stress	DSM IV ICD-10	AD	Mild-Mod	62	NR	69.8y (65-85y) 53%M	60 mg/d	3m	SAS-G SCAG WAIS WMS	No

EvTable42. Study results: Nicergoline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4m		Final: (specify) 6m	
Hermann 1997	ITT Analysis 1] Placebo 2] Nicergoline 30 mg BID 3] change from baseline of differences between Placebo and Nicergoline	SCAG	1] 63.17(12.73) 2] 63.74(11.31)		1] 59.59(16.88) 2] 52.50(12.41)	3] <0.0001	1] 57.48(17.16) 2] 49.54(12.82)	3]<0.0001
		MMSE	1] 20.28 (2.62) 2] 20.23 (2.06)		1] 4.50 (1.07) 2] 3.69 (0.70)	3] <0.0001	1] 21.56 (3.81) 2] 24.03 (3.06)	3]<0.0001
		CGI (Item 2)	1] 5.02 (0.22) 2] 4.97 (0.18)		1] 22.31(10.19) 2] 24.17(10.21)	3] 0.1292	1] 4.43 (1.18) 2] 3.46 (0.92)	3]<0.0001
		WAIS-DST	1] 19.73 (8.73) 2] 20.62 (7.93)		1] 8.02 (1.57) 2] 8.93 (1.53)	3] .0004	1] 22.44 (9.91) 2] 5.37(11.08)	3] 0.0602
		WAIS-DSPT	1] .67 (1.63) 2] 7.56 (1.64)		1] 12.00 (4.78) 2] 13.66 (4.11)	3] 0.0328	1] 8.13 (1.66) 2] 8.98 (1.86)	3] 0.0042
		WAIS-DTIC	1] 10.63 (4.58) 2] 11.33 (4.08)		1] 10.41 (3.48) 2] 8.63 (3.54)	3] 0.0017	1] 12.15 (4.52) 2] 14.57 (3.79)	3] 0.0026
		Blessed-A scale	1] 10.85 (3.32) 2] 10.69 (3.35)				1] 9.94 (3.98) 2] 7.69 (3.75)	3] 0.0010

EvTable43. Study results: Nicergoline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6m		Final: (specify) 12m	
Nappi 1997	OC Analysis							
	1] Placebo	<u>SCAG</u> Total Score	1] 49.14 (6.9) 2] 49.14 (9.2)		1] 50.12 (7.8) 2] 45.53 (10.4)	3] <0.05	1] 53.72 (10.2) 2] 45.37 (11.7)	3] <0.001
	2] Nicergoline 30 mg bid	<u>MMSE</u>	1] 21.98 (2.6) 2] 21.25 (2.9)		1] 20.35 (3.5) 2] 21.78 (4.0)	3] <0.05	1] 19.14 (3.8) 2] 21.27 (4.1)	3] <0.01
	3] Nicergoline difference from placebo	Physician Global Impression (% improved)					1] 10% 2] 53%	3] <0.001
		Patient- Global Impression (% improved)					1] 8% 2] 64%	3] <0.001

EvTable44. Study results: Nicergoline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify)		Final] (specify) 8 w	
Saletu 1995	OC Analysis	CGI(Item 1)	1] 4.29 (0.95)				1] 4.04 (0.95)	5] <0.05
	1] Placebo (SDAT)		2] 4.33 (1.01)				2] 3.54 (0.88)	6] <0.001
	2] Nicergoline 30 mg bid (SDAT)		3] 4.23 (1.03)				3] 4.00 (1.17)	7] <0.03
	3] Placebo (MID)		4] 4.42 (1.02)				4] 3.83 (1.09)	8] <0.001
	4] Nicergoline 30 mg bid (MID)	CGI(Item 2)						9] <0.025
	5] Placebo change from baseline (SDAT)							10] <0.05
	6] Nicergoline change from baseline (SDAT)						1] 3.75 (0.61)	9] 0.001
	7] Placebo change from baseline (MID)						2] 3.21 (0.78)	10] 0.04
	8] Nicergoline change from baseline (MID)	SCAG total					3] 3.81 (0.57)	
	9] Placebo vs Nicergoline change from baseline (SDAT)						4] 3.29 (0.62)	
	10] Placebo vs Nicergoline change from baseline (MID)							
		MMSE	1] 55.2 (13.4)				1] 51.5 (11.0)	5] <0.05
			2] 56.8 (15.5)				2] 46.6 (11.4)	6] <0.01
			3] 54.3 (13.8)				3] 50.1 (14.3)	7] NS
			4] 55.8 (16.1)				4] 49.1 (15.9)	8] NS
		HAM-D						9] <0.01
								10] <0.05
		NOSIE						

EvTable45. Study results: Nicergoline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3m		Final: (specify) 6m	
Winblad, 2001a	ITT Analysis	<u>CGIC</u>			1] 4.26		1] 4.39	5] NS
		<u>Informant rated</u>			2] 4.08		2] 4.32	
		CGIC			1] 4.07		1] 4.16	5] NS
		Patient rated			2] 3.91		2] 4.09	
		<u>ADAS-Cog</u>					3] 1.38 (0.57)*	3] <0.01
							4]-0.17 (0.55)*	4] NS
							5] 1.55	5] <0.05
		ADAS-Noncog					3] 0.75 (0.26)*	3] <0.01
							4] 0.14 (0.26)*	4] NS
							5] 0.61	5] <0.1
		ADAS-Total					3] 1.93 (0.68)*	3] NS
							4]-0.08 (0.65)*	4] <0.01
							5] 2.01	5] <0.05
		IADL	1] 18.6 (6.6)				3] 1.47 (0.34)	5] NS
			2] 18.3 (6.9)				4] 1.13 (0.33)	
		PSMS	1] 9.5 (3.8)				3] 1.07 (0.19)	5] NS
			2] 9.6 (3.5)				4] 0.84 (0.19)	

*SEM

EvTable46. Study results: Nicergoline - Antagonic Stress.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Schneider 1994	OC Analysis	SCAG	1] 66.5 (11.5) 2] 71.2 (7.7)				1] 50.5 (8.6) 2] 46.1 (6.5)	3] <0.001 4] <0.001 5] 0.002 favors AS
	1] Nicergoline 20 mg tid							
	2] Antagonic Stress 3 capsules tid	SASG	1] 65.8 (9.5) 2] 68.4 (8.9)				1] 52.1 (9.4) 2] 47.3 (6.5)	3] <0.001 4] <0.001 5] 0.000 favors AS
	3] Nicergoline change from baseline	WAIS Digit symbol	1] 7.8 (1.2) 2] 7.5 (1.6)				1] 9.6 (1.6) 2] 11.5 (2.4)	3] <0.001 4] <0.001 5] 0.000 favors AS
	4] Antagonic Stress change from baseline							
	5] Difference between Nicergoline and Antagonic Stress in change from baseline							

EvTable47. Adverse Events: Nicergoline.

Adverse events (AE) identified in included studies	Hermann, 1997	Nappi, 1997	Saletu, 1995	Winblad, 2001a	NICERGOLINE ANTAGONIC STRESS Schneider, 1994
Withdrawn (%) due to AE	T: 1 C: 3	T: 0 C: 0	T: 2 C: 2	T: 9 C: 8	T: NR C: NR
AE Checklist (Max 5)	2	4	5	3	0
None Reported					x
Balance					
Accidental Injury					
Dizziness	x		x		
Falls					
Behavioral					
Agitation	x			x	
Cardiovascular	x				
Arrhythmia	x		x		
Hypotension					
Hypertension					
Extrapyramidal	x				
Tremor					
Gastrointestinal	x	x		x	
Abdominal pain					
Constipation	x		x		
Diarrhea	x		x		
Dyspepsia	x				
Nausea, vomiting	x				
Metabolic/nutritional				x	
Eating disorder	x				
Weight Change			x		
Neurological				x	
Asthenia					
Psychiatric				x	
Anxiety	x				
Confusion, delirium					
Depression	x		x		
Respiratory					
Cough, cold, infection	x				
Rhinitis	x		x		
Other	x	x	x		
Aberrant hematology					
Fatigue, weakness	x				
Fever, flu, pneumonia	x				
Headache	x	x	x	x	
Hepatic abnormality					
Muscle/joint disorder	x				
Pain	x				
Rash, skin disorder	x		x		
Sleep disorder	x		x	x	
Urinary disorder				x	

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable48. Key characteristics: Physostigmine.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Moller 1999	NR	6	Placebo Physostigmine	DSM-III-R NINCDS	AD	Mild-Mod	181	143	69.3y (50-85y) 48%M	30 and 60 mg patch	24w	ADAS CGIC HAM-D HIS MMSE NOSGER	No
Thal 1996b	IS PI	6	Placebo Physostigmine	NINCDS	AD	Probable	366	333	68.6y (47-87y) 51%M 97% White	Titration: 9, 12, 15 mg bid, increased weekly	6w	ADAS-Cog CGI-C IADL MMSE PSMS	No
Thal 1999	IF	5	Placebo Physostigmine	NINCDS	AD	Probable	475	210	73.4y range 51-92 39.8% M 91% White 80% High School Grads Community	2w washout Titration: 9mg bid, 12mg bid, 15mg bid 24w drug in increased every 3w week 9 to 24 30mg/d or 36mg/d	24w	ADAS-Cog GERRI IADL CIBIC+ CGIC	No

EvTable48. Key characteristics: Physostigmine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	#Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
VanDyck 2000	IF	7	Placebo Physostigmine	NINCDS	AD	Probable Mild-Mod	176	148	71.5y (44-88y) 63%M 95 % White 4 %Black 1% Hispanic	30 mg/d (15 mg bid)	12w	ADAS-Cog CIBIC+ CIBIC CGIC IADL MMSE	No

EvTable49. Study results: Physostigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24 w	
Moller, 1999	OC Analysis							
	1] Placebo	ADAS-Cog	1] 28.5 2] 29.8 3] 30.8				1] 27.1 2] 31.4 3] 31.3	
	2] Physostigmine 30 mg patch	ADAS-Noncog	1] 7.7 2] 7.5 3] 6.9				1] 6.6 2] 7.0 3] 6.5	
	3] Physostigmine 60 mg patch	ADAS-Total	1] 36.2 2] 37.3 3] 37.6				1] 33.7 2] 38.4 3] 37.9	
	4] Treatment vs Placebo from baseline	NOSGER	1] 68.8 2] 67.5 3] 66.6				1] 68.8 2] 72.0 3] 38.8	
		CGIC % improved					1] 47% 2] 36% 3] 21%	4] NS

EvTable50. Study results: Physostigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Thal 1996b	ITT Analysis	ADAS-Cog					1] 0.63 2] -1.12	3] 0.003
		CGIC					1] -0.04 2] 0.22	3] 0.0120
		MMSE					1] -0.60 2] 0.05	3] 0.132
		PSMS					1] 0.14 2] 0.33	3] 0.383
		IADL					1] 4.13 2] 1.67	3] 0.101

EvTable51. Study results: Physostigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24 w	
Thal 1999	ITT Analysis (LOCF) 1] Placebo 2] Physostigmine Controlled-release 15mg bid DIFFERENCE FROM PLACEBO 3] Physostigmine Controlled-release 18mg bid DIFFERENCE FROM PLACEBO	ADAS-cog					2] -2.9 3] -2.9	2] 0.002 3] 0.001
		CIBIC+					2] 0.31 3] 0.26	2] 0.019 3] 0.042
		CGIC					2] 0.16 3] 0.24	2] NS 3] NS
		GERRI					2] -0.07 3] -0.03	2] NS 3] NS
		IADL					2] -3.85 3] -0.10	2] NS 3] NS

EvTable52. Study results: Physostigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Van Dyck 2000	ITT Analysis 1] Placebo change from baseline 2] Physostigmine 12 or 15 mg BID change from baseline 3] difference between Placebo and Physostigmine in change from baseline	<u>ADAS-Cog</u>					1] 1.06 (5.17) 2] -0.96 (5.22)	3] 0.01
		<u>CIBIC+</u>					1] -0.33 (0.82) 2] 0.00 (0.88)	3] 0.02
		<u>CGIC</u>					1] -0.30 (0.84) 2] -0.12 (0.87)	3] 0.17
		MMSE					1] -0.87 (3.20) 2] -0.25 (2.98)	3] 0.22
		IADL					1] 3.51 (12.54) 2] 1.28 (12.48)	3] 0.25

EvTable53. Adverse Events: Physostigmine.

Adverse events (AE) identified in included studies	Moller, 1999	Thal, 1996b	Thal, 1999	Van Dyck, 2000
Withdrawn (%) due to AE	T: 12 C: 1	T: 16 C: 5	T: 55 C: 5	T: NR C: NR
AE Checklist (Max 5)	1	2	2	2
None Reported				
Balance				X
Accidental Injury			X	X
Dizziness		X		S
Falls				
Behavioral				
Agitation			X	X
Cardiovascular				
Arrhythmia				
Hypotension				
Hypertension				
Extrapyramidal				
Tremor	X	X		S
Gastrointestinal	X		X	X
Abdominal pain	X	X	X	X
Constipation				
Diarrhea		X	X	X
Dyspepsia			X	X
Nausea, vomiting	X	X	X	X
Metabolic/nutritional		X		
Eating disorder		X	X	X
Weight Change				S
Neurological				
Asthenia		X	X	S
Psychiatric				
Anxiety				X
Confusion, delirium				S
Depression				
Respiratory				S
Cough, cold, infection				
Rhinitis				
Other	X	X	X	S
Aberrant hematology				
Fatigue, weakness				
Fever, flu, pneumonia				
Headache	X	X		X
Hepatic abnormality				
Muscle/joint disorder				X
Pain				X
Rash, skin disorder	X			
Sleep disorder			X	X
Urinary disorder				

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable54. Key characteristics: Posatirelin.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Ferrari 1998	NR	5	Placebo Posatirelin	DSM-III-R ICD-10 NINCDS	AD PDD VaD	Probable	213	172	78.8y (≥65y) 42%M	10 mg/d	3m	GBS GDS HDRS HIS MMSE RMT	AD vs Vad
Gasbarrini 1997	NR	6	Placebo Posatirelin	NINCDS NINDS- AIREN	AD VaD	Probable Mild-Mod	360	357	77.6y (≥60y) 38%M	10mg/d	3m	GBS Laboratory tests Rey Memory Test	No
Parnetti 1996	PI	7	Placebo Posatirelin	NINDS- AIREN	VaD	Probable	136	105	69.4y (60-75y) 66%M comorbidity: hypertension	10 mg ml/d	12w	GBS Laboratory tests RMT TPAT	No
Parnetti 1995	NR	6	Placebo Posatirelin Citicoline	NINCDS	AD	Probable Mild-Mod	222	214	74.9y (65-85y) 34%M	10 mg/d 500 mg/d	3m	GBS GDS HDRS MMSE Laboratory tests	No

EvTable55. Study results: Posatirelin.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Ferrari, 1998	ITT population (LOCF)							
	1] Placebo change from baseline (ALL)	GBS Rating Scale Total score					1] 1.1 (2.0)* 2] -9.2 (3.4)* 4] 0.3 (3.2) 5] -2.2 (3.6)	3] <0.001
	2] Posatirelin 10mg/ml IM change from baseline (ALL)	GBS Rating Scale motor function				7] 1.5 (2.5) 8] -17.4 (5.8)	1] 0.9 (0.5)* 2] -1.5 (0.7)* 4] -0.4 (1.6) 5] -0.6 (0.9)	3] 0.010
	3] Posatirelin vs. Placebo change from baseline	GBS Rating Scale intellectual				7] -0.4 (1.2) 8] -8.8 (3.1)	1] -0.4 (0.9)* 2] -4.5 (1.8)* 4] -0.4 (1.6) 5] -0.9 (1.9)	3] 0.005
	4] Placebo change from baseline (AD)							
	5] Posatirelin 10mg/ml IM change from baseline (AD)	GBS Rating Scale emotional				7] -0.4 (1.2) 8] -8.8 (3.1)	1] -0.04 (0.3)* 2] -0.7 (0.4)* 4] 0.0 (0.4) 5] -0.03 (0.4) 7] -0.1 (0.4) 8] -1.6 (0.7)	3] 0.078
	6] Posatirelin Placebo change from baseline (AD)							
	7] Placebo change from baseline (VaD)							
	8] Posatirelin 10 mg/ml IM change from baseline (VaD)							
	9] Posatirelin vs. Placebo change from baseline (VaD)							

* SEM

EvTable56. Study results: Posatirelin.

Author Year	Outcome Measures	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 45d		Final: (specify) 90d	
Gasbarrini 1997	ITT Population							
	1] Placebo	<u>GBS motor impairment</u>	1] 12.2(7.7) 2] 12.1(7.7)		1] 11.8(7.3) 2] 11.1(7.4)		1] 11.9(7.6) 2] 10.5(7.6)	3] <0.001
	2] Posatirelin 10mg/d IM	<u>GBS intellectual impairment</u>	1] 25.9(10.7) 2] 25.7(10.7)		1] 25(10.7) 2] 23.6(10.2)		1] 24.5(11.4) 2] 21.9(10.7)	3] <0.001
	3] difference between Placebo and Posatirelin from baseline	<u>GBS emotional impairment</u>	1] 7.4(3.6) 2] 7.4(3.6)		1] 7.1(3.6) 2] 6.6(3.4)		1] 6.9(3.8) 2] 6.1(3.5)	3] <0.001
		GBS Factor I	1] 11.6 (5.1) 2] 11.6 (5.0)		1] 11.3 (5.2) 2] 10.7 (4.6)		1] 11.2 (5.6) 2] 9.9 (4.9)	3] <0.001
		GBS Factor II	1] 12.2 (7.7) 2] 12.1 (7.7)		1] 11.8 (7.3) 2] 11.1 (7.4)		1] 11.9 (7.6) 2] 10.5 (7.6)	3] <0.001
		GBS Factor III	1] 6.1 (3.2) 2] 6.4 (3.2)		1] 6.1 (3.2) 2] 5.8 (3.3)		1] 5.9 (3.2) 2] 5.4 (3.3)	3] <0.001
		GBS Factor IV	1] 19.3 (8.3) 2] 19.4 (8.3)		1] 18.6 (8.4) 2] 17.5 (8.3)		1] 18.1 (8.9) 2] 16.2 (8.6)	3] <0.001

EvTable57. Study results: Posatirelin.

Author Year	Analysis Groups	Outcome Measures	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 45d		Final: (specify) 90d	
Parnetti, 1996	OC Population 1] Placebo 2] Posatirelin 10mg/ml 3] Posatirelin 10mg/ml vs Placebo change from baseline	GBS-ADL	1] 7.8 (5.1) 2] 8.5 (5.1)		1] 8.0 (5.2) 2] 7.5 (5.8)		1] 7.9 (5.0) 2] 7.0 (5.1)	3] 0.001
		GBS Emotional Impairment	1] 6.1 (2.3) 2] 5.6 (2.4)		1] 5.9 (2.4) 2] 4.8 (2.3)		1] 5.7 (2.6) 2] 4.3 (2.1)	3] 0.066
		GBS Motor impairment	1] 7.9 (5.1) 2] 8.5 (5.1)		1] 8.0 (5.2) 2] 7.5 (4.8)		1] 7.9 (5.0) 2] 7.1 (5.1)	3] 0.001
		GBS Intellectual impairment	1] 19.8 (8.9) 2] 20.3 (8.8)		1] 19.9 (9.3) 2] 16.1 (7.2)		1] 20.4 (9.6) 2] 14.7 (7.5)	3]<0.001
		TP Global	1] 0.6 (0.4) 2] 0.7 (0.4)		1] 0.7 (0.4) 2] 0.8 (0.5)		1] 0.6 (0.4) 2] 0.9 (0.5)	3] 0.076
		Randt acquisition	1] 65.1 (13.0) 2] 64.2 (10.0)		1] 63.4 (2.3) 2] 69.0 (10.1)		1] 63.1 (12.9) 2] 70.3 (11.2)	3] <0.001
		Randt recall	1] 64.8 (23.7) 2] 66.7 (25.5)		1] 61.4 (24.3) 2] 67.5 (23.1)		1] 63.0 (26.9) 2] 72.2 (26.8)	3] 0.058
		Randt memory	1] 60.8 (20.8) 2] 60.5 (18.8)		1] 57.0 (18.9) 2] 63.5 (16.0)		1] 57.7 (21.4) 2] 67.1 (19.0)	3] <0.001

EvTable58. Study results: Citicoline - Posatirelin.

Author Year	Analysis Groups	Outcome s Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 45 d		Final: (specify) 90 d	
Parnetti 1995	OC Population 1] Placebo (Ascorbic Acid 100 mg/d) 2] Citicoline 500 mg/d 3] Posatirelin 10 mg/d 4] Ascorbic Acid change from baseline 5] Citicoline change from baseline 6] Posatirelin change from baseline 7] Posatirelin vs Citicoline change from baseline	<u>GBS</u> Emotional impairment	1] 1.9 (1.0) 2] 1.9 (1.0) 3] 1.9 (1.2)		1] 1.8 (1.0) 2] 1.9 (1.1) 3] 1.7 (1.0)		1] 1.9 (1.0) 2] 1.7 (1.0) 3] 1.6 (0.9)	4] NS 5] NS 6] <0.025
		<u>GBS</u> Impaired orientation & memory	1] 2.2 (0.9) 2] 2.2 (1.0) 3] 2.1 (1.0)		1] 2.1 (1.0) 2] 2.1 (1.0) 3] 2.0 (1.0)		1] 2.1 (1.1) 2] 2.1 (1.0) 3] 1.8 (1.0)	4] NS 5] NS 6] NS 7] 0.038 favors Posatirelin
		<u>GBS</u> Impaired ability ADL	1] 1.2 (0.8) 2] 1.4 (1.1) 3] 1.2 (1.0)		1] 1.3 (0.8) 2] 1.3 (1.0) 3] 1.1 (1.0)		1] 1.3 (0.9) 2] 1.4 (1.0) 3] 1.1 (1.0)	4] NS 5] NS 6] <0.025
		<u>GBS</u> Depression / Anxiety	1] 1.5 (0.9) 2] 1.5 (0.8) 3] 1.6 (1.1)		1] 1.5 (0.9) 2] 1.5 (0.9) 3] 1.5 (1.0)		1] 1.4 (0.9) 2] 1.4 (0.9) 3] 1.4 (0.9)	4] NS 5] NS 6] NS 7] 0.031 favors Posatirelin
		<u>GBS</u> Impaired attention & motivation	1] 2.2 (0.9) 2] 2.1 (1.0) 3] 2.1 (1.1)		1] 2.1 (0.9) 2] 2.0 (1.0) 3] 1.9 (0.9)		1] 2.1 (1.0) 2] 1.9 (1.0) 3] 1.8 (0.8)	4] NS 5] NS 6] <0.025
		<u>GBS</u> Intellectual impairment	1] 2.2 (0.8) 2] 2.1 (0.9) 3] 2.0 (0.9)		1] 2.1 (0.9) 2] 2.0 (0.9) 3] 1.9 (0.9)		1] 2.1 (1.0) 2] 2.0 (0.9) 3] 1.8 (0.8)	5] <0.025 7] 0.037 favors Posatirelin
		<u>GBS</u> Motor impairment	1] 1.2 (0.8) 2] 1.4 (1.1)		1] 1.3 (0.8) 2] 1.3 (1.0) 3] 1.1 (1.0)		1] 1.3 (0.9) 2] 1.4 (1.0) 3] 1.1 (1.0)	5] <0.025
		MMSE	1] 16.4 (2.7) 2] 16.5 (2.6) 3] 16.6 (2.5)				1] 17.1 (4.1) 2] 17.6 (3.9) 3] 17.8 (3.4)	4] NS 5] NS 6] NS
		HDRS	1] 13.0 (5.0) 2] 11.4 (4.9) 3] 12.6 (5.0)				1] 11.4 (4.9) 2] 11.3 (5.2) 3] 11.1 (5.3)	4] NS 5] NS 6] NS

EvTable59. Adverse Events: Posatirelin.

Adverse events (AE) identified in included studies	Ferrari, 1998	Gasbarrini, 1997	Parnetti, 1995	Parnetti, 1996
Withdrawn (%) due to AE	T: 4 C: 3	T: 3 C: 1	T: 0 C: 0	T: NR C: NR
AE Checklist (Max 5)	3	4	4	2
None Reported				
Balance	X			
Accidental Injury				
Dizziness	X	X		
Falls				
Behavioral				
Agitation	X	X	X	X
Cardiovascular				
Arrhythmia	X		X	X
Hypotension				
Hypertension		X		
Extrapyramidal				
Tremor			X	X
Gastrointestinal				X
Abdominal pain	X			
Constipation				
Diarrhea		X		
Dyspepsia			X	
Nausea, vomiting	X	X		X
Metabolic/nutritional		X		
Eating disorder				
Weight Change				
Neurological				
Asthenia				X
Psychiatric				
Anxiety				
Confusion, delirium			X	
Depression				
Respiratory				
Cough, cold, infection				
Rhinitis				
Other		X	X	X
Aberrant hematology		X		
Fatigue, weakness				
Fever, flu, pneumonia				
Headache	X	X	X	
Hepatic abnormality				
Muscle/joint disorder		X		
Pain				
Rash, skin disorder	X	X	X	
Sleep disorder	X		X	X
Urinary disorder		X	X	

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable60. Key characteristics: Rivastigmine.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Agid 1998	IF	7	Placebo Rivastigmine	DSM-III-R NINCDS	Dementia AD	Mild-Mod Probable	386	357	69.4y (50-90y) 44%M	Titration: 3 w Then 4 or 6 mg/d	15w	Benton Visual Retention CGIC Digit Symbol Substitution ECG Fuld Object-Memory Laboratory tests MMSE NOSGER Trail Making	No
Corey-Bloom 1998 Auxiliary: Farlow 2001 Farlow 2000 Kumar 2000 Del Ser 2000 Doraiswamy 2002	IF	8	Placebo Rivastigmine	DSM-IV NINCDS	AD	Mild-Modly Sev	699	545	74.5y (45-89y) 39%M 93% of patients were taking medications for other conditions: cardiovascular (43%), gastro (59%) analgesics (56%) 95% White	Titration: w 1-7; Then dose range: 1 –6 mg/d	26w	ADAS-Cog CIBIC+ GDS MMSE PDS	Vascular Risk (HIS)

EvTable60. Key characteristics: Rivastigmine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Forette 1999	IF	6	Placebo Rivastigmine	NINCDS DSM-III-R	DAT	Mild-Mod	114	85	70.6y (NR) %M NR	Titration: 2 mg/d, 3 mg/d at d 4, increment of 0.5 mg q4d until d 28 and increment of 1 mg qw 12 mg/d	18w	ADAS-cog CIBIC+ Digit span test ECG Laboratory tests NOSGER Wechsler logical memory test	No
McKeith 2000	IF	8	Placebo Rivastigmine	Consensus Criteria	Lewy- body Dementi a	Mild-Mod	120	92	73.9y (57-87y) 56%M	Titration: 1.5 mg bid for 2 w up to 6 mg bid	20w	CGC-plus ECG Laboratory tests MMSE NPI NPI-10 NPI-4 UPDRS	No
Potkin 2001	PI	5	Placebo Rivastigmine	DSM IV NINCDS	AD	Mild-Modly Sev	27	27	75.9y (64-89y) %M NR	Titration over a period of 12 w then 3, 6, or 9 mg/d	26w	CIBIC+ FDG-PET scan MMSE MRI Snodgrass Picture Naming Task	No

EvTable60. Key characteristics: Rivastigmine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Rosler 1999 AUXILIARY Rosler 2001 Farlow 2000 Rosler 1998 Doraiswamy 2002	IF	8	Placebo Rivastigmine	DSM IV NINCDS	AD	Mild-Modly Sev	725	581	72y (45-95y) 41%M Community. Comorbidity: diabetes, hypertension, arthritis	Titration on w 1 to 12 with increments of 1.5 mg/d4 mg/d 12 mg/d	14w	ADAS-Cog CIBIC CIBIC Progressive Deterioration scale Global Deterioration scale MMSE	No

EvTable61. Study results: Rivastigmine.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 7w		Final: (specify) 13w	
AGID 1998	Observed Cases	CGIC					1] 29.91%	5] <0.05
	1] Placebo	Successful outcome					2] 31.53%	
	2] Rivastigmine 4 mg BID	Digit Symbol Substitution			6] 0.1(7.4) 7] 2.1(5.8) 8] 2.0(5.4)	5] <0.005	6] 0.5(6.9) 7] 1.7(5.1) 8] 2.8(8.1)	4] NS 5] <0.05
	3] Rivastigmine 6 mg BID							
	4] Rivastigmine 4 mg bid versus Placebo change from baseline	MMSE					6] 0.0(2.6) 7] 0.0(3.3) 8] 0.3(3.1)	
	5] Rivastigmine 6 mg bid versus Placebo change from baseline	Trail Making			6] -0.6(31.2) 7] -4.3(36.9) 8] -5.4(45.3)		6] 0.5(28.7) 7] -1.6(39.0) 8] -7.3(48.9)	
	6] Placebo change from baseline	NOSGER IADL					6] -0.2 (3.3) 7] 0.0 (3.3) 8] -0.7 (3.5)	4] NS 5] NS
	7] Rivastigmine 4 mg bid change from baseline							
	8] Rivastigmine 6 mg bid change from baseline	NOSGER Mood, memory, self- care, social behavior, disturbing behavior						4] NS 5] NS

EvTable61. Study results: Rivastigmine cont'd.

REF ID#	Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify) 7w		Final: (specify) 13w	
			FULD Object-Memory Evaluation (TS)			6] 0.0 (6.2) 7] 2.2 (7.3) 8] 2.0 (6.6)	4] <0.01 5] <0.05	6] -0.9 (5.5) 7] -0.4 (6.2) 8] 0.7 (6.2)	4] <0.05 5] <0.05
			FULD Object-Memory Evaluation (TR)			6] 0.5(4.6) 7] 1.7 (5.3) 8] 2.4 (4.8)	4] <0.05 5] <0.005	6] 0.1 (4.3) 7] 0.8 (4.6) 8] 1.1 (4.2)	4] NS 5] <0.05

EvTable62. Study results: Rivastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 26w	
Corey-Bloom 1998	ITT Analysis 1] Placebo	<u>ADAS-Cog</u>	1] 21.7 2] 22.4 3] 22.3				4] - 4.09 5] - 2.36 6] - 0.31	8] <0.001
Farlow 2001	2] Rivastigmine 1-4 mg/ d	<u>CIBIC+</u>					4) 0.49 5) 0.23 6) 0.20	8] < 0.01
Farlow 2000	3] Rivastigmine 6-12 mg/ d	<u>PDS</u>					4] -4.90 5] -5.19 6] - 1.52	8] < 0.001
Kumar 2000	4] Placebo change from baseline	<u>GDS</u>	1] 3.9 2] 4.0 3] 4.0				4] -0.32 5] -0.16 6] -0.13	7] <0.05 8] < 0.030
Delser 2000	5] Rivastigmine low dose change from baseline							
Dorais- wamy 2002	6] Rivastigmine high dose change from baseline	MMSE	1] 20 2] 19.5 3] 19.6	9] 0.481			4] -7.9 5] -0.35 6] 0.30	8]< 0.05
	7] Placebo vs Rivastigmine low dose							
	8] Placebo vs Rivastigmine high dose							
	9] Difference among groups							

EvTable63. Study results: Rivastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 18 w	
Forette 1999	Completers Analysis	ADAS-Cog	1] 21.7 (8) 2] 24.1 (11.6) 3] 23.2 (8.5)				4] 2 5] -2.75 6] 0.25	7] 0.054
	1] Placebo							
	2] Rivastigmine 12 mg d variable bid	NOSGER Self-care					4] -0.3 (2.5) 5] -0.4 (2.0) 6] -0.6 (2.4)	8] NS
	3] Rivastigmine 12 mg d variable tid	NOSGER Disturbing behaviour					4] 0.1 (3.1) 5] -0.3 (2.1) 6] -0.7 (3.4)	8] NS
	4] Placebo mean change from baseline	NOSGER IADL					4] 0.8 (4.0) 5] 0.4 (3.1) 6] -0.7 (4.0)	8] NS
	5] Rivastigmine bid mean change from baseline	NOSGER Memory					4] 1.3 (3.7) 5] -0.7 (2.9) 6] -1.0 (2.7)	5] 0.037 6] 0.014 8] 0.032
	6] Rivastigmine tid mean change from baseline	NOSGER Mood					4] -0.6 (3.2) 5] 0.7 (3.0) 6] -0.4 (3.4)	8] NS
	7] Rivastigmine bid vs. placebo	NOSGER Social behaviour					4] 0.3 (3.3) 5] 0.0 (2.6) 6] -1.1 (3.8)	8] NS
	8] Rivastigmine bid & tid vs. placebo	CIBIC+ Improved					1] 16% 2] 57% 3] 36%	7] 0.027
		Wechsler logical memory test					5] 1.8 (2.3) 6] 0.1 (2.3)	8] 0.012 immediate recall only

EvTable64. Study results: Rivastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 20w	
McKeith 2000	ITT Analysis							
	1] Placebo	<u>NPI-4</u>	1] 11.7 (8.6) 2] 12.2 (8.2)				3] 0.8 (7.3) 4] 2.5 (8.4)	5] 0.088 CI (-1.1 – 4.6)
	2] Rivastigmine 12 mg d	<u>NPI-10</u>					3] 1.2 (10.7) 4] 5.0 (16.2)	5] 0.048 CI (-1.6 – 9.2)
	3] Placebo mean change from baseline	<u>CCASSS</u>				5] 0.01		5] 0.048
	4] Rivastigmine mean change from baseline	CGC+						5] NS
	5] Rivastigmine versus placebo change from baseline	MMSE	1] 17.8 (4.4) 2] 17.9 (4.7)					5] NS

EvTable65. Study results: Rivastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 26 w	
Potkin 2000	OC Population 1] Placebo 2] Rivastigmine 9 mg/d 3] Placebo vs. Rivastigmine 4] Rivastigmine metabolism change from baseline	Snodgrass Picture Naming task CIBIC+ Ratio stabilized					1] 2/7 2] 15/20	3] NS 3] <0.03*

* Tested the difference between the placebo and treatment group that had deteriorated (CIBIC+ score 5-7) only

EvTable66. Study results: Rivastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 18w		Final: (specify) 26w	
Rosler 1999	ITT Analysis	ADAS-Cog	1] 23.4 2] 23.9 3] 23.6				4] - 1.34 5] - 1.37 6] 0.26	7] > 0.05 8] > 0.011
Rosler 2001	1] Placebo	ADAS COG >4 point improvement					1] 16% 2] 15% 3] 24%	7] >0.05 8] <0.1
Farlow 2000	2] Rivastigmine 1-4 mg/ d							
Rosler 1998	3] Rivastigmine 6-12 mg/ d							
Dorais- wamy 2002	4] Placebo change from baseline	CIBIC					4] 4.38 5] 4.24 6] 3.91	7] > 0.05 8] < 0.001
	5] Rivastigmine low dose change from baseline	CIBIC+			4] 4.09 5] 4.06 6] 3.85		4] 4.34 5] 4.20 6] 3.93	8] < 0.05
	6] Rivastigmine high dose change from baseline	PDS						
	7] Placebo vs Rivastigmine low dose		1] 54.1 2] 53.8 3] 55.2				4] -2.18 5] -3.37 6] 0.05	7] > 0.05 8] < 0.07
	8] Placebo versus Rivastigmine high dose	GDS					4] -0.26 5] -0.22 6] -0.06	8] < 0.05
		MMSE					4] -0.47 5] -0.62 6] 0.21	8] < 0.05

EvTable67. Adverse Events: Rivastigmine.

Adverse events (AE) identified in included studies	Agid, 1998	Corey-Bloom, 1998	Forette, 1999	McKeith, 2000	Potkin, 2001	Rosler, 1999
Withdrawn (%) due to AE	T: 11 C: 4	T: 18 ⁺ C: 7	T: 27 C: 4	T: 12 C: 11	T: NR C: NR	T: 15 ⁺ C: 7
AE Checklist (Max 5)	4	3	5	4	2	2
None Reported					X	
Balance						
Accidental Injury						
Dizziness	X	S*	X			X
Falls						
Behavioral						
Agitation				X		
Cardiovascular			NS	NS		
Arrhythmia						
Hypotension						
Hypertension		S*				
Extrapyramidal						
Tremor						
Gastrointestinal		S*				
Abdominal pain	X		X			X
Constipation						
Diarrhea	X					X
Dyspepsia		S*				
Nausea, vomiting	X	S*	X	S		S*
Metabolic/nutritional						
Eating disorder		S*	X	S		X
Weight Change		S*	X	X		
Neurological		S*	X			
Asthenia		S*				
Psychiatric						
Anxiety						
Confusion, delirium						
Depression						
Respiratory						
Cough, cold, infection						
Rhinitis						
Other		S*	X			X
Aberrant hematology	NS*		NS	NS		
Fatigue, weakness		S*				X
Fever, flu, pneumonia						
Headache	X		X			X
Hepatic abnormality	NS*					
Muscle/joint disorder						
Pain						
Rash, skin disorder						
Sleep disorder		S*		S		
Urinary disorder						

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable68. Key characteristics: Tacrine.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	#Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Allain 1999	NR	7	Tacrine + Placebo Tacrine + Silymarin	NINCDS DSM-III-R	AD	Mild-Mod	222	194	74.2y (NR) 39%M Community All subjects + Tacrine	420 mg/d	15w	MMSE SKT	No
Gutzmann 2002	PI	7	Idebenone Tacrine	NINCDS DSM-III-R	AD PDD	Mild-Mod	203	44	71.2y (44-90y) 36%M 100% White 100% Community	360 mg/d 160 mg/d	60w	ADAS-Cog ADAS-Noncog ADAS-Total CGI CT EIS HIS MRI NOSGER-IADL	No

EvTable68. Key characteristics: Tacrine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Knapp 1994b Auxiliary: Farlow 1998, Schneider 1997, Raskind 1997, Henke 1997, Schneider 1996, Knopman 1996, Gracon 1996, Smith 1996, Knapp 1994	IF	7	Placebo Tacrine	NINCDS	AD	Probable Mild-Mod	663	279	72.8y (49-95y) 48%M	Titration: Group1-40 mg/d for 6w then 80 mg/d for 24w Group2-40 mg/d for 6w then 80 mg/d for 6w then 120 mg/d for 18w Group3-40 mg/d for 6w then 80 mg/d for 6w then 120 mg/d for 6w then 160 mg/d for 12w	30w	ADAS-Cog ADAS-Noncog ADAS-Total GDS CIBI FCCA GDS IADL MMSE PDS PSMS	ERT APOE Genotyp e Gender

EvTable68. Key characteristics: Tacrine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Maltby 1994	NI	6	Placebo Tacrine + Lecithin	NINCDS DSM-III-R	AD	Probable Mild-Mod	41	32	68.8y (52-84y) 51%M 100% Commu nity	Tacrine: Started at 25 mg/d and doses increased by 25 mg q2w up to 100 mg/d	36w	Activities of daily living Carer Stress Assessment Cholinergic sensitive test Extrapyramidal score Digit Span Face recognition GDS LFT-Liver function test London psychogeriatric rating scale MMSE Mood states scale National adult reading test Neurological exam Selective reminding test Symptoms of stress Verbal Fluency Walsh tests	No
Prentice 1996	NI PI	5	Placebo Tacrine	DSM-III-R NINCDS	AD	Probable	23	19	68.0y (NR) 13%M	40 mg/d for 6 w, then 80 mg/d for 6 w	13w	CAMCOG CAMTOT MMSE Rivermead Behavioral Memory Test – Profile RPT Score SPET Scan	No

EvTable68. Key characteristics: Tacrine cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Weinstein 1991 Auxiliary: Goad 1991	NI IS	7	Placebo THA & Lecithin	DSM-III-R	AD	Probable	13	12	74.6y (56-79y) 50%M 100% Community	Titration: increments of 25 mg/d for 4 w, then 100 mg/d 10 g/d	12w	Burden Scale CAMCOG CAMDEX CT HDRS IDDD Laboratory tests MMSE	No
Wong 1999	IS	5	Placebo Tacrine	NINCDS	AD	Probable Mild-Mod	100	94	73.8y (52-94y) 50%M	30 mg/d for 6 w, 60 mg/d for 6 w, 90 mg/d for 6 w, then 120 mg/d	30w	CGIC CASI IQCODE ADS MMSE FCCA HIS	No
Wood 1994	IF	6	Placebo Tacrine	NINCDS	AD	Mild-Mod	154	131	75y (NR) 54%M 100% Community	Titration: 20 mg/bid for 2 d, 20 mg/tid for 2 d 40 mg bid for 7 d. Then the dose could be increased or decreased in 20 mg amount to reach optimum dose. Max dose: 120 mg/d	12w	ADAS-Noncog AMTS Blessed Scale CGRS GBS LFT-Liver function test Mann-Whitney test MMSE RGRS Rosen	No

EvTable69. Study results: Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 15w	
Allain 1999	ITT Analysis	<u>SKT</u>	1] 17.2 (6.1) 2] 17.6 (6.1)				1] 17.4 (6.3) 2] 18.2 (6.6)	1] NS 2] NS
		<u>MMSE</u>	1] 17.8 (4.4) 2] 17.1 (4.2)				1] 18.3 (5.4) 2] 17.3 (5.3)	1] NS 2] NS
	1] Placebo + Tacrine 80mg qid							
	2] Silymarin 140mg tid + Tacrine 80 mg qid							

EvTable70. Study results: Idebenone-Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 60 w	
Gutzmann 2002	ITT Analysis 1] Idebenone 360 mg d 2] Tacrine 160 mg d variable 3] Difference between Idebenone and Tacrine	<u>EIS%</u>					1] 28.9%	3] <0.0001 favours Idebenone
		Rating = -1					2] 9.0%	
		Rating = 0					1] 54.8%	
		Rating = 1					2] 83.8%	
		Rating = 2					1] 16.3%	
		Rating = 3					2] 7.1%	3] NS
		ADAS-Total	1] 41.55(16.46) 2] 41.52(14.92)				1] 34.51(17.43) 2] 30.44(16.32)	
		ADAS-Cog	1] 30.23(11.59) 2] 30.93(10.59)				1] 26.40(16.67) 2] 24.81(14.92)	
		ADAS-Noncog	1] 11.32 (6.79) 2] 10.55(5.86)				1] 8.11(7.56) 2] 5.63(6.10)	
		CGI-S	1] 5.22 (0.46) 2] 5.19 (0.44)				1] 4.43 (1.58) 2] 4.53 (1.45)	
		NOSGER-IADL	1] 13.88(4.43) 2] 13.78(4.55)				1] 13.13 (5.49) 2] 12.5 (6.25)	3] NS

EvTable71. Study results: Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 30w	
Knapp 1994	ITT Analysis	<u>CIBI</u>					1] 19% improved 4] 41% improved 5] -0.01 CI (-0.4-0.1) 6] -0.02 CI (-0.4 -0.006) 7] 0.2 CI (-0.4 - -0.01)	5] 0.33 6] 0.04 7] 0.04
	1] Placebo							
	2] Tacrine 80mg/d							
	3] Tacrine 120mg/d	<u>ADAS-Cog</u>					1] 29.2 (11.8) 2] 30.9 (13.4) 3] 28.5 (11.1) 4] 28.0 (11.8) 5] -1.4 CI (-3.5-0.7) 6] -2.0 CI (-3.5 - -0.5) 7] -2.2 CI (-3.5 - -0.8)	5] 0.20 6] 0.008 7] 0.002
	4] Tacrine 160mg/d variable							
	5] Tacrine 80mg/d versus Placebo	<u>FCCA</u> (% improved)					1] 16% 4] 42%	7] <0.002
	6] Tacrine 120mg/d versus Placebo	GDS					5] 0.07 CI (-0.1-0.3) 6] -.05 CI (-0.2-0.08) 7] -0.2 CI (-0.3 - -0.04)	5] 0.48 6] 0.47 7] 0.01
	7] Tacrine 160mg/d variable versus Placebo	ADAS-Noncog					5] -0.8 CI (-2.2-0.6) 6] -0.3 CI (-1.3-0.7) 7] -0.7 CI (-1.6-0.2)	5] 0.26 6] 0.52 7] 0.12
		ADAS-Total					5] -2.4 CI (-5.2-0.4) 6] -2.2 CI (-4.2 - -0.3) 7] -3.0 CI (-4.8- -1.1)	5] 0.09 6] 0.03 7] 0.002
		MMSE					1] 18.2 (5.0) 2] 17.1 (4.6) 3] 18.7 (4.6) 4] 18.8 (4.5) 5] 0.6 CI (-0.6-1.7) 6] 0.4 CI (-0.4 - -1.2) 7] 0.9 CI (0.1-1.6)	5] 0.33 6] 0.37 7] 0.02

EvTable72. Study results: Tacrine & Lethicin.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6 m		Final: (specify) 9 m	
Maltby 1994	OC Population 1] Placebo & Lethicin 9X1200mg d 2] Tacrine 100 mg/d variable & Lethicin 9X1200mg d 3] Treatment effect Tacrine & Lethicin vs. Placebo & Lethicin	MMSE	1] 17.3 (6.7) 2] 16.6 (6.8)		1] 16.4 (6.8) 2] 16.1 (6.5)		1] 15.2 (8.0) 2] 14.4 (6.7)	3] 0.9444
		IADL	1] 10.3 (2.6) 2] 8.5 (2.5)		1] 9.5 (2.6) 2] 8.3 (2.8)		1] 9.6 (2.7) 2] 7.6 (3.4)	3] 0.622
		LPRS	1] 14.1 (7.7) 2] 15.2 (6.5)		1] 15.7 (10.9) 2] 18.2 (9.6)		1] 18.9 (12.1) 2] 19.5 (10.9)	3] 0.638
		Verbal (words)	1] 35.5 (15.9) 2] 34.9 (15.1)		1] 37.5 (16.0) 2] 38.7 (16.5)		1] 34.9 (16.3) 2] 38.5 (16.3)	3] 0.200
		Visual (objects)	1] 36.8 (21.2) 2] 30.7 (13.3)		1] 32.8 (22.8) 2] 31.4 (18.1)		1] 30.8 (22.7) 2] 26.2 16.9)	3] 0.359
		Digit forward	1] 5.9 (2.7) 2] 6.1 (2.8)		1] 5.2 (2.6) 2] 5.9 (2.4)			3] 0.723
		Verbal fluency	1] 22.4 (13.2) 2] 23.1 (15.0)		1] 19.8 (11.8) 2] 24.3 (14.8)			3] 0.198
		Face recognition	1] 3.8 (1.4) 2] 3.9 (2.0)		1] 3.4 (1.8) 2] 3.4 (1.2)			3] 0.651
		Carer stress assess.	1] 2.4 (4.4) 2] 4.0 (6.3)		1] 2.6 (4.9) 2] 2.0 (2.2)		1] 2.2 (3.9) 2] 3.1 (4.8)	3] 0.397

EvTable73. Study results: Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Prentice 1996	OC Population	MMSE	1] 18.4 (6.7) 2] 14.9 (5.0)				1] 19.1 (7.0) 2] 15.7 (5.2)	3] NS
	1] Placebo 2] Tacrine 40 mg/d 6w, then 80 mg/d 6w.	CAMTOT	1] 67.0 (12.5) 2] 53.5 (17.3)				1] 68.4 (18.0) 2] 54.7 (20.7)	3] NS
	3] Placebo vs. Tacrine	RPT	1] 4.1 (3.7) 2] 4.1 (3.0)				1] 5.6 (4.7) 2] 3.6 (3.1)	3] NS

EvTable74. Study results: Tetrahydroaminoacridine & Lethicin.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Weinstein 1990	OC Population 1] Placebo 2] Tetrahydro- aminoacridine (THA) 100 mg/d & Lethicin 10 g/d 3] THA vs. Placebo	CAMCOG IDDD Burden scale MMSE	1] 63 (12) 2] 44 (26)				1] 64 (16) 2] 45 (29)	3] NS 3] NS 3] NS 3] NS

EvTable75. Study results: Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 18w		Final: (specify) 30w	
Wong 1998	ITT Analysis	<u>CGIC</u>					3] 0.05 4] 0.02	5] 0.802
	1] Placebo							
	2] Tacrine 120 mg/d	<u>CASI</u>					3] -3.60 4] -0.85	5] 0.050
	3] Placebo mean change from baseline	<u>IQCODE</u>					3] 2.95 4] 3.13	5] 0.835
	4] Tacrine mean change from baseline	ADS					3] -1.00 4] -0.66	5] 0.978
	5] Tacrine vs. Placebo mean change from baseline	MMSE	1] 17.9 (4.8) 2] 16.2 (4.8)				3] 1.50 4] -0.21	5] 0.057

EvTable76. Study results: Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12 w	
Wood 1994	ITT (LOCF) Analysis 1] Placebo 2] Tacrine 80 mg/d variable 3] Placebo least squares mean 4] Tacrine least squares mean 5] Tacrine vs. Placebo	MMSE					3] 18.44 (0.54)* 4] 18.82 (0.55)*	5] 0.55
		CGRS						5] 0.012
		RGRS						5] 0.03
		Blessed					3] 4.89 (0.24)* 4] 4.73 (0.25)*	5] 0.60
		GBS					3] 31.70 (1.70)* 4] 29.10 (1.71)*	5] 0.20
		Adas- noncog					3] 8.11 (0.48)* 4] 7.97 (0.51)*	5] 0.53
		AMTS					3] 4.87 (0.21)* 4] 5.21 (0.22)*	5] 0.18

* SEM

EvTable77. Adverse Events: Tacrine.

Adverse events (AE) identified in included studies	Knapp, 1994	Maltby, 1994	Prentice 1996	Weinstein, 1991	Wong, 1999	Wood, 1994	Tacrine+Silymarin (T) Tacrine+Placebo(C) Allain 1988	TACRINE (T) IDEBENONE (C) Gutzmann, 2002
Withdrawn (%) due to AE	T: 55 C:11	T: 43 C: 0	T: 0 C: 0	T: 14 C: 0	T: 27 C:12	T: 16 C: 5	T: 8 C: 8	T: 41 C: 17
AE Checklist (Max 5)	3	3	1	3	2	3	3	3
None Reported								
Balance								
Accidental Injury								
Dizziness	X				X	X		
Falls								
Behavioral		X					NS	
Agitation	X							
Cardiovascular								
Arrhythmia								
Hypotension								
Hypertension								
Extrapyramidal								
Tremor					X			
Gastrointestinal	X							S*
Abdominal pain	X				X			
Constipation								
Diarrhea	X				X	X	NS	
Dyspepsia	X							
Nausea, vomiting	X	X		X	X	X	NS	S*
Metabolic/nutritional								
Eating disorder	X				X		NS	
Weight Change	X							
Neurological			X					
Asthenia							NS	
Psychiatric								
Anxiety								
Confusion, delirium								
Depression		X						
Respiratory							NS	
Cough, cold, infection								
Rhinitis	X							
Other		X			X	X		
Aberrant hematology								
Fatigue, weakness							NS	
Fever, flu, pneumonia								
Headache	X					X		
Hepatic abnormality	X	X	X	X	X	X		S*
Muscle/joint disorder		X						
Pain								
Rash, skin disorder								
Sleep disorder							NS	
Urinary disorder		X				X		

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable78. Key characteristics: Velnacrine.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Antuono 1995	IF	7	Placebo Velnacrine	NINCDS	AD	Probable	449	280	72.8y (47-90y) 38%M	150 or 225 mg/d	24 w	ADAS-Cog CATS CGI-C PGR PSMS RAGS	No
Huff 1991	PI	6	Placebo HP 128 (Velnacrine)	NINCDS	AD	Probable	16	15	70.5y (46-84y) 31%M	100 mg bid	13d	ADAS Benton MAE CGI Plasma Levels RAGS Sentence Repetition Token Test Verbal Fluency Visual Naming	No
Zemlan 1996	IF	6	Placebo Velnacrine	NINCDS	AD	Probable	309	225	71.6y (51-89y) 41%M	10 mg, 25 mg, 50 mg or 75 mg tid	15w	ADAS-Cog CGI-C IADL PGIR PSMS	No

EvTable79. Study results: Velnacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12w		Final: (specify) 24w	
Antuono 1995	LOCF Analysis							
	1] Placebo	<u>ADAS-Cog estimate</u>			4] -0.5 5] -0.5 6] -2.0	4] NS 5] NS 6] <0.001	4] 1.5 5] 1.5 6] -1.0	7] NS 8] NS 9] <0.05
	2] Velnacrine 150 mg/d	<u>CGI-C estimate</u>			1] 4.1 2] 4.0 3] 3.8	4] 0.001 5] 0.001 6] 0.001	1] 4.3 2] 4.1 3] 0.0	4] 0.001 5] 0.001 6] 0.001 7] 0.001
	3] Velnacrine 225 mg/d							
	4] Placebo change from baseline	PGIR			1] 4.1 2] 3.9 3] 3.8		1] 4.4 2] 4.3 3] 3.9	7] <0.05
	5] Velnacrine 150 mg/d change from baseline	PSMS			4] 0.62 5] -0.01 6] 0.30		4] 1.07 5] 0.49 6] 0.43	7] <0.05
	6] Velnacrine 225 mg/d change from baseline	RAGS			4] 1.12 5] 0.46 6] 0.10		4] 2.68 5] 2.87 6] 1.55	7] NS 8] NS 9] NS
	7] Combined Velnacrine vs Placebo change from baseline	CATS						7] <0.05 9] <0.01
	8] Velnacrine 150 mg/d vs Placebo change from baseline							
	9] Velnacrine 225 mg/d vs Placebo change from baseline							

EvTable80. Study results: HP 128 (Velnacrine).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 10 d	
Huff, 1991	OC Population	ADAS					4] 0.3	5] NS
	1] Placebo	MAE					4] 2.6	5] NS
	2] HP 128 (Velnacrine) 100 mg bid	Controlled oral word association						
	3] Placebo mean change from baseline	Visual naming					4] 3.0	5] NS
	4] HP 128 mean change from baseline	Token test					4] 0.9	5] NS
	5] HP 128 vs. placebo	Sentence repetition test					4] 0.0	5] NS
		RAGS					3] 4.0 4] 3.3	5] NS
		CGI						5] >0.05

EvTable81. Study results: Velnacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 15w	
Zemlan 1996	OC Analysis 1] Placebo 2] Velnacrine 75 mg tid best dose improvement relative to placebo	ADAS-Cog					2] 2.15	2] <0.001
		CGI-Improvement					2] 0.25	2] 0.046
		ADAS-Total					2] 2.36	2] 0.001
		ADAS-Noncog					2] 0.03	2] 0.934
		PGIR					2] 0.24	2] 0.073
		CGI-Severity					2] -0.01	2] 0.776
		PSMS					2] 0.19	2] 0.294
		IADL					2] -0.28	2] 0.453

EvTable82. Adverse Events: Velnacrine.

Adverse events (AE) identified in included studies	Antuono, 1995	Huff, 1991	Zemlan, 1996
Withdrawn (%) due to AE	T: 5 C: 4	T: 8 C: 0	T: 33 C: 22
AE Checklist (Max 5)	3	3	3
None Reported			
Balance			
Accidental Injury			
Dizziness	X	X	
Falls			
Behavioral			
Agitation	X		
Cardiovascular			X
Arrhythmia		X	
Hypotension		X	
Hypertension		X	
Extrapyramidal			
Tremor			
Gastrointestinal		X	X
Abdominal pain			X
Constipation		X	
Diarrhea	X	X	X
Dyspepsia			
Nausea, vomiting	X	X	X
Metabolic/nutritional			
Eating disorder	X		X
Weight Change			
Neurological			
Asthenia			X
Psychiatric			
Anxiety			
Confusion, delirium			
Depression			
Respiratory			X
Cough, cold, infection			
Rhinitis			
Other	X		
Aberrant hematology	X		X
Fatigue, weakness			
Fever, flu, pneumonia			
Headache	X		
Hepatic abnormality	X		X
Muscle/joint disorder			
Pain			
Rash, skin disorder	X		X
Sleep disorder	X		
Urinary disorder			

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable83. Key Characteristics: Various cholinergic neurotransmitter modifying agents.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population (ethnicity, setting, comorbidity)	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Canal 1996	IF	6	Placebo Eptastigmine	NINCDS	AD	Mild-Mod	103	94	67.5y (48-85y) 49%M	20 mg bid (if ≤ 65 kg) 20 mg tid (if >65 kg)	4w	ADL CGI-C CGCI HAM-D IADL LMT Semantic Word Fluency Test Trail Making	No
Imbimbo 1999	IF	8	Placebo Eptastigmine	NINCDS DSM IV	AD	Mod-Modly Sev	491	424	71.0y (52-90y) 37%M 99% White	5 mg tid (start) 4-week step-wise increase to 15 mg tid & 20 mg tid	24w	ADAS-Cog CIBIC+ GDS HAM-D IADL	No
Xu 1995	IS	6	Placebo Huperzine	DSM-III-R	AD	Mild-Sev	103	103	66.0y (54-90y) 55%M Asian	1.6 mg/d	8w	ADL HDS HIS MMS MQ	No
Rockwood 1997 Auxiliary: Rockwood 2000	IF	7	Placebo Linopirdine	DSM-III-R NINCDS	AD	Mild-Mod	382	311	71.6y (NR) 44%M 98% White	30 mg tid	6m	ADAS-Total ADAS-Noncog ADAS-Cog CGI DBDS IADL MMSE PSMS A SKT	No
VanDyck 1997	PI	5	Placebo Linopirdine	NINCDS	AD	Mild-Mod	37	34	68.7y (NR) 54%M	40 mg tid	4w	ADAS-Cog CGIC DBDS	No

EvTable83. Key Characteristics: Various cholinergic neurotransmitter modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population (ethnicity, setting, comorbidity)	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Mohr 1997	NR	5	Placebo Sabeluzole	DSM-III-R NINCDS	AD	Mild- Mod	39	15	68.8y (53-79y) 49%M	10 mg bid	48w	ADAS-Cog CT Scan MMSE COG9 Memory	No
Popa 1994	IS	5	Meclofenoxate (MF) Antagonic Stress	DSM-III-R	AD	Mild- Mod	63	NR	69.7y (65-87y) 52%M	260 mg tid	3m	SAS-G SCAG WAIS WMS	No
Schneider 1994	NR	5	Nicergoline Antagonic Stress	DSM IV ICD-10	AD	Mild- Mod	62	NR	69.8y (65-85y) 53%M	60 mg/d	3m	SAS-G SCAG WAIS WMS	No
Xu 1999	IS	7	Huperzine capsules Huperzine tablets	NINCDS DSM-III-R	AD	Mild- Sev	60	60	72.0y (54-80y) 43%M Asian	400 µg/d	60d	CGI CGI-C GBS-SDS HDS-R IADL Memory Quotient MMSE TESS	No

EvTable84. Study results: Eptastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 29d	
Canal 1996	OC Population 1] Placebo change from baseline 2] Eptastigmine 20 mg BID or TID change from baseline 3] Difference between Placebo and Eptastigmine in change from baseline	ADL					1] 0.35 (0.32)* 2] -0.10 (0.10)*	3] 0.383
		IADL					1] 0.65 (0.40)* 2] -0.39 (0.20)*	3] 0.020
		LMT					1] 0.00 (0.65)* 2] 1.46 (0.43)*	3] 0.157
		SWFT					1] -0.75 (0.54)* 2] 0.65 (0.43)*	3] 0.087
		TMT					1] -4.60 (5.13)* 2] -17.15(4.68)*	3] 0.247
		CGI					1] 0.05 (0.05)* 2] -0.14 (0.05)*	3] 0.258
		CGIC Physician					1] 4.10 (0.07)* 2] 3.68 (0.06)*	3] 0.006
		CGIC Caregiver					1] 3.75 (0.10)* 2] 3.54 (0.08)*	3] 0.180

*SEM

EvTable85. Study results: Eptastigmine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24 w	
Imbimbo 1999	ITT Population							
	1] Placebo	<u>ADAS-Cog</u>					4] 2.62 (7.58) 5] 1.05 (6.79) 6] 0.41 (6.88)	7] 0.04 8] 0.005
	2] Eptastigmine 15 mg TID							
	3] Eptastigmine 20 mg TID	<u>CIBIC+</u>					4] 4.36 (0.89) 5] 4.21 (0.86) 6] 4.03 (0.33)	7] 0.138 8] 0.001
	4] Placebo change from baseline	IADL					4] 1.23 (2.55) 5] 0.83 (2.20) 6] 0.58 (1.78)	7] 0.088 8] 0.005
	5] Eptastigmine 15 mg TID change from baseline							
	6] Eptastigmine 20 mg TID change from baseline							
	7] Eptastigmine 15 mg TID change from placebo							
	8] Eptastigmine 20 mg TID change from placebo							

EvTable86. Study results: Haboyin (Huperzine-A).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 30d		Final: (specify) 60d	
XU 1999	OC Analysis 1] Haboyin (Huperzine-A) 400ug capsules bid 2] Haboyin (Huperzine-A) 400ug tablets bid 3] Capsules difference from baseline 4] Tablets difference from baseline 5] Difference between capsules and tablets	Memory Quotient	1] 40(13) 2] 43(12)	5] >0.05			1] 48(17) 2] 49(16)	3] <0.01 4] <0.01 5] >0.05
		MMSE	1] 13(5) 2] 15(4)	5] >0.05	1] 14(7) 2] 16(5)	3] <0.01 4] <0.01 5] >0.05	1] 18(8) 2] 19(6)	3] <0.01 4] <0.01 5] >0.05
		HDS-R	1] 11(5) 2] 13(5)	5] >0.05	1] 13(6) 2] 15(6)	3] <0.01 4] <0.01 5] >0.05	1] 16(8) 2] 17(7)	3] <0.01 4] <0.01 5] >0.05
		IADL	1] 21(5) 2] 22(6)	5] >0.05	1] 20(5) 2] 21(6)	3] <0.01 4] <0.01 5] >0.05	1] 19(6) 2] 19(6)	3] <0.01 4] <0.01 5] >0.05
		GBS-SDS	1] 52(23) 2] 51(21)	5] >0.05	1] 47(22) 2] 44(22)	3] <0.01 4] <0.01 5] >0.05	1] 40(25) 2] 36(24)	3] <0.01 4] <0.01 5] >0.05

EvTable87. Study results: Linopirdine.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Rockwood 1997	ITT Analysis (Endpoint)	<u>ADAS-cog</u>	1] 20.5 (8.44)* 2] 20.3 (8.58)*				1] 22.5 2] 20.2	3] .001
Rockwood 2000	1] Placebo	<u>CGI</u>	1] 3.92 (0.87)* 2] 3.91 (0.86)*				1] 3.69 (0.72) 2] 3.81 (0.74)	
	2] Linopirdine 30 mg tid	<u>CGI % improved</u>					1] 13% (est) 2] 15% (est)	3] 0.558
	3] Linopirdine vs Placebo	ADAS-noncog	1] 2.4 (2.6)* 2] 2.3 (2.6)*				1] 3.4 (0.9)* 2] 3.0 (0.8)*	3] NS
		ADAS total	1] 22.8 (9.5)* 2] 22.6 (10.0)*				1] 25.4 (2.6)* 2] 23.2 (0.7)*	3] <0.05
		SKT	1] 38.5 (6.6)* 2] 37.8 (6.3)*				1] 40.8 (2.4)* 2] 39.5 (1.5)*	3] NS
		IADL	1] 19.8 (5.1)* 2] 19.1 (5.2)*				1] 21.2 (1.4)* 2] 20.1 (1.0)*	3] NS
		Dementia behavior disturbance scale	1] 14.7 (9.7)* 2] 14.3 (9.6)*				1] 15.7 (1.1)* 2] 14.9 (0.6)*	3] NS
		MMSE	1] 19.6 (4.51) 2] 19.4 (4.05)					

*SEM

EvTable88. Study results: Linopirdine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
VanDyck 1997	OC Analysis	ADAS						3] 0.12
	1] Placebo	DBDS						3] 0.13
	2] Linopirdine 40 mg tid	CGIC						3] 0.07
	3] Linopirdine vs Placebo							

EvTable89. Study results: Sabeluzole.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 48w	
Mohr, 1997	OC Analysis	ADAS-cog	1] 20.4 (2.0)*				1] 27.1 (2.8)*	4] <0.01
	1] Placebo		2] 17.8 (1.8)*				2] 22.3 (3.0)*	5] <0.05
	2] Sabeluzole 5 mg		3] 23.3 (2.8)*				3] 28.4 (3.5)*	6] <0.05
	3] Sabeluzole 10 mg	MMSE	1] 18.5 (1.1)					7] NS
			2] 18.9 (0.9)					
	4] Placebo change from baseline	Cog-9	3] 17.9 (0.6)					
	5] Sabeluzole 5mg change from baseline		1] 11.0 (1.8)				1] 17.3 (2.9)	4] <0.01
	6] Sabeluzole 10mg change from baseline	Memory	2] 7.2 (1.2)				2] 11.3 (2.3)	5] <0.05
3] 13.4 (2.4)			3] 17.9 (2.9)	6] <0.05				
7] Between groups		1] 10.5 (0.6)	1] 11.7 (0.6)	4] <0.05				
		2] 10.7 (0.7)	2] 11.0 (0.8)	5] NS				
		3] 11.5 (0.5)	3] 12.2 (0.5)	6] NS				
				7] NS				

*SEM

EvTable90. Study results: Antagonic Stress - Meclofenoxate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Popa, 1994	OC Population							
	1] Meclofenoxate (MF) 260 mg tid	SCAG	1] 68.3 (7.2) 2] 71.2 (7.7)				1] 52.0 (7.6) 2] 46.1 (6.5)	3] <0.001 4] <0.001 5] <0.001 favors AS
	2] Antagonic Stress (AS)							
	3] MF final v baseline	SASG	1] 67.4 (10.6) 2] 68.4 (8.9)				1] 53.3 (13.1) 2] 47.3 (6.5)	3] <0.001 4] <0.001 5] <0.01 favors AS
	4] AS final v baseline							
	5] AS vs MF final	WMS (MQ)	1] 82.0 (6.5) 2] 81.3 (9.8)				1] 100.0 (11.3) 2] 108.6 (11.4)	3] <0.001 4] <0.001 5] <0.05 favors AS
		WAIS (DI)	1] 13.8 (4.4) 2] 15.4 (4.3)				1] 9.8 (4.1) 2] 6.7 (3.9)	3] <0.001 4] <0.001 5] <0.010 favors AS

EvTable91. Study results: Nicergoline - Antagonic Stress.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Schneider 1994	OC Analysis	SCAG	1] 66.5 (11.5) 2] 71.2 (7.7)				1] 50.5 (8.6) 2] 46.1 (6.5)	3] <0.001 4] <0.001 5] 0.002 favors AS
	1] Nicergoline 20 mg tid							
	2] Antagonic Stress 3 capsules tid	SASG	1] 65.8 (9.5) 2] 68.4 (8.9)				1] 52.1 (9.4) 2] 47.3 (6.5)	3] <0.001 4] <0.001 5] 0.000 favors AS
	3] Nicergoline change from baseline	WAIS Digit symbol	1] 7.8 (1.2) 2] 7.5 (1.6)				1] 9.6 (1.6) 2] 11.5 (2.4)	3] <0.001 4] <0.001 5] 0.000 favors AS
	4] Antagonic Stress change from baseline							
	5] Difference between Nicergoline and Antagonic Stress in change from baseline							

EvTable92. Study results: Huperzine A.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify)		Final] (specify) 8w	
Xu 1995	OC Analysis							
	1] Placebo	MQ	1] 48 (21) 2] 56 (21)	3] >0.05			1] 52 (26) 2] 64 (26)	3] <0.05 4] <0.01 5] <0.01
	2] Huperzine A							
	3] Difference between Placebo and Huperzine A	MMS	1] 14 (5) 2] 16 (5)	3] >0.05			1] 15 (6) 2] 19 (6)	3] <0.01 4] >0.05 5] <0.01
	4] Placebo difference from baseline	HDS	1] 16 (5) 2] 16 (6)	3] >0.05			1] 15 (7) 2] 20 (6)	3] <0.01 4] >0.05 5] <0.01
	5] Huperzine A difference from baseline	ADL	1] 31 (9) 2] 33 (10)	3] >0.05			1] 31.9 (0.7) 2] 29 (9)	3] >0.05 4] >0.05 5] <0.01

EvTable93. Adverse Events: Neurotransmitters - Various Cholinergic neurotransmitter modifying agents.

Adverse events (AE) identified in included studies	EPTASTIGMINE Canal, 1996	EPTASTIGMINE Imbimbo, 1999	HUPERZINE Xu, 1995	HUPERZINE Xu, 1999	LINOPIRDINE Van Dyck, 1997	LINOPIRDINE Rockwood, 1997	SABELUZOLE Mohr, 1997	MECLOFENOXATE ANTAGONIC STRESS Popa, 1994	NICERGOLINE ANTAGONIC STRESS Schneider, 1994	HUPERZINE Xu, 1999
Withdrawn (%) due to AE	T: 4 C: 0	T: 8 C: 7	T: 0 C: 0	T: 0 C: 0	T: 0 C: 0	T: 21 C: 2	T: 0 C: 0	T: NR C: NR	T: NR C: NR	T: NR C: NR
AE Checklist (Max 5)	4	4	3	5	0	3	0	1	0	5
None Reported					x		x	X	X	
Balance				NS						NS
Accidental Injury										
Dizziness			NS							
Falls		x								
Behavioral			NS							
Agitation			NS	NS						NS
Cardiovascular		x								
Arrhythmia	x	x								
Hypotension										
Hypertension		x								
Extrapyramidal										
Tremor										
Gastrointestinal		x								
Abdominal pain		x								
Constipation										
Diarrhea		x	NS							
Dyspepsia										
Nausea, vomiting	x	x	NS	NS						NS
Metabolic/nutritional										
Eating disorder		x	NS	NS						NS
Weight Change										
Neurological										
Asthenia										
Psychiatric										
Anxiety		x								
Confusion, delirium		x								
Depression		x								
Respiratory		x								
Cough, cold, infection										
Rhinitis			NS							
Other		x		NS						NS
Aberrant hematology		x								
Fatigue, weakness										
Fever, flu, pneumonia		x								
Headache		x								
Hepatic abnormality						S				
Muscle/joint disorder		x								
Pain		x								
Rash, skin disorder		x								
Sleep disorder		x	NS	NS						NS
Urinary disorder		x								

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable94. Key characteristics: Haloperidol.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Allain 2000	NR	6	Placebo Tiapride Haloperidol	DSM-III-R	AD	Mild-Mod	306	259	79.6y (55-94y) 36%M	300 mg/d 6 mg/d	21d	CGI Global Improvement MMSE MOSES UKU	No
Auchus 1997	NI	6	Placebo Haloperidol Fluoxetine	NINCDS	AD	Probable	15	12	75.6y (NR) 33%M 100% Community	3 mg/d 20 mg/d	6w	BEHAVE-AD CMAI CSI	No
DeDeyn 1999	PI	7	Placebo Risperidone Haloperidol	DSM IV	PDD VaD Mixed	Severe	344	223	81.0y (median) (56-97y) 44%M 99.9% white 100% Institution	0.25 mg with increments of 0.25 every 4d then: 4 mg/d 4 mg/d	12w	BEHAVE-AD CGI CMAI CMH ECG EPS ESRS FAST Heart Rate Laboratory tests MMSE	VaD vs all

EvTable94. Key characteristics: Haloperidol cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Petrie 1982	PI	6	Placebo Loxapine Haloperidol	DSM III	PDD MID	Mod-Sev	64	37	72.7y (60-95y) 49%M 100% institution	Gradually increased with a fixed- flexible dosage for 4 w 50 mg/d 10 mg/d variable	10w	BPRS CGI CGIC EKG Laboratory tests NOSIE SCAG	No
Teri 2000	NI IS	6	Placebo Haloperidol Trazodone BMT	NINCDS	AD	Probable – Possible	149	91	74.8y (NR) 45%M 86% white	3 mg/d 300 mg/d	16w	ABID ADCS-CGIC BRSD-CERAD Caregiver Burden Screen CMAI IADL MMSE PSM RMBPC SCB	No

EvTable94. Key characteristics: Haloperidol cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
DRUG VS DRUG (NO PLACEBO)													
Carlyle 1993	NR	5	Loxapine Haloperidol	DSM-III- R	PDD AD MID	Mod-Sev	40	31	79.0y (65-91y) 55%M 100% Institution	50 mg tid 10 mg tid	28d	Aggression Chart Blood count Electrolytes ESR Renal & Liver Function Test	No
Coccaro 1990	NI IS	6	Haloperidol Oxazepam Diphenhydramine	DSM III	Dementia	NR	59	52	75.3y (58-99y) 59%M 100% Institution	Started with 0.5 or 1.0 mg/d (haloperidol), 10 or 20mg/d (oxazepam), 25 or 50 mg/d (diphenhydrami- ne) and increased to max doses of 5 mg/d 60 mg/d 200 mg/d respectively	8w	ADAS BPRS CDRS NOSIE PSMS Treatment emergent	No
Chan 2001	NI	6	Haloperidol Risperidone	DSM IV	AD VaD	Severe	58	55	80.5y (≥55y) 28%M 100% Asian	2 mg/d	12w	BEHAVE-AD CMAI CMMSE FAST Simpson-Angus Scale	No

EvTable95. Study results: Haloperidol - Tiapride.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 21 d	
Allain 2000	ITT Endpoint Analysis	MOSES					1] 49%	8] 0.04
	1] Placebo	% responders					2] 63%	9] 0.004
	2] Tiapride 100- 300 mg/d	(% with 25% decrease in irritability/ aggressiven ess					3] 69%	10] 0.38
	3] Haloperidol 2-6 mg/d	subscore)						
	4] Across treatment	MOSES					1] 15.53 (5.25)	5] 0.0009
	5] Tiapride vs Placebo		1] 20.28 (2.85)				2] 13.33 (4.20)	6] 0.008
	6] Haloperidol vs Placebo	Global Improvement very improved	2] 19.90 (2.92)				3] 13.75 (4.59)	7] 0.53
	7] Tiapride vs Haloperidol		3] 20.52 (3.27)				1] 14%	4] NS
	8] Placebo vs Tiapride change from baseline	Global Improvement no change					2] 24%	
	9] Placebo vs Haloperidol change from baseline	CGI					3] 31%	
	10] Triapride vs Haloperidol change from baseline	MMSE					1] 21%	8] 0.03
							2] 12%	9] 0.02
							3] 12%	10] NS
								8] NS
								9] NS
								10] NS

EvTable96. Study results: Haloperidol - Fluoxetine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify)		Final] (specify) 6 w	
Auchus 1996	OC Population 1] Placebo 2] Fluoxetine 20 mg/d 3] Haloperidol 3mg/d 4] Across group treatment effect	CMAI	1] 34.4 (8.2) 2] 33.8 (3.0) 3] 37.4 (4.4)				1] 33.0 (3.5) 2] 35.2 (10.3) 3] 35.0 (11.2)	4] 0.82
		BEHAVE-AD	1] 5.6 (3.4) 2] 7.0 (4.2) 3] 11.8 (4.9)				1] 6.6 (3.5) 2] 8.8 (3.5) 3] 9.2 (7.1)	4] 0.35
		CSI	1] 116.2 (57.0) 2] 160.4 (121.8) 3] 165.4 (50.3)				1] 134.8 (62.1) 2] 143.6 (79.3) 3] 179.4 (91.9)	4] 0.67

EvTable97. Study results: Risperidone - Haloperidol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
DeDeyn 1999	ITT Population	BEHAVE-AD total	1] 16.6 2] 16.5 3] 16.3				1] -4.2 2] -6.6 3] -5.2	4] 0.19 6] 0.01
	1] Placebo							
	2] Haloperidol 1.2mg/d	Behave-AD Aggressiveness	1] 5.0 2] 4.7 3] 5.0				1] -0.8 2] -1.6 3] -1.7	4] 0.004 6] 0.01 7] 0.05 favors Risperidone
	3] Risperidone 1.1 mg/d							
	4] Risperidone vs Placebo							
	5] Risperidone vs Placebo change from baseline	CMAI total aggressive	1] 27.5 2] 26.3 3] 25.6				1] -1.6 2] -3.3 3] -3.9	4] 0.01 7] 0.02
	6] Haloperidol vs Placebo change from baseline	CMAI physical aggressive	1] 19.7 2] 19.3 3] 18.9				1] -0.7 2] -0.3 3] -2.7	4] 0.01 7] 0.01
	7] Risperidone vs Placebo change from baseline	CMAI verbal aggressive	1] 7.7 2] 7.0 3] 6.8				1] -0.8 2] -1.0 3] -1.2	4] 0.01
		CGI						5] <0.05
		MMSE						5] NS
		FAST						5] NS
		Behave-AD % with > 30% reduction from baseline					1] 47% 2] 63% 3] 54%	5] 0.25

EvTable98. Study results: Loxapine - Haloperidol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: 10w	
Petrie, 1982	Efficacy Analysis Population	CGIC (marked or moderate improvement)					1] 9% 2] 35% 3] 32%	
	1] Placebo							
	2] Haloperidol 10 mg/d (max)	BPRS total	1] 46.36 2] 46.35 3] 50.79				1] 48.90 2] 39.60 3] 43.84	4] < 0.05 5] < 0.05 6] <0.05 7] <0.05
	3] Loxapine 50 mg/d (max)	SCAG total	1] 61.0 2] 55.9 3] 62.9				1] 60.9 2] 47.3 3] 54.4	4] < 0.05 5] < 0.05
	4] Haloperidol vs baseline							
	5] Loxapine vs baseline	NOSIE	1] 157.2 2] 184.0 3] 155.0				1] 151.2 2] 192.0 3] 171.4	5] < 0.05
	6] Placebo vs Haloperidol change from baseline							
	7] Placebo vs Loxapine change from baseline							

EvTable99. Study results: Haloperidol - Trazodone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 16w	
Teri, 2000	Efficacy Analysis	ADCS-CGIC %improvement					1] 32% 2] 32% 3] 41% 4] 31%	6] 0.99 7] 0.81 8] 0.65 9] 0.75 10] 0.52 11] 0.86
	1] BMT							
	2] Haloperidol mean dose 1.8 mg/d							
	3] Trazodone mean dose 200 mg/d	BRSD Change score					1] -3.56 (2.85) 2] -5.35 (22.41) 3] -6.95 (20.87) 4] -5.28 (24.36)	5] NS
	4] Placebo							
	5] Group Effect	MMSE Change score					1] -0.05 (2.58) 2] -0.61 (2.69) 3] -1.97 (3.15) 4] -0.28 (3.35)	6] NS 7] NS 8] NS 9] NS 10] <0.05 favours BMT 11] NS
	6] Placebo vs Trazodone							
	7] Placebo vs Haloperidol							
	8] Placebo vs BMT	Lawton-Brody ADL Physical Change score					1] -0.27 (1.96) 2] 2.53 (4.00) 3] 1.62 (2.56) 4] 1.31 (2.47)	6] <0.05 favours placebo 7] <0.05 favours placebo
	9] Traxodon vs Haloperidol							
	10] Trazodone vs BMT	Lawton-Brody ADL Instrumental Change score					1] 0.17 (1.84) 2] 1.79 (3.20) 3] 1.81 (3.32) 4] 0.89 (3.32)	6] <0.05 7] <0.05 favours placebo
	11] Haloperidol vs BMT							

EvTable 99. Study results: Haloperidol - Trazodone cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 16w	
			SCB Subjective					1] -2.95 (7.29) 2] -1.88 (8.89) 3] -1.97 (10.06) 4] -2.58 (9.67)	5] NS
			Screen for Caregiver Burden Subjective					1] -2.95 (7.29) 2] -1.88 (8.89) 3] -1.97 (10.06) 4] -2.58 (9.67)	
			Objective					1] -1.23 (3.32) 2] -0.44 (3.22) 3] -1.14 (4.04) 4] -1.25 (4.02)	

EvTable100. Study results: Loxapine - Haloperidol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 14d		Final: (specify) 28d	
Carlyle 1993	OC Analysis 1] Haloperidol 7.0 mg/d (mean) 2] Loxapine 36.0 mg/d (mean) 3] Difference between Haloperidol and Loxapine	Mean Aggression Score for responders Mean depression score Response rate	1] 6.0 2] 8.6		1] 4.8 2] 6.6	3] NS	1] 2.5 2] 4.2 1] 11/14 2] 14/17	3] NS 3] NS

EvTable101. Study results: Haloperidol – Oxazepam - Diphenhydramine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point:		Final: 8w	
Coccaro 1990	Completers Analysis	CDRS mean score	1] 2.78 2] 2.76 3] 2.66	4] > 0.10				
	1] Haloperidol 5 mg/d (max)							
	2] Oxazepam 60 mg/d (max)	ADAS	1] 11.00 (5.95) 2] 11.50 (4.90) 3] 9.82 (3.68)				1] 8.39 (6.09) 2] 9.12 (4.33) 3] 6.12 (4.78)	4] NS 5] < 0.001 6] NS
	3] Diphenhydramine 200 mg/d (max)	BPRS	1] 6.33 (3.01) 2] 5.81 (2.17) 3] 5.67 (2.72)				1] 4.78 (2.44) 2] 5.50 (2.71) 3] 4.47 (2.85)	4] NS 5] < 0.02 6] NS
	4] Between groups, change from baseline	PSMS	1] 42.17 (12.95) 2] 45.75 (11.02) 3] 39.35 (10.36)				1] 37.89 (15.36) 2] 43.68 (11.47) 3] 34.76 (9.94)	4] NS 5] < 0.001 6] NS
	5] Change from baseline						1] 78.31 (9.45) 2] 80.69 (9.89) 3] 73.00 (11.53)	
	6] Between groups at timepoint	NOSIE	1] 78.19 (7.67) 2] 80.69 (9.10) 3] 73.47 (5.88)					4] NS 5] NS 6] <0.02

EvTable102. Study results: Haloperidol - Risperidone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12 w	
Chan 2001	OC Analysis 1] Haloperidol 0.5-2 mg/d 2] Risperidone 0.5-2 mg/d 3] Haloperidol change from baseline 4] Risperidone change from baseline 5] Between treatments	CMAI BEHAVE-AD (Aggressive- ness) FAST CMMSE	1] 46.4 (10.5) 2] 48.9 (14.5) 1] 2.1 (2.0) 2] 2.2 (2.5) 1] 8.2 (5.0) 2] 7.9 (6.0)				1] 36.3 (10.4) 2] 40.8 (16.9) 1] 0.8 (1.5) 2] 0.9 (2.0) No data extracted 3] -0.15 4] -0.42	3] 0.000 4] 0.002 5] 0.95 3] 0.011 4] 0.019 5] 0.56 3] 0.84 4] 0.70

EvTable103. Adverse Events: Haloperidol.

Adverse events (AE) identified in included studies	TIAPRIDE (T1) HALOPERIDOL (T2) PLACEBO (C) Allain, 2000	HALOPERIDOL(T1) FLUOXETINE (T2) PLACEBO (C) Auchus, 1997	HALOPERIDOL (T1) RISPERIDONE (T2) PLACEBO (C) De Deyn, 1999	HALOPERIDOL (T) LOXAPINE (T2) PLACEBO (C) Petrie, 1982	HALOPERIDOL (T1) TRAZODONE (T2) PLACEBO (C) Teri, 2000
Withdrawn (%) due to AE	T1: 5 T2: 17 C: 6	T1: 33 T2: 0 C: 17	T1: NR T2: NR C: NR	T1: 18 T2: 21 C: 5	T1: NR T2: NR C: NR
AE Checklist (Max 5)	3	3	1	5	2
None Reported					
Balance		x			S*
Accidental Injury			NS*		
Dizziness					NS*
Falls			NS*		
Behavioral	x			NS*	
Agitation			NS*		
Cardiovascular	x			x	
Arrhythmia					
Hypotension	x			x	
Hypertension	x				
Extrapyramidal	S*			x	S*
Tremor	x	x			NS*
Gastrointestinal				x	
Abdominal pain					
Constipation	x				
Diarrhea	x				
Dyspepsia					
Nausea, vomiting	x				
Metabolic/nutritional					
Eating disorder					
Weight Change					
Neurological				x	
Asthenia	x				
Psychiatric					
Anxiety	x	x			
Confusion, delirium		x			
Depression		x			
Respiratory					
Cough, cold, infection					
Rhinitis					
Other	NS*			S*	NS*
Aberrant hematology					
Fatigue, weakness					NS*
Fever, flu, pneumonia					
Headache					
Hepatic abnormality					
Muscle/joint disorder					
Pain					
Rash, skin disorder					
Sleep disorder	x		NS*		
Urinary disorder	x				

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable103. Adverse Events: Haloperidol (Drug vs Drug Trials) cont'd

Adverse events (AE) identified in included studies	HALOPERIDOL (T) RISPERIDONE (C) Chan, 2001	HALOPERIDOL(T) LOXAPINE (C) Carlyle, 1993	DIPHENHYDRAMINE (T1) HALOPERIDOL (T2) OXAZEPAM (T3) Coccaro, 1990
Withdrawn (%) due to AE	T: 4 C: 7	T: 20 C: 15	T1: 5 T2: 10 T3: 11
AE Checklist (Max 5)	1	5	3
None Reported			
Balance			
Accidental Injury			
Dizziness			
Falls			
Behavioral			
Agitation		x	
Cardiovascular			
Arrhythmia			
Hypotension	NS*	x	
Hypertension			
Extrapyramidal	S*	x	x
Tremor			
Gastrointestinal			
Abdominal pain			
Constipation	x		
Diarrhea			
Dyspepsia			
Nausea, vomiting	x		
Metabolic/nutritional			
Eating disorder			
Weight Change			
Neurological			
Asthenia			
Psychiatric			
Anxiety			
Confusion, delirium		x	
Depression			
Respiratory			
Cough, cold, infection			
Rhinitis			
Other		x	x
Aberrant hematology			
Fatigue, weakness			
Fever, flu, pneumonia			
Headache			
Hepatic abnormality			
Muscle/joint disorder			
Pain			
Rash, skin disorder			
Sleep disorder	x		
Urinary disorder			

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable104. Key characteristics: Memantine.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Orgogozo 2002	IF	6	Placebo Memantine	NINDS- AIREN	VaD	Mild-Mod	321	234	76.4y (NR) 53%M 100% white	5 mg/d on w 1 10 mg/d on w 2 15 mg/d on w 3 then 20 mg/d	28w	ADAS-cog CGIC CIBIC-plus ECG GBC GBS MIS MMSE NOSGER Laboratory Tests	No
Wilcock 2002	IS	7	Placebo Memantine	DSM-III-R NINDS- AIREN	VaD	Mild-Mod	579	464	77.4y (54-97y) 51%M 100% Community	5 mg/d increasing of 5 mg/d each week for 4 w then 20 mg/d	28w	ADAS-Cog CIBIC+ CGI-C CGI-S GBS MMSE NOSGER	MMSE Type of VaD Gender
Winblad 1999	NR	6	Placebo Memantine	DSM-III-R	DAT VaD PDD	Modly Sev- Severe	166	151	71.2y (60-80y) 42%M 100% Institution	5 mg/d on w 1 then 10 mg/d	12w	BGP CGI-C CGI-S CT Scan Ferm's D-test GDS HAM-D HIS MMSE	AD/VaD Care Dependence

EvTable105. Study results: Memantine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 28w	
Orgogozo 2002	ITT/OC Analysis							
	1] Placebo change from baseline	<u>ADAS-Cog</u>					1] 1.58 (6.42) 2] -1.25 (1.39)	3] 0.0016
	2] Memantine 20mg/d change from baseline	<u>CIBIC-plus</u>					1] 4.11 (1.48) 2] 3.82 (1.39)	3] 0.284
	3] Memantine vs Placebo	MMSE					1] 0.52 (4.07) 2] 1.75 (3.38)	3] 0.0121
	ITT for Primary outcomes	CGI-C Clinician					1] 3.85 (1.19) 2] 3.58 (1.09)	3] 0.0938
	Per protocol For Secondary outcomes	CGI-C Caregiver					1] 3.82 (1.31) 2] 3.52 (1.26)	3] 0.0921
		NOSGER					1] 3.26 (12.95) 2] 2.73 (11.67)	3] 0.8119
		GBS					1] 3.38 (16.34) 2] -0.36(15.38)	3] 0.1194

EvTable106. Study results: Memantine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 28w	
Wilcock 2002	ITT and PP Analysis	<u>ADAS-Cog</u>					3] -2.28 (7.77) 4] -0.53 (7.02)	5] 0.0005
	1] Placebo							
	2] Memantine 20mg/d	<u>CGI-C</u>						5] 0.292
	3] Placebo change from baseline							
	4] Memantine change from baseline	MMSE					3] 0.51 (3.9) 4] 0.24 (3.8)	5] NS
	5] Memantine vs Placebo	NOSGER	1] 67.69 (14.21) 2] 68.90 (15.84)				3] 3.45 (11.08) 4] 2.32 (11.12)	5] 0.22
		GBS	1] 32.15 (14.58) 2] 33.83 (14.26)				3] 2.48 (15.95) 4] 1.65 (12.00)	5] 0.02

EvTable107. Study results: Memantine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 2h		Final: (specify) 24h	
Winblad 1999	ITT Analysis 1] Placebo change from baseline 2] Memantine 10mg/d change from baseline 3] Difference between placebo and memantine in change from baseline	<u>CGI-C</u>			1] 40% 2] 48%		1] 46% 2] 73%	3] <0.001
		<u>BGP</u> (care dependence)					1] -1.1 (11.8) 2] -3.1 (12.2)	3] 0.016
		CGI-S					1] 53% 2] 78%	
		BGP (total)					1] -4.6 (7.0) 2] -7.2 (7.1)	3] 0.015

EvTable108. Adverse Events: Memantine.

Adverse events (AE) identified in included studies	Orgogozo, 2002	Wilcock, 2002	Winblad, 1999
Withdrawn (%) due to AE	T: 12 C: 13	T: 9 C: 7	T: NR C: NR
AE Checklist (Max 5)	4	4	3
None Reported			
Balance			
Accidental Injury		x	
Dizziness	NS	x	
Falls		x	
Behavioral			
Agitation	NS	x	
Cardiovascular	NS		x
Arrhythmia			
Hypotension			
Hypertension			
Extrapyramidal		x	
Tremor			
Gastrointestinal			
Abdominal pain			x
Constipation		x	
Diarrhea		x	
Dyspepsia			
Nausea, vomiting		x	x
Metabolic/nutritional			x
Eating disorder			
Weight Change			
Neurological	NS	x	
Asthenia			
Psychiatric		x	
Anxiety			
Confusion, delirium	NS	x	
Depression			
Respiratory		x	x
Cough, cold, infection		x	
Rhinitis			
Other		x	x
Aberrant hematology			
Fatigue, weakness			
Fever, flu, pneumonia			x
Headache		x	
Hepatic abnormality			
Muscle/joint disorder		x	x
Pain		x	
Rash, skin disorder			
Sleep disorder		x	
Urinary disorder		x	

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable109. Key characteristics: Selegiline.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	#Completing Trial	Mean age (range) % Male (M)	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Agnoli 1991	NR	5	Placebo L-Deprenyl	NINCDS	PDD	Probable Mild-Mod	10	10	68.6y (NR) 40%M	5 mg bid	60d	Cerebral Blood Flow GBS RMT SPECT-Tc-HMPAO TP	No
Burke 1993a Auxiliary: Burke 1993b	PI IS	6	Placebo L-Deprenyl	NINCDS	DAT	Mild	39	33	73.1y (NR) 74%M	5 mg bid	15m	BDS BNT BPRS CDR-SB COWA CS DDS DSCS GERRI MMSE Neuropsychological Battery WAIS-R Block design WAIS-R Digit Span WSM-R	No
Filip 1998	NR	6	Placebo Selegiline (L-Deprenyl)	DSM III	PDD AD	Mild-Mod	173	142	83.0y (≥60y) 29%M 100% Institution	10 mg/d	24w	CGI Clock Drawing Test ECG EEG MMSE NOSIE Sternberg's Memory Scanning	Clock drawing test result

EvTable109. Key characteristics: Selegiline cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M)	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Freedman 1998	IS NI	5	Placebo Deprenyl	NINCDS	AD	Probable	51	41	70.4y (50-80y) 47%M	5 mg/d for 7 d then 10 mg/d	6m	ADAS-Noncog BPRS BSRT GDS COWATT MCPT CSDD MMSE RAGS-E	No
Mangoni 1991 Auxiliary: Smirne 1993	NR	7	Placebo L-Deprenyl	NINCDS DSM III	DAT PDD	Probable Mild-Mod	119	112	68.8y (NR) 38%M	10 mg/d	3m	Blessed-D Digit Span Drawing test IPSC-E Short Story TP Word Fluency WMS	GDS result
Sano 1997 Auxiliary: Thal 1996	NI IS	5	Placebo Vitamin E Selegiline Selegiline + VitaminE	NINCDS	AD	Moderate	341	341	73.4y (NR) 35%M	Vitamin E 1000 IU bid Selegiline 5mg bid	2y	ADAS-Cog Blessed Dementia Scale CDR MMSE Time to end-point (event free survival)	No

EvTable110. Study results: Selegiline.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 60d	
Agnoli 1992	OC Analysis 1] Placebo 2] L-Deprenyl 3] Difference between placebo and L-Deprenyl	RMT A&R	1] 62 (17.2) 2] 54.8 (8.6)	3] <0.05			1] 54 (16.1) 2] 64.2 (15)	3] <0.05
		RMT DR	1] 45.8 (7.3) 2] 36.6 (7.2)	3] NS			1] 40.8 (5.4) 2] 47.2 (8.0)	3] NS
		RMT MI	1] 47 (13.7) 2] 37.2 (6.1)	3] <0.05			1] 39 (12.7) 2] 49 (12.8)	3] <0.05
		TP time	1] 6.7 (2.7) 2] 6.2 (2.6)	3] <0.05			1] 7.9 (1.5) 2] 5.4 (2.7)	3] <0.05
		TP omissions	1] 11.8 (6.7) 2] 8.5 (3.9)	3] <0.05			1] 14 (6.8) 2] 5.2 (2.7)	3] <0.05
		TP errors	1] 10.4 (7.0) 2] 10.2 (14)				1] 5.8 (4.9) 2] 1.0 (1.4)	3] <0.05
		GBS Intellectual function	1] 22.4 (8.4) 2] 20 (8.9)	3] <0.05			1] 23 (11.1) 2] 18 (9.6)	3] <0.05
		GBS verb fluency	1] 8.6 (3.5) 2] 5.5 (2.5)	3] <0.05			1] 6.9 (2.5) 2] 7.1 (3.2)	3] <0.05
		GBS picture copying	1] 16 (2.9) 2] 16.2 (3)	3] <0.05			1] 13 (4.2) 2] 17.2 (3)	3] <0.05

EvTable111. Study results: Selegiline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 8m		Final: (specify) 15m	
Burke 1993a	OC Analysis	CDR	1] 1.0 (0.0) 2] 1.0 (0.0)		1] 1.2 (0.5) 2] 1.3 (0.5)	3] <0.01	1] 1.3 (0.5) 2] 1.3 (0.6)	3] 0.31
Burke 1993	1] Placebo							
	2] L-Deprenyl	CDRS-sum of boxes	1] 5.7 (1.1) 2] 6.2 (1.4)		1] 6.7 (2.3) 2] 7.4 (2.4)	3] <0.01	1] 8.3 (3.2) 2] 7.8 (3.1)	3] NS
	3] Placebo vs. L-Deprenyl vs. baseline	MMSE	1] 19.6 (4.5) 2] 18.8 (5.0)		1] 17.7 (7.8) 2] 16.1 (6.2)	3] <0.01	1] 13.1 (7.4) 2] 15.5 (6.3)	3] 0.25
		BDS	1] 5.4 (2.5) 2] 7.7 (4.0)		1] 6.1 (3.9) 2] 7.5 (4.2)	3] NS	1] 7.7 (3.9) 2] 9.6 (4.8)	3] NS
		GERRI	1] 101.2 (21.4) 2] 110.0 (30.4)		1] 106.9 (23.6) 2] 112.1 (30.7)	3] NS	1] 118 (23.6) 2] 113.9 (33.6)	3] NS
		DSS	1] 1.8 (1.4) 2] 0.8 (1.0)		1] 1.5 (1.7) 2] 0.7 (1.1)	3] NS	1] 0.9 (1.8) 2] 0.5 (0.9)	3] NS
		DSCS	1] 2.1 (1.5) 2] 2.5 (1.6)		1] 2.8 (2.0) 2] 2.6 (1.9)	3] NS	1] 2.1 (2.4) 2] 2.4 (1.6)	3] NS
		CS	1] 4.3 (4.0) 2] 3.2 (2.3)		1] 3.2 (3.1) 2] 3.7 (3.9)	3] NS	1] 4.7 (4.7) 2] 2.4 (1.9)	3] NS
		BPRS	1] 24.4 (2.9) 2] 25.6 (3.5)		1] 24.6 (3.7) 2] 28.4 (9.0)	3] NS	1] 28.2 (6.2) 2] 25.1 (5.9)	3] NS

EvTable112. Study results: Selegiline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12w		Final: (specify) 24 w	
Filip 1999	OC Analysis							
	1] Placebo	SMST-intercept	1] 87.8 (56.4)		1] 82.0 (56.2)		1] 77.8 (54.2)	8] NS
			2] 97.6 (54.8)		2] 67.5 (42.2)		2] 69.2 (51.5)	9] 0.011
	2] Selegiline		3] 98.3 (53.0)		3] 86.7 (64.7)		3] 73.5 (55.7)	
			4] 103.6(53.4)		4] 70.4 (34.9)		4] 71.8 (65.2)	
	3] Placebo (normal subgroup by CDT)		5] 79.6 (58.3)		5] 78.3 (49.3)		5] 81.1 (53.5)	
			6] 94.7 (55.8)		6] 66.1 (45.6)		6] 68.0 (44.3)	
	4] Selegiline (normal subgroup by CDT)	SMST-slope	1] 6.2 (13.8)		1] 4.8 (9.3)		1] 4.6 (8.3)	8] NS
			2] 2.5 (10.8)		2] 7.4 (8.4)		2] 6.9 (9.5)	9] 0.047
	5] Placebo (pathologic subgroup by CDT)		3] 6.3 (12.8)		3] 4.9 (12.7)		3] 5.8 (10.3)	
			4] 4.2 (8.0)		4] 9.8 (10.6)		4] 9.7 (11.6)	
	6] Selegiline (pathologic subgroup by CDT)		5] 6.1 (14.7)		5] 4.7 (5.7)		5] 3.7 (6.4)	
			6] 1.7 (11.9)		6] 6.2 (7.0)		6] 5.6 (8.1)	
	7] Selegiline vs. Placebo (ALL)	SMST-total error	1] 7.4 (5.7)		1] 11.1 (8.7)		1] 11.3 (10.1)	8] NS
			2] 8.3 (6.8)		2] 9.8 (7.8)		2] 9.2 (8.5)	9] 0.029
			3] 6.6 (3.9)		3] 9.8 (6.4)		3] 9.5 (7.1)	
	8] Selegiline vs. Placebo (normal subgroup)		4] 5.8 (5.1)		4] 8.2 (7.8)		4] 5.9 (7.3)	
			5] 8.0 (6.8)		5] 12.1 (10.2)		5] 12.8 (11.8)	
	9] Seileline vs. Placebo (pathologic subgroup)	CDT	6] 9.5 (7.2)		6] 105. (7.7)		6] 10.8 (8.6)	
			1] 3.1 (1.4)				1] 3.0 (1.3)	8] 0.001
			2] 3.2 (1.2)				2] 2.8 (1.4)	9] NS
			3] 1.8 (0.7)				3] 2.7 (1.3)	
			4] 2.1 (0.7)				4] 1.9 (1.1)	
			5] 4.1 (1.0)				5] 3.3 (1.3)	
			6] 3.7 (0.9)				6] 3.3 (1.31)	

EvTable112. Study results: Selegiline cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify) 12w		Final: (specify) 24 w	
			MMSE	1] 3.9 (1.1) 2] 3.4 (1.1) 3] 3.8 (1.1) 4] 3.5 (0.8) 5] 3.9 (1.1) 6] 3.4 (1.3)				1] 3.7 (1.4) 2] 3.9 (1.1) 3] 3.6 (1.4) 4] 4.2 (1.1) 5] 3.7 (1.3) 6] 3.8 (1.1)	7] 0.004 8] 0.041 9] NS
			CGI						7] <0.005

EvTable113. Study results: Selegiline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Freedman 1998	ITT OC Analysis 1] Placebo 2] L-Deprenyl 3] Change from baseline placebo 4] Change from baseline L-Deprenyl 5] Difference between placebo and L-Deprenyl in change from baseline	BPRS	1] 24.0 (3.3) 2] 23.8 (3.5)				1] 25.8 (6.0) 2] 24.8 (4.0) 3] 1.79 (4.5) 4] 1.02 (2.9)	5] 0.6
		MMSE	1] 18.4 (4.4) 2] 17.3 (3.7)				1] 18.5 (6.2) 2] 17.3 (5.1)	5] NS
		GDS	1] 3.9 (0.8) 2] 4.3 (0.8)				1] 4.0 (0.8) 2] 4.4 (0.9)	5] NS
		ADAS-Noncog	1] 3.7 (3.1) 2] 3.4 (2.3)				1] 4.3 (4.0) 2] 2.7 (2.3)	5] NS
		CSDD	1] 3.3 (2.3) 2] 3.1 (1.9)				1] 3.2 (3.0) 2] 2.6 (1.9)	5] NS
		BSRT	1] 24.5 (11.7) 2] 20.4 (9.4)				1] 23.2 (12.6) 2] 20.4 (10.5)	5] NS
		RAGS-E	1] 39.3 (8.8) 2] 38.1 (7.9)				1] 39.0 (11.1) 2] 37.6 (9.6)	5] NS
		COWAT	1] 28.4 (15.3) 2] 26.6 (17.2)				1] 28.0 (18.4) 2] 22.4 (15.8)	5] NS
		MCPT	1] 24.6 (3.0) 2] 23.4 (4.4)				1] 24.6 (1.8) 2] 24.6 (2.0)	5] NS

EvTable114. Study results: Selegeline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 60d		Final: (specify) 90d	
Mangoni 1991	OC Analysis	BDS-1 daily living	1] 8.73 (4.34) 2] 8.76 (3.98)		1] 8.94 (4.57) 2] 7.33 (3.49)		1] 9.18 (4.72) 2] 6.86 (3.43)	4] <0.01
Smirne 1993	1] Placebo	BDS-II	1] 23.57 (8.69) 2] 23.35 (7.40)		1] 20.80 (9.17) 2] 25.52 (7.06)		1] 21.24 (9.21) 2] 26.69 (6.42)	4] <0.01
	2] L-Deprenyl 10 mg/d	Total						
	3] ANOVA within treatment	IPSC-E Psychic Total	1] 104.98 (26.10) 2] 116.32 (32.39)		1] 107.49 (27.64) 2] 101.71 (21.43)		1] 113.80 (31.27) 2] 96.16 (17.81)	4] <0.01
	4] Multivariate ANOVA between treatments including all four test occasions	IPSC-E Somatic Total	1] 24.71 (9.94) 2] 24.68 (9.24)		1] 25.04 (9.75) 2] 23.08 (7.00)		1] 25.37 (9.38) 2] 22.27 (7.36)	4] <0.05
	5] Placebo GDS=3	Digit Span	1] 4.45 (1.35) 2] 4.19 (1.64)		1] 4.24 (1.58) 2] 4.81 (1.23)		1] 3.93 (1.50) 2] 4.90 (1.33)	4] <0.01
	6] Placebo GDS=4	WMS short story – int.	1] 3.73 (3.66) 2] 2.87 (2.65)		1] 2.89 (2.68) 2] 4.60 (3.69)		1] 2.33 (2.88) 2] 4.94 (3.83)	4] <0.01
	7] Placebo GDS=5	WMS short story-del	1] 2.01 (2.77) 2] 3.10 (3.08)		1] 2.49(2.78) 2] 3.30 (3.86)		1] 1.78 (2.45) 2] 3.9 (3.49)	4] <0.01
	8] L-Deprenyl GDS=3	Word fluency	1] 9.41 (5.37) 2] 8.26 (5.38)		1] 7.13 (4.91) 2] 9.75 (5.71)		1] 7.35 (5.84) 2] 10.47 (5.62)	4] <0.01
	9] L-Deprenyl GDS=4	Drawing test	1] 12.33 (5.98) 2] 13.76 (4.97)		1] 11.73 (6.14) 2] 14.78 (4.64)		1] 11.39 (6.02) 2] 15.82 (4.23)	4] <0.05
	10] L-Deprenyl GDS=5							

EvTable114. Study results: Selegiline cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify) 60d		Final: (specify) 90d	
			TPAT	5] 25 (9) 6] 19 (8) 7] 11 (8) 8] 24 (8) 9] 20 (9) 10] 13 (13)		5] 25 (8) 6] 15 (11) 7] 8 (10) 8] 25 (7) 9] 24 (7) 10] 16 (13)		5] 25 (8) 6] 16 (10) 7] 8 (10) 8] 27 (5) 9] 26 (5) 10] 20 (10)	

EvTable115. Study results: Selegiline - Vitamin E.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 10m		Final: (specify) 20m	
Sano 1997	ITT Analysis	<u>Event-free survival</u>			1] 79%		1] 40%	5] 0.077
	1] Placebo				2] 86%		2] 51%	6] 0.087
Thal 1996	2] Vitamin E 1000IU bid				3] 60%		3] 60%	7] 0.21
	3] Selegiline 5mg bid	Event-free survival with MMSE as covariate			4] 80%		4] 49%	
	4] Vitamin E 1000IU bid + Selegiline 5mg bid							5] 0.001
	5] Vitamin E 1000IU bid vs Placebo from baseline							6] 0.012
	6] Selegiline 5mg bid vs Placebo from baseline							7] 0.049
	7] Vitamin E 1000IU bid + Selegiline 5mg bid vs Placebo from baseline							

EvTable116. Adverse Events: Selegiline.

Adverse events (AE) identified in included studies	Agnoli, 1992	Burke, 1993	Filip, 1999	Freedman, 1998	Mangoni, 1991	Sano, 1997
Withdrawn (%) due to AE	T: 0 C: 0	T: 0 C: 0	T: 9 C: 4	T: 0 C: 0	T: 4 C: 2	T: 0 C: 0
AE Checklist (Max 5)	0	3	1	3	3	1
None Reported	X	X				
Balance						S*
Accidental Injury						
Dizziness				X	X	
Falls						S*
Behavioral						
Agitation				X		
Cardiovascular			X			NS*
Arrhythmia				X		
Hypotension						
Hypertension						
Extrapyramidal						NS*
Tremor				X		
Gastrointestinal				X	X	NS*
Abdominal pain						
Constipation						
Diarrhea						
Dyspepsia					X	
Nausea, vomiting					X	
Metabolic/nutritional				X		
Eating disorder						
Weight Change				X		
Neurological			X			NS*
Asthenia						
Psychiatric			X			
Anxiety					X	
Confusion, delirium				X		
Depression						
Respiratory						
Cough, cold, infection						
Rhinitis						
Other			X	X	X	S*
Aberrant hematology						
Fatigue, weakness						
Fever, flu, pneumonia			X			
Headache				X		
Hepatic abnormality						
Muscle/joint disorder						
Pain						
Rash, skin disorder				X		NS*
Sleep disorder				X		
Urinary disorder						

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Completing Trial Number	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Dehlin 1985	NR	6	Placebo Alaproclate	DSM III	PDD MID Mixed	Mild-Sev	43	40	82.0y (65-93y) 44%M 100%Institution	200 mg bid	4w	CPRS GBS	No
Alvarez 2000	PI	5	Placebo Anapsos	NINCDS DSM IV	AD VaD	Mild-Mod	45	42	Mean NR (≥50y) %M NR 100% Community	360 mg/d or 720 mg/d	4w	ADAS-Cog	Disease Severity
Cutler 1993	PI	6	Placebo BMY	DSM-III-R NINCDS	AD	Mild-Mod	69	54	72.0y (54-92y) 41%M	300 mg tid	12w + 4w washout	ADAS CGI CNTB GERRI MMSE WFT	No
Tariot 1998	NI IS	7	Placebo Carbamazepin e	DSM-III-R NINCDS	AD Mixed VaD	Probable	51	47	85.5y (≥60y) 20%M 98% White 100%Institution	100 mg/d (start) increase by 50 mg q2-5d; modal dose: 300 mg/d	6w	BPRS BRSD CGI MMSE Overt Aggression Scale PSMS	No
Olin 2001	NI	5	Placebo Carbamazepin e	NINCDS	AD	Mild-Sev	21	16	74.7y (63-86y) 33%M 71% White 100% Community Agitation	100 mg/d (Day 1-3) 100 mg bid (Day 4-7) 100 mg tid (Day 8-14) 100 mg qid (end)	6w	BPRS CBC/SMAC Levels CGIC Ham-D IADL MMSE PSMS	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Nyth 1990	NR	7	Placebo Citalopram	DSM III	AD, SDAT VaD PDD MID	Mild-Mod	98	61	77.6y (NR) 22%M	20 mg/d for 2 w 30 mg/d for 2 w	4w	CGI GBS MADRS UKU side-effect rating scale Laboratory tests	AD/SDA T vs VaD
Pollock 2002	NI	6	Placebo Citalopram Perphenazine	DSM IV NINCDS	AD VaD MIXED	Probable Possible	85	39	80.6y (NR) 35%M 90% White	10 mg/d (c) or 0.05 mg/kg (p) for 3 d 20 mg/d (c) 0.1 mg/kg/d (p) for 14 d	17d	BPRS Laboratory tests MMSE Neurobehavioural Rating Scale UKU Side effect scale	No
Porsteinsson 2001	IS PI	7	Placebo Divalproex	DSM IV NINCDS	AD VaD MIXED	Probable Possible	56	49	85.0y (>60y) 30%M 100% Institution	375 mg/d + 125 mg/q3d (until side effects)	6w	BPRS BRSD CGI CMAI MMSE Overt Aggression Scale PSM	No
Tariot 2000b	IF	6	Placebo Divalproex	DSM IV	DAT VaD	Probable Possible	173	100	83.4y (68-100y) 35%M 158 White 11 Black 3 Hispanic 100% Institution	125 mg bid + 125 mg/d until 20 mg/kg/d	6w	BPRS BRMS CGI CMAI Laboratory tests MMSE	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Olafsson 1992	NR	6	Placebo Fluvoxamine	DSM III	SDAT MID PDD	NR	46	29	81.0y (median) (65-93y) 41%M 100% Institution	50 mg/d (start) 150 mg/d (end)	6w	GBS Scale Neuropsychological Battery Trail Making Test	No
Reifler 1989	NI IS	6	Placebo Imipramine	DSM III	PDD AD	Mild-Mod	61	57	72.0y (NR) 41%M 100% Community Depression	25 mg/d + 25 mg/w until therapeutic response 83 mg/d (mean)	8w	DRS ECG Ham-D HDS MMSE OARSADL WAIS-R	Depression
Claus 1998	NR	5	Placebo Lisuride	NINCDS	AD	Mild- Modly Sev	22	22	74.1y (NR) 50%M	0.075 mg/d (start) increments of 0.075 mg/w until 0.3 mg/d	8w	CGI CVLT DMSE MMSE	No
Thal 2000b	PI	5	Placebo Lu 25-109	NINCDS	AD	Mild-Mod	496	303	75.5y (47-95y) 42%M 92% White 8% Other Community (100%)	2 weeks dose titration then fixed doses: 25, 50, or 100 mg bid	6m	ADAS-Cog ADCS-CGIC ADCS-ADL BEHAVE-AD	No
Fuchs 1992	NR	5	Placebo Maprotiline	DSM-III-R	PDD	NR	127	94	80.0y (median) (48-96y) 43%M 100% Institution Mild depression	From 25 to 75 mg tid	8w	Blood pressure GDS MMS Std Video rating of global impression	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Passeri 1987	NR	6	Placebo Minaprine	DSM III NINCDS	MID SDAT	Probable	122	122	Mean NR (60-80y) 30%M 100% Institution	100 mg bid	3m	CGI HDRS Neuropsychological Battery Nowlis MRS SHGRS SRT TP	SDA T vs MID
Roth 1996	NR	7	Placebo Moclobemide	DSM III	AD	Mild-Mod	511	NR	73.6y (60-90y) 25%M 22% Community 78% Institution, Depression	400 mg/d	6w	BGP CGAE CGI-T ECG HAM-D MMSE SCAG	No
Moller 2001	NR	7	Placebo Naftidrofuryl	NINDS- AIREN DSM-III-R	VaD MIXED	Mild-Sev	378	278	71.5y (50-85y) 45%M	600 mg/d or 600 mg/d	6m	ADAS-cog CGI CT HIS Laboratory tests MADRS MMSE MRI NOSGER SCAG Trage 8 test kit	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Street 2000 Auxiliary: Clark 2001 Kennedy 2001 Mintzer 2001 Street 2002	IF	8	Placebo Olanzapine	NINCDS	AD	Moderate	206	152	82.8y (61-97y) 39%M 100% Institution	Fixed doses: 5, 10 or 15 mg/d	6w	ADAS-Cog Barnes Akathisia BPRS ECG EPS scales MMSE NPI/NH Simpson-Angus Gait	Psychosis Cognitive impairment level
Amaducci 1988 Auxiliary: SMID 1987	IS	5	Placebo Phosphatidylserine	NINCDS	AD	Mild-Sev	142	115	62.1y (40-80y) 40%M 100% Institution	200mg/d	3m	BDS Block tapping BSR CASE RMT SCT test Self Test TK	Severity of illnesses
Crook 1992a	PI	5	Placebo Phosphatidylserine	NINCDS DSM III	AD PDD	Mild-Mod	51	49	71.0y (55-85y) 31%M 100% Community	100 mg tid	12w	CGI Concern of memory Facial recognition First-last name test Interviewer notices memory loss MMSE Name-Face association Recall Tests Verbal Selective Reminding WAIS	Severity of illnesses

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Magai 2000	PI	7	Placebo Sertraline	DSM IV NINCDS	AD	Late- stage	31	27	89.0y (NR) 0%M 84% White 16% Other 100% Institution Major depression	25 mg/d (week 1-2) 50 mg/d (week 3-4) 100 mg/d (week 5-6)	8w	AFBS CMAI CSDD Facial Behavior GS	No
Lyketsos 2000	NI	6	Placebo Sertraline	NINCDS DSM IV	AD	Mild-Mod	22	16	77.0y (NR) 41%M 77% White 23% Other 100% Community Depression	25 mg/d (start) increased by 50 mg/w to 150/d	13w	ADL CS HAM-D IADL MMSE PDRS	No
Petracca 2001	NR	7	Placebo Fluoxetine	NINCDS DSM IV	AD	Probable	41	35	70.8y (NR) 45%M 76% Major depression 24% minor depression	10 mg/d for w 1 20 mg/d for w 2 30 mg/d for w 3 40 mg/d for w 4 to 6	6w	CGI FIM HAM-A HAM-D MMSE	No
Auchus 1997	NI	6	Placebo Haloperidol Fluoxetine	NINCDS	AD	Probable	15	12	75.6y (NR) 33%M 100% Community	3 mg/d 20 mg/d	6w	BEHAVE-AD CMAI CSI	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Katz 1999 Auxiliary: Jeste 2000 Pryse- Phillips 2000	IF IS	6	Placebo Risperidone	DSM IV	AD VaD Mixed	Mod-Sev	625	435	82.7y (≥55y) 32%M 100% Institution	0.5, 1.0 or 2.0 mg/d	12w	BEHAVE-AD	Gender Age Race
Teri 2000	NI IS	6	Placebo Haloperidol Trazodone BMT	NINCDS	AD	Probable Possible	149	91	74.8y (NR) 45%M 85% White 15% Other Community	Haloperidol: 0.5 mg/d (start) 3 mg/d (end) Trazodone: 50 mg/d (start) 300 mg/d (end)	16w	ABID ADCS-CGIC BRSD-CERAD Caregiver Burden Screen CMAI IADL MMSE PSM RMBPC SCB	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
DeDeyn 1999	PI	7	Placebo Risperidone Haloperidol	DSM IV	PDD VaD Mixed	Severe	344	223	81.0y (median) (56-97y) 44%M	Titration: 0.25 mg q4d up to 1 mg bid, then if no therapeutic effect and no signs of EPS 4 mg/d 4 mg/d	12w	BEHAVE-AD CGI CMAI ECG EPS ESRS FAST Laboratory tests MMSE	No VaD vs ALL
Allain 2000	NR	6	Placebo Tiapride Haloperidol	DSM-III-R	AD	Mild-Mod	306	259	79.6y (55-94y) 36%M 100% White 100% Institution Irritability Aggressiveness	Tiapride: 100 mg/d (Day 1-3) 200 mg/d (Day 4-end) Haloperidol: 2 mg/d (Day 1-3) 4 mg/d (Day 4-end)	21d	CGI Global Improvement MMSE MOSES UKU	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Meehan 2002	IF	6	Placebo Olanzapine Lorazepam	NINCDS DSM IV	AD VaD MIXED	Possible- Probable	272	248	77.6y (54-97y) 39%M 92% White 100% Institution Agitation	Additional injections optional Olanzapine: 12.5 mg/d (max) Loxapine: 2.5 mg/d (max)	24h	ACES BPRS Positive BPRS Total CGI-S CMAI COSTART ECG MMSE Total NPI/NH PANSS-EC Simpson-Angus score	No
Barnes 1982	PI	6	Placebo Thioridazine Loxapine	DSM III	PDD MID	NR	60	34	83.0y (>65y) %M NR 100% Institution Irritability, agitation	Titration: 1 capsule every 2-5 days as needed Thioridazine: 25 mg/d (start) 62.5 mg/d (mean) Loxapine: 5 mg/d (start) 10.5 mg/d (mean)	8w	BPRS CGI NOSIE SCAG	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Pollock 2002	NI	6	Placebo Citalopram Perphenazine	DSM IV NINCDS	AD VaD MIXED	Mod-Sev	85	39	80.6y (NR) 35%M 89% White 100% Institution	Citalopram: 10 mg/d (Day 1-3) 20 mg/d (Day 4-17) Perphenazine: 0.05 mg/kg/d (Day 1-3) 0.1 mg/kg/d (Day 4-17)	17d	BPRS Laboratory tests MMSE Neurobehavioural Rating Scale UKU Side effect scale	No
Bodick 1997 Auxiliary: Veroff 1998 Satlin 1997	IF	6	Placebo Xanomeline	NINCDS	AD	Mild-Mod	343	205	75.0y (60-90y) 43%M 92% White 8% Other 100% Community	75mg/d, or 150 mg/d or 225 mg/d	6m	ADAS-Cog ADSS CIBIC+ CNTB IADL MMSE NOSGER	No
Chan 2001	NI	6	Haloperidol Risperidone	DSM IV	AD VaD	Severe	58	55	80.5y (≥55y) 28%M Community and institution All Chinese	Titration: increases of 0.5 mg/q2d 2 mg/d	12w	BEHAVE-AD CMAI CMMSE FAST Simpson-Angus Scale	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Taragano 1996	NR	7	Fluoxetine Amitriptyline	NINCDS	AD	Probable	37	25	72.1y (NR) 22%M	10 mg/d 25 mg/d	45d	HAM-D MMSE	No
Ancill 1991	IF	6	Lorazepam Alprazolam	DSM-III-R	AD	NR	40	27	78.9y (>65y) 50%M 100% Institution Agitation	0.5 mg tid 0.25 mg tid	28d	AE CGI	No
Karlsson 2000	PI	7	Citalopram Mianserin	DSM-III-R	AD	Mild-Mod	345 53 demented	289 50 demented	75.0y (64-95y) 21%M 58% Community 42% Institution Major depression	Citalopram: 20 mg/d (week 1-4) 40 mg/d (week 5-12) Mianserin: 30 mg/d (week 1-4) 60 mg/d (week 5-12)	12w	CGI GBS MADRS MMSE WHO Well-being	No
Coccaro 1990	NI IS	6	Haloperidol Oxazepam Diphen- hydramine	DSM III	PDD	Mild-Sev	59	52	75.3y (58-99y) 59%M 100% Institution Agitation	5 mg/d 60 mg/d 200 mg/d	8w	ADAS BPRS CDRS NOSIE PSMS	No
Carlyle 1993	NR	5	Loxapine Haloperidol	DSM-III-R	PDD AD MID	Mod-Sev	40	31	79.0y (65-91y) 55%M 100% Institution Aggression	Loxapine: 5 mg bid (start) 50 mg tid (end) Haloperidol: 1 mg bid (start) 10 mg tid (end)	28d	Aggression Chart Blood count Electrolytes ESR Renal & Liver Function Test	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Passeri 1993	NR	5	5'-MTHF Tradozone	DSM III-R	AD MID	Mild- Moderate	96	96	Mean NR (65-94y) 45%M Depression	50 mg/d 100 mg/d	8w	Blood levels HDRS RVM – immediate recall RVM – delayed recall	AD vs MID
Gutzmann 1997	NR	7	Tiapride Melperone	DSM-III-R	Mixed	Mild-Sev	176	156	73.8y (40-100y) 29%M 100% Institution	400 mg/d 100 mg/d	28d	AGGR AIMS BePU (German Test) Laboratory tests CGI CLEX MMSE NOSIE RAPSU (German Test) VAS-ADL	No
Katona 1998	NR	6	Paroxetine Imipramine	DSM-III-R	PDD	Mild-Mod	198	147	76.6y (59-98y) 22%M 99% White 1% Other Depression	Paroxetine: 20 mg/d (week 1-2) 30 mg/d (week 3-4) 40 mg/d (end) Imipramine: 25 mg/d (3d) 50 mg/d (11d) 75 mg/d (2w) 100 mg/d (end)	8w	CGI Cornell Rating Scale GBS MADRS	No

EvTable117. Key Characteristics. Various non-cholinergic neurotransmitter/neuropeptide modifying agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Petrie 1982	PI	6	Placebo Loxapine Haloperidol	DSM III	PDD MID	Mod-Sev	64	37	72.7y (60-95y) 49%M 100% insstitution	Gradually increased with a fixed-flexible dosage for 4 w 50 mg/d 10 mg/d variable	10w	BPRS CGI CGIC EKG Laboratory tests NOSIE SCAG	No

EvTable118. Study results: Alaproclate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Dehlin 1985	Endpoint Analysis		Estimated				Estimated	
	1] Placebo	GBS motor function	1] 13.2 2] 9.8)				1] 14.1 2] 8.9	3] <0.05
	2] Alaproclate 200 mg bid	GBS intellectual function	1] 25.7 2] 21.8				1] 26.4 2] 20.1	3] NS
	3] Alaproclate vs. Placebo	GBS emotional function	1] 6.0 2] 4.8				1] 5.3 2] 3.8	3] NS

EvTable119. Study results: Anapsos.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
Alvarez, 2000	OC Analysis	<u>ADAS-Cog</u>	1] 28.78 (4.06)*				1] 29.20 (3.84)*	2] <0.05 vs. baseline
	1] Placebo total population		2] 36.07 (4.17)*				2] 34.60 (4.48)*	5] <0.05 vs. baseline
	2] Anapsos 360 mg qid total population		3] 34.44 (3.16)*				3] 35.11 (3.45)*	5] <0.01 vs. placebo
	3] Anapsos 720mg qid total population		4] 21.00 (4.09)*				4] 22.51 (4.04)*	8] <0.05 vs. baseline
	4] Placebo mild dementia subpopulation		5] 27.98 (4.53)*				5] 25.42 (4.57)*	8] <0.05 vs. placebo
	5] Anapsos 360 mg qid mild dementia sub-population		6] 28.15 (3.59)*				6] 27.21 (3.60)*	
	6] Anapsos 720 mg qid mild dementia subpopulation		7] 29.34 (7.93)*				7] 30.88 (8.09)*	
	7] Placebo Alzheimer's Disease subpopulation		8] 30.56 (6.08)*				8] 28.04 (6.48)*	
			9] 39.23 (3.19)*				9] 40.31 (3.68)*	
			10] 8.47(4.92)*				10] 28.27 (4.32)*	
			11] 43.42 (4.19)*				11] 43.35 (4.07)*	
			12] 29.64 (5.04)*				12] 29.91(5.38)*	

EvTable119. Study results: Anapsos cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 4w	
		8] Anapsos 360 mg QID Alzheimer's Disease subpopulation							
		9] Anapsos 720 mg QID Alzheimer's Disease subpopulation							
		10] Placebo Vascular Dementia subpopulation							
		11] Anapsos 360 mg QID Vascular Dementia subpopulation							
		12] Anapsos 720 mg QID Vascular Dementia subpopulation							

* SEM

EvTable120. Study results: BMY-21,502.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Cutler, 1993	OC Analysis	MMSE	1] 22.5 2] 23.5	3] >0.05			4] -0.5 5] -1.5	
	1] Placebo							
	2] BMY-21,502 300mg tid	ADAS		3] NS				
	3] BMY-21,502 vs. Placebo							
	4] Placebo change from baseline							
5] BMY-21,502 change from baseline	1] 3.69 2] 3.64	3] >0.05						

EvTable121. Study results: Carbamazepine

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Tariot, 1998	ITT Analysis	<u>CGI improved</u>					1] 21% 2] 77%	3] 0.001
	1] Placebo							
	2] Carbamazepine 304 mg tid variable	<u>BPRS total</u>	1] 53.3 (8.8) 2] 55.1 (9.6)				1] 52.4 (9.8) 2] 47.4 (10.2)	3] 0.0003
	3] Difference in change between Placebo and Carbamazepine	OAS	1] 13.2 (6.3) 2] 15.0 (6.4)				1] 11.3 (7.3) 2] 8.3 (8.0)	3] 0.008
		BRS for Dementia	1] 63.1 (25.8) 2] 77.7 (34.8)				1] 55.0 (29.2) 2] 53.4 (32.0)	3] 0.03

EvTable122. Study results: Carbamazepine

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Olin, 2001	OC Population	<u>BPRS total</u>					1] -4.2 (8.2) 2] -4.0 (7.9)	3] 0.519
	1] Placebo change from baseline	<u>CGIC % improved or no change</u>					1] 58% 2] 89%	3] 0.055
	2] Carbamazepine 400 mg/d variable change from baseline	Ham-D					1] -1.4 (3.3) 2] -4.2 (4.3)	3] 0.150
	3] Difference between Placebo and Carbamazepine change from baseline	PSMRS					1] -0.6 (1.6) 2] -0.5 (1.6)	3] 1.00
		IADL					1] 0.3 (2.2) 2] 0.7 (2.2)	3] 0.408
		MMSE					1] -0.5 (2.9) 2] -0.1 (2.7)	3] 0.644

EvTable123 Study results: Citalopram.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
Nyth, 1990	OC Analysis 1] Placebo average rating scores AD/SDAT 2] Citalopram 30 mg/d variable average rating scores AD/SDAT 3] Improvement between group differences AD/SDAT 4] Placebo vs Citalopram VaD	CGI-SI	1] 3.909 2] 3.897				1] 4.032 2] 3.897	3] 0.284 4] NS
		CGI-ANS	1] 1.795 2] 1.607				1] 1.971 2] 1.643	3] 0.423 4] NS
		GBS-MI	1] 7.813 2] 6.667				1] 7.813 2] 6.667	3] 0.731 4] NS
		GBS-II	1] 20.063 2] 22.666				1] 19.875 2] 21.333	3] 0.321 4] NS
		GBS – EB	1] 4.406 2] 4.555				1] 3.781 2] 3.296	3] 0.384 4] NS
		GBS-Confus.	1] 1.188 2] 1.223				1] 1.063 2] 0.704	3] 0.148 4] NS
		GBS-Irritabiliy	1] 0.969 2] 1.297				1] 0.938 2] 0.667	3] 0.017 4] NS
		GBS-anxiety	1] 0.876 2] 1.408				1] 0.688 2] 0.889	3] 0.276 4] NS
		GBS-restless	1] 0.782 2] 0.888				1] 0.719 2] 0.444	3] 0.081 4] NS
		MADRS Total score	1] 7.690 2] 8.307				1] 7.690 2] 6.115	3] 0.358 4] NS

EvTable124. Study results: Citalopram.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) up to 17d	
Pollock, 2002	ITT Analysis							
	1] Placebo	<u>Neuro- Behavioural Rating Score</u>	1] 58.3 (11.9)				1] 56.0 (15.2)	4] 0.002
	2] Citalopram 20 mg/d		2] 53.5 (10.2)				2] 43.5 (12.1)	5] 0.14
	3] Perphenazine 0.1mg/kg/d		3] 57.1 (14.0)				3] 49.9 (14.2)	
	4] Citalopram vs. placebo							
	5] Perphenazine vs. placebo							

EvTable125. Study results: Divalproex.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Porsteinsson 2001	OC Analysis 1] Placebo 2] Divalproex 826 mg/d variable 3] Difference between Placebo and Divalproex change from baseline	BPRS-total	1] 55.4 (7.4) 2] 543.9 (8.7)				1] 49.5 (10.5) 2] 47.9 (12.4)	3] 0.61
		Overt Aggression Scale	1] 16.9 (9.0) 2] 14.8 (7.6)				1] 12.0 (8.5) 2] 10.0 (8.3)	3] 0.95
		CERAD BRSD Weighted	1] 53.9 (20.9) 2] 48.2 (16.2)				1] 45.3 (26.0) 2] 38.3 (9.9)	3] .73
		CMAI	1] 77.2 (21.1) 2] 77.2 (18.9)				1] 69.9 (22.9) 2] 67.7 (23.3)	3] 0.65
		MMSE	1] 6.7 (6.7) 2] 7.0 (6.6)				1] 5.1 (6.2) 2] 5.2 (6.9)	3] 0.91
		PSMS	1] 14.3 (4.8) 2] 15.4 (4.4)				1] 14.6 (4.7) 2] 15.2 (4.5)	3] 0.41
		CGI % with therapeutic effect					1] 52% 2] 68%	3] 0.07

EvTable125. Study results: Divalproex.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Porsteinsson 2001	OC Analysis 1] Placebo 2] Divalproex 826 mg/d variable 3] Difference between Placebo and Divalproex change from baseline	<u>BPRS-total</u>	1] 55.4 (7.4) 2] 543.9 (8.7)				1] 49.5 (10.5) 2] 47.9 (12.4)	3] 0.61
		Overt Aggression Scale	1] 16.9 (9.0) 2] 14.8 (7.6)				1] 12.0 (8.5) 2] 10.0 (8.3)	3] 0.95
		CERAD BRSD Weighted	1] 53.9 (20.9) 2] 48.2 (16.2)				1] 45.3 (26.0) 2] 38.3 (9.9)	3] .73
		CMAI	1] 77.2 (21.1) 2] 77.2 (18.9)				1] 69.9 (22.9) 2] 67.7 (23.3)	3] 0.65
		MMSE	1] 6.7 (6.7) 2] 7.0 (6.6)				1] 5.1 (6.2) 2] 5.2 (6.9)	3] 0.91
		PSMS	1] 14.3 (4.8) 2] 15.4 (4.4)				1] 14.6 (4.7) 2] 15.2 (4.5)	3] 0.41
		CGI % with therapeutic effect					1] 52% 2] 68%	3] 0.07

EvTable126. Study results: Divalproex.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Tariot 2001b	ITT Analysis							
	1] Placebo	BRMS	1] 17.7 (0.50)* 2] 17.2 (0.48)*				3] -3.9 (0.79)* 4] -3.9 (0.77)*	5] 0.941
	2] Divalproex Sodium 20 mg/kg/d variable	CMAI Total score	1] 81.8 (2.70)* 2] 86.8 (2.63)*				3] -7.3 (2.72)* 4] -14.3 (2.65)*	5] 0.035
	3] Placebo change from baseline	BPRS Total score	1] 41.7 (1.33)* 2] 43.3 (1.29)*				3] -7.1 (1.73)* 4] -8.0 (1.67)*	5] 0.690
	4] Divalproex change from baseline	CGI (Part II)					3] 3.4 (0.14)* 4] 3.9 (0.15)*	5] 0.035 favors placebo
	5] Divalproex vs. Placebo	MMSE	1] 7.7 (0.77)* 2] 7.1 (0.75)*					5] NS

*SEM

EvTable127. Study results: Fluvoxamine.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Olafsson	OC Analysis 1] Placebo 2] Fluvoxamine 150 mg/d 3] Placebo difference from baseline 4] Fluvoxamine difference from baseline 5] difference between changes in Placebo and Fluvoxamine	Trail-making test - Median (range)	1] 2.8 (0-16) 2] 8.3 (0-15)				1] 0.8 (0-8) 2] 5.0 (0-17)	3] NS 4] NS 5] NS
		GBS - Median (range)	1] 63 (22-104) 2] 78 (13-132)				1] 68 (17-102) 2] 68 (19-120)	3] NS 4] NS 5] NS
		<i>Picture recall</i> Immediate recall	1] 0.2 (0-5) 2] 0.4 (0-4)				1] 0.2 (0-4) 2] 0.6 (0-4)	3] NS 4] NS 5] NS
		Delayed recall	1] 0.1 (0-2) 2] 0.2 (0-3)				1] 0.1 (0-4) 2] 0.4 (0-4)	3] NS 4] NS 5] NS
		<i>Picture recognition</i> Correct recognition	1] 2.5 (0-8) 2] 3.7 (1-8)				1] 4.8 (0-8) 2] 4.8 (0-8)	3] NS 4] NS 5] NS
		Concept distracters	1] 1.5 (0-7) 2] 0.8 (0-4)				1] 2.5 (0-7) 2] 1.0 (0-8)	3] NS 4] NS 5] NS
		Other distracters	1] 2.2 (0-8) 2] 1.3 (0-6)				1] 3.0 (0-6) 2] 1.0 (0-8)	3] NS 4] NS 5] NS
		<i>Finger Tapping</i> Dominant index finger	1] 31 (12-59) 2] 20 (0-60)				1] 30 (5-67) 2] 14 (0-64)	3] NS 4] NS 5] NS

EvTable128. Study results: Fluxoxamine cont'd.

REF ID#	Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 6w	
			Non-dominant index finger	1] 25 (0-48) 2] 20 (0-50)				1] 31 (0-51) 2] 16 (0-54)	3] NS 4] NS 5] NS

EvTable127. Study results: Fluvoxamine.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Olafsson	OC Analysis 1] Placebo 2] Fluvoxamine 150 mg/d 3] Placebo difference from baseline 4] Fluvoxamine difference from baseline 5] difference between changes in Placebo and Fluvoxamine	Trail-making test - Median (range)	1] 2.8 (0-16) 2] 8.3 (0-15)				1] 0.8 (0-8) 2] 5.0 (0-17)	3] NS 4] NS 5] NS
		GBS - Median (range)	1] 63 (22-104) 2] 78 (13-132)				1] 68 (17-102) 2] 68 (19-120)	3] NS 4] NS 5] NS
		<i>Picture recall</i> Immediate recall	1] 0.2 (0-5) 2] 0.4 (0-4)				1] 0.2 (0-4) 2] 0.6 (0-4)	3] NS 4] NS 5] NS
		Delayed recall	1] 0.1 (0-2) 2] 0.2 (0-3)				1] 0.1 (0-4) 2] 0.4 (0-4)	3] NS 4] NS 5] NS
		<i>Picture recognition</i> Correct recognition	1] 2.5 (0-8) 2] 3.7 (1-8)				1] 4.8 (0-8) 2] 4.8 (0-8)	3] NS 4] NS 5] NS
		Concept distracters	1] 1.5 (0-7) 2] 0.8 (0-4)				1] 2.5 (0-7) 2] 1.0 (0-8)	3] NS 4] NS 5] NS
		Other distracters	1] 2.2 (0-8) 2] 1.3 (0-6)				1] 3.0 (0-6) 2] 1.0 (0-8)	3] NS 4] NS 5] NS
		<i>Finger Tapping</i> Dominant index finger	1] 31 (12-59) 2] 20 (0-60)				1] 30 (5-67) 2] 14 (0-64)	3] NS 4] NS 5] NS
		Non-dominant index finger	1] 25 (0-48) 2] 20 (0-50)				1] 31 (0-51) 2] 16 (0-54)	3] NS 4] NS 5] NS

EvTable128. Study results: Imipramine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Reifler 1998	OC Analysis	HDS	1] 18.6 (4.0) 2] 19.3 (3.5) 3] 6.8 (2.4) 4] 6.9 (2.9)				1] 10.8 (3.5) 2] 11.5 (3.7) 3] 6.5 (1.8) 4] 7.9 (3.1)	5] NS 6] NS
	1] Placebo Depressed group							
	2] Imaprine 83 mg/d Depressed group	MMSE	1] 18.0 (5.5) 2] 16.9 (4.6) 3] 14.8 (15.1) 4] 13.4 (6.9)				1] 19.3 (6.5) 2] 18.7 (5.4) 3] 15.1 (6.2) 4] 13.1 (7.7)	5] NS 6] NS
	3] Placebo Not Depressed Group	DRS	1] 115.9 (14.3) 2] 111.2 (14.3) 3] 98.6 (24.8) 4] 80.4 (44.6)				1] 117.4 (13.7) 2] 104.3 (20.9) 3] 98.1 (26.4) 4] 72.7 (43.8)	5] <0.01 6] <0.01
	4] Imaprine 83mn/d Not Depressed Group	OARS-ADL	1] 19.6 (3.9) 2] 19.5 (3.6) 3] 19.5 (3.8) 4] 16.7 (5.8)				1] 17.8 (4.1) 2] 18.0 (3.8) 3] 18.3 (3.5) 4] 15.5 (5.4)	5] NS 6] NS
	5] Imipramine vs. Placebo Depressed group							
	6] Imipramine vs. Placebo Not Depressed group							

EvTable129. Study results: Lisuride.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Claus, 1998	OC Analysis 1] Placebo 2] Lisuride 0.3 mg tid 3] Placebo vs. Lisuride in change from baseline	MMSE	1] 18.7 (5.5)* 2] 22.9 (3.8)*				1] 17.1 (5.7)* 2] 22.6 (4.9)*	3] 0.03
		Dementia Mood Assessment Scale	1] 23.3 (4.5)* 2] 19.9 (3.3)*				1] 25.9 (5.2)* 2] 21.6 (3.8)*	3] 0.71
		CGI Improved					1] 8.3% 2] 20%	3] 0.72
		CVLT Total recall	1] 17.6 (2.3)* 2] 23.5 (3.1)*				1] 16.3 (2.8)* 2] 26.6 (3.2)*	3] 0.06
		Verbal Fluency Total correct	1] 12.3 (2.0) 2] 17.3 (2.5)				1] 11.2 (1.9) 2] 20.0 (3.7)	3] 0.10
		Visuospatial Associative Learning	1] 18.4 (4.4) 2] 20.1 (3.8)				1] 11.1 (3.6) 2] 14.7 (2.4)	3] 0.73
		Delayed Matching to sample	1] 12.4 (2.2) 2] 15.9 (2.5)				1] 10.8 (2.0) 2] 16.3 (2.1)	3] 0.59
		Working Memory Test between errors	1] 27.1 (9.8) 2] 49.9 (9.4)				1] 28.9 (9.8) 2] 28.6 (6.6)	3] 0.22
		Working Memory Test within errors	1] 2.1 (0.8) 2] 3.1 (1.2)				1] 1.8 (0.8) 2] 1.6 (0.4)	3] 0.46
		Visual Vigilance Task	1] 18.7 (5.1) 2] 18.7 (2.9)				1] 22.5 (5.5) 2] 11.7 (3.1)	3] 0.09
		Pegboard	1] 104.3 (24.4) 2] 139.7 (8.1)				1] 118.3 (14.9) 2] 136.1 (14.4)	3] 0.28

EvTable130. Study results: Lu25-109.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24w	
Thal, 2000b	ITT Analysis							
	1] Placebo change from baseline	<u>ADAS-Cog</u>					1] 1.16 2] 1.04 3] 0.90 4] 1.90	5] 0.51
	2] Lu25-109 25 mg tid change from baseline	<u>ADCS-CGIC</u>					1] 0.25 2] 0.34 3] 0.22 4] 0.33	5] 0.63
	3] Lu25-109 50 mg tid change from baseline	ADCS-ADL					1] -2.62 2] -2.79 3] -2.40 4] -3.13	5] 0.91
	4] Lu25-109 100 mg tid change from baseline							
	5] General linear ANCOVA vs. baseline	BEHAVE-AD					1] 0.07 2] -0.72 3] -0.15 4] -0.26	5] 0.35

EvTable131. Study results: Maprotiline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 8w	
Fuchs 1993	OC Analysis	<u>Video rating of Global Impression</u>					3] 0.6 4] -0.2	5] 0.60
	1] Placebo							
	2] Maprotiline 75 mg/d variable	MMSE	1] 15.8 2] 15.0				1] 17.5 2] 15.3	5] 0.22
	3] Placebo change from baseline	GDS	1] 8.4 2] 8.2				1] 6.6 2] 5.3	5] 0.09
	4] Maprotiline change from baseline							
	5] Placebo vs. Maprotiline change from baseline							

EvTable132. Study results: Minaprine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 60d		Final: (specify) 90d	
Passeri, 1998	Endpoint Analysis							
	1] Placebo SDAT patients	HDRS	1] 13 (7) 2] 15 (6) 3] 15 (5) 4] 13 (5)		1] 12 (7) 2] 11 (5) 3] 14 (4) 4] 8.8 (4)	5] <0.01 6] <0.01 7] <0.05 8] <0.01	1] 12 (7) 2] 10 (5) 3] 13(5) 4] 8.4 (5)	5] <0.05 6] <0.05 7] <0.01 8] <0.01
	2] Minaprine 50mg bid SDAT patients	SRT Anxiety	1] 5.5 (4.7) 2] 4.8 (3.7) 3] 5.5 (3.1) 4] 4.3 (2.9)		1] 4.8 (3.6) 2] 4.3 (3.7) 3] 4.2 (2.5) 4] 3.0 (1.8)	6] <0.05	1] 4.7 (3.5) 2] 3.6 (3.8) 3] 4.2 (2.4) 4] 2.7 (1.9)	5] <0.05 6] <0.05
	3] Placebo MID patients							
	4] Minaprine 50 mg bid MID patients	SRT Depression	1] 6.9 (5.2) 2] 6.9 (3.2) 3] 7.0 (3.6) 4] 7.6 (3.8)		1] 5.6 (4.4) 2] 4.8 (3.1) 3] 6.8 (4.2) 4] 4.8 (3.1)	5] <0.01 8] <0.01	1] 5.5 (4.7) 2] 4.4 (3.1) 3] 6.4 (4.0) 4] 3.8 (3.8)	6] <0.1 8] <0.01
	5] Minaprine vs. Placebo (SDAT)	SHGRS	1] 35 (8) 2] 34 (10) 3] 42 (11) 4] 38 (11)		1] 34 (8) 2] 33 (10) 3] 41 (11) 4] 35 (10)	6] <0.01	1] 34(8) 2] 32 (10) 3] 40 (12) 4] 34 (10)	6] <0.5 7] <0.01 8] <0.01
	6] Minaprine vs. Placebo (MID)	Nowlis	1] 14 (3.7) 2] 15 (4) 3] 16 (3.7) 4] 14 (3.8)		1] 14 (4) 2] 13 (3.9) 3] 15 (3.4) 4] 12 (2.8)	6] <0.01	1] 14 (4.1) 2] 13 (39) 3] 14 (2.9) 4] 12 (3.4)	5] <0.01 6] <0.01 7] <0.01 8] <0.01
	7] Minaprine vs. baseline (SDAT)							
	8] Minaprine vs. baseline (MID)							

EvTable133. Study results: Moclobemide.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Roth 1996	ITT Analysis	HAM-D					1] 91 2] 12.6	3] 0.001
	1] Placebo Difference from baseline	MMSE					1] 1.9 2] 2.6	3] 0.05
	2] Moclobemide 400 mg/d Difference from baseline	SCAG					1] 14.8 2] 17.3	3] NS
	3] Placebo vs. Moclobemide Change from baseline	BGP					1] 1.2 2] 1.6	3] NS
		CGAE Any improvement					1] 59% 2] 72%	3] <0.001

EvTable134. Study results: Naftidrofuryl.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) Response Rate no deterioration	
Moller 2001	ITT Analysis							
	1] Placebo- response rate	<u>ADAS-cog & SCAG (positive response)</u>					1] 58% 2] 75% 3] 73%	4] 0.005 5] 0.015 6] 0.73
	2] Naftidrofuryl 400 mg/d –	<u>ADAS-Cog</u>	1] 28.6 (7.7) 2] 29.0 (8.5) 3] 29.5 (7.5)				1] 67% 2] 84% 3] 73%	4] 0.003 5] 0.013 6] 0.61
	3] Naftidrofuryl 600 mg/d – response rate							
	4] Placebo vs. Naftidrofuryl 400 mg	<u>SCAG</u>	1] 51.0 (8.3) 2] 52.3 (9.8) 3] 50.7 (8.6)				1] 65% 2] 84% 3] 84%	4] 0.001 5] 0.001 6] 0.97
	5] Placebo vs. Naftidrofuryl 600 mg/d	NOSGER	1] 69.6 (15.4) 2] 69.8 (14.3) 3] 66.2 (15.8)				1] 50% 2] 63% 3] 58%	4] 0.049 5] 0.21 6] 0.49
	6] Naftidrofuryl 400 mg/d vs. 600 mg/d	CGI					1] 55% 2] 73% 3] 66%	4] 0.004 5] 0.10 6] 0.21

EvTable135. Study results: Olanzapine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6w	
Street 2000	ITT Analysis	NPI/NH <u>Core total</u>					1] -3.7 (10.3) 2] -7.6 (7.7) 3] -6.1 (8.2) 4] -4.9 (7.8)	5] <0.001 6] 0.006 7] 0.24
Clarke 2001	1] Placebo change from baseline							
Kennedy 2001	2] Olanzapine 5mg/d change from baseline	BPRS total					1] -1.4 (11.1) 2] -6.8 (8.6) 3] -5.6 (10.0) 4] -4.0 (10.9)	5] 0.005 6] 0.06 7] 0.13
Mitzner 2001	3] Olanzapine 10mg/d change from baseline							
Street 2001	4] Olanzapine 15mg/d change from baseline	ADAS-Cog In sub-group with mild to moderate cognitive impairment					1] 1.38 (6.23) 2] -0.94 (8.10) 3] 4.00 (7.03) 4] 1.83 (8.98)	5] 0.703 6] 0.203 7] 0.695
	5] Olanzapine 5mg/d vs. placebo							
	6] Olanzapine 10mg/d vs. placebo							
	7] Olanzapine 15mg/d vs. placebo							

EvTable136. Study results: Phosphatidylserine (PS).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3m		Final: (specify) 3m & 3m post	
Amaducci 1988	OC Analysis	Set Test	1] 27.57 2] 16.33 3] 26.02 4] 10.75		2] 11 4] 13	5] <0.10 6] NS	1] 23.93 2] 9.22 3] 27.10 4] 15.63	5] <0.01
SMID Group 1987	1] Placebo moderate impairment	BDS Nonpersonal memory	1] 3.00 2] 2.10 3] 3.23 4] 1.25		2] 2.0 4] 2.0	5] <0.10 6] NS	1] 3.11 2] 1.20 3] 2.96 4] 1.25	5] 0.05
	2] Placebo severe impairment							
	3] PS 200 mg/d moderate impairment							
	4] PS 200mg/d severe impairment	BDS part 2	1] 19.40 2] 14.50 3] 19.27 4] 11.50			5] NS 6] NS	1] 18.51 2] 10.40 3] 17.77 4] 14.25	5] 0.005
	5] PS vs Placebo change from baseline severe impairment	Block tapping	1] 1.38 2] 0.00 3] 1.03 4] 0.44			5] < 0.10 6] NS	1] 1.38 2] 0.11 3] 0.89 4] 1.25	5] 0.05
	6] PS vs placebo change from baseline moderate impairment	BDS Personal Memory	2] 5.5 4] 4.0		2] 4.4 4] 4.4	5] NS 6] NS		
		BDS Daily Living Score	2] 15.6 4] 16.4		2] 16.3 4] 16.6	5] NS 6] NS		

EvTable137. Study results: Phosphatidylserine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 9w		Final: (specify) 12w	
Crook, 1992a	ITT Analysis 1] Placebo 2] Phosphatidylserine 100mg tid 3] Phosphatidylserine 100mg tid vs. Placebo	PRS-Concern of Memory F - Value					3] 5.73	3] 0.02
		PRS-Recall of Interviewer and staff F - Value					3] 4.92	3] 0.04
		PRS-Recall of past day F - Value					3] 6.36	3] 0.02
		PRS-Recall of past week F - Value					3] 9.76	3] 0.01
		PRS-Interviewer Notices memory Loss F - Value					3] 13.21	3] 0.00
		MAC-F First-last name test					3] 12.29	3] <0.00
		MAC-F Name-face association						3] <0.05
		Memory for names of familiar persons			3] 6.12	3] <0.02		
		Ability to recall the location of misplaced objects						3] <0.05

EvTable138. Study results: Sertraline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 8w	
Magai, 2000	ITT Analysis 1] Placebo 2] Sertraline 100 mg escalating 3] Difference between Placebo and Sertraline change from baseline	CSDD	1] 6.36 (2.13) 2] 5.76 (1.89)				1] 4.43 (1.95) 2] 3.53 (2.07)	3] NS
		GS	1] 0.93 (0.91) 2] 0.47 (0.87)				1] 0.57 (0.64) 2] 0.06 (0.24)	3] NS
		CMAI	1] 27.71(19.49) 2] 28.05(21.45)				1] 21.57(11.52) 2] 23.24(20.00)	3] NS
		AFBS	1] 2.29 (2.23) 2] 3.88 (2.64)				1] 1.71 (2.23) 2] 3.06 (2.73)	3] NS
		Knit-brow face	1] 4.86 (2.71) 2] 3.94 (3.07)				1] 4.43 (3.63) 2] 1.94 (2.65)	3] <0.1
		Sad face	1] 2.57 (2.27) 2] 2.65 (3.75)				1] 1.43 (1.74) 2] 2.18 (2.35)	3] NS

EvTable139. Study results: Sertraline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6w		Final: (specify) 13w	
Lyketsos 2000	ITT Analysis 1] Placebo change from baseline 2] Sertraline 150 mg (escalating) change from baseline 3] Placebo vs. Sertraline change from baseline	Psychiatrist rating of responders					1] 10% 2] 75%	3] <0.05
		CSDD			1] 2.7 (6.2) 2] -12.5 (7.0)	1] <0.05	1] -2.6 (5.9) 2] -9.7 (8.3)	3] 0.03
		HAM-D			1] -4.4 (4.9) 2] -14.6 (10.0)	1] <0.05 2] <0.05	1] -3.4 (5.5) 2] -8.9 (12.4)	3] 0.20
		PDRS ADL subscale			1] 0.7 (5.2) 2] -0.9 (4.7)		1] 3.4 (3.5) 2] -0.8 (5.2)	3] 0.09
		MMSE			1] 0.3 (1.6) 2] -2.1 (5.0)		1] 0.8 (2.3) 2] -1.2 (4.7)	3] 0.18

EvTable140. Study results: Fluoxetine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify) 3m		Final] (specify) 6w	
Petracca 2001	ITT Analysis							
	1] Placebo	Ham-D	1] 17.2 (3.6) 2] 15.8 (2.1)				1] 10.0 (5.1) 2] 9.4 (5.7)	3] <0.001 4] <0.001 5] NS 6] NS
	2] Fluoxetine 40 mg/d dose escalation	CGI Severity	1] 3.0 (0.8) 2] 2.7 (0.9)				1] 2.4 (0.8) 2] 2.1 (0.7)	3] <0.001 4] <0.001
	3] Placebo change from baseline							
	4] Fluoxetine change from baseline	Ham-A	1] 9.1 (5.8) 2] 8.3 (5.1)				1] 6.2 (4.4) 2] 7.0 (5.6)	3] <0.001 4] <0.001 5] NS 6] NS
	5] Difference between Placebo and Fluoxetine change from baseline	MMSE	1] 23.2 (5.3) 2] 23.2 (4.5)				1] 23.9 (5.9) 2] 23.1 (6.8)	3] NS 4] NS 5] NS 6] NS
	6] Difference between Placebo and Fluoxetine	FIM	1] 64.2 (8.9) 2] 68.5 (3.4)				1] 67.1 (7.3) 2] 69.8 (2.8)	3] <0.01 4] <0.01 5] NS 6] NS

EvTable141. Study results: Haloperidol - Fluoxetine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify)		Final] (specify) 6 w	
Auchus 1996	OC Population 1] Placebo 2] Fluoxetine 20 mg/d 3] Haloperidol 3mg/d 4] Across group treatment effect	CMAI	1] 34.4 (8.2) 2] 33.8 (3.0) 3] 37.4 (4.4)				1] 33.0 (3.5) 2] 35.2 (10.3) 3] 35.0 (11.2)	4] 0.82
		BEHAVE-AD	1] 5.6 (3.4) 2] 7.0 (4.2) 3] 11.8 (4.9)				1] 6.6 (3.5) 2] 8.8 (3.5) 3] 9.2 (7.1)	4] 0.35
		CSI	1] 116.2 (57.0) 2] 160.4 (121.8) 3] 165.4 (50.3)				1] 134.8 (62.1) 2] 143.6 (79.3) 3] 179.4 (91.9)	4] 0.67

EvTable142. Study results: Risperidone.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Katz, 1999	ITT Endpoint Analysis	BEHAVE-AD						
Jeste, 2000	1] Placebo		1] 15.9				8] -4.2 (0.6)*	5] 0.37
Camilleri 2000	2] Risperidone 0.5 mg/d	CMAI Total	2] 15.9				9] -4.8 (0.7)*	6] 0.002
	3] Risperidone 1 mg/d	CMAI Verbal	3] 16.0				10] -6.5 (0.7)*	7] 0.001
	4] Risperidone 2 mg/d		4] 15.4				11] -6.4 (0.6)*	
	5] Risperidone 0.5 mg/d vs Placebo	CMAI Physical aggressive						6] 0.006
	6] Risperidone 1 mg/d vs Placebo	CGIC						7] <0.001
	7] Risperidone 2 mg/d vs Placebo		1] 4.2					
	8] Placebo change from baseline		3] 4.2					
	9] Risperidone 0.5 mg/d change from baseline		4] 4.2					
	10] Risperidone 1 mg/d change from baseline						8] -0.50	
	11] Risperidone 2 mg/d change from baseline						9] -1.25	
							10] -1.80	
							11] -2.30	
							8] -2.1	
							9] -3.0	
							10] -5.4	
							11] -6.4	
							1] 3.7	6] 0.002
							2] 3.3	7] <0.001
							3] 3.2	

*SEM

EvTable143. Study results: Haloperidol - Trazodone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 16w	
Teri, 2000	Efficacy Analysis	ADCS-CGIC %improvement					1] 32% 2] 32% 3] 41% 4] 31%	6] 0.99 7] 0.81 8] 0.65 9] 0.75 10] 0.52 11] 0.86
	1] BMT							
	2] Haloperidol mean dose 1.8 mg/d							
	3] Trazodone mean dose 200 mg/d	BRSD Change score					1] -3.56 (2.85) 2] -5.35 (22.41) 3] -6.95 (20.87) 4] -5.28 (24.36)	5] NS
	4] Placebo							
	5] Group Effect	MMSE Change score					1] -0.05 (2.58) 2] -0.61 (2.69) 3] -1.97 (3.15) 4] -0.28 (3.35)	6] NS 7] NS 8] NS 9] NS 10] <0.05 favours BMT 11] NS
	6] Placebo vs Trazodone							
	7] Placebo vs Haloperidol							
	8] Placebo vs BMT	Lawton-Brody ADL Physical Change score					1] -0.27 (1.96) 2] 2.53 (4.00) 3] 1.62 (2.56) 4] 1.31 (2.47)	6] <0.05 favours placebo 7] <0.05 favours placebo
	9] Traxodon vs Haloperidol							
	10] Trazodone vs BMT	Lawton-Brody ADL Instrumental Change score					1] 0.17 (1.84) 2] 1.79 (3.20) 3] 1.81 (3.32) 4] 0.89 (3.32)	6] <0.05 7] <0.05 favours placebo
	11] Haloperidol vs BMT							

EvTable143. Study results: Haloperidol - Trazodone cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 16w	
			SCB Subjective					1] -2.95 (7.29) 2] -1.88 (8.89) 3] -1.97 (10.06) 4] -2.58 (9.67)	5] NS
			Screen for Caregiver Burden Subjective					1] -2.95 (7.29) 2] -1.88 (8.89) 3] -1.97 (10.06) 4] -2.58 (9.67)	
			Objective					1] -1.23 (3.32) 2] -0.44 (3.22) 3] -1.14 (4.04) 4] -1.25 (4.02)	

EvTable144. Study results: Risperidone - Haloperidol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
DeDeyn 1999	ITT Population	BEHAVE-AD total	1] 16.6 2] 16.5 3] 16.3				1] -4.2 2] -6.6 3] -5.2	4] 0.19 6] 0.01
	2] Haloperidol 1.2mg/d	Behave-AD Aggressiveness	1] 5.0 2] 4.7 3] 5.0				1] -0.8 2] -1.6 3] -1.7	4] 0.004 6] 0.01 7] 0.05 favors Risperidone
	3] Risperidone 1.1 mg/d							
	4] Risperidone vs Placebo							
	5] Risperidone vs Placebo change from baseline	CMAI total aggressive	1] 27.5 2] 26.3 3] 25.6				1] -1.6 2] -3.3 3] -3.9	4] 0.01 7] 0.02
	6] Haloperidol vs Placebo change from baseline	CMAI physical aggressive	1] 19.7 2] 19.3 3] 18.9				1] -0.7 2] -0.3 3] -2.7	4] 0.01 7] 0.01
	7] Risperidone vs Placebo change from baseline	CMAI verbal aggressive	1] 7.7 2] 7.0 3] 6.8				1] -0.8 2] -1.0 3] -1.2	4] 0.01
		CGI						5] <0.05
		MMSE						5] NS
		FAST						5] NS
		Behave-AD % with > 30% reduction from baseline					1] 47% 2] 63% 3] 54%	5] 0.25

EvTable145. Study results: Haloperidol - Tiapride.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 21 d	
Allain 2000	ITT Endpoint Analysis	MOSES					1] 49%	8] 0.04
	1] Placebo	% responders					2] 63%	9] 0.004
	2] Tiapride 100-300 mg/d	(% with 25% decrease in irritability/aggressiveness subscore)					3] 69%	10] 0.38
	3] Haloperidol 2-6 mg/d							
	4] Across treatment	MOSES					1] 15.53 (5.25)	5] 0.0009
	5] Tiapride vs Placebo		1] 20.28 (2.85)				2] 13.33 (4.20)	6] 0.008
	6] Haloperidol vs Placebo	Global Improvement very improved	2] 19.90 (2.92)				3] 13.75 (4.59)	7] 0.53
	7] Tiapride vs Haloperidol		3] 20.52 (3.27)				1] 14%	4] NS
	8] Placebo vs Tiapride change from baseline	Global Improvement no change					2] 24%	
	9] Placebo vs Haloperidol change from baseline	CGI					3] 31%	
							1] 21%	
							2] 12%	
							3] 12%	
								8] 0.03
								9] 0.02
								10] NS
								8] NS
								9] NS
								10] NS
	10] Triapride vs Haloperidol change from baseline	MMSE						

EvTable146. Study results: Olanzapine - Lorazepam.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value		
			Baseline		Mid-Point: (specify) 2h		Final: (specify) 24h			
Meehan, 2002	OC Analysis	PANSS-EC			1] -5.27 (6.87)	5]<0.05	1] -3.81 (6.20)	5]<0.05		
					2] -7.86 (6.05)	6]<0.01	2] -6.44 (6.00)	6]<0.05		
					3] -8.67 (6.97)	7]<0.01	3] -6.29 (6.75)	7] NS		
					4] -8.49 (6.55)		4] -5.75 (5.99)			
		CMAI			1] -2.78 (3.4)	5] NS	1] -2.21 (3.7)	5] NS		
					2] -3.77 (2.93)	6]<0.05	2] -2.82 (3.21)	6] NS		
					3] -3.97 (3.89)	7]<0.05	3] -3.36 (3.92)	7] NS		
					4] -4.18 (3.52)		4] -2.82 (3.08)			
		ACES			1] 1.04 (1.66)	5]<0.05	1] 0.63 (1.14)	5] NS		
					2] 1.80 (1.61)	6]<0.01	2] 0.90 (1.19)	6] <0.01		
					3] 1.88 (1.86)	7]<0.01	3] 1.29 (1.49)	7] <0.05		
					4] 2.19 (1.83)		4] 1.07 (1.12)			
		BPRS Total					1]-10.29(11.72)	5] NS		
							2]-10.51(11.50)	6] NS		
							3]-10.59(11.31)	7] NS		
							4]- 9.12(10.27)			
		BPRS Positive					1] -2.09 (3.80)	5] NS		
							2] -1.72 (3.50)	6] NS		
							3] -1.86 (3.39)	7] NS		
							4] -1.32 (3.32)			
		CGI-S					1] -0.59 (0.92)	5] NS		
							2] -0.38 (0.80)	6] NS		
							3] -0.47 (0.89)	7] NS		
							4] -0.46 (0.80)			
		MMSE Total					1] 0.37 (3.62)	5] NS		
							2] 0.31 (2.29)	6] NS		
							3] 0.10 (3.01)	7] NS		
							4] 0.08 (3.04)			

EvTable147. Study results: Thoridazine - Loxapine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 8w	
Barnes 1981	Endpoint Analysis							
	1] Placebo	BPRS Total					1] 39.74 2] 39.73 3] 36.89	4] <.05 5] <.05 6] <.01 7] NS 8] NS
	2] Thoridazine 62.5mg/d							
	3] Loxapine 10.5mg/d	SCAG					1] 59.24 2] 58.82 3] 53.58	4] <.05 5] <.05 6] <.01 7] NS 8] NS
	4] Improvement from baseline Placebo							
	5] Improve from baseline Thoridazine	CGI- Improvement					1] 3.41 2] 3.18 3] 3.11	7] NS 8] NS
	6] Improve from baseline Loxapine	NOSIE						7] NS 8] NS 9] NS
	7] Thioridazine vs Placebo from baseline							
	8] Loxapine vs Placebo from baseline	CGI Severity					1] 3.17 2] 3.07 3] 3.05	4] NS 5] <0.01 6] <0.01
	9] Loxapine vs Thioridazine from baseline							

EvTable148. Study results: Citalopram.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) up to 17d	
Pollock, 2002	ITT Analysis							
	1] Placebo	<u>Neuro- Behavioural Rating Score</u>	1] 58.3 (11.9)				1] 56.0 (15.2)	4] 0.002
	2] Citalopram 20 mg/d		2] 53.5 (10.2)				2] 43.5 (12.1)	5] 0.14
	3] Perphenazine 0.1mg/kg/d		3] 57.1 (14.0)				3] 49.9 (14.2)	
	4] Citalopram vs. placebo							
	5] Perphenazine vs. placebo							

EvTable149. Study results: Xanomeline Tartrate (XT).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Bodick 1997 Veroff, 1998 Satlin 1997	Endpoint analysis	<u>ADAS-Cog</u>					1] 1.42 2] 1.03 3] 0.38 4] -1.42	5] 0.935 6] 0.367 7] 0.045 8] 0.033
	1] Placebo change from baseline							
	2] XT 25 mg tid change from baseline	<u>CIBIC+</u>					1] 4.33 2] 4.44 3] 4.09 4] 4.00	5] 0.846 6] 0.036 7] 0.022 8] 0.005
	3] XT 50 mg tid change from baseline	ADSS						8] 0.002
	4] XT 75 mg tid change from baseline	NOSGER					1] 1.15 2] 0.06 3] -1.49 4] -1.69	5] 0.457 6] 0.078 7] 0.032 8] 0.018
	5] XT 25 mg tid difference from Placebo	IADL					1] 0.48 2] 0.58 3] -0.29 4] -0.12	5] 0.786 6] 0.088 7] 0.026 8] 0.010
	6] XT 50 mg tid difference from Placebo							
	7] XT 75 mg tid difference from Placebo	CNTB					4] 2.77	7] 0.039
	8] Dose response							

EvTable150. Study results: Haloperidol - Risperidone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12 w	
Chan 2001	OC Analysis							
	1] Haloperidol 0.5-2 mg/d	CMAI	1] 46.4 (10.5) 2] 48.9 (14.5)				1] 36.3 (10.4) 2] 40.8 (16.9)	3] 0.000 4] 0.002 5] 0.95
	2] Risperidone 0.5-2 mg/d	BEHAVE-AD (Aggressive- ness)	1] 2.1 (2.0) 2] 2.2 (2.5)				1] 0.8 (1.5) 2] 0.9 (2.0)	3] 0.011 4] 0.019 5] 0.56
	3] Haloperidol change from baseline	FAST					No data extracted	
	4] Risperidone change from baseline	CMMSE	1] 8.2 (5.0) 2] 7.9 (6.0)				3] -0.15 4] -0.42	3] 0.84 4] 0.70
	5] Between treatments							

EvTable151. Study results: Fluoxetine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 30d		Final: (specify) 45d	
Taragano 1997	OC Analysis 1] Fluoxetine 10 mg/d 2] Amitriptyline 25 mg/d 3] Between treatments	Ham-D	1] 25.3 (3.8) 2] 26.3 (4.0)	3] 0.10	1] 19.3 (3.2) 2] 17.8 (2.5)	3] 0.10	1] 16.7 (2.9) 2] 15.6 (3.2)	3] 0.10
		MMSE	1] 20.0 (3.2) 2] 18.8 (4.2)	3] 0.10			1] 21.4 (2.9) 2] 21.5 (3.5)	3] 0.10

EvTable152. Study results: Lorazepam, Alprazolam.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 28d	
Ancill, 1991	OC Analysis 1] Lorazepam mean 3.1 mg/d 2] Alprazolam mean 1.5 mg/d 3] Lorazepam vs Alprazolam	CGI % improved					1] 29% 2] 42%	3] NS

EvTable153. Study results: Citalopram.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6w		Final: (specify) 12w	
Karlsson, 2000	OC Analysis 1] Citalopram 40 mg variable 2] Mianserin 60 mg variable 3] Citalopram vs. Mianserin from baseline	<u>MADRS</u>	1] 26 2] 27		1] 18 2] 18		1] 15 2] 16	3] >0.7

EvTable154. Study results: Haloperidol – Oxazepam - Diphenhydramine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point:		Final: 8w	
Coccaro 1990	Completers Analysis	CDRS mean score	1] 2.78 2] 2.76 3] 2.66	4] > 0.10				
	1] Haloperidol 5 mg/d (max)							
	2] Oxazepam 60 mg/d (max)	ADAS	1] 11.00 (5.95) 2] 11.50 (4.90) 3] 9.82 (3.68)				1] 8.39 (6.09) 2] 9.12 (4.33) 3] 6.12 (4.78)	4] NS 5] < 0.001 6] NS
	3] Diphenhydramine 200 mg/d (max)	BPRS	1] 6.33 (3.01) 2] 5.81 (2.17) 3] 5.67 (2.72)				1] 4.78 (2.44) 2] 5.50 (2.71) 3] 4.47 (2.85)	4] NS 5] < 0.02 6] NS
	4] Between groups, change from baseline	PSMS	1] 42.17 (12.95) 2] 45.75 (11.02) 3] 39.35 (10.36)				1] 37.89 (15.36) 2] 43.68 (11.47) 3] 34.76 (9.94)	4] NS 5] < 0.001 6] NS
	5] Change from baseline						1] 78.31 (9.45) 2] 80.69 (9.89) 3] 73.00 (11.53)	
	6] Between groups at timepoint	NOSIE	1] 78.19 (7.67) 2] 80.69 (9.10) 3] 73.47 (5.88)					4] NS 5] NS 6] <0.02

EvTable155. Study results: Loxapine - Haloperidol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 14d		Final: (specify) 28d	
Carlyle 1993	OC Analysis 1] Haloperidol 7.0 mg/d (mean) 2] Loxapine 36.0 mg/d (mean) 3] Difference between Haloperidol and Loxapine	Mean Aggression Score for responders	1] 6.0 2] 8.6		1] 4.8 2] 6.6	3] NS	1] 2.5 2] 4.2	3] NS
		Mean depression score Response rate					1] 11/14 2] 14/17	3] NS

EvTable156. Study results: 5'-MTHF - Trazodone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4w		Final: (specify) 8w	
Passeri 1992	OC Analysis	HDRS	1] 23 (5)		1] 20(6)	3] <0.01	1] 18 (6)	3] <0.01
	1] 5'-MTHF 50 mg/d		2] 23 (3)		2] 21 (4)	4] <0.05	2] 19 (5)	4] <0.01
			5] 23 (5)		5] 21 (6)	7] <0.01	5] 18 (6)	7] <0.01
	2] Trazodone 100mg/d		6] 23 (4)		6] 21 (5)	8] <0.01	6] 19 (6)	8] <0.01
			9] 21 (5)		9] 17 (7)		9] 18 (5)	11] <0.01
	3] 5'-MTHF change from baseline		10] 23 (3)		10] 22 (2)		10] 20 (3)	
	4] Trazodone change from baseline	RVM immediate recall	1] 20 (7)				1] 23 (8)	3] <0.01
			2] 22 (9)				2] 22 (9)	7] <0.01
	5] 5'-MTHF subgroup AD		5] 20 (7)				5] 23 (7)	
			6] 22 (9)				6] 22 (8)	
	6] Trazodone subgroup AD		9] 20 (8)				9] 22 (7)	
			10] 20 (8)				10] 22 (11)	
	7] 5'-MTHF change from baseline subgroup AD	RVM delayed recall	1] 2 (2)				1] 3 (2)	
			2] 3 (2)				2] 3 (2)	
			5] 3 (2)				5] 3 (2)	
			6] 3 (2)				6] 3 (2)	
			9] 2 (2)				9] 3 (2)	
	8] Trazodone change from baseline subgroup AD		10] 4 (2)				10] 3 (2)	
	9] 5'-MTHF subgroup MID							
	10] Trazodone subgroup MID							
	11] Trazodone change from baseline subgroup MID							

EvTable157. Study results: Tiapride - Melperone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 28d	
Gutzmann 1997	ITT Analysis 1] Melperone 400 mg/d 2] Tiapride 100 mg/d 3] Melperone vs Tiapride change from baseline	CGI (item 1) Severity of illness % Severely III or markedly III	1] 68% 2] 77%				1] 35.9% 2] 52.6%	
		CGI (Item 2) global change % responders					1] 72.5% 2] 73.4%	3] 0.675
		NOSIE social competence	1] 28.5 2] 29.2				1] 32.0 2] 31.4	
		NOSIE irritability	1] 30.9 2] 32.1				1] 25.7 2] 26.8	
		AGGR	1] 10.6 2] 11.0				1] 5.7 2] 4.8	
		VAS-ADL	1] 31.1 2] 32.5				1] 41.8 2] 39.9	
		VAS- verbal aggression	1] 28.5 2] 30.9				1] 14.7 2] 15.8	
		VAS-aggressive behaviour	1] 26.9 2] 38.4				1] 16.5 2] 18.3	

*SEM

EvTable158. Study results: Imipramine - Paroxetine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 8w	
Katona, 1998	ITT Analysis 1] Paroxetine 50 mg bid variable change from baseline 2] Imipramine 50 mg bid variable change from baseline 3] Difference between Paroxetine group and Imipramine group	<u>MADRS</u>					1] -12.6 (10.0) 2] -11.8 (10.0)	3] >0.368
		<u>CGI severity of Illness</u>					1] -1.3 (1.5) 2] -1.3 (1.5)	3] >0.286
		CGI improvement Score					1] 2.7 (1.5) 2] 2.7 (1.6)	
		CSDD					1] -8.9 (6.7) 2] -7.1 (7.5)	3] <0.049 favours Paroxetine
		GBS total score					1] -11.7 (18.1) 2] -12.0 (19.6)	3] >0.651

EvTable159. Study results: Loxapine - Haloperidol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: 10w	
Petrie, 1982	Efficacy Analysis Population	CGIC (marked or moderate improvement)					1] 9% 2] 35% 3] 32%	
	1] Placebo							
	2] Haloperidol 10 mg/d (max)	BPRS total	1] 46.36 2] 46.35 3] 50.79				1] 48.90 2] 39.60 3] 43.84	4] < 0.05 5] < 0.05 6] <0.05 7] <0.05
	3] Loxapine 50 mg/d (max)	SCAG total	1] 61.0 2] 55.9 3] 62.9				1] 60.9 2] 47.3 3] 54.4	4] < 0.05 5] < 0.05
	4] Haloperidol vs baseline							
	5] Loxapine vs baseline	NOSIE	1] 157.2 2] 184.0 3] 155.0				1] 151.2 2] 192.0 3] 171.4	5] < 0.05
	6] Placebo vs Haloperidol change from baseline							
	7] Placebo vs Loxapine change from baseline							

EvTable160. Adverse events. Various cholinergic neurotransmitter modifying agents.

Adverse events (AE) identified in included studies	ALAPROCLATE Dehlin, 1985	ANAPSOS Alvarez, 2000	BMV Cutler, 1993	CARBAMAZEPINE Olin, 2001	CARBAMAZEPINE Tariot, 1998	CITALOPRAM Nyth, 1990	DIVALPROEX Porsteinsson, 2000	DIVALPROEX Tariot, 2001b	FLUOXETINE Petracca, 2001	FLUOXETINE Taragano, 1997
Withdrawn (%) due to AE	T: 0 C: 13	T: 7 C: 7	T: 24 C: 9	T: 0 C: 25	T: 15 C: 0	T: 24 C: 14	T: 7 C: 14	T: 22 C: 4	T: 6 C: 4	T: 58 C: 22
AE Checklist (Max 5)	2	3	1	4	4	3	5	5	5	3
None Reported			x							
Balance	x	x			x					
Accidental Injury								S		
Dizziness	x	x				X			X	
Falls		x			NS					
Behavioral		x								
Agitation	x						x			
Cardiovascular							x	S		
Arrhythmia		x				NS				
Hypotension	x				S					
Hypertension										
Extrapyramidal					NS					
Tremor									X	
Gastrointestinal					NS			x	X	
Abdominal pain										
Constipation									X	X
Diarrhea	x			x			x			X
Dyspepsia									X	
Nausea, vomiting		x		x			x	NS		X
Metabolic/nutritional								NS		
Eating disorder								NS		
Weight Change								NS		
Neurological							x			
Asthenia										
Psychiatric	x	x				X				
Anxiety										
Confusion, delirium	x				NS		x		X	X
Depression	x					X				
Respiratory							x			
Cough, cold, infection										
Rhinitis										
Other	x					X	S	NS	X	
Aberrant hematology				NS			NS	S		
Fatigue, weakness					NS	X	x			
Fever, flu, pneumonia					NS		x			
Headache										
Hepatic abnormality										
Muscle/joint disorder	x						x			
Pain										
Rash, skin disorder		x			NS	X	x	NS		
Sleep disorder	x				NS	X	x	S		
Urinary disorder					NS		x	NS		

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable160. Adverse events. Various cholinergic neurotransmitter modifying agents cont'd.

Adverse events (AE) identified in included studies	FLUVOXAMINE Olafsson, 1992	IMIPRAMINE Reifler, 1989	LISURIDE Claus, 1998	LU 25-109 Thal, 2000b	MAPROTILINE Fuchs, 1993	MINAPRINE Passeri, 1987	MOCLOBEMIDE Roth, 1996	NAFTIDROFURYL Moller, 2001
Withdrawn (%) due to AE	T: 18 C: 33	T: 15 C: 3	T: 0 C: 0	T: 30 C: 12	T: 2 C: 2	T: 0 C: 0	T: NR C: NR	T: 0 C: 0
AE Checklist (Max 5)	2	2	1	1	3	5	2	2
None Reported								
Balance						x		
Accidental Injury								
Dizziness		NS	x	x	NS		NS	
Falls								
Behavioral	x							
Agitation				x		x	NS	
Cardiovascular		x						x
Arrhythmia							NS	
Hypotension								
Hypertension					NS		NS	
Extrapyramidal								
Tremor								
Gastrointestinal							NS	
Abdominal pain				x			NS	
Constipation					x		NS	
Diarrhea				x			NS	
Dyspepsia						x		
Nausea, vomiting	x			x	x		NS	
Metabolic/nutritional		x		x				
Eating disorder				x				
Weight Change				x	NS		NS	
Neurological								
Asthenia				x				
Psychiatric	x				x			
Anxiety		x				x	x	
Confusion, delirium				x				
Depression	x			x				
Respiratory								
Cough, cold, infection								
Rhinitis								
Other		x		x	x		NS	x
Aberrant hematology								
Fatigue, weakness			x	x	x		NS	
Fever, flu, pneumonia		x						
Headache			x	x		x	x	
Hepatic abnormality								
Muscle/joint disorder							NS	
Pain								
Rash, skin disorder						x		
Sleep disorder	x	NS		x			NS	
Urinary disorder				x	x			

NR = Withdrawals due to AE Not Reported

+ = Dose response effect on AE

x = Reported adverse event/side effect but not tested for significant differences between groups

S or NS = Reported and tested for statistical differences between placebo and treatment group

S* or NS* = Reported and tested for statistical differences between two (three) treatment groups

[] = Symptom NOT reported in the paper

EvTable160. Adverse events. Various cholinergic neurotransmitter modifying agents cont'd.

Adverse events (AE) identified in included studies	OLANZEPINE Street, 2000	PHOSPHATIDYL- SERINE Amaducci, 1988	PHOSPHATIDYL- SERINE Crook, 1992a	RISPERIDONE Katz, 1999	SERTRALINE Magai, 2000	SERTRALINE Lyketsos, 2000	XANOMELINE Bodick, 1997
Withdrawn (%) due to AE	T: 12 C: 4	T: 0 C: 0	T: 0 C: 0	T: 16 ⁺ C: 12	T: 12 C: 14	T: 0 C: 0	T: 40 ⁺ C: 35
AE Checklist (Max 5)	2	3	4	4	5	2	2
None Reported		x	x				
Balance	S*						
Accidental Injury	NS*			X			
Dizziness							
Falls				X			
Behavioral							
Agitation	NS*			X	x	NS	
Cardiovascular							
Arrhythmia							
Hypotension							
Hypertension							
Extrapyramidal				X			
Tremor						NS	
Gastrointestinal						NS	
Abdominal pain							
Constipation							
Diarrhea							
Dyspepsia							S*
Nausea, vomiting							S*
Metabolic/nutritional							S*
Eating disorder	NS*						
Weight Change	NS*						
Neurological							
Asthenia							
Psychiatric							
Anxiety	NS*						
Confusion, delirium					x	x	
Depression							
Respiratory				X			
Cough, cold, infection	NS*			X			
Rhinitis				X			
Other	NS*			X	x		S*
Aberrant hematology							
Fatigue, weakness							
Fever, flu, pneumonia	NS*			X			
Headache							
Hepatic abnormality							
Muscle/joint disorder							
Pain	NS*			X			S*
Rash, skin disorder	NS*			X			
Sleep disorder	S*			X			
Urinary disorder				X			

NR = # Withdrawals due to adverse events Not Reported; + = Dose Effect on Adverse Events

x = Reported adverse event/side effect but not tested for significant differences between groups

S or NS = Reported and tested for statistical differences between placebo and treatment group

S* or NS* = Reported and tested for statistical differences between two (three) treatment groups

[] = Symptom NOT reported in the paper

EvTable160. Adverse events. Various cholinergic neurotransmitter modifying agents cont'd.

Adverse events (AE) identified in included studies	LORAZEPAM (T) ALPRAZOLAM (C) Ancill, 1991	HALOPERIDOL (T1) OXAZEPAM (T2) DIPHENHYD (T3) Coccaro, 1990	HALOPERIDOL (T) LOXAPINE (C) Carlyle, 1993	HALOPERIDOL (T1) LOXAPINE (T2) PLACEBO (C) Patrie, 1982	5'-MTHF (T) TRAZODONE (C) Passeri, 1993	TIAPRIDE (T) MELPERONE (C) Gutzmann, 1997	HALOPERIDOL (T1), BMT (T2), TRAZODONE (T3) Teri, 2000
Withdrawn (%) due to AE	T: 19 C: 11	T1: 10 T2: 11 T3: 5	T: 20 C: 15	T1: 21 T2: 15 C: 5	T: 0 C: 0	T: 11 C: 6	T1: NR T2: NR T3: NR
AE Checklist (Max 5)	4	3	5	5	3	5	2
None Reported							
Balance					x		S*
Accidental Injury							
Dizziness						x	NS*
Falls			x				
Behavioral	NS*			NS*			
Agitation	NS*	x				x	
Cardiovascular				x			
Arrhythmia							
Hypotension	S*		x	x			
Hypertension						x	
Extrapyramidal		x	x	x		x	S*
Tremor			x				NS*
Gastrointestinal				x		x	
Abdominal pain							
Constipation			x				
Diarrhea							
Dyspepsia							
Nausea, vomiting						x	
Metabolic/nutritional						x	
Eating disorder							
Weight Change						x	
Neurological	S*			x		x	
Asthenia							
Psychiatric							
Anxiety							
Confusion, delirium	S*		x				
Depression						x	
Respiratory							
Cough, cold, infection							
Rhinitis							
Other	S*	x	x	S*	x	x	NS*
Aberrant hematology							
Fatigue, weakness						x	NS*
Fever, flu, pneumonia						x	
Headache							
Hepatic abnormality							
Muscle/joint disorder							
Pain							
Rash, skin disorder							
Sleep disorder						x	
Urinary disorder			x			x	

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable160. Adverse events. Various cholinergic neurotransmitter modifying agents cont'd.

Adverse events (AE) identified in included studies	HALOPERIDOL (T1) FLUOXETINE (T2) PLACEBO (C) Auchus, 1997	CITALOPRAM (T) MIANSERIN (C) Karlsson, 2000	PERPHENAZINE CITALOPRAM Pollock, 2002	TIAPRIDE (T1) HALOPERIDOL (T2) PLACEBO (C) Altain, 2000	HALOPERIDOL (T) RISPERIDONE (C) Chan, 2001	HALOPERIDOL (T1) RISPERIDONE (T2) PLACEBO (C) De Devn, 1999	CITALOPRAM (T1) PERPHENAZINE (T2) PLACEBO
Withdrawn (%) due to AE	T1: 33 T2: 0 C: 17	T: 5 C: 9	T:NR C:NR	T1: 5 T2: 17 C: 6	T: 4 C: 7	T1: NR T2: NR C: NR	
AE Checklist (Max 5)	3	3	2	3	1	1	
None Reported							
Balance	x						
Accidental Injury						NS*	
Dizziness		NS*					
Falls						NS*	
Behavioral				x			
Agitation		NS*	S*			NS*	
Cardiovascular				x			
Arrhythmia							
Hypotension				x	NS*		
Hypertension				x			
Extrapyramidal			NS*	S*	S*		
Tremor	x			x			
Gastrointestinal							
Abdominal pain							
Constipation		NS*		x	x		
Diarrhea				x			
Dyspepsia		x					
Nausea, vomiting		NS*		x	x		
Metabolic/nutritional							
Eating disorder							
Weight Change							
Neurological			NS*				
Asthenia				x			
Psychiatric							
Anxiety	x	NS*		x			
Confusion, delirium	x						
Depression	x						
Respiratory							
Cough, cold, infection							
Rhinitis							
Other				NS*			
Aberrant hematology							
Fatigue, weakness		S*					
Fever, flu, pneumonia							
Headache		NS*					
Hepatic abnormality							
Muscle/joint disorder							
Pain		NS*					
Rash, skin disorder							
Sleep disorder		S*		x	x	NS*	
Urinary disorder		x		x			

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable160. Adverse events. Various cholinergic neurotransmitter modifying agents cont'd.

Adverse events (AE) identified in included studies	PAROXETINE (T) IMIPRAMINE (C) Katona, 1998	OLANZEPINE (T) LORAZEPAM (C) Meehan, 2002	THIORIDAZINE (T1) LOXAPINE (T2) PLACEBO (C) Barnes, 1982
Withdrawn (%) due to AE	T: 18 C: 17	T: 0 C: 0	T1: 18 T2: 21 C: 12
AE Checklist (Max 5)	4	3	3
None Reported			
Balance			
Accidental Injury		NS*	
Dizziness			
Falls			
Behavioral			
Agitation			x
Cardiovascular		NS*	x
Arrhythmia		NS*	
Hypotension			x
Hypertension		NS*	x
Extrapyramidal		NS*	x
Tremor			
Gastrointestinal			
Abdominal pain			
Constipation			
Diarrhea			
Dyspepsia			
Nausea, vomiting	x		
Metabolic/nutritional			
Eating disorder			
Weight Change			
Neurological			
Asthenia	x		
Psychiatric	x		
Anxiety			
Confusion, delirium	x		
Depression			
Respiratory	x		
Cough, cold, infection			
Rhinitis			
Other	x		x
Aberrant hematology			
Fatigue, weakness			x
Fever, flu, pneumonia			
Headache		NS*	
Hepatic abnormality			
Muscle/joint disorder			
Pain			
Rash, skin disorder			
Sleep disorder	x	NS*	
Urinary disorder			

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable161. Key characteristics: Cerebrolysin (CERE).

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Bae 2000	IF	7	Placebo Cerebrolysin	NINCDS	AD	Mild-Mod	53	53	71.0y (NR) 34%M	30 ml/d 5 d/w	4w	ADAS-Cog Katz-ADL CGIS/C GDS Lawton-IADL MMSE	No
Panisset 2002	NI	8	Placebo Cerebrolysin	NINCDS	AD	Probable Mild-Mod	192	171	74.1y (59-81y) 42%M 93.5% White	30 ml/d 5 d/w	6m	ADAS-cog CIBIC+ MMSE DAD CORNELL PSMS IADL Behave-AD CDR CMH Trail Making Test	APOE Genotype
Ruether 1994 Auxiliary: Ruether 2000	NR	7	Placebo Cerebrolysin	DSM-III-R	AD	Mild-Mod	120	120	71.5y (NR) 34%M	30 ml 5d/wk	28d	BF-S GDS Ham-D MMSE NAI SCAG Trail Making ZVT-G	No

EvTable161. Key characteristics: Cerebrolysin (CERE) cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Ruether 2001 Auxiliary: Ruether 2002	IF	8	Placebo Cerebrolysin	NINCDS ICD 10	AD	Probable Mild-Mod	149	136	73.0y (50-85y) 42%M	30ml Cere 70ml Saline 5d/w	16w	ADAS-Cog ADAS-Noncog CGI MADR-S NAI SKT	MMSE
Xiao 1999	IF	6	Placebo Cerebrolysin	DSM IV	VaD	Mild-Modly Sev	148	147	69.7y (55-85y) 69%M	30 ml/d 5d/wk	4w	ADL CGI Ham-D MMSE NAI SCAG Trail Making ZVT	No
Xiao 2000	IF	7	Placebo Cerebrolysin	DSM-III-R NINCDS	AD	Mild-Modly Sev	157	155	70.3y (55-85y) 50%M	30 ml 5d/w	4w	ADL CGI HAM-D MMSE NAI SCAG ZVT	No

EvTable162. Study results: Cerebrolysin (CERE).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
Bae, 2000	ITT Analysis							
	1] Placebo	ADAS-Cog	1] 33.51 (13.35) 2] 32.52 (14.65)				3] -0.36 (3.59) 4] -3.23 (4.75)	5] 0.02
	2] CERE 30 ml IV qid	CGIS/C (Improved)					1] 21.1% 2] 61.8%	5] 0.01
	3] Placebo change from baseline	MMSE	1] 14.6 (5.5) 2] 16.3 (4.8)				3] 0.21 4] 1.68	5] 0.04
	4] CERE change from baseline	GDS	1] 7.1 (3.4) 2] 6.1 (2.9)					5] NS
	5] CERE vs Placebo change from baseline	KATZ ADL	1] 9.9 (3.2) 2] 9.1 (3.4)					5] NS
		LAWTON IADL	1] 25.2 (5.9) 25.5 (6.4)					5] NS

EvTable163. Study results: Cerebrolysin (CERE).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3m		Final: (specify) 6m	
Panisset, 2002	OC Analysis				Endpoint 2 months after end of therapy		5 months after end of therapy	
	1] Placebo mean change from baseline	<u>CIBIC+</u>			1] 4.29 (0.11)* 2] 4.08 (0.10)	3] 0.033	1] 4.46 (0.12) 2] 4.42 (0.12)	
	2] CERE 30 ml 5d/w for 4 weeks mean change from baseline	<u>ADAS-Cog</u>	1] 23.63 (1.53) 2] 24.20 (1.68)		1] -0.88 (0.61) 2] 0.04 (0.62)	3] 0.284	1] 1.02 (0.69) 2] 2.83 (0.68)	
		DAD			1] -2.34 (1.4) 2] -1.54 (1.38)	3] 0.680	1] -4.08(1.63) 2] -6.04 (1.60)	
	3] CERE vs. Placebo change from baseline	MMSE	1] 20.93 (0.33) 2] 20.22 (0.34)		1] 0.17 (0.34) 2] -0.06 (0.34)	3] 0.620	1] -0.34 (0.39) 2] -0.93 (0.38)	

EvTable164. Study results: Cerebrolysin (CERE).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
Ruether, 1994 Ruether, 2000	ITT Analysis	<u>ZVT-G</u>	1] 184.4 (39.3) 2] 184.1 (32.4)				1] 185.6 (21.5) 2] 161.5 (22.8)	3] <0.05 4] <0.0001
	1] Placebo							
	2] CERE 30 ml 5d/w for 4 weeks	<u>SCAG</u>	1] 66.9 (7.5) 2] 66.5 (6.5)				1] 65.8 (6.1) 2] 49.8 (5.2)	3] <0.05 4] <0.0001
	3] CERE 30 ml vs baseline	<u>CGI % with improvement</u>					1] 20% 2] 100%	3] 0.0001 4] <0.0001
	4] Cere vs. Placebo change from baseline	<u>t</u>	1] 48.2 (2.7) 2] 48.1 (2.2)				1] 45.6 (3.3) 2] 34.5 (2.1)	3] <0.05 4] NS
		NAI						
		Bf-S	1] 44.7 (3.6) 2] 43.7 (4.3)				1] 41.6 (5.7) 2] 26.9 (6.7)	3] <0.05

EvTable165. Study results: Cerebrolysin (CERE).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 16w Endpoint		Final: (specify) 28w Follow-up	
Ruether 2001	ITT Analysis	<u>CGI</u>	1] 5.16 (0.07) 2] 5.24 (0.07)		3] 4.60 (0.11) 4] 4.18 (0.11)	6] .004	3] 4.86 (0.12) 4] 4.81 (0.12)	6] <0.024
Ruether 2002	1] Placebo							
	2] CERE 30 ml/d 5d/w for 4 weeks (repeat after 8w washout)	<u>ADAS-Cog</u>	1] 30.21 (1.57) 2] 32.01 (1.44)		3] 1.1 (0.59) 4] -2.1(0.69)	5] .001	3] 1.6 (0.59) 4] 0.0 (0.65)	5]<0.025 6] 0.024
		NAI			3] 0.0 (0.21) 4] -0.5 (0.29)	5] 0.07	3] 0.4(0.30) 4] 0.0 (0.32)	5] 0.071
	3] Placebo change from baseline	ADAS- Noncog			3] -0.2 (0.29) 4] -1.2 (0.45)	6] .003	3] 0.9 (0.37) 4] -0.1(0.38)	5]<0.025 6] 0.003
	4] CERE 30 ml/d change from baseline	MADR-S				5] NS		5] 0.102
	5] CERE vs Placebo change from baseline	SKT				5] NS		5] 0.161
	6] Mean treatment difference between CERE and Placebo							

EvTable166. Study results: Cerebrolysin (CERE).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 2w		Final: (specify) 4w	
Xiao, 1999	ITT Analysis 1] Placebo 2] CERE30 ml QID 5d/w 3] CERE vs Placebo change from baseline	MMSE	1] 20.08 (3.58) 2] 19.61 (3.31)		1] 21.01 (4.57) 2] 20.96 (3.74)	3] 0.252	1] 21.80 (4.17) 2] 22.29 (4.19)	3] 0.028
		CGI improved			1] 61% 2] 48%	3] 0.19	1] 71% 2] 73%	3] 0.11
		HAM-D	1] 8.47 (4.83) 2] 9.68 (4.95)		1] 7.33 (4.73) 2] 7.56 (5.24)	3] 0.078	1] 6.33 (3.94) 2] 6.68 (5.86)	3] 0.179
		SCAG	1] 43.55(12.23) 2] 44.06(12.16)		1] 40.20(11.20) 2] 39.74(11.47)	3] 0.359	1] 36.94(11.63) 2] 37.00(12.68)	3] 0.767
		ADL	1] 30.50 (8.25) 2] 30.92 (9.05)		1] 29.40 (8.82) 2] 30.45 (9.82)	3] 0.429	1] 28.34 (8.95) 2] 29.53(10.34)	3] 0.377
		NAI	1] 45.94 (6.39) 2] 46.44 (6.51)		1] 45.60 (6.90) 2] 45.88 (6.58)	3] 0.756	1] 45.36 (6.92) 2] 45.11 (7.09)	3] 0.355
		ZVT-1	1] 182.96 (95.77) 2] 204.69 (122.27)		1] 166.81 (84.30) 2] 165.00 (92.05)	3] 0.125	1] 170.45 (93.54) 2] 159.63 (85.33)	3] 0.017
		ZVT-2	1] 177.88 (107.90) 2] 209.23 (163.85)		1] 166.28 (92.93) 2] 170.74 (114.50)	3] 0.193	1] 170.05 (93.86) 2] 159.00 (95.88)	3] 0.016

EvTable167. Study results: Cerebrolysin (CERE).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
Xiao, 2000	ITT Analysis 1] Placebo IV 2] CERE IV 30 ml qid, 4d/w 3] CERE IV vs Placebo change from baseline	<u>CGI</u> <u>% improved</u>			1] 52% 2] 39%	3] 0.03 favours placebo	1] 60% 2] 72%	3] 0.02
		<u>MMSE</u>	1] 19.3 (3.2) 2] 18.81 (2.32)		1] 20.12(4.16) 2] 20.05(3.37)	3] 0.417	1] 20.57 (4.61) 2] 21.26 (4.08)	3] 0.043
		HAM-D	1] 7.16 (5.02) 2] 7.04 (4.93)		1] 6.26 (4.48) 2] 6.50 (4.63)	3] 0.403	1] 5.56 (4.16) 2] 5.29 (4.97)	3] 0.783
		SCAG	1] 42.07(12.44) 2] 41.91(11.99)		1] 39.73(12.73) 2] 37.87(12.12)	3] 0.025	1] 38.66(12.64) 2] 35.71(13.48)	3] 0.014
		ADL	1] 29.48 (7.59) 2] 29.73 (7.88)		1] 29.30 (7.72) 2] 28.77 (8.24)	3] 0.105	1] 28.40 (7.97) 2] 27.23 (9.17)	3] 0.061
		NAI	1] 46.25 (6.44) 2] 46.62 (7.16)		1] 46.52 (6.36) 2] 45.19 (7.00)	3] 0.007	1] 6.40 (7.02) 2] 43.05 (9.12)	3] 0.003
		ZVT(1)	1] 204.09 (119.73) 2] 217.05 (111.54)		1] 198.72 (144.79) 2] 189.85 (110.92)	3] 0.141	1] 186.08 (124.78) 2] 173.14 (134.02)	3] 0.023
		ZVT(2)	1] 208.50 (139.69) 2] 210.61 (129.57)		1] 196.30 (142.64) 2] 189.29 (109.76)	3] 0.295	1] 196.53 (150.97) 2] 173.05 (119.22)	3] 0.071

EvTable168. Adverse Events: Cerebrolysin (CERE).

Adverse events (AE) identified in included studies	Bae, 2000	Panisset, 2002	Ruether, 1994	Ruether, 2001	Xiao, 1999	Xiao, 2000
Withdrawn (%) due to AE	T: 0 C: 0	T: 1 C: 0	T: 0 C: 0	T: 1 C: 1	T: 0 C: 0	T: NR C: NR
AE Checklist (Max 5)	5	4	5	4	3	2
None Reported	X		X			
Balance				X		
Accidental Injury		NS				
Dizziness		NS			NS	NS
Falls						
Behavioral						
Agitation					NS	
Cardiovascular		NS			NS	NS
Arrhythmia				X		
Hypotension					NS	NS
Hypertension					NS	NS
Extrapyramidal						
Tremor						NS
Gastrointestinal		NS				NS
Abdominal pain						
Constipation						
Diarrhea						
Dyspepsia						
Nausea, vomiting		NS		X		
Metabolic/nutritional						
Eating disorder						
Weight Change		S				
Neurological						
Asthenia		NS				
Psychiatric						
Anxiety		S				
Confusion, delirium						
Depression					NS	
Respiratory						
Cough, cold, infection				X		
Rhinitis				X		
Other		NS		X	NS	NS
Aberrant hematology		NS		NS		
Fatigue, weakness						
Fever, flu, pneumonia				X		
Headache		S		X		
Hepatic abnormality						
Muscle/joint disorder						
Pain		NS				
Rash, skin disorder					NS	
Sleep disorder					NS	NS
Urinary disorder						

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable169. Key characteristics: Estrogens.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Asthana 2001	PI	5	Placebo 17b-estradiol	NINCDS	AD	Probable Mild-Mod	20	20	80.0y (61-90y) 0%M	0.10 mg/d	8w	BMICT Boston Naming Test BPRS BSRT CIBIC DPRS FCMT IADL MMSE OMDR PSMS SCWIT Story Recall Trail-Making Test Treisman Visual Search Visual Paired- Associates	No
Henderson 2000	IF	6	Placebo Conjugated equine estrogens	NINCDS	AD	Probable Mild-Mod	42	36	78.0y (NR) 0%M	1.25 mg/d	16w	ADAS-Cog ADL CGIC CSGDS IADL MADRS Neuropsychological battery	No

EvTable169. Key characteristics: Estrogens cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Kyomen 1999 Auxiliary: Kyomen 2002	PI IS	8	Placebo Conjugated Equine Estrogens	DSM-III-R	Dementia	Mod-Sev	15	14	83.8y (>60y) 13%M 100% Institution	2.5 mg/d	4w	ABSR CSDD DSSS Katz ADL scale OAS (modified) TSI	No
Mulnard 2000	IF	7	Placebo Estrogen	NINCDS	AD	Probable Mild-Mod	120	97	75.0y (56-91y) 0%M	1.25mg/ d	1y	ADAS-Cog ADCS-CGIC ADL ADL-BDRS Blessed-D CDRS Dependency Scale DST EFR HAM-D HDRS MAACLR MMSE NDT Neuropsychological Battery Trail-Making Test A	No

EvTable169. Key characteristics: Estrogens cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Wang 2000	IS PI	7	Placebo Premarin	NINCDS	AD	Mild-Mod	50	47	71.8y (NR) 0%M	1.25 mg/d	12w	BEHAVE-AD CASI (Chinese) CDR CIBIC+ HAM-D HDRS MMSE-CE HARS	No

EvTable170. Study results: Estradiol.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 8w	
Asthana 2001	OC Analysis 1] Placebo	<u>SCWIT</u>					3] 110 (13)* 4] 90 (10)*	5] 0.02
	2] Estradiol (17-β)	<u>BSRT</u>					3] 7.0 (0.8)* 4] 8.2 (0.8)*	5] 0.049
	3] Placebo change over baseline	FCMT					3] 22 (2.5)* 4] 26 (4.0)*	5] 0.03
	4] Estrogen change over baseline	CIBIC						5] 0.30
	5] Placebo vs estrogen in change from baseline	DPRS (mood) Functional Assessment (PSMS, IADL)						5] NS 5] NS

*SEM

EvTable171. Study results: Estrogen.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify) 4w		Final] (specify) 16w	
Henderson 2000	OC Analysis	<u>ADAS-Cog</u>			3] 1.2 (1.5) 4] -0.2 (1.1)	5] >0.1	3] 0.5 (1.7) 4] 1.8 (1.2)	5] >0.1
	1] Placebo							
	2] Estrogen (unopposed conjugated equine, Premarin)	CGIC			3] 3.8 (0.1) 4] 3.9 (0.1)	5] >0.1	3] 4.2 (0.1) 4] 4.2 (0.2)	5] >0.1
	3] Placebo change from baseline	ADL/IADL			3] -1.1 (1.1) 4] 0.3 (0.8)	5] >0.1	3] 2.9 (1.5) 4] 2.9 (1.1)	5] >0.1
	4] Estrogen change from baseline	CSGDS			3] 1.2 (0.8) 4] 0.1 (1.2)	5] >0.1	3] -0.7 (1.2) 4] -1.4 (1.4)	5] >0.1
	5] Placebo vs Estrogen change from baseline	MADRS			3] -2.0 (1.2) 4] -2.6 (1.3)	5] >0.1	3] 1.1 (1.4) 4] 0.2 (1.6)	5] >0.1

EvTable172. Study results: Estrogen.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify)		Final] (specify) 4w	
Kyomen 1999 Kyomen, 2002	OC Analysis							
	1] Placebo	ABSR					3] 4.74 (4.07) 4] 2.05 (1.73)	5] 0.03
	2] Estrogen (conjugated equine)	CSDD					3] -2.17 (4.00) 4]- 3.75 (4.17)	5] 0.387
	3] Placebo change from baseline	KATZ ADL					3] 0.71 (1.71) 4] 0.41 (1.56)	5] 0.677
	4] Estrogen change from baseline	DSSS	1] 31.7 (9.0) 2] 31.6 (8.7)				1] 31.7 (9.0) 2] 19.6 (6.5)	3] <0.05 4] <0.03
	5] Placebo vs Estrogen in change from baseline	TSI					3] 0.86 (3.24) 4] -0.71 (3.18)	5] 0.143

EvTable173. Study results: Estrogen.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12m	
Mulnard 2000	ITT Analysis							
	1] Placebo	ADCS-CGIC					3] 5.0 (1.1) 4] 5.1 (0.9) 5] 5.2 (0.9) 6] 74% 7] 80% 8] 80%	9] 0.46 10] 0.36 11] 0.73 12] 0.73
	2] Estrogen							
	3] Placebo change from baseline	MMSE					3] -3.1(4.1) 4] -2.7(3.5) 5] -2.7(3.9)	9] 0.48 10] 0.64
	4] Estrogen (0.625 mg/d) change from baseline	ADAS-Cog					3] 3.6 (4.7) 4] 6.3 (8.7) 5] 4.8 (5.4)	9] 0.09 10] 0.32
	5] Estrogen (1.25 mg/d) change from baseline	CDRS					3] 0.2 (0.4) 4] 0.4 (0.7) 5] 0.5 (0.6)	9] 0.03 10] 0.01
	6] Placebo (worsened %)	ADL-BDRS part 1					3] 1.2 (1.5) 4] 1.0 (1.2) 5] 1.0 (1.2)	9] 0.80 10] 0.61
	7] Estrogen (0.625mg/d, worsened %)	ADL-BDRS part 2					3] 0.8 (1.6) 4] 1.0 (1.4) 5] 0.92 (1.4)	9] 0.56 10] 0.73
	8] Estrogen(1.25 mg/d, worsened %)	ADL- Dependency					3] 0.4 (1.1) 4] 0.4 (0.8)	9] 0.59 10] 0.21

EvTable173. Study results: Estrogen cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 12m	
		9] Placebo vs estrogen 0.65mg/d change from baseline	Score					5] 0.5 (1.0)	
			Mood scores(HDRS)					3] 0.03 (3.9)	9] 0.69
								4] 0.5 (3.7)	10] 0.69
								5] -1.1 (4.3)	
		10] Placebo vs estrogen 1.25mg/d change from baseline	Memory scores (EFR)					3] -5.7 (22.4)	9] 0.08
								4] -11.1 (15.2)	10] 0.41
								5] -8.2 (13.2)	
		11] Placebo vs estrogen .065mg/d % worse	Memory scores (NDT)					3] -0.9 (3.1)	9] 0.57
								4] -0.9 (3.5)	10] 0.19
								5] -2.1 (2.6)	
		12] Difference between Place and estrogen 1.25mg/d % worse	Attention scores (letter cancellation)					3] -1.3 (5.5)	9] 0.90
								4] -0.6 (8.7)	10] 0.45
								5] -2.3 (6.0)	
			Attention scores Trail-making test-A					3] 18.6 (43.4)	9] 0.89
								4] 19.0 (54.2)	10] 0.98
								5] 18.8 (42.8)	
			Attention scores DST					3] -3.9 (6.8)	9] 0.47
								4] -2.4 (6.8)	10] 0.99
								5] -4.5 (8.5)	
			Language scores (category fluency)					3] -2.9 (6.6)	9] 0.06
								4] -6.3 (9.0)	10] 0.13
								5] -5.0 (5.7)	
			Language scores (letter fluency)					3] -1.7 (6.8)	9] 0.32
								4] -3.1 (5.8)	10] 0.71

EvTable173. Study results: Estrogen cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 12m	
			Motor scores (Grooved Pegboard Test)					5] -2.1 (7.1) 3] -5.2 (42.4) 4] -0.6 (2.7) 5] -5.9 (5.5)	9] 0.90 10] 0.86
			Motor scores(Finger Tapping Test)					3] 4.0 (9.6) 4] -1.3 (10.2) 5] 1.7 (6.9)	9] 0.04 10] 0.25

EvTable174. Study results: Estrogen.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point] (specify) 6w		Final] (specify) 12w	
Wang 2000	ITT Analysis							
	1] Placebo	<u>CASI-Total</u>			3] -0.7 (8.2) 4] 0.4 (5.2)	5] NS	3] 0.5 (8.2) 4] 1.0 (8.0)	5] NS
	2] Estrogen (Conjugated, Premarin)	<u>CDR</u>			3] 0.0 (0.4) 4] 0.0 (0.3)	5] NS	3] 0.1 (0.4) 4] 0.0 ±0.4	5] NS
	3] Placebo change from baseline	<u>CIBIC+</u>			3] -0.2 (0.8) 4] -0.2 (0.9)	5] NS	3] -0.2 (0.8) 4] -0.2 (1.0)	5] NS
	4] Estrogen change from baseline	BEHAVE- AD					3] -0.8 (5.0) 4] -0.4 (3.8)	5] NS
	5] Placebo vs Estrogen in change from baseline	HARS					3] 0.4 (2.6) 4] -0.8 (4.7)	5] NS
		HDRS					3] 0.4 (4.8) 4] -1.2 (5.8)	5] NS

EvTable175. Adverse Events: Estrogens.

Adverse events (AE) identified in included studies	Asthana, 2001	Henderson, 2000	Kyomen, 1999	Mulnard, 2000	Wang, 2000
Withdrawn (%) due to AE	T: 0 C: 0	T: 5 C: 5	T: 0 C: 0	T: 14 C: 5	T: 4 C: 0
AE Checklist (Max 5)	1	3	5	3	2
None Reported			x		
Balance					
Accidental Injury					
Dizziness					
Falls				NS	
Behavioral					
Agitation					
Cardiovascular					
Arrhythmia					
Hypotension					
Hypertension					NS
Extrapyramidal					
Tremor					
Gastrointestinal					
Abdominal pain					
Constipation					
Diarrhea					
Dyspepsia					
Nausea, vomiting					NS
Metabolic/nutritional					
Eating disorder					
Weight Change					
Neurological					
Asthenia					
Psychiatric				NS	
Anxiety					
Confusion, delirium					
Depression					
Respiratory					
Cough, cold, infection					
Rhinitis					
Other	x	x		NS	S
Aberrant hematology					
Fatigue, weakness					
Fever, flu, pneumonia					
Headache					NS
Hepatic abnormality					
Muscle/joint disorder					
Pain					
Rash, skin disorder	x				NS
Sleep disorder					
Urinary disorder					

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
 x = Reported adverse event/side effect but not tested for significant differences between groups
 S or NS = Reported and tested for statistical differences between placebo and treatment group
 S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
 [] = Symptom NOT reported in the paper

EvTable176. Key characteristics: Gingko Biloba.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	#Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Kanowski 1996	PI	8	Placebo Ginkgo Biloba (EGb 761)	DSM-III-R	DAT MID	Mild-Mod	216	156	69.6y (≥55y) 33%M 100% Community	240 mg/d	24w	CGI NAB MADRS SKT	DAT vs MID
Le Bars 1997 Auxiliary: Le Bars 2002 Le Bars 2000 Pors 1998	IF	8	Placebo Ginkgo Biloba	DSM-III-R ICD-10	AD MID	Mild-Modly Sev	327	137	69.0y (45-90y) 46%M	40 mg tid	52w	ADAS-Cog CGIC GERRI	AD vs MID+AD MMSE
Maurer 1997	NR	6	Ginkgo Biloba (Egb761)	DSM-III-R NINCDS	DAT PDD	Mild-Mod	20	18	64.6y (50-80y) 50%M	240 mg/d	3m	ADAS-Cog ADAS-Noncog CGI EEG SKT Trail Making	No

EvTable177. Study results: Ginkgo Biloba (EGb761).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24w	
Kanowski 1996	ITT Analysis							
	1] Placebo	<u>SKT</u>	1] 11.2 (3.4) 2] 10.2 (3.0) 3] 12.2 (3.3)				1] 10.4 (4.9) 2] 8.0 (4.3) 3] 11.8 (3.9)	7] <0.05
	2] Ginkgo Biloba (EGb761) 240 mg/d		4] 10.9 (3.7) 5] 11.0 (3.4) 6] 10.2 (2.8)				4] 9.4 (4.4) 5] 10.1 (5.0) 6] 7.6 (4.2)	
	3] Placebo probable MID	<u>NAB</u>	1] 21.1 (3.7) 2] 21.0 (3.6) 3] 23.3 (4.0)				1] 20.5 (3.6) 2] 20.1 (3.6) 3] 22.3 (3.9)	7] <0.10
	4] Ginkgo Biloba probable MID		4] 21.2 (3.3) 5] 20.7 (3.5) 6] 21.0 (3.7)				4] 20.6 (3.3) 5] 20.2 (3.5) 6] 19.9 (3.7)	
	5] Placebo probable DAT	<u>CGI(numbers improved</u>	3] 4.8 (0.4) 4] 4.9 (1.0)				1] 42/77 2] 57/79	7] <0.05
	6] Ginkgo Biloba probable DAT	<u>compared with baseline)</u>	5] 4.9 (0.4) 6] 5.0 (0.4)				3] 3.8 (1.0) 4] 3.7 (0.8) 5] 4.6 (1.0) 6] 4.0 (0.8)	
	7] Placebo vs. Ginkgo Biloba change from baseline	<u>MADRS</u>	1] 16.1 (7.2) 2] 16.5 (6.8)				1] 13.6 (7.3) 2] 13.2 (6.5)	7] NS

EvTable178. Study results: Ginkgo Biloba (EGb761).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 52w	
Le Bars 1997	ITT Analysis 1] Placebo change from baseline AD & MID 2] Ginkgo Biloba (EGb761) 40 mg tid change from baseline AD & MID 3] Placebo change from baseline AD only 4] Ginkgo Biloba (EGb761) 40 mg tid change from baseline AD only 5] Placebo vs Ginkgo Biloba in change from baseline AD & MID 6] Placebo vs Ginkgo Biloba in change from baseline AD only	<u>ADAS-Cog</u>					1] 1.5 CI (0.4 to2.5) 2] 0.1 CI (-1.8 to1.0) 3] 1.5 CI (0.3 to 2.6) 4] -0.2 CI (-1.2 to 0.8)	5] 0.04 6] 0.02
Le Bars 2002								
Le Bars 2000								
Por 1998		<u>GERRI</u>					1] 0.08 CI (0.01 to 0.14) 2]-0.06 CI(-0.13 to 0.01) 3] 0.09 CI (0.02 to 0.17) 4] -0.09 CI (-0.16 to -0.02)	5] 0.004 6] <0.001
		<u>CGIC</u> <u>(Rating</u>					1] 4.2 CI (4.1-4.3) 2] 4.2 CI (4.1-4.4) 3] 4.2 CI (4.1 to 4.4) 4] 4.2 CI (4.1 to 4.4)	5] 0.77 6] 0.21
		<u>mean)</u>						

EvTable179. Study results: Ginkgo Biloba (EGb761, Tebonin forte).

Author Year	Analysis Groups	Outcome Measures	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Maurer 1997	SKT: ITT Analysis Others: OC Analysis	<u>SKT</u>	1] 18.11 (9.43) 2] 19.67 (6.31)				1]18.89(9.13) 2]16.78(6.87)	3] <0.013
	1] Placebo	ADAS-Cog	1]36.10(15.23) 2]31.21(12.63)				1]36.13(15.56) 2]30.33(14.77)	3] NS
	2] Ginkgo Biloba (EGb761, Tebonin forte) 240 mg/d	CGI (numbers improved compared with baseline)					1] 1/9 2] 5/9	4] 0.069
	3] Difference between placebo and Ginkgo Biloba in change from baseline							
	4] Difference between placebo and Ginkgo Biloba in numbers improved							

EvTable180. Adverse Events: Gingko Biloba.

Adverse events (AE) identified in included studies	Kanowski, 1996	Le Bars, 1997	Maurer, 1997
Withdrawn (%) due to AE	T: 0 C: 0	T: 6 C: 6	T: 0 C: 0
AE Checklist (Max 5)	4	3	5
None Reported		x	x
Balance			
Accidental Injury			
Dizziness			
Falls			
Behavioral			
Agitation			
Cardiovascular			
Arrhythmia			
Hypotension			
Hypertension			
Extrapyramidal			
Tremor			
Gastrointestinal	x		
Abdominal pain			
Constipation			
Diarrhea			
Dyspepsia			
Nausea, vomiting			
Metabolic/nutritional			
Eating disorder			
Weight Change			
Neurological			
Asthenia			
Psychiatric			
Anxiety			
Confusion, delirium			
Depression			
Respiratory			
Cough, cold, infection			
Rhinitis			
Other			
Aberrant hematology			
Fatigue, weakness			
Fever, flu, pneumonia			
Headache	x		
Hepatic abnormality			
Muscle/joint disorder			
Pain			
Rash, skin disorder	S		
Sleep disorder			
Urinary disorder			

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable181. Key characteristics: Idebenone.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Bergamasco 1994	NR	6	Placebo Idebenone	DSM III NINCDS	DAT	Mild – Modly Sev Probable	92	83	70.0y (55-80y) 47%M	30 mg tid	90d	BDRS GBS Laboratory tests Rey's 15 Word Test Rey's A Figure Test SCAG Token Test Word Fluency Test	No
Gutzmann 1998 Auxiliary: Weyer 1996	NR	6	Placebo Idebenone	DSM-III-R NINCDS	AD PDD	Mild-Modly Sev	450	379	69.9y (58-82y) 34%M	120 mg tid	12m	ADAS (Cog/NODCOS) Adverse Events Caregiver observation CGI Laboratory tests ECG IADL NOSGER SKT	Disease severity
Gutzmann 2002	PI	7	Idebenone Tacrine	NINCDS DSM-III-R	AD PDD	Mild-Mod	203	44	71.2y (44-90y) 36%M 100% White 100% Community	360 mg/d 160 mg/d	60w	ADAS-Cog ADAS-Noncog ADAS-Total CGI CT EIS HIS MRI NOSGER-IADL	No

EvTable181. Key characteristics. Idebenone cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Marigliano 1992	NR	7	Placebo Idebenone	DSM-III-R	MID	Mild-Mod	108	108	73.6y (65-80y) 49%M 31% Institution 69% Community	45 mg bid	120d	ECG GBS HIS HRSD Laboratory tests MMSE Randt Memory Test Token Test TP	No
Weyer 1997	NR	5	Placebo Idebenone	DSM-III-R NINCDS	PDD DAT	Mild-Mod	300	247	70.0y (54-90y) 34%M	30 mg/d 90 mg/d	6m	ADAS- Total ADAS-Cog ADAS-Noncog CGI DSS Greene's Assessment HAMD MMSE NAA NAB Laboratory tests	Disease severity

EvTable182. Study results: Idebenone.

Author Year	Outcomes Measured	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 45d		Final: (specify) 90d	
Bergamasco 1994	OC Analysis	SCAG total score	1] 60.00(2.02)* 2] 57.10(1.99)*		1] 59.20(1.64)* 2] 55.24(1.75)*	4] <0.05	1] 57.62(1.70)* 2] 53.00(2.18)*	4] <0.05 5] <0.05
	1] Placebo							
	2] Idebenone 30 mg tid	Rey's 15 Word Test				4] <0.05		4] <0.05
	3] Placebo difference from baseline		1] 14.07(0.91)* 2] 17.31(1.41)*		1] 14.95(0.87)* 2] 20.14(1.64)*		1] 15.24(1.02)* 2] 21.28(1.78)*	5] NS
	4] Idebenone difference from baseline	TK	1] 22.84(0.82)* 2] 23.80(0.89)*		1] 22.79(0.83)* 2] 24.48(0.92)*	4] <0.05	1] 23.22(0.81)* 2] 24.68(0.89)*	3] <0.05 4] <0.05 5] <0.05
	5] Idebenone vs Placebo change from baseline	BDRS	1] 17.07(1.01)* 2] 17.50(1.22)*		1] 16.83(1.06)* 2] 18.12(1.23)*		1] 16.03(1.18)* 2] 19.17(1.37)*	

*SEM

EvTable183. Study results: Idebenone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6m		Final: (specify) 12m	
Gutzman, 1998	ITT Analysis	ADAS Total			4] -5.6 (8.2) 5] -7.2 (7.4) 6] -8.4 (8.1)	7] <0.0027	4] -4.9 (8.5) 5] -7.1 (8.6) 6] -8.8 (9.5)	7] 0.0001
Weyer, 1996	1] Placebo 90mg							
	2] Idebenone 90 mg tid	ADAS-Cog	1] 34.3 (9.3) 2] 35.3 (9.3) 3] 32.7 (8.0)		4] -3.4 5] -4.9 6] -6.0	7] <0.001	4] -3.0 5] -5.0 6] -7.0	7] <0.0005
	3] Idebenone 120 mg tid	CGI	1] 5.2 (0.4) 2] 5.2 (0.5) 3] 5.1 (0.4)				1] 63.3 2] 73.4 3] 87.3	7] 0.0000
	4] Placebo change from baseline							
	5] Idebenone 90 mg tid change from baseline	NOSGER- IADL	1] 16.0 (4.8) 2] 16.0 (4.9) 3] 15.3 (4.9)				1] 36.7 2] 41.1 3] 48.4	7] 0.0298
	6] Idebenone 120 mg tid change from baseline							
	7] Dose trend							

EvTable184. Study results: Idebenone-Tacrine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 60 w	
Gutzmann 2002	ITT Analysis 1] Idebenone 360 mg d 2] Tacrine 160 mg d variable 3] Difference between Idebenone and Tacrine	<u>EIS%</u>					1] 28.9%	3] <0.0001 favours Idebenone
		Rating = -1					2] 9.0%	
		Rating = 0					1] 54.8%	
		Rating = 1					2] 83.8%	
		Rating = 2					1] 16.3%	
		Rating = 3					2] 7.1%	
		ADAS-Total	1] 41.55(16.46) 2] 41.52(14.92)				1] 34.51(17.43) 2] 30.44(16.32)	3] NS
		ADAS-Cog	1] 30.23(11.59) 2] 30.93(10.59)				1] 26.40(16.67) 2] 24.81(14.92)	
		ADAS-Noncog	1] 11.32 (6.79) 2] 10.55(5.86)				1] 8.11(7.56) 2] 5.63(6.10)	
		CGI-S	1] 5.22 (0.46) 2] 5.19 (0.44)				1] 4.43 (1.58) 2] 4.53 (1.45)	
		NOSGER-IADL	1] 13.88(4.43) 2] 13.78(4.55)				1] 13.13 (5.49) 2] 12.5 (6.25)	3] NS

EvTable185. Study results: Idebenone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 60d		Final: (specify) 150d	
Marigliano 1992	OC Analysis							
	1] Placebo	RMT- Delayed Recall	1] 62 2] 60		1] 59 2] 64	4] <0.04	1] 55 2] 61	5] <0.02
	2] Idebenone 45 mg bid	RMT- Figure Recognition	1] 7 2] 6.5		1] 6 2] 6.4	5] <0.05	1] 5.5 2] 6.5	
	3] Placebo change from baseline	RMT Paired	1] 6 2] 5.5		1] 6 2] 5		1] 2.5 2] 6	5] <0.02
	4] Idebenone 45 mg bid change from baseline	Word- Acquisition						
	5] Idebenone 45 mg bid vs. placebo	GBS Intellectual Functions	1] 16.5 2] 16.5		1] 16 2] 15		1] 13 2] 11	5] <0.02
		GBS Motor functions	1] 7 2] 7		1] 7 2] 6		1] 4.5 2] 5	5] <0.05

EvTable186. Study results: Idebenone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3m		Final: (specify) 6m	
Weyer, 1997	ITT Analysis	ADAS-Total			1] -1.6 2] -5 3] -6.9 4] -6.1 5] -6.5 6] -9.5		1] -5.7 2] -5.6 3] -8.0 4] -7.0 5] -7.6 6] -11.1	7] 0.037 8] 0.009
	1] Placebo change from baseline							
	2] Idebenone 30 mg tid change from baseline	ADAS-Cog			1] -3.6 2] -3.8 3] -5.5 4] -4.8 5] -5.0 6] -7.3		1] -4.5 2] -4.4 3] -6.5 4] -5.8 5] -5.9 6] -8.9	7] 0.031 8] 0.006
	3] Idebenone 90 mg tid change from baseline							
	4] Placebo ADAS-Total >20 change from baseline	ADAS- Noncog			1] -0.9 2] -1.3 3] -1.3 4] -1.5 5] -1.6 6] -2.3		1] -1.2 2] -1.9 3] -2.0 4] -1.8 5] -2.2 6] -3.2	7] 0.035 8] 0.014
	5] Idebenone30 ADAS-Total >20 change from baseline	CGI % improved						
	6] Idebenone 90 ADAS-Total >20 change from baseline						1] 67.5 2] 72.2 3] 81.6 4] 47.9 5] 66.7 6] 88.5	7] 0.018 8] 0.000
	7] Treatment effect 0 to 6m							
	8] Treatment effect 0 to 6m sub-group ADAS-Cog >20							

EvTable187. Adverse Events: Idebenone.

Adverse events (AE) identified in included studies	Bergamasco, 1994	Gutzmann, 1998	Marigliano, 1992	Weyer, 1997	IDEBENONE(C) TACRINE (T) Gutzmann, 2002
Withdrawn (%) due to AE	T: 4 C: 2	T: NR C: NR	T: 0 C: 0	T: 5 C: 5	T: 41 C: 17
AE Checklist (Max 5)	4	1	5	3	3
None Reported					
Balance			x		
Accidental Injury					
Dizziness		x		NS*	
Falls			x		
Behavioral	x				
Agitation					
Cardiovascular				x	
Arrhythmia			x		
Hypotension		NS*			
Hypertension					
Extrapyramidal					
Tremor					
Gastrointestinal			x		S*
Abdominal pain	x				
Constipation					
Diarrhea					
Dyspepsia				x	
Nausea, vomiting	x			x	S*
Metabolic/nutritional					
Eating disorder					
Weight Change					
Neurological		NS*			
Asthenia					
Psychiatric					
Anxiety	x				
Confusion, delirium			x		
Depression					
Respiratory		x		NS*	
Cough, cold, infection					
Rhinitis					
Other			x		
Aberrant hematology		NS*			
Fatigue, weakness					
Fever, flu, pneumonia		x			
Headache				x	
Hepatic abnormality				NS*	S*
Muscle/joint disorder					
Pain		x			
Rash, skin disorder			x		
Sleep disorder	x				
Urinary disorder					

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable188. Key characteristics: Oxiracetam.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Bottini 1992	PI	6	Placebo Oxiracetam	DSM III NINCDS	PDD MID Mixed	NR	65	58	71.0y (54-82y) 43%M	800 mg bid	12w	DSPT QoL RPM RT TK WLM Verbal Fluency Short Story	No
Maina 1989	NR	6	Placebo Oxiracetam	DSM III	PDD MID MIXED	NR	289	272	73.0y (<85y) 50%M	800 mg bid	12w	BDS Global Evaluation of efficacy IPSE-E NMICS IPAX-E AE	PDD vs MID
Mangoni 1988	PI	6	Placebo Oxiracetam	NINCDS	AD	Probable- Possible Mild-Mod	30	30	62.0y (52-79y) 67%M	800 mg bid	24w	IPSC-E LNNB Sensory motor left scale Frontal right scale	No

EvTable188. Key characteristics: Oxiracetam cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	#Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Rozzini 1992 Auxiliary: Rozzini 1993	NR	4	Placebo Oxiracetam	DSM-III-R	PDD	Mild-Mod	110	94	73.8y (>65y) 34%M	800 mg bid	26w	AMI BI BDI BMDI IADL MMSE RT	No
Villardita 1992	PI	6	Placebo Oxiracetam	NINCDS DSM III	AD MID	Mild-Mod	60	60	69.8y (52-83y) 60%M	800 mg bid	90d	ACPT BTT DS IADL-E IPSC-E LAS MMSE MWF Rosy's WT	No

EvTable189. Study results: Oxiracetem.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6w		Final: (specify) 12w	
Bottini 1992	Completer Analysis 1] Placebo 2] Oxiracetem 800 mg bid 3] Placebo vs baseline 4] Oxiracetem vs baseline 5] Oxiracetem vs Placebo	Reaction Time milli second	1] 783 (634) 2] 1018 (873)				1] 697 (350) 2] 996 (1101)	5] NS
		Verbal Fluency phonemic	1] 12.6 (8.21) 2] 13.3(12.37)				1] 9.8 (7.74) 2] 14.2 (10.2)	5] 0.009
		Verbal Fluency semantic	1] 13.8 (13.0) 2] 12.9 (7.31)				1] 13.0 (7.02) 2] 14.7 (7.31)	5] <0.02
		Short Story	1] 7.8 (12.54) 2] 5.2 (6.24)				1] 6.1 (5.942) 2] 9.7 (11.67)	5] 0.02
		RPM	1] 13.9 (7.34) 2] 13.1 (6.15)				1] 12.7 (6.08) 2] 14.5 (6.5)	5] 0.007
		DSPT	1] 4.3 (0.94) 2] 4.2 (0.95)				1] 4.5 (1.03) 2] 4.7 (1.16)	3] <0.01 4] <0.01 5] NS
		WLM	1] 4.0 (3.71) 2] 3.6 (2.98)				1] 4.7 (3.97) 2] 5.1 (3.73)	3] <0.01 4] <0.01 5] <0.05
		TK	1] 22.6 (5.83) 2] 23.3 (6.11)				1] 23.8 (6.16) 2] 23.1 (7.17)	5] NS
		Italian QoL	1] 2.7 (0.55) 2] 2.8 (0.56)		1] 2.7 (0.54) 2] 3.1 (0.53)		1] 2.5 (0.62) 2] 3.2 (0.50)	5] <0.001

EvTable190. Study results: Oxiracetem.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Maina, 1989	Per Protocol Analysis 1] Placebo 2] Oxiracetem 800 mg bid 3] Difference between treatments in favor of Oxiracetem 4] Time by treatment interaction	BDS NMICS IPSE-E Global evaluation of efficacy	1] 10.5 (4.20) 2] 10.7 (4.25) 1] 17.3 (7.06) 2] 18.4 (7.34) 1] 1.9 (0.35) 2] 2.0 (0.34)	3] NS			1] 10.3 (4.54) 2] 9.0 (4.17) 1] 17.7 (20.9) 2] 20.9 (6.93) 1] 1.8 (est) 2] 1.51 (est) Good or very good 1] 21% 2] 64%	3] <0.01 3] <0.01 4] <0.01 3] <0.01 3] <0.01

EvTable191. Study results: Oxiracetem.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12w		Final: (specify) 24w	
Mangoni 1998	Completer Analysis 1] Placebo 2] Oxiracetem 800 bid 3] Between treatment in favor of Oxiracetem 4] Time by treatment interaction	IPSC-E LNNB Sensory-motor left scale Frontal right scale	1] 2.1 (0.75) 2] 2.3 (0.68)	3] NS	1] 1.94 (est) 2] 1.67 (est)		1] 2.1 (est) 2] 1.75 (est) 3] 7/14 tests 4] 7/14 tests	3] <0.01 4] <0.001 3] <0.01 4] <0.01 4] <0.01 4] <0.01

EvTable192. Study results: Oxiracetem.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 2m		Final: (specify) 6m	
Rozzini, 1992	OC Analysis	BI	1] 94.89 (9.79) 2] 94.38(10.69)		1] 94.67(10.02) 2] 94.17(10.77)		1] 94.46(10.28) 2] 93.96(11.32)	4] NS
Rozzini 1993	1] Placebo							
	2] Oxiracetem 800 mg bid	IADL	1] 1.52 (1.64) 2] 1.48 (1.62)		1] 1.57(1.70) 2] 1.50 (1.72)		1] 1.63 (1.85) 2] 1.56 (1.73)	3] NS 4] NS
	3] Oxiracetem vs baseline	MMSE	1] 21.98 (2.77) 2] 22.08 (2.78)		1] 22.22(3.01) 2] 23.35(3.18)	3] <0.05	1] 22.04(3.28) 2] 22.71 (3.58)	
	4] Placebo vs. Oxiracetam change from baseline	RT score	1] 1.84 (1.95) 2] 8.67 (2.05)		1] 8.84 (1.79) 2] 7.64 (2.47)	3] <0.05	1] 8.85(2.04) 2] 7.54(2.44)	3] <0.02
		BDI	1] 20.43(14.49) 2] 23.79(17.96)		1] 21.04(15.92) 2] 20.75(16.21)		1] 22.25(16.82) 2] 23.58(16.4)	
		Test of Attention Matrix	1] 18.83 (8.72) 2] 18.46 (9.14)		1] 19.39 (7.21) 2] 19.27 (8.96)	3] NS	1]18.85 (6.92) 2] 20.31 (8.92)	3] NS

EvTable193. Study results: Oxiracetem.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 90d	
Villardita 1992	OC Analysis	MMSE	1] 16.7 (0.8)* 2] 15.5 (0.9)*				1] 15.6 (0.8)* 2] 21.6 (1.0)*	3] NS 4] <0.01 5] <0.01
	1] Placebo							
	2] Oxiracetem 800 mg bid	ACPT	1] 18.5 (0.8)* 2] 17.2 (0.9)*				1] 17.3 (1.0)* 2] 21.6 (1.0)*	3] NS 4] <0.05 5] <0.05
	3] Placebo vs baseline							
	4] Oxiracetem vs baseline	R15WT immediate	1] 18.8 (1.1)* 2] 18.5 (1.1)*				1] 18.4 (1.1)* 2] 20.6 (1.1)*	3] NS 4] <0.05 5] <0.05
	5] Oxiracetem vs Placebo	R15WT delayed	1] 2.4 (0.3)* 2] 2.3 (0.3)*				1] 2.5 (0.2)* 2] 2.8(0.2)*	3] NS 4] <0.05 5] NS
		DS forward	1] 4.5 (0.3)* 2] 4.5 (0.2)*				1] 4.1 (0.2)* 2] 4.5 (0.2)*	5] NS
		DS backward	1] 3.1 (0.2)* 2] 3.0 (0.2)*				1] 3.1 (0.2)* 2] 2.9 (0.2)*	5] NS
		IPSC-E total	1] 152.7 (4.1)* 2] 149.9(4.4)*				1] 155.9 (4.2)* 2] 157.6 (5.2)*	5] NS
		IADL-E total	1] 20.9(1.2)*				1] 19.6 (1.0)* 2] 26.0 (1.0)*	3] <0.05 4] <0.01 5] <0.01

*SEM

EvTable194. Adverse Events: Oxiracetam.

Adverse events (AE) identified in included studies	Bottini, 1992	Maina, 1989	Mangoni, 1988	Rozzini, 1992	Villardita, 1992
Withdrawn (%) due to AE	T: 6 C: 0	T: 2 C: 2	T: 0 C: 0	T: 4 C: 9	T: 0 C: 0
AE Checklist (Max 5)	4	5	5	2	5
None Reported			x		
Balance					
Accidental Injury					
Dizziness	x				
Falls					
Behavioral					
Agitation		x			
Cardiovascular					
Arrhythmia					
Hypotension					
Hypertension					
Extrapyramidal	x				
Tremor					
Gastrointestinal				x	
Abdominal pain	x	x		x	
Constipation		x			
Diarrhea					
Dyspepsia				x	
Nausea, vomiting					
Metabolic/nutritional					
Eating disorder	x				
Weight Change					
Neurological					
Asthenia					
Psychiatric					
Anxiety				x	x
Confusion, delirium		x			
Depression					
Respiratory					
Cough, cold, infection					
Rhinitis					
Other	x				
Aberrant hematology		NS			
Fatigue, weakness					
Fever, flu, pneumonia					
Headache					
Hepatic abnormality					
Muscle/joint disorder					
Pain					
Rash, skin disorder	x				
Sleep disorder	x				x
Urinary disorder	x				

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable195. Key characteristics. Pentoxifylline.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Knezevic 1996	IF	6	Placebo Pentoxifylline	DSM-III-R	MID	Mild-Mod	289	239	69.7y (45-90y) 56%M	1200 mg/d	9m	GBS MMSE Neuropsychological SCAG	No
Black 1992	PI	6	Placebo Pentoxifylline	DSM III	MID	Nild-Mod	64	38	75.4y (55-98y) 52%M	400 mg tid	36w	ADAS-Cog ADAS-Noncog HIS MMSE	Vascular change vs Discrete strokes
Ghose 1987	IF	6	Placebo Pentoxifylline	DSM III	PDD MID	Mild-Mod	36	28	77.0y (60-88y) 50%M	400 mg tid	12w	Digit Span Test DSPT MMSE RT SCAG Shopping list	MID vs PDD
DRUG VS DRUG													
Parnetti 1997	NR	5	Sulodexide Pentoxifylline	NINDS- AIREN	VaD	Probable Mild-Mod	93	86	75.0y (65-80y) 40%M	100 mg/d 1200 mg/d	6m	GBS Laboratory tests MMSE	No

EvTable196. Study results: Pentoxifylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 9m	
Knezevic, 1996	ITT/LOCF Analysis 1] Placebo 2] Pentoxifylline 1200 mg/day 3] Placebo change from baseline 4] Pentoxifylline change from baseline 5] Placebo vs Pentoxifylline change from baseline	GBS total	1] 37.2 (17.2) 2] 37.9 (17.5)				3] 0.0 (13.9) 4] -2.1 (14.7)	5] 0.065
		GBS Intellectual					3] 0.0 (7.0) 4] -0.9 (7.6)	5] 0.060
		GBS Motor					3] 0.2 (5.0) 4] 0.0 (9.5)	5] 0.275
		GBS emotional					3] 0.3 (2.3) 4] -0.3 (2.8)	5] 0.072
		MMSE						5] NS
		SCAG Total	1] 43.9 (11.8) 2] 44.8 (11.7)				3] -1.9 (9.0) 4] -3.1 (10.4)	5] 0.034
		SCAG cognitive					3] -0.4 (3.5) 4] -1.3 (3.6)	5] 0.007

EvTable197. Study results: Pentoxifylline.

Author Year	Analysis Groups	Outcome Measures	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 36w	
Black 1992	Efficacy Analysis	ADAS-total	1] 30.08				1] 37.03	3] 0.058
			2] 26.5				2] 28.19	6] 0.023
			4] 29.17				4] 34.98	9] 0.002
			5] 22.94				5] 20.85	
			7] 30.77				7] 41.99	
			8] 26.72				8] 25.36	
		ADAS-Cog	1] 25.39				1] 30.41	3] 0.064
			2] 21.69				2] 23.10	6] 0.020
			4] 24.81				4] 28.83	9] 0.005
			5] 18.34				5] 17.11	
			7] 25.72				7] 33.24	
			8] 21.72				8] 21.14	
		ADAS-Noncog	1] 4.69				1] 6.63	3] 0.23
			2] 4.81				2] 5.09	6] 0.12
			4] 4.36				4] 6.16	9] 0.017
			5] 4.60				5] 3.73	
			7] 5.05				7] 8.73	
			8] 5.00				8] 4.22	
		ADAS without cognitive memory	1] 7.78				1] 10.16	3] 0.036
			2] 6.63				2] 6.72	6] 0.005
			4] 7.20				4] 8.96	9] 0.001
			5] 5.07				5] 3.47	
			7] 7.78				7] 12.84	
			8] 6.00				8] 5.28	
		9] Placebo vs Pentoxifylline change from baseline subgroup with discrete strokes						

EvTable198. Study results: Oxpentifylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Ghose, 1987	Per protocol analysis	<u>MMSE</u>	1] 19.4 2] 19.6 3] 16.6 4] 16.5 5] 19.9 6] 16.6				1] 20.4 2] 21.3 3] 18.3 4] 20.3 5] 21.3 6] 19.5	7] <0.05 8] NS
	1] Placebo PDD							
	2] Placebo MID							
	3] Oxpentifylline 400 mg tid PDD	<u>SCAG</u>	1] 42.1 (14.3) 2] 44.1 (16.3) 3] 53.5 (15.6) 4] 43.0 (16.7) 5] 42.1 (14.3) 6] 50 (16)				1] 37.3 (14.0) 2] 39.9 (14.8) 3] 44.3 (13.0) 4] 41.0 (10.4) 5] 37.4 (14.4) 6] 43.51 (12.9)	7] NS 8] NS
	4] Oxpentifylline 400 mg tid MID							
	5] Placebo all							
	6] Oxpentifylline all	<u>DSST</u>	5] 5.5 (1.4) 6] 5.4 (1.3)				5] 6.2 (1.4) 6] 5.7 (1.2)	7] NS 8] NS
	7] Oxpentifylline vs Placebo MID							
	8] Oxpentifylline vs Placebo PDD							

EvTable199. Study results: Sulodexide - Pentoxifylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4m		Final: (specify) 6m	
Parnetti, 1997	OC Analysis							
	1] Sulodexide 50 mg bid	<u>GBS motor impairment</u>	1] 1.64 (0.14)* 2] 1.59 (0.13)*		1] 1.58 (0.14)* 2] 1.53 (0.14)*	3] <0.01	1] 1.54 (0.16)* 2] 1.46 (0.17)*	3] <0.01
	2] Pentoxifylline 400 mg tid	<u>GBS intellectual impairment</u>	1] 2.09 (0.09)* 2] 1.98 (0.08)*		1] 1.88 (0.09)* 2] 1.95 (0.1)*	3] <0.01	1] 1.79 (0.1)* 2] 1.87 (0.12)*	3] <0.01
	3] sulodexide change vs baseline	<u>GBS emotional impairmen</u>	1] 2.1 (0.12)* 2] 1.89 (0.1)*		1] 1.88 (0.12)* 2] 1.9 (0.1)*	3] <0.12	1] 1.76 (0.12)* 2] 1.75 (0.11)*	3] <0.01
		MMSE	1] 17.6 (0.4)* 2] 18 (0.4)*				1] 20 (0.6)* 2] 20 (0.4)*	

*SEM

EvTable200. Adverse Events. Pentoxifylline.

Adverse events (AE) identified in included studies	Black, 1992	Ghose, 1987	Knezivic, 1996	PENTOXIFYLLINE SULODEXIDE Parnetti, 1997
Withdrawn (%) due to AE	T: 22 C: 25	T: 12 C: 6	T: 0 C: 0	T: 7 C: 6
AE Checklist (Max 5)	1	2	3	2
None Reported			X	
Balance				
Accidental Injury				
Dizziness				
Falls				
Behavioral				
Agitation				
Cardiovascular	X			X
Arrhythmia				
Hypotension				
Hypertension				
Extrapyramidal				
Tremor				
Gastrointestinal	X	X		
Abdominal pain				X
Constipation				
Diarrhea				
Dyspepsia				
Nausea, vomiting	X	X		
Metabolic/nutritional				
Eating disorder				
Weight Change				
Neurological	X			
Asthenia				X
Psychiatric				
Anxiety				
Confusion, delirium				
Depression				
Respiratory				
Cough, cold, infection				
Rhinitis				
Other				
Aberrant hematology				
Fatigue, weakness				
Fever, flu, pneumonia				
Headache				X
Hepatic abnormality				
Muscle/joint disorder				
Pain				
Rash, skin disorder				
Sleep disorder				
Urinary disorder				

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable201. Key characteristics: Propentofylline.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Marcusson 1997	NR	5	Placebo Propentofylline	DSM-III-R	AD VaD	Mild-Mod	260	187	72.4y (NR) %M NR	300 mg tid	12m	ADAS-Cog BfS CGI DSST ECG GBS Laboratory tests MMSE NAA NAB SKT Syndrome Short Test Zerssen Adjective Mood Scale	AD vs VaD
Mielke 1996	IS	5	Placebo Propentofylline	DSM-III-R	VaD	Mild-Mod	30	25	68.7y (55-79y) 58%M	300 mg tid	3m	DSST Fragmented Picture Task Memory Tasks MMSE PET Physiological tests	No
Mielke 1998	NR	5	Placebo Propentofylline	NINCDS DSM-III-R	AD	Probable Mild-Mod	30	28	64.8y (52-78y) 57%M	300 mg tid	3m	DSST FAST MMSE PET BSRT SRT-DR Verbal Fluency	No

EvTable201. Key characteristics: Propentofylline cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male Population	Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Saletu 1990 Auxiliary: Moller 1994	NR	6	Placebo HWA 285 (Propentofylline)	DSM-III-R	Dementia	Mild	190	165	68.5y (NR) 49%M	300 mg tid	12w	Alphabetical Cross-out test Benton Test CGI-CGC CGI-S Cognitive Difficulties Scale CT DST EEG Folstein Mini-Mental State GBS Gruenberger Verbal Memory Test MMSE MRI Pictures of Objects Task and Learning Trail Making Test	MMSE

EvTable202. Study results: Propentofylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6m		Final: (specify) 12m	
Marcusson 1997	ITT Analysis 1] Placebo change from baseline 2] Propentofylline 300 mg tid change from baseline 3] Change from baseline in treatment vs placebo	<u>GBS</u>			1] 1.5 (12.5) 2] -2.2 (11.1)	3] 0.003	1] 4.5 (15.5) 2] -0.4 (14.2)	3] 0.001
		<u>CGI item II</u>			1] -0.4 (2.3) 2] -1.0 (2.7)	3] 0.028	1] -0.3 (4.0) 2] -1.2 (5.1)	3] 0.072
		<u>SKT</u>			1] -0.4 (3.9) 2] -1.9 (3.4)	3] 0.001	1] -0.1 (3.8) 2] -1.5 (3.9)	3] 0.002
		CGI item I			1] 0.03 (0.6) 2] -0.14 (0.6)	3] 0.014	1] 0.09 (0.7) 2] -0.12 (0.8)	3] 0.004
		MMSE			1] 0.4 (3.5) 2] 0.9 (3.1)	3] 0.072	1] -0.6 (4.2) 2] 0.6 (3.9)	3] 0.001
		DSST			1] -0.5 (6.0) 2] 0.6 (6.9)	3] 0.222	1] -0.9 (6.5) 2] 1.1 (8.2)	3] 0.062
		NAB			1] 1.1 (3.4) 2] 0.1 (3.4)	3] 0.021	1] 6.8 (3.7) 2] 0.6 (3.8)	3] 0.007
		NAA			1] -0.3 (4.7) 2] -0.0(4.3)	3] 0.395	1] 0.9 (4.9) 2] -0.2 (5.1)	3] 0.698
		BfS			1] 0.1 (10.6) 2] -1.5 (9.4)	3] 0.588	1] -0.1 (10.1) 2] -1.2 (9.9)	3] 0.893

EvTable203. Study results: Propentofylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Mielke, 1996	OC Analysis	MMSE		3] NS			1] 21.9 (3.1) 2] 21.4 (3.2)	3] <0.1
	2] Propentofylline 300 mg tid change from baseline	DSST		3] NS			1] 19.6 (8.1) 2] 15.6 (7.5)	3] <0.1
	3] Propentofylline 300 mg tid vs Placebo							

EvTable204. Study results: Propentofylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3m	
Mielke, 1998	OC Analysis 1] Placebo change from baseline 2] Propentofylline 300 mg tid change from baseline 3] Change from baseline in treatment vs placebo	MMSE	1] 21.3 (3.4) 2] 19.5 (3.5)				1] -1.0 (1.6) 2] -1.4 (3.8) 3] NS	1] NS 2] 0.02 3] NS
		BSRT					1] 0.1 (0.7) 2] -0.1 (0.5) 3] NS	1] NS 2] NS 3] NS
		SRT DR					1] 0.4 (1.1) 2] 0.5 (1.3) 3] NS	1] NS 2] NS 3] NS
		DSST					1] -2.1 (7.5) 2] 3.2 (6.2) 3] <0.06	1] NS 2] NS 3] <0.06
		Verbal Fluency					1] 0.6 (7.5) 2] -1.9 (7.3) 3] NS	1] NS 2] NS 3] NS

EvTable205. Study results: Propentofylline.

Author Year	Analysis Groups	Test Used	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Saletu, 1990	OC Analysis	GBS Total	1] 26.2 (11.1) 2] 29.4 (14.8)	5] 0.1632			3] -2.5 (10.4) 4] -11.6 (11.8)	3]<0.05 4] <0.05 5] 0.0001
Moller 1994	1] Placebo							
	2] Propentofylline 300mg tid	MMSE	1] 20.9 (2.3) 2] 21.0 (2.7)	5] 0.8794			3] 2.3 (4.0) 4] 4.2 (3.2)	3]<0.05 4]<0.05 5] 0.0038
	3] Placebo change from baseline	CGI –S	1] 5.06 (.71) 2] 5.19 (.74)	5] 0.619			3] -0.50 (1.20) 4] -0.82 (1.00)	3] <0.05 4] <0.05 5] 0.014
	4] Propentofylline change from baseline	CGI-CGC					3] 4.02 (1.37) 4] 3.33 (0.97)	3] <0.05 4] <0.05 5] 0.003
	5] Placebo vs Propentofylline	CGI-efficacy					3] 2.70 (1.05) 4] 2.31 (1.04)	3] <0.05 4] <0.05 5] 0.032
		8 Psychometric tests				5] NS		3] 0.05 4] 0.05 5] NS

EvTable206. Adverse Events: Propentofylline.

Adverse events (AE) identified in included studies	Marcusson{1984}, 1997	Mielke{2645}, 1996	Mielke{1855}, 1998	Saletu{4169}, 1990
Withdrawn (%) due to AE	T: 12 C: 8	T: 0 C: 0	T: 8 C: 0	T: 0 C: 13
AE Checklist (Max 5)	2	4	2	1
None Reported		X		
Balance				
Accidental Injury				
Dizziness	X			X
Falls				
Behavioral				
Agitation				
Cardiovascular				
Arrhythmia				
Hypotension				
Hypertension				
Extrapyramidal				
Tremor				
Gastrointestinal	X			X
Abdominal pain	X		X	
Constipation				
Diarrhea				
Dyspepsia				
Nausea, vomiting	X			
Metabolic/nutritional				
Eating disorder				
Weight Change				
Neurological				
Asthenia				
Psychiatric				
Anxiety				X
Confusion, delirium				
Depression				
Respiratory				
Cough, cold, infection				
Rhinitis				
Other				
Aberrant hematology				
Fatigue, weakness				
Fever, flu, pneumonia				
Headache	X			X
Hepatic abnormality				
Muscle/joint disorder				
Pain				
Rash, skin disorder			X	
Sleep disorder				
Urinary disorder				

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable207. Key characteristics: Other agents.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Senin 1991	PI	8	Placebo Aniracetam	NINCDS	AD	Mild-Mod	109	109	72.4y (65-80y) 44%M 100% Community	750 mg bid	6m	Blessed-D Corsi Test Gibson Spiral Maze MMSE Raven Colored Progressive Matrices Rey's 15 SCAG TP	No
Ban 1991b	NR	5	Placebo Ateroid	DSM III	PDD MID SDAT	NR	155	148	73.0 y (NR) 56% M 30% Community 70% Institution	200 LRU tid	12w	SCAG HDS Laboratory tests	PDD vs MID
Shrotriya 1996	IF	6	Placebo BMY (Nootropic)	NINCDS DSM-III-R	AD	Mild-Mod	69	54	72.0 y (54-92 y) 41% M	300 mg tid	12w + 4w washout	ADAS ADAS-Cog AE CGI-S CNTB MMSE	No
Cucinotta 1992	NR	6	Placebo Buflomedil	DSM-III-R	VaD	Mild-Mod	88	73	74.0 y (NR) 47% M	300 mg bid	270d	Birren test CGI Dementia rating scale HDS HIS MMSE Nowlis mood test SHGRS	No

EvTable207. Key characteristics: Other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Danielczyk 1988	NR	8	Placebo CBM 36-733 (Ergokryptine)	DSM III	PDD	Mild- Mod	125	117	73.8 y (60-90 y) 41% M Community Institution	0.4 mg/d (Day 1-3) 0.8 mg/d (Day 4-6) 1.2 mg/d (Day 7-9) 2 mg/d (Day 10-end)	8w	Digit span Laboratory tests Neuropsychological Battery NOSIE Psychometric tests SCAG Trail Making Test WAIS	No
Shimada 1994	NI IS	5	Placebo Choto-san	DSM-III-R	VaD	NR	60	57	78.9 y (NR) 15% M 100% Asian 77% Cardiovascular 5% Diabetes 5% Parkinson's 5% Liver/Renal	2.5 g tid	12w	Global improvement rating Hasegawa's dementia scale Overall safety rating Utility rating	No
Terasawa 1997	PI	5	Placebo Choto-san	DSM-III-R	VaD	NR	139	119	76.6 y (NR) 36 100% Asian; 80% Cardiovascular 8% Diabetes 5% Parkinson's 3% Liver/Renal	7.5 g tid	12w	HDS-R Various Global Rating Scales	No
Schellenberg 1997	PI	5	Placebo Cyclandelate	DSM-III-R	AD	NR	139	92	75.0 y (62-85 y) 21% M Community	400 mg qid	16w	ADAS FIGT MEMT NCT NSL NST SCAG	No

EvTable207. Key characteristics: Other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Weyer 2000	PI	7	Placebo Cyclandelate	DSM-III-R	PDD VaD Mixed	Mild- Modly Sev	196	147	78.6 y (54-91y) 27% M Community	800 mg bid	24w	ADAS-Cog CGI-C NOSGER-IADL	MMSE ADAS- Cog Treatme nt Centre
Peabody 1986	NI IS	6	Placebo Vasopressin (DAVP)	DSM III	PDD	Mild- Mod	14	NR	NR	30 µg/d (start) 180 µg/d	3w	BUSCHKE-S&L SCAG HAM-D POMS VAMS	No
Cucinotta 1996 Auxiliary: Cucinotta 1998	IF	6	Placebo Ergokryptine (DEK)	NINCDS	AD	Mild- Mod	215	155	74.0 y (60-85 y) 33% M Italian, Institution	5 mg bid (week 1-2) 10 mg bid (week 3-4) 20 mg bid (until end)	1y	AE GBS HAM-D MDB MMSE Rey	No
Treves 1999	NR	5	Placebo Denbufylline	DSM-III-R	VaD AD MIXED	Mild- Mod	336	229	74.0 y (>60 y) 42% M	100 mg bid	16w	MMSE WAIS-DSST WAIS-VDC	No
Crapper- McLachlan 1991	NI	7	Placebo DFO	NINCDS	AD	Proba ble	48	39	63.1y (57-69 y) 49% M 100% Community	500 mg bid (start) 125 mg bid (Day 17- end)	2y	VHB Wechsler Adult Intelligence- revised verbal Wechsler Memory Scale Western Aphasia Battery	No

EvTable207. Key characteristics: Other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Scharf 1999	NI IS	7	Placebo Diclofenac & Misoprostol	DSM IV	AD	Mild-Mod	41	26	73.0 y (≥50 y) 44% M Community	50 mg & 200 µg bid	25w	ADAS-Cog ADAS-Noncog ADAS-total CGIC GCIC GDS IADL MMSE PSMS	No
Ban 1991a	IS	7	Placebo Glycosamino- glycan Polysulphate	DSM III	PDD MID	Mod-Sev	155	148	73.5 y (56-98 y) 57% M Hispanic, Italian 30% Community 70% Institution	200 LRU tid	12w	ADL BPRS CGI-GI CGI-SI CT EEG EKG GDS HDS HIS MMSE SCAG WMS-RR	No

EvTable207. Key characteristics: Other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population (Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Ala 1990	PI	5	Placebo GM-1 Monosialotetrahexosylgan	NINCDS	AD	Mild-Mod	46	42	71.0 y (63-79 y) 39% M	100 mg/d	12w	BCRS Benton Racial Recognition Test BPRS Buschke Selective Reminding Test Clock drawing Complex Figure Test Grooved pegboard test HAM-D IADL Letter cancellation Lipids MMSE NOSIE Selective Reminding Symbol-Digit Modalities test Verbal fluency	No
Crook 1992b	NR	5	Placebo Guanfacine	NINCDS DSM III	AD PDD	Mild-Mod	29	26	71.0 y (60-81y) 45% M Community	0.5 mg/d	13w	Benton Visual Retention CGI variables MMSE Neuropsychiatric rating scale WAIS Vocabulary Wechsler Paired Assoc.	No
Thompson 1990	IF	7	Placebo Hydergine-LC	DSM III NINCDS	PDD AD	Probable	80	68	71.0 y (55-79 y) 59% M	1 mg tid	24w + 2w wash out	GERRI Ham-D IPSCE SCAG WMS WAIS-DSST	No

EvTable207. Key characteristics: Various other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total # Randomized	# Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Van Gool 2001	NI IS	8	Placebo Hydroxychloro- quine	NINCDS	AD	Min-Mild	168	155	70.6 y (NR) 42% M	400 mg/d if ≥ 65 kg 200 mg/d if ≤ 65 kg	18m	ADAS-Cog CAMDEX IDDD RMBPC	No
Rogers 1993	NR	5	Placebo Indomethacin	NINCDS	AD	Mild-Mod	44	28	78.0 y (NR) 54% M Community	100 mg/d if ≤ 100 lbs 125/d if 101-150 lbs 150 mg/d if ≥ 150 lbs	6m	ADAS BNT MMSE TK	No
Adair 2001	NI	7	Placebo N- acetylcysteine	NINCDS	AD	Mild-Mod	47	43	NR Community	50 mg/kg/d	6m	ADL BNT CT Scan MMSE Neuropsychological Battery	No
Aisen 2002	NI	7	Placebo Nimesulide	DSM III	AD	Probable	40	38	74.0 y (NR) 58% M	100 mg bid	12w	ADAS-Cog ADL BPRS CDR HAM-D MMSE	No

EvTable207. Key characteristics: Various other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Ban 1990	IS	7	Placebo Nimodipine	DSM III	PDD MID	Mild- Modly Sev	178	175	75.4 y (55-95 y) 42% M Italian, 19% Community 79% Institution	90 mg/d	12w	CBC & platelet count CGI Severity of illness CGI-global improvement EEG GDS HAM-D HIS Laboratory tests MMSE Neurologic Evaluation Plutchik Geriatric Rating Scale SCAG Wechsler Memory Scale	No
Pantoni 2000a Auxiliary: Pantoni 2000	IF	6	Placebo Nimodipine	DSM-III-R	MID	Mild-Mod	259	251	74.3 y (45-85 y) 47% M White, Community & Institution	30 mg tid	26w	ADL CDR CGI Digit Span Fuld Object Memory GBS IADL MMSE RDS Word Fluency ZVT-G	VaD vs MID

EvTable207. Key characteristics: Various other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Carlson 2001 Auxiliary: Breitner 1999	IS	7	Placebo Nizatidine	DSM-III- R NINCDS NINDS- AIREN	AD VaD	Probable – Possible	51	42	80.7 y (67-96 y) 38% M	75 mg bid	1y	Benton Visual Retention Test Boston Naming Test Category Fluency CERAD Constructional Praxis COWA CPT IADL Logical Memory I & 2 MMSE MMSE Trail-Making Test WLM WMS WMS-R	No
Kragh- Sorensen 1986	NI	6	Placebo ORG 2766	DSM III	PDSD	Mild-Mod	156	126	82.5 y (>65 y) 23% M	80 mg bid	28d	GAGS Laboratory tests LPRS RDS SCAG	No

EvTable207. Key characteristics: Various other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Soininen 1985 Auxiliary Partanen 1986 Soininen 1984	IS	7	Placebo ORG 2766	DSM III	AD	Mild-Sev	77	73	73.2 y (NR) 32% M Institution	20 mg bid	6m	AGS-E GPI-E LPRS McGBRS SCAG	No
Aisen 2000b Auxiliary: Aisen 2000a	NI IS	7	Placebo Prednisone	NINCDS	AD	Mild-Mod	138	92	72.3 y (>50 y) 69% M	20 mg/d (week 1-4) 10 mg/d (week 5-end)	1y	ADAS-Cog BDRS BPRS CDR-SB HAM-D MMSE	No
Simmons 2002	PI IS	7	Placebo Simvastatin	NINCDS	AD	Mild-Mod	44	37	68.3 y (60-77 y) 43% M	80 mg/d	26w	ADAS-cog Laboratory tests MMSE	MMSE

EvTable207. Key characteristics: Various other agents cont'd.

Author Y ear	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Nolan 1991	NI	5	Placebo Thiamine	NINCDS	AD	Mild-Mod	15	10	76.3 y (59-87 y) 33% M Community	3 g/day	12m	Boston Naming test CERAD Constructional praxis test Delayed recall and recognition MMSE Verbal learning score Word list learning test	No
Fischhof 1996	NR	5	Placebo Vincamine	DSM-III-R	PDD VaD	Mild-Mod	152	142	Mean NR (50-85 y) % M NR Institution	30 mg bid	12w	BGP CGI SCAG SKT	MID vs DAT
Croisile 1993	IF	4	Placebo Piracetam	NINCDS	AD	Mild-Mod	33	30	42% M Community	8 gr/day per oz	1y	Aphasia Battery Blessed A CT EEG Laboratory tests Logical Digit Span MADRS MMSE Neuropsychological Battery	No

EvTable207. Key characteristics: Various other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Kanowski 1990	PI	8	Placebo Xantinol nicotinate	DSM-III-R	AD MID	Mild-Mod	313	297	82.0 y (≥60 y) 19% M	1g tid	12w	BGP CGI NAI SCAG Digit Correction Test Digit Symbol Substitution Test	MID vs DAT
Thomas 2001	NR	7	Donepezil Vitamin E Rivastigmine (open label)	NINCDS	AD	Mild-Mod	60	54	66.1y (57-78 y) 44% M	Donepezil: 5 mg/d (1 month) 10 mg/d (until end) Vitamin E: 2000 IU (fixed)	6m	ADAS-cog CT/MRI ERP scalp topography GBS GDS MMSE NPI WAIS	No
Taragano 1997	NR	7	Fluoxetine Amitriptyline	NINCDS DSM III	AD	Mild-Mod	37	25	72.0 y (NR) 22% M Community, Major depression	10 mg/d 25 mg/d	45d	HAM-D MMSE	No
Passeri 1993	NR	5	5'-MTHF (folate) Tradozone	DSM III-R	AD MID	Mild-Mod	96	96	Mean NR (65-94 y) 45% M Depression	50 mg/d 100 mg/d	8w	Blood levels HDRS RVM – immediate recall RVM – delayed recall	AD vs MID

EvTable207. Key characteristics: Various other agents cont'd.

Author Year	Funding Source	Quality Score	Interventions	Criteria for Diagnosis	Diagnosis	Disease Severity	Total Number Randomized	Number Completing Trial	Mean age (range) % Male (M) Population	Highest Dose	Treatment Period	Outcomes Measured	Outcome reports stratified
Parnetti 1995	NR	6	Placebo Posatirelin Citicoline	NINCDS	AD	Mild-Mod	222	214	74.9 y (65-85 y) 34% M 100% Community	10 mg/d 500 mg/d	3m	GBS GDS HDRS MMSE Laboratory tests	No
Spilich 1996	PI	6	Pyritinol Hydergine	NINCDS	AD	Mild-Mod	102	100	73.0 y (NR) 31% M 100% Hispanic	600 mg/d 4.5 mg/d	12w	CETM SCAG	No
Cucinotta 1988	PI	5	Sulfomuco- polysaccharid es CDP-choline	DSM III	MID	Mild-Mod	30	23	79.4 y (NR) 27% M 100% Institution	500 LRU 1000 mg	4w	Blessed-D Digit-Span Digit-Symbol NMS SCAG TP	No
Parnetti 1997	NR	5	Sulodexide Pentoxifylline	NINDS- AIREN	VaD	Mild-Mod	93	86	75.0 y (65-80 y) 40% M Institution	100 mg/d 1200 mg/d	6m	GBS Laboratory tests MMSE	No
Sano 1997 Auxiliary: Thal 1996	NI IS	5	Placebo Vitamin E Selegiline Selegiline + VitaminE	NINCDS	AD	Mod	341	341	73.4 y (NR) 35% M	Vitamin E 1000 IU bid Selegiline 5mg bid	2y	ADAS-Cog Blessed Dementia Scale CDR MMSE Time to end-point (event free survival)	No

EvTable208. Study results: Aniracetam

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4m		Final: (specify) 6m	
Senin 1991	OC Analysis 1] Placebo 2] Aniracetam 1500 mg/d 3] Placebo time x baseline 4] Aniracetam time x baseline 5] Placebo vs. Aniracetam	SCAG total	1] 48.2 (1.3)* 2] 50.5 (0.9)*		1] 52.2 (1.6)* 2] 42.3 (1.5)*	3] <0.01 4] <0.001 5] <0.001	1] 56.5 (2.3) 2] 39.4 (1.5)	3] <0.01 4] <0.001 5] <0.001
		Blessed I	1] 7.2 (0.4)* 2] 7.9 (0.4)*		1] 7.9 (0.4)* 2] 6.3 (0.5)*	3] <0.001 4] <0.001 5] <0.02	1] 8.8 (0.5) 2] 5.7 (0.5)	3] <0.001 4] <0.001 5] <0.001
		Blessed II	1] 22.9 (0.5)* 2] 21.6 (0.6)*		1] 21.5 (0.8)* 2] 24.4 (0.8)*	3] <0.001 4] <0.001 5] <0.01	1] 20.2 (1) 2] 26.1 (0.6)	3] <0.001 4] <0.001 5] <0.001
		Toulouse-Pieron (global)	1] 0.4 (0.1)* 2] 0.3 (0.08)*		1] 0.1 (0.1)* 2] 0.7 (0.08)*	3] <0.02 4] <0.02 5] <0.001	1] 0.2 (.01) 2] 0.8 (.01)	3] <0.02 4] <0.001 5] <0.001
		Rey immediate recall	1] 18.4 (0.9)* 2] 16.9 (0.9)*		1] 16.2 (0.9)* 2] 20.8 (1)*	5] <0.001	1] 16.2 (1.1) 2] 23.4 (1)	3] 0.001 4] <0.001 5] <0.001
		Rey delayed recall	1] 1.9 (0.2)* 2] 1.8 (0.2)*		1] 1.3 (0.2)* 2] 2.6 (0.3)*	5] <0.02	1] 1.6 (0.3) 2] 2.9 (0.4)	3] <0.001 4] <0.001 5] <0.02

*SEM

EvTable209. Study results: Ateroid.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Ban 1991b	OC Analysis							
	1] Placebo	SCAG total					1] 17.3 2] 24.1	3] <0.05
	2] Ateroid 200LRU tid	SCAG cognitive					1] 13.0 2] 19.2	3] <0.05
	3] Ateroid vs Placebo	SCAG agitation					1] 17.77 2] 25.4	3] NS
		SCAG overall impression					1] 13.5 2] 20.2	3] <0.10
		SCAG depression					1] 22.0 2] 27.5	3] <0.10
		HDS						3] <0.04

EvTable210. Study results: BMJ 21,502.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Shrotrya, 1996	OC Analysis 1] Placebo	ADAS-Cog	1] 16.2 2] 15.1	3] 0.81			4] -0.5 5] -1.5	3] >0.05
	2] BMJ 21,502 300 mg tid	CGI severity	1] 3.49 2] 3.29	3] 0.11				3] NS
	3] BMJ 21,502 300 mg tid vs Placebo	MMSE	1] 22.5 2] 23.5	3] 0.26				
	4] Placebo changes from baseline	CGI % Improvement					1] 28% 2] 37%	3] >0.05
	5] BMJ changes from baseline							

EvTable211. Study results: Buflomedil.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 90d		Final: (specify) 270d	
Cucinotta, 1992	OC Analysis	SHGRS	1] 43.00		1] 37.00		1] 29.00	
			2] 35.50		2] 30.50		2] 30.00	
			3] 29.00		3] 25.40		3] 23.75	
			4] 41.00		4] 39.00		4] 34.00	
		Median Rating scale & Neuro- psychological test scores	1] 16.00		1] 10.75		1] 6.25	
			2] 18.50		2] 16.50		2] 11.50	
			3] 21.50		3] 9.50		3] 7.00	
			4] 29.00		4] 22.50		4] 22.00	
		Dementia Rating scale	1] 39.00		1] 28.00		1] 15.25	
			2] 22.50		2] 18.50		2] 17.50	
			3] 27.00		3] 22.00		3] 15.75	
			4] 37.25		4] 30.25		4] 28.00	
		Birren test	1] 1.19		1] 1.64			
			2] 1.38		2] 2.50			
			3] 1.83		3] 1.93			
			4] 1.27		4] 1.50			
		Nowlis Mood test	1] 13.38		1] 12.12			
			2] 13.17		2] 12.00			
			3] 15.83		3] 12.50			
			4] 15.50		4] 15.75			
		Clinical Global Impression	1] 0.25		1] 0.82		1] 1.33	
			2] 0.00		2] 1.00		2] 0.67	
			3] 0.00		3] 0.12		3] 0.67	
			4] 0.03		4] 0.05		4] 0.00	

EvTable212. Study results: CBM 36-733 (Ergokryptine).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 8w	
Danielczyk, 1997	OC Analysis 1] Placebo 2] CBM 36-733 2.0 mg/d 3] Differences between groups	SCAG- Overall Impression	1] 5.0 2] 5.0				1] 5.0 2] 4.0	3] 0.016
		SCAG- Cognitive Dysfunction	1] 18.0 2] 16.0				1] 17.0 2] 15.0	3] 0.276
		SCAG- Interpersonal Relationship	1] 12.0 2] 10.0				1] 11.0 2] 8.0	3] 0.421
		SCAG-Apathy	1] 14.0 2] 15.0				1] 14.0 2] 13.0	3] 0.011
		SCAG- Affect	1] 10.0 2] 9.0				1] 10.0 2] 8.0	3] 0.385
		SCAG- Somatic Dysfunction	1] 8.0 2] 7.0				1] 7.0 2] 7.0	3] 0.679
		Psychometric Test Battery -tests with significant difference					2/9	

EvTable213. Study results: Choto-san.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 8w		Final: (specify) 12w	
Shimada 1994	OC Analysis 1] Placebo 2] Choto-san 2.5g tid 3] Difference between Placebo and Choto-san 4] Choto-san vs baseline	Global Improvement rating (% improved)			1] 19% 2] 61%	3] <0.01	1] 23% 2] 71%	3] <0.01
		Utility rating (% useful)					1] 25% 2] 78%	3] <0.01
		Global improvement -subjective symptoms			1] 12% 2] 59%	3] <0.01	1] 21% 2] 55%	3] <0.01
		-neurological symptoms			1] 13% 2] 5%	3] NS	1] 25% 2] 14%	3] NS
		-psychiatric symptoms			1] 12% 2] 58%	3] <0.01	1] 24% 2] 58%	3] <0.01
		Disturbance in daily living			1] 12% 2] 29%		1] 19% 2] 38%	3] <0.05
		Hasegawa dementia	1] 15 (3.84) 2] 15 (3.76)	4] NS	1] 16 (5.82) 2] 18 (4.79)	4] <0.01	1] 17 (5.97) 2] 19 (5.71)	4] <0.01

EvTable214. Study results: Choto-san.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 8w		Final: (specify) 12w	
Terasawa 1997	OC Analysis 1] Placebo 2] Choto-san 2.5g tid 3] difference between Placebo and Choto-san	Global severity rating %improved			1] 42.9% 2] 70.9%	3] 0.001	1] 48.4% 2] 34.4%	3] <0.001
		HDS-R			1] 17.3(5.3) 2] 18.0(6.4)	3] NS	1] 17.4(6.0) 2] 19.3(6.6)	3] NS
		Utility Rating (% Useful)					1] 33% 2] 44%	3] <0.001

EvTable215. Study results: Cyclandelate.

Author Year	Analysis Groups	Ooutcomes Meassured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 8w		Final: (specify) 16w	
Schellenberg 1997	ITT Analysis 1] Placebo 2] Cyclandelate 1,600 mg d 3] Cyclandelate vs. placebo	SCAG	1] 44.0 (39.0, 53.0) 2] 39.0 (35.0, 51.0)		1] 36.0 (28.0, 53.0) 2] 33.0 (28.0, 38.0)		1] 38.0 (29.0, 67.0) 2] 32.0 (26.0, 37.0)	3] 0.0004
		ADAS	1] 24.0 (15.0, 39.0) 2] 20.0 (14.0, 37.0)		1] 20.0 (13.0, 34.0) 2] 17.0 (9.0, 22.0)		1] 17.0 (11.0, 42.0) 2] 16.0 (8.0, 20.0)	3] 0.0006
		NCT	1] 16.5 (8.0, 35.0) 2] 20.2 (10.2, 32.2)		1] 25.8 (10.8, 51.2) 2] 37.2 (22.6, 54.0)		1] 34.9 (11.4, 52.8) 2] 26.4 (19.4, 53.4)	3] >0.2
		NST	1] 39.0 (19.0, 59.0) 2] 48.0 (27.0, 57.0)		1] 50.0 (23.0, 65.0) 2] 56.0 (39.0, 68.0)		1] 54.0 (27.0, 65.0) 2] 59.5 (36.0, 66.0)	3] 0.0051
		FIGT	1] 14.5 (7.0, 34.0) 2] 22.0 (13.0, 34.0)		1] 25.5 (11.0, 45.0) 2] 33.5 (18.0, 49.0)		1] 30.0 (11.0, 45.0) 2] 31.5 (18.0, 45.0)	3] >0.2
		MEMT	1] 17.0 (12.0, 19.0) 2] 18.0 (14.0, 19.0)		1] 18.0 (14.0, 19.0) 2] 18.0 (17.0, 19.0)		1] 18.0 (15.0, 19.0) 2] 18.0 (16.0, 19.0)	3] >0.2
		NSL	1] 53.0 (45.0, 61.0) 2] 49.0 (43.0, 58.0)		1] 51.0 (38.0, 63.0) 2] 41.0 (36.0, 50.0)		1] 55.0 (39.0, 69.0) 2] 41.0 (27.0, 49.0)	3] .000002

EvTable216. Study results: Cyclandelate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24w	
Weyer, 2000	ITT Analysis 1] Placebo change from baseline 2] Cyclandelate 800 mg bid change from baseline 3] Difference between Placebo and Cyclandelate in change from baseline	<u>ADAS-Cog% responders < -10 points</u>					1] 12.8% 2] 18.6%	3] NS
		<u>NOSGER-IADL % responders < -5 points</u>					1] 5.3% 2] 10.3%	3] NS
		<u>CGI-C % responders at least minimal improvement</u>					1] 51.1% 2] 58.8%	3] NS
		ADAS-Cog					1] -1.5 (7.4) 2] -2.7 (8.8)	3] 0.320
		NOSGER-IADL					1] -0.2 (2.7) 2] -0.6 (3.9)	3] 0.181

EvTable217. Study results: Desamino-D-Arginine-Vasopressin (DDAVP).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 3w	
Peabody 1986	OC Analysis 1] Placebo 2] DDAVP 45mg qid 3] DDAVP vs Placebo	HAM-D	1] 4.4 (4.6) 2] 3.2 (3.6)				1] 3.6 (2.8) 2] 4.5 (3.9)	3] 0.02
		SCAG affect	1] 4.1 (1.3) 2] 4.6 (1.8)				1] 4.4 (1.2) 2] 4.7 (2.6)	3] 0.01
		SCAG interpersonal	1] 4.8 (1.2) 2] 4.6 (1.0)				1] 4.5 (0.4) 2] 5.0 (1.5)	3] 0.02
		SCAG 19 (overall)	1] 4.4 (0.7) 2] 3.4 (1.0)				1] 4.5 (0.6) 2] 3.3 (1.1)	3] 0.06
		SCAG total	1] 36 (5) 2] 29 (6)				1] 35 (3) 2] 30 (65)	3] NS
		Buschke total recall	1] 6.1 (1.8) 2] 7.4 (2.1)				1] 5.8 (2.0) 2] 7.7 (2.5)	3] NS
		POMS Vigor	1] 55 (9) 2] 59 (7)				1] 56 (9) 2] 63 (6)	3] NS
		POMS Fatigue	1] 42 (5) 2] 42 (5)				1] 44 (6) 2] 41 (5)	3] NS
		POMS Depression	1] 37 (4) 2] 38 (5)				1] 37 (3) 2] 38 (5)	3] NS

EvTable218. Study results: Dihydroergokryptine (DEK).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6m		Final: (specify) 12m	
Cucinotta, 1996	Endpoint Analysis 1] Placebo	<u>GBS Factor</u> <u>1</u>	1] 8.7 (4.4) 2] 9.6 (4.3)		1] 9.4 (4.4) 2] 8.4 (4.2)		1] 10.5 (4.8) 2] 8.0 (4.8)	3] 0.000
Cucinotta, 1998	2] DEK 20mg bid	<u>GBS Factor</u> <u>2</u>	1] 5.2 (6.0) 2] 4.9 (5.7)		1] 5.4 (6.2) 2] 4.1 (4.6)		1] 7.2 (7.7) 2] 3.8 (4.7)	3] 0.001
	3] DEK 20mg bid vs Placebo	<u>GBS Factor</u> <u>3</u>	1] 3.5 (2.7) 2] 3.3 (2.4)		1] 3.1 (2.3) 2] 2.7 (2.3)		1] 3.0 (2.8) 2] 2.5 (2.3)	3] 0.229
		<u>GBS Factor</u> <u>4</u>	1] 13.1 (7.5) 2] 12.6 (7.6)		1] 13.5 (7.8) 2] 11.1 (6.7)		1] 15.3 (7.5) 2] 10.3 (6.7)	3] 0.000
		MMSE	1] 19.6 (2.6) 2] 19.7 (2.6)	3] 0.645			1] 8% reduction 2] same mean	3] <0.001
		HAM-D	1] 9.1 (4.9) 2] 9.4 (4.7)	3] 0.621				
		REY Test Immediate Recall	1] 14.4 (7.0) 2] 14.3 (7.9)		1] 12.2 (7.2) 2] 14.2 (8.3)		1] 11.7 (7.0) 2] 14.5 (8.1)	3] 0.022
		REY Test Delayed Recall	1] 3.0 (2.9) 2] 3.4 (2.5)		1] 3.0 (3.0) 2] 3.8 (3.4)		1] 2.4 (2.7) 2] 4.0 (3.7)	3] 0.044
		GBS-C	1] 4.7 (2.8) 2] 4.9 (5.7)		1] 4.9 (3.3) 2] 4.1 (2.6)		1] 5.4 (3.3) 2] 3.6 (2.6)	3] 0.004
		GBS – M	1] 5.2 (6.0) 2] 4.9 (5.7)		1] 5.4 (6.2) 2] 4.1 (4.6)		1] 7.2 (7.7) 2] 3.8 (4.7)	3] 0.001
		GBS - I	1] 18.8 (8.9) 2] 19.1 (9.2)		1] 19.8 (8.8) 2] 16.8 (8.2)		1] 22.3 (8.9) 2] 16.6 (9.0)	3] 0.000

EvTable219. Study results: Denbufylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 16w	
Treves, 1999	OC Analysis 1] Placebo 2] Denbufylline 25 mg bid 3] Denbufylline 50 mg bid 4] Denbufylline 100 mg bid 5] Denbufylline Overall doses 6] Treatment effect vs Placebo	<u>MMSE</u>	1] 17.3 (4.2)				1] 16.9 (5.1)	6] 0.7
			2] 18.4 (4.6)				2] 19.5 (6)	
			3] 18.6 (4.7)				3] 19.1 (5.7)	
			4] 18.2 (4.8)				4] 19.1 (5.5)	
			5] 18.4 (4.7)				5] 19.3 (5.7)	
		<u>DSST</u>	1] 6.6 (6.8)				1] 7.1 (7.1)	6] NS
			2] 7.2 (6.5)				2] 9.4 (9)	
			3] 8.5 (8.6)				3] 9.7 (8.6)	
			4] 7.5 (5.4)				4] 8.3 (7.6)	
			5] 7.7 (6.9)				5] 9.2 (8.5)	
		WAIS-voc	1] 28.9 (15.5)				1] 26.1 (16)	6] NS
			2] 28.8 (17.2)				2] 30.9 (19.6)	
			3] 26.6 (17.1)				3] 29.4 (19.3)	
			4] 27.7 (16.1)				4] 29.6 (19.4)	
			5] 27.7 (16.8)				5] 30.0 (19.4)	

EvTable220. Study results: Desferrioxamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 18m	
Crapper McLachlan 1991	OC Analysis 1] Placebo slope of change from baseline 2] Desferrioxamine 500 mg BID slope of change from baseline 3] Difference from baseline between Desferrioxamine and Placebo	<u>VHB</u>	1] -1.717 (1.689)				1] -0.866 (0.932) 3] 0.0375	

EvTable221. Study results: Diclofenac and Misoprostol (D/M).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 25w	
Scharf 1999	ITT Analysis 1] Placebo 2] Diclofenac 50mg and Misoprostol 200µg (D/M) 3] Placebo mean change from baseline 4] D/M group mean change from baseline 5] mean treatment differences placebo vs. D/M	ADAS-cog					3] 1.93 (5.55) 4] 0.25 (4.50)	5] 0.571
		GDS					3] 0.57 (0.51) 4] 0.35 (0.49)	5] 0.384
		CGIC					3] 4.57 (0.51) 4] 4.29 (0.69)	5] 0.340
		MMSE					3] -0.86 (3.21) 4] 0.41 (2.69)	5] 0.237
		ADAS-noncog					3] 1.36 (3.93) 4] -0.59 (3.89)	5] 0.319
		ADAS-Total					3] 3.24 (8.85) 4] -0.75 (1.34)	5] 0.125
		IADL					3] 1.86 (2.03) 4] 0.06 (2.95)	5] 0.161
		PSMS					3] 0.21 (0.89) 4] 0.53 (1.84)	5] 0.340
		CGIC-C Caregiver rated					3] 4.79 (1.05) 4] 4.47 (1.01)	5] 0.768

EvTable222. Study results: Glycoaminoglycan.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Ban, 1991a	OC Analysis 1] Placebo 2] Glycoaminoglycan polysulphate 200 LRU tid 3] Improvement in scores Glycoaminoglycan polysulphate 200 LRU tid vs. placebo	CGI- SI	1] 4.6 (0.69) 2] 4.6 (0.63)				1] 4.0 (0.89) 2] 3.7 (0.77)	3] <0.18
		CGI –GI	1] 3.8 (0.41) 2] 3.7 (0.71)				1] 3.5 (0.71) 2] 3.0 (0.78)	3] <0.001
		BPRS Total scores	1] 31.4 (11.75) 2] 28.9 (12.12)				1] 25.2 (10.73) 2] 20.6 (10.52)	3] <0.01
		ADL	1] 32.9 (9.10) 2] 33.0 (9.77)				1] 32.2 (9.11) 2] 30.5 (9.57)	3] <0.16
		SCAG	1] 56.6 (10.66) 2] 56.1 (13.47)				1] 46.8 (11.78) 2] 42.6 (12.72)	3] <0.05
		MMSE	1] 14.3 (4.64) 2] 14.9 (4.93)				1] 16.2 (5.16) 2] 18.2 (6.01)	3] <0.04
		GDS	1] 4.8 (0.77) 2] 4.8 (0.72)				1] 4.4 (0.95) 2] 4.3 (0.93)	3] <0.50
		HDS	1] 11.8 (3.87) 2] 12.1 (3.98)				1] 11.1(4.28) 2] 10.6 (4.53)	3] <0.04
		WMS-RR PAL	1] 5.9 (3.98) 2] 6.8 (4.7)				1] 5.9 (4.79) 2] 7.7 (4.93)	3] <0.081
		Logical Memory	1] 1.4 (1.42) 2] 1.8 (1.54)				1] 1.5 (1.46) 2] 2.1 (2.01)	3] <0.18
		Immediate Recall	1] 1.5 (1.38) 2] 1.6 (1.55)				1] 1.3 (1.24) 2] 2.0 (2.01)	3] <0.03
		Delayed Recall	1] 0.5 (.22) 2] 0.8 (.23)				1] 0.8 (.36) 2] 1.1 (.31)	3] NS

EvTable223. Study results: Monosialotetrahexosylgan (GM-1).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 12w		Final: (specify) 24w	
Ala 1990	OC Analysis 1] Placebo 2] GM-1 100mg IM d 3] Placebo vs. GM-1 4] Placebo change from baseline 5] GM-1 change from baseline	HAM-D	1] 7.04 (4.01) 2] 5.58 (4.07)		1] 6.61 (4.19) 2] 5.79 (4.61)		1] 6.43 (2.90) 2] 7.21 (4.20)	3] NS 4] >0.05 5] >0.05
		PSM	1] 8.13 (1.63) 2] 8.32 (1.42)		1] 8.09 (1.76) 2] 8.16 (1.80)		1] 9.22 (2.39) 2] 9.42 (2.67)	3] NS 4] 0.032 5] 0.049
		IADL	1] 17.1 (3.2) 2] 18.5 (3.7)		1] 17.4 (3.5) 2] 19.5 (4.1)		1] 18.4 (3.6) 2] 20.6 (4.7)	3] NS 4] 0.012 5] 0.027
		BPRS Total score	1] 8.65 (5.47) 2] 6.42 (3.81)		1] 8.65 (6.32) 2] 8.74 (7.49)		1] 9.22 (4.00) 2] 9.79 (6.33)	3] NS 4] >0.05 5] 0.024
		NOSIE Total score	1] 178 (26) 2] 184 (15)		1] 177 (28) 2] 184 (21)		1] 169 (32) 2] 173 (28)	3] NS 4] 0.031 5] 0.036
		MMSE	1] 17.5 (3.2) 2] 17.5 (3.3)		1] 17.7 (4.4) 2] 18.3 (4.0)		1] 17.0 (4.9) 2] 16.5 (4.5)	3] NS
		BCRS	1] 19.7 (3.3) 2] 19.7 (2.4)		1] 18.9 (2.7) 2] 19.2 (2.6)		1] 19.6 (3.2) 2] 20.3 (2.9)	3] NS
		Complex figure – copy accuracy	1] 10.8 (7.2) 2] 11.6 (7.4)		1] 10.7 (6.9) 2] 10.6 (6.5)		1] 9.8 (6.2) 2] 10.7 (7.3)	3] NS

EvTable223. Study results: Monosialotetrahexosylgan (GM-1) cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify) 12w		Final: (specify) 24w	
			Verbal fluency	1] 25.1 (9.6) 2] 25.6 (11.1)		1] 26.0 (10.3) 2] 24.4 (11.2)		1] 23.6 (9.4) 2] 22.4 (10.0)	3] NS
			Selective reminding-Retrieval	1] 7.30 (6.96) 2] 9.58 (8.31)		1] 6.35 (6.11) 2] 8.42 (5.69)		1] 9.52 (9.40) 2] 8.95 (6.79)	3] NS
			Symbol digit	1] 11.9 (11.1) 2] 14.6 (8.3)		1] 13.9 (10.8) 2] 15.8 (6.3)		1] 13.8 (10.5) 2] 10.6 (7.4)	3] NS 5] 0.017
			8 other cognitive tests						3] NS

EvTable224. Study results: Guanfacine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 13w	
Crook, 1992b	ITT Analysis	CGI						5] <.05
	1] Placebo							
	2] Guanfacine .5mg/d	Wechsler Paired Associates	1] 6.54 (2.04) 2] 6.90 (2.94)				3] 6.08 (1.77) 4] 4.97 (2.72)	
	3] Placebo change from baseline	Benton Visual Retention- Number Correct	1] 2.69 (1.49) 2] 3.20 (1.66)				3] 2.15 (1.57) 4] 1.67 (1.18)	
	4] Guanfacine .5mg/d change from baseline							
	5] Guanfacine .5mg/d change from baseline relative to placebo	Benton Visual Retention- Errors	1] 15.31 (4.48) 2] 12.73 (6.22)				3] 15.69 (5.06) 4] 17.80 (4.28)	5] <.03

EvTable225. Study results: Hydergine-LC.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 24w	
Thompson 1990	OC Population 1] Placebo change from baseline 2] Hydergine-LC 1mg tid change from baseline 3] Placebo vs Hydergine-LC change from baseline	WAIS DSST					1] 1.76(1.64)* 2] 8.85(2.07)*	3] <0.01
		WMS- Logical					1] -0.02(0.23)* 2] -0.21(0.23)*	3] >0.30
		WMS-Visual					1] -0.07(0.12)* 2] -0.10(0.10)*	3] >0.50
		GERRI					1] 0.04(0.07)* 2] -0.23(0.09)*	3] <0.02
		IPSCE					1] -0.11(1.36)* 2] -1.43(1.65)*	3] >0.50
		HAM-D					1] 0.46(0.45)* 2] 0.39(0.39)*	3] >0.50
		SCAG					1] -3.60(1.35)* 2] -2.37(1.69)*	3] >0.50

*SEM

EvTable226. Study results: Hydroxychloroquine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 9m		Final: (specify) 18m	
Van Gool, 2001	ITT Analysis							
	1] Placebo	IDDD	1] 10.9 (7.0) 2] 11.6 (7.1)	3] NS	1] 15.9 (9.2) 2] 17.0 (10.0)	3] NS	1] 21.3 (10.5) 2] 22.6 (11.4)	3] NS
	2] Hydroxchloroquine 200 or 400 mg QID	ADAS-Cog	1] 17.6 (9.1) 2] 18.0 (9.4)	3] NS	1] 20.0 (9.70) 2] 21.7 (12.8)	3] NS	1] 25.7 (14.3) 2] 26.4 (14.9)	3] NS
	3] Difference between Placebo and Hydroxchloroquine	RMBPC	1] 27.8 (10.8) 2] 28.6 (11.0)	3] NS	1] 30.2 (11.7) 2] 32.0 (11.5)	3] NS	1] 34.2 (12.4) 2] 36.3 (12.0)	3] NS

EvTable227. Study results: Indomethacin.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Rogers 1993	Completer Analysis 1] Placebo % change from baseline 2] Indomethacin 150mg/d % change from baseline 3] Difference between Placebo and Indomethacin change from baseline	ADAS					1] -13.3 (5.6)* 2] 1.4 (4.9)*	3] 0.061
		MMSE					1] -13.4 (4.4)* 2] -0.9 (4.8)*	3] 0.069
		BNT					1] -6.6 (5.5)* 2] 4.4 (3.7)*	3] 0.120
		TK					1] -0.4 (2.9)* 2] 0.5 (1.0)*	3] 0.773
		Overall all tests					1] -8.4 (2.3)* 2] 1.3 (1.8)*	3] <0.003

*SEM

EvTable228. Study results: N-acetylcysteine (NAC).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3m		Final: (specify) 6m	
Adair 2001	ITT Analysis 1] Placebo 2] NAC 50 mg/kg/d 3] Difference between Placebo and NAC	<u>MMSE</u>	1] 18.0 (3.6) 2] 19.8 (3.7)		1] 17.5 (3.6) 2] 20.5 (4.7)	3] 0.056	1] 16.8 (4.6) 2] 19.8 (5.3)	3] NS
		<u>ADL</u>	1] 16.2 (5.0) 2] 14.2 (4.3)		1] 18.5 (4.3) 2] 15.2 (4.7)		1] 20.1 (4.9) 2] 16.1 (5.0)	3] NS
		BNT	1] 18.9 (6.5) 2] 20.9 (5.6)		1] 19.6 (6.0) 2] 21.1 (5.3)		1] 18.0 (6.9) 2] 21.2 (6.2)	3] NS

EvTable229. Study results: Nimesulide.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Aisen 2001	ITT Analysis							
	1] Placebo	MMSE	1] 22.7 (1.0)* 2] 21.8 (1.1)	6] 0.54				
	2] Nimesulide 100mg bid	ADAS-Cog	1] 21.4 (2.6)* 2] 19.8 (1.5)*	6] 0.59			3] -0.5 (1.0)* 4] 0.9 (1.0)*	5] 0.49
	3] Placebo change from baseline	CDR -SB	1] 4.6 (0.5)* 2] 4.7 (0.7)*	6] 0.91			3] 0.2 (0.3)* 4] 0.7(0.3)*	5] 0.70
	4] Nimesulide change from baseline	ADL	1] 3.8 (0.5)* 2] 3.3 (.03)*	6] 0.34			3] -0.3 (0.2)* 4] -0.2 (0.3)*	5] 0.73
	5] Nimesulide change from baseline relative to placebo	HAM-D	1] 3.8 (0.8)* 2] 4.6 (0.8)*	6] 0.52			3] 1.0 (0.6)* 4] -0.2 (0.9)*	5] 0.30
	6] Placebo vs Nimesulide	BPRS	1] 29.6 (1.4)* 2] 28.5 (1.5)*	6] 0.59			3] 0.4 (0.9)* 4] 0.4 (1.4)*	5] 0.99

*SEM

EvTable230. Study results: Nimodipine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 90d	
Ban, 1990	OC Analysis 1] Placebo 2] Nimodipine 30 mg bid 3] Nimodipine 30 mg bid vs. placebo	CGI-SI	1] 3.8 (1.44) 2] 3.8 (1.51)				1] 3.6 (1.3) 2] 3.2 (1.28)	3] <0.018
		CGI-GI	1] 3.7 (0.59) 2] 3.3 (0.77)				1] 3.3 (.86) 2] 2.7 (0.86)	3] <0.001
		HAM-D total	1] 13.1 (6.20) 2] 14.4(7.67)				1] 12.1 (6.42) 2] 10.4 (5.63)	3] <0.001
		MMSE total	1] 18.4 (5.57) 2] 17.6 (5.47)				1] 19.2 (5.74) 2] 21.0 (5.14)	3] <0.001
		GDS total	1] 4.0 (.79) 2] 4.1 (.83)				1] 3.8 (.90) 2] 3.5 (.85)	3] <0.001
		SCAG total	1] 53.0 (14.40) 2] 57.5 (15.70)				1] 46.9 (14.47) 2] 44.8 (12.79)	3] <0.001
		PLUTCHIK Total	1] 19.2 (7.14) 2] 20.3 (6.87)				1] 17.6 (6.26) 2] 16.9 (7.41)	3] <0.013
		WMS	1] 74.9 (16.67) 2] 71.6 (13.19)				1] 80.7 (19.07) 2] 85.4 (18.72)	3] <0.001

EvTable231. Study results: Nimodipine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point:		Final: 26w	
Pantoni, 2000a	ITT Analysis	<u>GBS total</u>	1] 0.85 (0.55) 2] 0.88 (0.58)				1] 0.23 (0.49) 2] 0.85 (0.55)	3] 0.67
		<u>GBS-i</u>	1] 1.06 (0.65) 2] 1.13 (0.73)				1] 0.25 (0.58) 2] 0.21 (0.64)	3] 0.60
	2] Nimodipine 90 mg/d change from baseline	CDR					1] 1.16 (0.55) 2] 1.12 (0.60)	3] 0.67
		MMSE	1] 21.46 (4.24) 2] 21.24 (4.07)				1] -0.83 (3.29) 2] -0.87 (3.66)	3] 0.94 favours placebo
	3] Placebo vs Nimodipine change from baseline	CGI (item 1)					1] 0.14 (0.61) 2] 0.21 (0.57)	3] 0.35
		CGI (item 2)					1] 4.02 (1.06) 2] 4.02 (0.99)	3] 0.95 favours placebo
		FOM (total recall)					1] -2.56 (5.53) 2] -2.28 (6.54)	3] 0.67
		IADL					1] 0.14 (0.39) 2] 0.11 (0.32)	3] 0.41

EvTable232. Study results: Nizatidine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12m	
Carlson, 2002 Breitner, 1999	ITT Analysis 1] Placebo change from baseline 2] Nizatidine 75 mg bid change from baseline 3] Difference between placebo and Nizatidine change from baseline	COWA Letter fluency					1] 1.4 2] -0.2	3] 0.460
		Category Fluency					1] 0.5 2] 2.0	3] 0.611
		Boston Naming test					1] 1.4 2] 0.6	3] 0.231
		WMS Immediate Recall					1] 0.4 2] 2.1	3] 0.147
		WMS Delayed Recall					1] -0.8 2] 0.9	3] 0.087
		Word List Immediate Recall					1] 1.8 2] 2.1	3] 0.413
		Word list Delayed Recall					1] 0.3 2] 0.3	3] 0.916
		Constructional Praxis test					1] 1.6 2] 1.9	3] 0.704
		Constructional Praxis, recall IADL					1] 1.1 2] 0.3	3] 0.224

EvTable233. Study results: ACTH4-9 (Org 2766).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 4w	
Kragh-Sorensen 1986	OC Analysis							
	1] Org 2766 5 mg vs Placebo	GAGS						1] <0.05
	2] Org 2766 20 mg vs Placebo	SCAG total						1] <0.01 2] <0.09 3] <0.01 4] <0.01
	3] Org 2766 40 mg vs Placebo							
	4] Org 2766 80 mg vs Placebo	LPRS						4] <0.05

EvTable234. Study results: ACTH 4-9 (Org 2766).

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 6m	
Soininen 1985	OC Analysis 1] Placebo vs ACTH 4-9 (Org 2766)	SCAG					1] 0.8 CI (-6.8 to 5.1)	1] NS
Partanen 1986		AGS-E					1] -0.2 CI (-3.1 to 2.7)	1] NS
Soininen 1984		LPRS					1] 1.2 CI (- 3.4 to 5.8)	1] NS
		McGBRS					1] -0.0 CI (-2.4 to 2.4)	1] NS
		GPI-E					1] -0.1 CI (-3.8 to 3.7)	1] NS

EvTable235. Study results: Prednisone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 1y	
Aisen 2000b	ITT Analysis	ADAS-Cog					1] 6.3 (6.4) 2] 8.2 (7.8)	3] 0.16
Aisen 2000a	1] Placebo change from baseline	CDR-SB					1] 2.2 (1.8) 2] 2.9 (2.5)	3] 0.07
	2] Prednisone 20 mg decreasing change from baseline	BDRS					1] 1.7 (1.9) 2] 1.7 (2.1)	3] 0.60
		HAM-D					1] 0.7 (3.6) 2] 1.7 (4.5)	3] 0.25
	3] Difference between Placebo and Prednisone in change from baseline	BPRS					1] 2.0 (6.6) 2] 5.4 (8.2)	3] 0.003

EvTable236. Study results: Simvastatin.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 26w	
Simons 2002	OC Analysis							
	1] Placebo	MMSE	1] 17.1 (4.9)				1] 14.4 (5.6)	3] <0.05
	2] Simvastatin 80 mg/d		2] 17.8 (5.0)				2] 17.2 (4.8)	4] NS 5] <0.02
	3] Placebo change from baseline	ADAS-cog	1] 33.2 (11.3)				3] 3.4 (7.0)	3] NS
	4] Simvastatin change from baseline		2] 29.4 (10.4)				4] 4.1 (6.5)	4] NS 5] NS
	5] Simvastatin vs. Placebo							

EvTable237. Study results: Thiamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6m		Final: (specify) 12m	
Nolan, 1990	Completers Analysis	<u>MMSE</u>	1] 16.0 (5.7) 2] 16.6 (5.73)		1] 16.4 (7.7) 2] 13.4 (7.2)	3] <.05	1] 14.6 (7.09) 2] 10.4 (9.13)	3] <0.05
	1] Lactose placebo	Verbal Learning Score						3] <0.05
	2] Thiamine 3 g/ d	BNT						3]<0.05
	3] Change from baseline across groups							

EvTable238. Study results: Vincamine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6w		Final: (specify) 12w	
Fischhof 1996	OC Population							
	1] Placebo DAT	CGI	1] 5.0 (0.3)			5] <0.05	1] 4.6 (0.8)	
	2] Placebo MID		2] 4.9 (0.4)				2] 4.9 (0.7)	
			3] 5.0 (0.0)				3] 2.8 (1.0)	
			4] 5.0 (0.2)				4] 3.7 (1.1)	
	3] Vincamine 30 mg/d DAT	CGI Total Improvement					6] 25%	5] <0.05
							7] 72%	8] <0.05
								9] <0.05
	4] Vincamine 30 mg/d MID	CGI Worse					6] 7%	
							7] 3%	
	5] Vincamine vs Placebo	SCAG	1] 69.0 (7.4)			5] <0.05	1] 67.9 (8.0)	
			2] 70.1 (6.5)				2] 68.1 (7.5)	
	6] All Placebo		3] 68.8 (7.2)				3] 62.1 (8.5)	
			4] 68.3 (7.0)				4] 63.4 (6.7)	
	7] All Vincamine							
	8] Placebo vs Vincamine DAT	BGP Need for help	1] 15.0 (9.1)				1] 14.2 (9.0)	
			2] 12.5 (9.0)				2] 12.3 (8.7)	
			3] 12.3 (8.8)				3] 10.4 (8.8)	
			4] 12.9 (9.1)				4] 11.0 (8.3)	
	9] Placebo vs Vincamine MID	SKT	1] 17.5 (3.3)			5] <0.05	1] 17.5 (3.6)	
			2] 17.6 (2.8)				2] 17.2 (3.1)	
			3] 17.8 (2.6)				3] 14.8 (4.4)	
			4] 18.1 (2.6)				4] 14.8 (4.4)	

EvTable239. Study results: Piracetam.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12m	
Croisile 1993	OC Analysis 1] Placebo 2] Piracetam 8g/d 3] Placebo difference from baseline 4] Piracetam difference from baseline 5] Placebo vs Piracetam	MMSE	1] 19.31(3.32) 2] 19.21 (3.98)				1] 16.4(6.60) 2] 18.10(5.70)	3] <0.05 4] NS 5] NS
		MADRS	1] 5.75(3.45) 2] 9.50(6.82)				1] 7.88(5.68) 2] 10.14(7.61)	3] NS 4] NS 5] NS
		Blessed A	1] 4.28(3.34) 2] 4.96(3.72)				1] 7.72(5.47) 2] 6.46(4.51)	3] <0.01 4] <0.05 5] NS
		Aphasia Battery	1] 1.76(1.94) 2] 1.67(1.75)				1] 6.15(7.68) 2] 3.41(4.88)	3] <0.01 4] NS 5] NS
		Logical Digit Span	1] 14.69(4.21) 2] 12.36(4.65)				1] 10.88(6.16) 2] 11.50(5.37)	3] <0.05 4] NS 5] NS

EvTable240. Study results: Xantinolnicotinate.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify)		Final: (specify) 12w	
Kanowski 1990	OC Population	<u>CGI</u>					1] 5.6 2] 4.6 3] 5.3 4] 4.5	5] <0.001 6] <0.002 7] <0.001 8] <0.001
	1] Placebo SDAT subgroup							
	2] Xantinolnicotinate 1g tid SDAT subgroup	SCAG Decrease in mean value					1] 2.1% 2] 10.1% 3] 1.7% 4] 9.8%	5] <0.0001 6] <0.0002
	3] Placebo MID subgroup	BGP						
	4] Xantinolnicotinate 1g tid MID subgroup	Digit Connection Test						5] NS 6] NS
	5] difference between Placebo and Xantinolnicotinate SDAT subgroup	Digit Symbol Substitution test						5] <0.001 6] <0.001
	6] difference between Placebo and Xantinolnicotinate MID subgroup							5] <0.01 6] <0.03

EvTable240. Study results: Xantinolnicotinate cont'd.

REF ID#	Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify)		Final: (specify) 12w	
		7] Xantinolnicotinate SDAT change from baseline							
		8] Xantinolnicotinate MID change from baseline							

EvTable241. Study results: Donepezil (DPZ) - Vitamin E.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 3 m		Final: (specify) 6 m	
Thomas 2001	OC Analysis							
	1] DPZ 10 mg d	WAIS	1] 72 (2.0)* 2] 72 (2.0)*		1] 74 (2.0)* 2] 72 (2.0)*		1] 75 (2.0)* 2] 71 (2.1)*	3] 0.15 4] 0.43
	2] Vitamin E 2,000 IU							
	3] change from baseline with DPZ	MMSE	1] 16 (0.5)* 2] 16 (0.5)*		1] 16 (0.6)* 2] 15 (0.5)*	5] <0.001 favors DZP	1] 16 (0.5)* 2] 15 (0.6)*	3] 0.06 4] 0.07 5] <0.001 favors DPZ
	4] change from baseline with Vitamin E	ADAS-cog	1] 33.34 (2.7)* 2] 33.45 (2.6)*		1] 31.55 (2.7)* 2] 36.09 (2.8)*		1] 31.84 (2.7)* 2] 39.07 (2.7)*	3] <0.001 4] <0.01
	5] DPZ vs Vitamin E change from baseline	NPI	1] 21.9 (0.5)* 2] 21.9 (0.5)*				1] 16.8 (0.2)* 2] 22.8 (1.2)*	

*SEM

EvTable242. Study results: Fluoxetine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 30d		Final: (specify) 45d	
Taragano 1997	OC Analysis 1] Fluoxetine 10 mg/d 2] Amitriptyline 25 mg/d 3] Between treatments	Ham-D	1] 25.3 (3.8) 2] 26.3 (4.0)	3] 0.10	1] 19.3 (3.2) 2] 17.8 (2.5)	3] 0.10	1] 16.7 (2.9) 2] 15.6 (3.2)	3] 0.10
		MMSE	1] 20.0 (3.2) 2] 18.8 (4.2)	3] 0.10			1] 21.4 (2.9) 2] 21.5 (3.5)	3] 0.10

EvTable243. Study results: 5'-MTHF - Trazodone.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4w		Final: (specify) 8w	
Passeri 1992	OC Analysis	HDRS	1] 23 (5)		1] 20(6)	3] <0.01	1] 18 (6)	3] <0.01
	1] 5'-MTHF 50 mg/d		2] 23 (3)		2] 21 (4)	4] <0.05	2] 19 (5)	4] <0.01
			5] 23 (5)		5] 21 (6)	7] <0.01	5] 18 (6)	7] <0.01
	2] Trazodone 100mg/d		6] 23 (4)		6] 21 (5)	8] <0.01	6] 19 (6)	8] <0.01
		9] 21 (5)		9] 17 (7)		9] 18 (5)	11] <0.01	
	3] 5'-MTHF change from baseline	10] 23 (3)		10] 22 (2)		10] 20 (3)		
	4] Trazodone change from baseline	RVM immediate recall	1] 20 (7)				1] 23 (8)	3] <0.01
			2] 22 (9)				2] 22 (9)	7] <0.01
			5] 20 (7)				5] 23 (7)	
	5] 5'-MTHF subgroup AD		6] 22 (9)				6] 22 (8)	
	9] 20 (8)				9] 22 (7)			
6] Trazodone subgroup AD	10] 20 (8)				10] 22 (11)			
7] 5'-MTHF change from baseline subgroup AD	RVM delayed recall	1] 2 (2)				1] 3 (2)		
		2] 3 (2)				2] 3 (2)		
		5] 3 (2)				5] 3 (2)		
		6] 3 (2)				6] 3 (2)		
8] Trazodone change from baseline subgroup AD		9] 2 (2)				9] 3 (2)		
	10] 4 (2)				10] 3 (2)			
9] 5'-MTHF subgroup MID								
10] Trazodone subgroup MID								
11] Trazodone change from baseline subgroup MID								

EvTable244. Study results: Citicoline.

Author Year	Analysis Groups	Outcome s Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 45 d		Final: (specify) 90 d	
Parnetti 1994	OC Population 1] Placebo (Ascorbic Acid 100 mg/d) 2] Citicoline 500 mg/d 3] Posatirelin 10 mg/d 4] Ascorbic Acid change from baseline 5] Citicoline change from baseline 6] Posatirelin change from baseline 7] Posatirelin vs Citicoline change from baseline	<u>GBS</u> Emotional impairment	1] 1.9 (1.0) 2] 1.9 (1.0) 3] 1.9 (1.2)		1] 1.8 (1.0) 2] 1.9 (1.1) 3] 1.7 (1.0)		1] 1.9 (1.0) 2] 1.7 (1.0) 3] 1.6 (0.9)	4] NS 5] NS 6] <0.025
		<u>GBS</u> Impaired orientation & memory	1] 2.2 (0.9) 2] 2.2 (1.0) 3] 2.1 (1.0)		1] 2.1 (1.0) 2] 2.1 (1.0) 3] 2.0 (1.0)		1] 2.1 (1.1) 2] 2.1 (1.0) 3] 1.8 (1.0)	4] NS 5] NS 6] NS 7] 0.038 favors Posatirelin
		<u>GBS</u> Impaired ability ADL	1] 1.2 (0.8) 2] 1.4 (1.1) 3] 1.2 (1.0)		1] 1.3 (0.8) 2] 1.3 (1.0) 3] 1.1 (1.0)		1] 1.3 (0.9) 2] 1.4 (1.0) 3] 1.1 (1.0)	4] NS 5] NS 6] <0.025
		<u>GBS</u> Depression / Anxiety	1] 1.5 (0.9) 2] 1.5 (0.8) 3] 1.6 (1.1)		1] 1.5 (0.9) 2] 1.5 (0.9) 3] 1.5 (1.0)		1] 1.4 (0.9) 2] 1.4 (0.9) 3] 1.4 (0.9)	4] NS 5] NS 6] NS 7] 0.031 favors Posatirelin
		<u>GBS</u> Impaired attention & motivation	1] 2.2 (0.9) 2] 2.1 (1.0) 3] 2.1 (1.1)		1] 2.1 (0.9) 2] 2.0 (1.0) 3] 1.9 (0.9)		1] 2.1 (1.0) 2] 1.9 (1.0) 3] 1.8 (0.8)	4] NS 5] NS 6] <0.025
		<u>GBS</u> Intellectual impairment	1] 2.2 (0.8) 2] 2.1 (0.9) 3] 2.0 (0.9)		1] 2.1 (0.9) 2] 2.0 (0.9) 3] 1.9 (0.9)		1] 2.1 (1.0) 2] 2.0 (0.9) 3] 1.8 (0.8)	5] <0.025 7] 0.037 favors Posatirelin
		<u>GBS</u> Motor impairment	1] 1.2 (0.8) 2] 1.4 (1.1)		1] 1.3 (0.8) 2] 1.3 (1.0) 3] 1.1 (1.0)		1] 1.3 (0.9) 2] 1.4 (1.0) 3] 1.1 (1.0)	5] <0.025

EvTable244. Study results: Citicoline cont'd.

REF ID#	Author Year	Analysis Groups	Outcome s Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
				Baseline		Mid-Point: (specify) 45 d		Final: (specify) 90 d	
			MMSE	1] 16.4 (2.7) 2] 16.5 (2.6) 3] 16.6 (2.5)				1] 17.1 (4.1) 2] 17.6 (3.9) 3] 17.8 (3.4)	4] NS 5] NS 6] NS
			HDRS	1] 13.0 (5.0) 2] 11.4 (4.9) 3] 12.6 (5.0)				1] 11.4 (4.9) 2] 11.3 (5.2) 3] 11.1 (5.3)	4] NS 5] NS 6] NS

EvTable245. Study results: Pyritinol - Hydergine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 6w		Final: (specify) 12w	
Spilich 1996	OC Analysis	SCAG	1] 27.0 2] 28.0		1] 22.5 2] 25.0	3] 0.14	1] 16.5 2] 22.0	3] 0.008 favors Pyritinol
	2] Hydergine 4.5 mg/d	CETM	1] 13.0 2] 14.8		1] 17.0 2] 17.8		1] 19.8 2] 17.0	4] <0.001 5] <0.002
	3] Between drugs vs baseline							
	4] Pyritinol change from baseline							
	5] Hydergine change from baseline							

EvTable246. Study results: Sulfomucopolysaccharides - Cytidine Diphosphocholine.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 2w		Final: (specify) 4w	
Cucinotta 1987	OC Analysis 1] Sulphomucopolysaccharides 500 units 2] CDP-choline 1.0g 3] difference between Sulphomucopolysaccharides and CDP-choline	SCAG	1] 35.35(1.32)* 2] 36.00(1.55)*		1] 32.28(1.28)* 2] 35.00(1.38)*		1] 31.61(1.94)* 2] 35.58(1.56)*	3] <0.05 favors Sulphomucopolysaccharides
		NMS	1] 16.71(0.59)* 2] 17.93(0.56)*		1] 16.14(0.54)* 2] 17.80(0.49)*		1] 15.45(0.66)* 2] 18.08(0.71)*	3] <0.02 favors Sulphomucopolysaccharides
		Digit Symbol	1] 44.66(6.84)* 2] 47.41(4.99)*				1] 65.77(9.46)* 2] 52.00(6.44)*	3] <0.05 favors Sulphomucopolysaccharides
		Digit Span	1] 6.72 (0.33)* 2] 5.91 (0.42)*				1] 6.90 (0.34)* 2] 6.25 (0.44)*	3] NS

*SEM

EvTable247. Study results: Sulodexide - Pentoxifylline.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 4m		Final: (specify) 6m	
Parnetti, 1997	OC Analysis 1] Sulodexide 50 mg bid 2] Pentoxifylline 400 mg tid 3] sulodexide change vs baseline	<u>GBS motor impairment</u>	1] 1.64 (0.14)* 2] 1.59 (0.13)*		1] 1.58 (0.14)* 2] 1.53 (0.14)*	3] <0.01	1] 1.54 (0.16)* 2] 1.46 (0.17)*	3] <0.01
		<u>GBS intellectual impairment</u>	1] 2.09 (0.09)* 2] 1.98 (0.08)*		1] 1.88 (0.09)* 2] 1.95 (0.1)*	3] <0.01	1] 1.79 (0.1)* 2] 1.87 (0.12)*	3] <0.01
		<u>GBS emotional impairment</u>	1] 2.1 (0.12)* 2] 1.89 (0.1)*		1] 1.88 (0.12)* 2] 1.9 (0.1)*	3] <0.12	1] 1.76 (0.12)* 2] 1.75 (0.11)*	3] <0.01
		MMSE	1] 17.6 (0.4)* 2] 18 (0.4)*				1] 20 (0.6)* 2] 20 (0.4)*	

*SEM

EvTable248. Study results: Selegiline - Vitamin E.

Author Year	Analysis Groups	Outcomes Measured	Result Value	P Value	Result Value	P Value	Result Value	P Value
			Baseline		Mid-Point: (specify) 10m		Final: (specify) 20m	
Sano 1997	ITT Analysis	<u>Event-free survival</u>			1] 79%		1] 40%	5] 0.077
	1] Placebo				2] 86%		2] 51%	6] 0.087
Thal 1996	2] Vitamin E 1000IU bid				3] 60%		3] 60%	7] 0.21
	3] Selegiline 5mg bid	Event-free survival with MMSE as covariate			4] 80%		4] 49%	
	4] Vitamin E 1000IU bid + Selegiline 5mg bid							5] 0.001
	5] Vitamin E 1000IU bid vs Placebo from baseline							6] 0.012
	6] Selegiline 5mg bid vs Placebo from baseline							7] 0.049
	7] Vitamin E 1000IU bid + Selegiline 5mg bid vs Placebo from baseline							

EvTable249. Adverse Events: Other agents.

Adverse events (AE) identified in included studies	ANIRACETAM Senin, 1991	ATEROID Ban, 1991b	BMV 21,502 Shrotriya, 1996	CBM 36-733 Danielczyk, 1988	CHOTO-SAN Shimada, 1994	CHOTO-SAN Terasawa, 1997	DEK Cucinotta, 1996	VASOPRESSIN (DDAVP) Peabody, 1986	DENBUFYLLINE Treves, 1999
Withdrawn (%) due to AE	T: 0 C: 0	T: 1 C: 2	T: 24 C: 9	T: 3 C: 5	T: 3 C: 0	T: 7 C: 3	T: 4 C: 2	T: 0 C: 0	T: 21 C: 16
AE Checklist (Max 5)	3	2	3	4	3	2	3	2	2
None Reported									
Balance									x
Accidental Injury									
Dizziness			x	x					
Falls									
Behavioral	x								
Agitation							x		
Cardiovascular	x	x		NS			x		
Arrhythmia			x	x			x		
Hypotension							x		
Hypertension				NS		x			
Extrapyramidal									
Tremor									
Gastrointestinal		x					x		
Abdominal pain	x								
Constipation									
Diarrhea						x	x		
Dyspepsia			x			x	x		x
Nausea, vomiting	x						x		x
Metabolic/nutritional						x			
Eating disorder									
Weight Change									
Neurological		x	x	x		x			x
Asthenia									
Psychiatric									
Anxiety			x	x			x		
Confusion, delirium			x				x	x	
Depression									
Respiratory				x					x
Cough, cold, infection									
Rhinitis			x						
Other			x				x		x
Aberrant hematology					x				
Fatigue, weakness									
Fever, flu, pneumonia						x			
Headache			x				x		
Hepatic abnormality	x		x			x			
Muscle/joint disorder									
Pain							x		
Rash, skin disorder							x		
Sleep disorder	x								
Urinary disorder				x		x	x		x

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable249. Adverse Events: Other agents cont'd.

Adverse events (AE) identified in included studies	NIMODIPINE Ban, 1990	NIMODIPINE Pantoni, 2000a	NIZATIDINE Carlson, 2002	ORG 2766 Soininen, 1985	ORG 2766 Kragh-Sorensen, 1986	PIRACETAM Croisile, 1993	THIAMINE Nolan, 1991
Withdrawn (%) due to AE	T: 1 C: 0	T: 0 C: 0	T: NR C: NR	T: 0 C: 0	T: NR C: NR	T: 0 C: 0	T: 0 C: 0
AE Checklist (Max 5)	5	4	3	2	5	3	0
None Reported					x		x
Balance							
Accidental Injury							
Dizziness							
Falls							
Behavioral							
Agitation							
Cardiovascular	NS	NS	x				
Arrhythmia	NS						
Hypotension	NS						
Hypertension							
Extrapyramidal		x					
Tremor							
Gastrointestinal	NS	x				x	
Abdominal pain							
Constipation						x	
Diarrhea	NS						
Dyspepsia							
Nausea, vomiting	NS						
Metabolic/nutritional							
Eating disorder	NS						
Weight Change				NS		NS	
Neurological		S	x				
Asthenia							
Psychiatric		x					
Anxiety							
Confusion, delirium							
Depression							
Respiratory		x					
Cough, cold, infection							
Rhinitis							
Other		x					
Aberrant hematology		x		S		NS	
Fatigue, weakness							
Fever, flu, pneumonia							
Headache	NS						
Hepatic abnormality							
Muscle/joint disorder							
Pain							
Rash, skin disorder		x					
Sleep disorder	NS						
Urinary disorder		x					

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable249. Adverse Events: Other agents cont'd.

Adverse events (AE) identified in included studies	BUFLOMEDIL Cucinotta, 1992	CYCLANDELA Weyer, 2000	CYCLANDELA Schellenberg, 1997	GUANFACINE Crook, 1992b	HYDERGINE Spilich, 1996	HYDERGINE Thompson, 1990	VINCAMINE Fischhof, 1996	XANITOLNICOTINAT E Kanowski, 1990	PENTOXIFYLLINE Parnetti, 1997
Withdrawn (%) due to AE	T:NR C:NR	T: 7 C: 6	T: 9 C: 6	T: 0 C: 0	T: C:	T: 0 C: 0	T:NR C:NR	T: 0 C: 0	T: 7 C: 6
AE Checklist (Max 5)	1	3	3	0	4	1	3	3	2
None Reported	x			x					
Balance							x	x	
Accidental Injury									
Dizziness								x	
Falls									
Behavioral									
Agitation									
Cardiovascular					x				X
Arrhythmia									
Hypotension							x		
Hypertension									
Extrapyramidal									
Tremor									
Gastrointestinal			x						
Abdominal pain									X
Constipation									
Diarrhea			x				x		
Dyspepsia									
Nausea, vomiting			x				x	x	
Metabolic/nutritional		x							
Eating disorder									
Weight Change									
Neurological									
Asthenia									X
Psychiatric									
Anxiety							x		
Confusion, delirium								x	
Depression			x						
Respiratory									
Cough, cold, infection									
Rhinitis									
Other			x			x		x	
Aberrant hematology									
Fatigue, weakness								x	
Fever, flu, pneumonia									
Headache			x					x	X
Hepatic abnormality									
Muscle/joint disorder									
Pain									
Rash, skin disorder								x	
Sleep disorder							x		
Urinary disorder		x							

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable249. Adverse Events: Other agents cont'd.

Adverse events (AE) identified in included studies	N- ACETYLCYSTEINE Adair 2001	GM-1 Ala, 1990	LYCOSAMINO GLYCAN- OLYSULPHATE Ban, 1991a	DEFERRIOXAMINE Crapper, McLachlan, 1991	SIMVASTATIN Simmons, 2002	POSATIRELIN Parnetti, 1995	5'-MTHF (T) TRADOZONE (C) Passeri, 1993
Withdrawn (%) due to AE	T: 0 C: 0	T: 10 C: 0	T: 1 C: 4	T: 0 C: 0	T: 8 C: 0	T: 0 C: 0	T: 0 C: 0
AE Checklist (Max 5)	2	3	5	3	2	4	3
None Reported							
Balance							x
Accidental Injury							
Dizziness	X						
Falls							
Behavioral	X						
Agitation		X				X	
Cardiovascular		X	X				
Arrhythmia						X	
Hypotension							
Hypertension							
Extrapyramidal							
Tremor						X	
Gastrointestinal			X				
Abdominal pain							
Constipation			X				
Diarrhea	X						
Dyspepsia						X	
Nausea, vomiting							
Metabolic/nutritional	X		X				
Eating disorder				X			
Weight Change				X			
Neurological							
Asthenia							
Psychiatric							
Anxiety							
Confusion, delirium		X				X	
Depression							
Respiratory							
Cough, cold, infection							
Rhinitis							
Other	X		X		X	X	x
Aberrant hematology							
Fatigue, weakness	X						
Fever, flu, pneumonia							
Headache	X	X				X	
Hepatic abnormality							
Muscle/joint disorder	X				X		
Pain		X					
Rash, skin disorder	X	X				X	
Sleep disorder	X	X				X	
Urinary disorder		X				X	

NR = Withdrawals due to AE Not Reported + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper

EvTable249. Adverse Events: Other agents cont'd.

Adverse events (AE) identified in included studies	PREDNISON Aisen, 2000b	NIMESULIDE Aisen, 2002	DICLOFENAC ISOPROSTOL Scharf, 1999	INDOMETHACIN Rogers, 1993	HYDROXY- CHLOROQUINE Van Gool, 2001
Withdrawn (%) due to AE	T: 0 C: 0	T: 5 C: 5	T: 46 C: 6	T: 42 C: 30	T: 0 C: 0
AE Checklist (Max 5)	3	5	2	2	0
None Reported					
Balance					
Accidental Injury					
Dizziness		X			
Falls					
Behavioral		X	X	X	
Agitation					
Cardiovascular					
Arrhythmia					
Hypotension					
Hypertension					
Extrapyramidal					
Tremor					
Gastrointestinal		X		X	
Abdominal pain		S	X		
Constipation	X	S			
Diarrhea					
Dyspepsia					
Nausea, vomiting		X			X
Metabolic/nutritional					
Eating disorder					
Weight Change					
Neurological				X	
Asthenia					
Psychiatric					
Anxiety					
Confusion, delirium					
Depression					
Respiratory					
Cough, cold, infection					
Rhinitis					
Other	S			X	X
Aberrant hematology		X	X		
Fatigue, weakness					
Fever, flu, pneumonia					
Headache	X			X	X
Hepatic abnormality	S	X	X		
Muscle/joint disorder					
Pain					
Rash, skin disorder		S			
Sleep disorder					
Urinary disorder	S	X			

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
 x = Reported adverse event/side effect but not tested for significant differences between groups
 S or NS = Reported and tested for statistical differences between placebo and treatment group
 S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
 [] = Symptom NOT reported in the paper

EvTable249. Adverse Events: Other agents cont'd.

Adverse events (AE) identified in included studies	FLUOXETINE AMITRIPTYLINE Taragano, 1997	SULFOMUCO- POLYSACCHARIDE S CDP-CHOLINE Cucinotta, 1988	DONEPEZIL Thomas, 2001	SELEGILINE VITAMIN E Sano, 1997
Withdrawn (%) due to AE	T: 58 C: 22	T: C:	T: 0 C: 0	T: 0 C: 0
AE Checklist (Max 5)	3		3	1
None Reported			X	
Balance				S*
Accidental Injury				
Dizziness				
Falls				S*
Behavioral				
Agitation				
Cardiovascular				NS*
Arrhythmia				
Hypotension				
Hypertension				
Extrapyramidal				NS*
Tremor				
Gastrointestinal				NS*
Abdominal pain				
Constipation	X			
Diarrhea	X			
Dyspepsia				
Nausea, vomiting	X			
Metabolic/nutritional				
Eating disorder				
Weight Change				
Neurological				NS*
Asthenia				
Psychiatric				
Anxiety				
Confusion, delirium	X			
Depression				
Respiratory				
Cough, cold, infection				
Rhinitis				
Other				S*
Aberrant hematology				
Fatigue, weakness				
Fever, flu, pneumonia				
Headache				
Hepatic abnormality				
Muscle/joint disorder				
Pain				
Rash, skin disorder				NS*
Sleep disorder				
Urinary disorder				

NR = Withdrawals due to AE Not Reported; + = Dose response effect on AE
x = Reported adverse event/side effect but not tested for significant differences between groups
S or NS = Reported and tested for statistical differences between placebo and treatment group
S* or NS* = Reported and tested for statistical differences between two (three) treatment groups
[] = Symptom NOT reported in the paper